



Kurzke, Matthias and Marzuola, Jeremy L. and Spirn, Daniel (2017) Gross-Pitaevskii vortex motion with critically scaled inhomogeneities. *SIAM Journal on Mathematical Analysis*, 49 (1). pp. 471-500. ISSN 0036-1410

Access from the University of Nottingham repository:

<http://eprints.nottingham.ac.uk/40895/1/M104901.pdf>

Copyright and reuse:

The Nottingham ePrints service makes this work by researchers of the University of Nottingham available open access under the following conditions.

This article is made available under the University of Nottingham End User licence and may be reused according to the conditions of the licence. For more details see: http://eprints.nottingham.ac.uk/end_user_agreement.pdf

A note on versions:

The version presented here may differ from the published version or from the version of record. If you wish to cite this item you are advised to consult the publisher's version. Please see the repository url above for details on accessing the published version and note that access may require a subscription.

For more information, please contact eprints@nottingham.ac.uk

GROSS–PITAEVSKII VORTEX MOTION WITH CRITICALLY SCALED INHOMOGENEITIES*

MATTHIAS KURZKE[†], JEREMY L. MARZUOLA[‡], AND DANIEL SPIRN[§]

Abstract. We study the dynamics of vortices in an inhomogeneous Gross–Pitaevskii equation $i\partial_t u = \Delta u + \frac{1}{\varepsilon^2}(\rho_\varepsilon^2(x) - |u|^2)u$. For a unique scaling regime $|\rho_\varepsilon(x) - 1| = O(\log \varepsilon^{-1})$, it is shown that vortices can interact both with the background perturbation and with each other. Results for associated parabolic and elliptic problems are discussed.

Key words. vortices, Gross–Pitaevskii, critical scaling

AMS subject classification. 35Q60

DOI. 10.1137/15M1049014

1. Introduction. We consider the behavior of vortices in an inhomogeneous Gross–Pitaevskii (IGP) equation

$$(1.1) \quad \begin{cases} i\partial_t u = \Delta u + \frac{1}{\varepsilon^2}(p^2(x) - |u|^2)u & \text{in } \Omega, \\ \nu \cdot \nabla u = 0 & \text{on } \partial\Omega, \end{cases}$$

where Ω is a bounded, simply connected domain in \mathbf{R}^2 .

An important feature of solutions of (1.1) is the potential appearance of vortices, points where $u(x, t) : \mathbf{R}^2 \rightarrow \mathbb{C}$ has a zero with nontrivial topological degree. We concentrate on the case where there are n vortices of degree $d_j \in \{-1, 1\}$ at positions $a_j(t) \in \mathbf{R}^2$ for $j = 1, \dots, n$. We will show below that solutions take the form

$$u(x, t) \approx \prod_{j=1}^n \rho_\varepsilon(x) \left(\frac{x - a_j(t)}{|x - a_j(t)|} \right)^{d_j}.$$

Here $\frac{x - a_j(t)}{|x - a_j(t)|}$ is interpreted in the complex sense as $\frac{z - a_j(t)}{|z - a_j(t)|}$ for $z - a_j = x_1 - a_{j,1} + i(x_2 - a_{j,2})$ with $\rho_\varepsilon(x) \rightarrow 0$ as $|x - a_j| \rightarrow 0$ and $|u(x, t)|^2 \approx p^2(x)$ away from the $a_j(t)$'s. In a sense identifying the location and degrees of vortices is sufficient to describe the solution $u(x, t)$.

The IGP equation (1.1) arises in several contexts, including the behavior of superfluid ^4He near a boundary, Bose–Einstein condensation (BEC) under a very weak confinement potential, and nonlinear optics. In the first case it has been observed experimentally that anomalous behavior of superfluid ^4He vortex filament motion indicates that vortex tubes can become pinned to sites on the material boundary; see [46, 38, 45]. Such pinning can be incorporated heuristically by the introduction of

*Received by the editors November 20, 2015; accepted for publication (in revised form) November 17, 2016; published electronically February 16, 2017.

<http://www.siam.org/journals/sima/49-1/M104901.html>

Funding: The work of the second author was supported in part by NSF applied math grant DMS-1312874 and NSF CAREER grant DMS-1352353. The work of the third author was supported in part by NSF CAREER grant DMS-0955687 and NSF grant DMS-1516565.

[†]School of Mathematical Sciences, University of Nottingham, Nottingham NG7 2RD, United Kingdom (matthias.kurzke@nottingham.ac.uk).

[‡]Mathematics Department, UNC, Chapel Hill, NC 27599 (marzuola@math.unc.edu).

[§]Mathematics Department, University of Minnesota, Minneapolis, MN 55455 (spirn@math.umn.edu).

a potential $p(x)$ in the Gross–Pitaevskii equation (see [34]) or complex Ginzburg–Landau equations (see [15]). In the second case, IGP is a useful model in a BEC in which the vortices interact both with each other and with the trap potential. In the third case, IGP is useful in modeling optical vortices. See, for instance [3, 2, 34], which are related to motion of vortices with nonvanishing total charge in inhomogeneous potentials and confinement.

The dynamics of vortices in Gross–Pitaevskii equations have been the subject of much study in the past few decades. The connection between the vortex motion in superfluids, where $p(x) \equiv 1$, and simplified ODEs was made by Fetter in [11], and it was shown that vortices interact with each other through a Coulomb potential. On the other hand, in BECs with nontrivial trapping potentials, $p(x) = O(1)$, it was shown by Fetter and Svidzinsky in [12] that vortices interact solely with the background potential and are carried along level sets of the Thomas–Fermi profile.

On the other hand, it has been observed, both experimentally [14, 32] and numerically [29], that vortices do interact with other vortices and with the background potential. In one particularly interesting direction, experimentalists have generated dipoles by dragging nonradially symmetric BECs through laser obstacles in [32] and by the Kibble–Zurek mechanism in [14]. These bound dipoles interact in nontrivial ways with each other and the background potentials. Reduced ODE models for the dynamics of these vortex configurations can be found in [30, 41, 42, 44] and are in good agreement with numerical simulations of the full equation (1.1) and with physical experiments [30]. The objective of our work is to place these widely studied models on a rigorous footing.

We are interested in the critical asymptotic regime where vortices interact with both the background potential $p(x)$ and each other. In the following, we will take $p(x) = p_\varepsilon(x)$ with

$$(1.2) \quad |1 - p_\varepsilon^2(x)| \lesssim \frac{1}{|\log \varepsilon|}$$

and show that asymptotic scalings of $p_\varepsilon(x)$ outside of the range (1.2) induce dynamics that are dominated by only the background potential or by only vortex-vortex interactions.

1.1. Background potential. Following Lassoued and Mironescu [26] we write

$$u(x, t) = \eta_\varepsilon(x)w(x, t),$$

where η_ε is a minimizer of an energy (A.2) and consequently a nontrivial solution to the elliptic PDE

$$(1.3) \quad \begin{cases} 0 = \Delta \eta_\varepsilon + \frac{1}{\varepsilon^2} \eta_\varepsilon (p_\varepsilon^2 - \eta_\varepsilon^2) & \text{in } \Omega, \\ \nu \cdot \nabla \eta_\varepsilon = 0 & \text{on } \partial\Omega, \end{cases}$$

then (1.1) is equivalent to the Schrödinger equation

$$(1.4) \quad \begin{cases} i\eta_\varepsilon^2 \partial_t w = \operatorname{div}(\eta_\varepsilon^2 \nabla w) + \frac{\eta_\varepsilon^4}{\varepsilon^2} (1 - |w|^2)w & \text{in } \Omega, \\ \nu \cdot \nabla w = 0 & \text{on } \partial\Omega. \end{cases}$$

The local and global well-posedness of (1.4) for fixed ε can be established using a variation of the argument in Brézis and Gallouet [7].

Associated to (1.4) is a weighted Hamiltonian,

$$e_\varepsilon^{\eta_\varepsilon}(w) := \frac{1}{2}\eta_\varepsilon^2 |\nabla w|^2 + \frac{\eta_\varepsilon^4}{4\varepsilon^2} (1 - |w|^2)^2;$$

consequently, $e_\varepsilon^1(w)$ is the standard gaugeless Ginzburg–Landau energy density as in [4]. We set

$$E_\varepsilon^{\eta_\varepsilon}(w) := \int_\Omega e_\varepsilon^{\eta_\varepsilon}(w) dx$$

to be the total energy of the mapping w associated to background potential η_ε . We will be interested in an asymptotic range of $p_\varepsilon(x)$'s such that the resulting energy feels both the background potential and the other vortices at the same order. As will be elicited below, our regime corresponds to having the background η_ε expanded in the following way: $\eta_\varepsilon^2(x) = 1 + \frac{Q_\varepsilon(x)}{|\log \varepsilon|}$ with $Q_\varepsilon(x) \rightarrow Q_0(x)$ as $\varepsilon \rightarrow 0$.

1.2. Conservation laws. There is a set of conservation laws for (1.4), and to write them down efficiently, we introduce some notation. For $b, c \in \mathbb{C}$ set $(\cdot, \cdot) = \frac{1}{2}(b\bar{c} + \bar{b}c)$ and let us define

$$j(u) := (iu, \nabla u) \equiv \text{supercurrent},$$

$$J(u) := \det \nabla u = \frac{1}{2} \operatorname{curl} j(u) \equiv \text{Jacobian};$$

then solutions satisfy the following differential identities:

$$(1.5) \quad \frac{\eta_\varepsilon^2}{2} \partial_t (|w|^2 - 1) = \operatorname{div} (\eta_\varepsilon^2 j(w)),$$

$$(1.6) \quad \partial_t e_\varepsilon^{\eta_\varepsilon}(w) = \operatorname{div}(\eta_\varepsilon^2(\nabla w, \partial_t w)).$$

A third conservation law holds for the Jacobian, we which we now define, tensorially. If the matrix \mathbb{J} is given by \mathbb{J}_{ij} with $\mathbb{J}_{21} = -\mathbb{J}_{12} = 1$ and $\mathbb{J}_{11} = \mathbb{J}_{22} = 0$, then $\operatorname{curl} F = \mathbb{J}_{k\ell} \partial_\ell F_k$, and if $J(w) = \frac{1}{2} \operatorname{curl} j(w)$, then following Jerrard and Smets [16] we have after a lengthy calculation

$$(1.7) \quad \partial_t J(w) = \mathbb{J}_{\ell j} \partial_j \left[\frac{1}{\eta_\varepsilon^2} \partial_k (\eta_\varepsilon^2 \partial_\ell w, \partial_k w) \right] - \mathbb{J}_{\ell j} \partial_j \left[\eta_\varepsilon^2 \partial_\ell \frac{(1 - |w|^2)^2}{4\varepsilon^2} \right].$$

Multiplying by $\phi \in C_0^2(\Omega)$ yields a conservation law observed in [16],

$$(1.8) \quad \int_\Omega \phi \partial_t J(w) dx = \int_{\mathbb{R}^2} \mathbb{J}_{\ell j} \partial_j \partial_k \phi (\partial_k w, \partial_\ell w) dx$$

$$- \int_\Omega \mathbb{J}_{\ell j} \partial_j \phi \left[\frac{\partial_k \eta_\varepsilon^2}{\eta_\varepsilon^2} (\partial_\ell w, \partial_k w) + \partial_\ell \eta_\varepsilon^2 \frac{(1 - |w|^2)^2}{4\varepsilon^2} \right] dx.$$

The dynamics of a vortex can be inferred by choosing suitable test functions ϕ in (1.8) which elicit the vortex positions. Since we expect the Jacobian $J(w)$ to be roughly equal to a delta function, localized at the site of each vortex α , we can choose $\phi = x_m \chi(x - \alpha)$, $m \in \{1, 2\}$, for smooth cutoff function $\chi(x) = 1$ in B_r and compactly supported in B_{2r} . Integrating the left-hand side over the support yields, formally, $\pi \dot{\alpha}$. On the right-hand side, the first term has support in an annulus about each vortex, due to the structure of the test function, and since the energy

concentrates outside of the support of the annulus, one sees that it generates an $O(1)$ vortex-vortex interaction term. The second term on the right-hand side has support inside of a ball about each vortex, and since the energy tensor is large in this region [24], it generates an $O(|\log \varepsilon| |\nabla \log \eta_\varepsilon|)$ vortex-potential interaction term. Therefore, for a vortex to interact with both the background potential and other vortices, it requires that $|\nabla \log \eta_\varepsilon| = O(|\log \varepsilon|^{-1})$ for both terms to be of the same order.¹

In fact if $|\nabla \log \eta_\varepsilon| \ll |\log \varepsilon|^{-1}$, then the vortex dynamics will be dominated by the interaction with the background potential, as in [16], whereas if $|\nabla \log \eta_\varepsilon| \gg |\log \varepsilon|^{-1}$, then vortex dynamics will be driven by its interaction with the current generated by fellow vortices; see [9].

1.3. Weak topology. In order to measure the distance of the vortices to the site of the expected location of the vortices given by the ODE (1.11), we use the flat norm $\dot{W}^{-1,1}(\Omega)$. We denote the norm $\dot{W}^{-1,1}$ unless this topology is used on a subdomain, such as

$$\|\mu\|_{\dot{W}^{-1,1}(B_r(\alpha))} = \sup_{\|\nabla \phi\|_{L^\infty(B_r(\alpha))} \leq 1} \left\{ \int_{B_r(\alpha)} \phi d\mu \text{ such that } \phi \in W_0^{1,\infty}(B_r(\alpha)) \right\}.$$

We use a helpful estimate for concentrations in the flat norm; see Brezis, Coron, and Lieb [6]. Define

$$(1.9) \quad r_\alpha := \frac{1}{8} \min \left\{ \min_{i \neq j} |\alpha_i - \alpha_j|, \min_j \text{dist}(\alpha_j, \partial\Omega) \right\};$$

then the following holds.

LEMMA 1.1. *Suppose α_k, ξ_k are points in Ω . If*

$$\left\| \sum d_k \delta_{\alpha_k} - d_k \delta_{\xi_k} \right\|_{\dot{W}^{-1,1}} \leq \frac{1}{4} r_\alpha,$$

then

$$\left\| \sum d_k \delta_{\alpha_k} - d_k \delta_{\xi_k} \right\|_{\dot{W}^{-1,1}} = \sum_k |\alpha_k - \xi_k|.$$

Proof. For a proof see [16], for example. □

Consequently, the flat norm provides a good way to measure the location of singularities. Since the Jacobian, $J(w)$, converges to a sum of weighted delta functions, we will use $\dot{W}^{-1,1}$ topology as a convenient metric to track vortex trajectories.

Associated to a set of locations $\alpha_j \in \Omega$ and degrees $d_j \in \{\pm 1\}$ we can define a *canonical harmonic map*, w_* , that describes the limiting harmonic map with prescribed singularities,

$$w_*(\alpha, d) \equiv e^{i\psi_*} \prod_{j=1}^n \left(\frac{x - \alpha_j}{|x - \alpha_j|} \right)^{d_j},$$

where harmonic ψ_* is chosen so that $\partial_\nu w_*(\alpha, d) = 0$ on $\partial\Omega$. We will use w_* when there is no ambiguity.

¹The third term on the right-hand side of (1.8) is negligible.

1.4. Energy expansion. Given a collection of n vortices with centers $\alpha = \{\alpha_1, \dots, \alpha_n\}$, the Hamiltonian $E_\varepsilon^{\eta_\varepsilon}(w)$ can be expanded out to second order with $E_\varepsilon^{\eta_\varepsilon}(w) = H_\varepsilon(\alpha) + o_\varepsilon(1)$, where

$$H_\varepsilon(\alpha) := \sum_{j=1}^n (\pi|\log \varepsilon| + \pi Q_0(\alpha_j) + \gamma_0) + W(\alpha, d),$$

which will be established in the sense of Γ -convergence in Proposition 4.3. Here n will be used throughout to represent the number of vortices, $Q_0(x)$ is the limiting rescaled background perturbation, and $W(\alpha, d)$ is the renormalized energy that arises in the work of Bethuel, Brezis, and Hélein [4],² given by

$$\begin{aligned} W(\alpha, d) &= \lim_{r \rightarrow 0} \left[\min_{\substack{v \in H^1(\Omega_r(\alpha); S^1) \\ \deg(v; \partial B_r(\alpha_j)) = d_j}} \int_{\Omega_r(\alpha)} \frac{1}{2} |\nabla v|^2 dx - \sum_{j=1}^n \pi \log \frac{1}{r} \right] \\ &= -\pi \sum_{j \neq k} d_j d_k \log |\alpha_j - \alpha_k| + \text{boundary terms,} \end{aligned}$$

where $\Omega_r(\alpha) \equiv \Omega \setminus \cup_j B_r(\alpha_j)$. In order to express the boundary terms, we define $G(\cdot, \alpha, d) = \sum_{j=1}^n G(\cdot, \alpha_j, d_j)$, where

$$\Delta G(\cdot, \alpha_j, d_j) = 2\pi d_j \delta_{\alpha_j} \text{ in } \Omega \text{ with } G = 0 \text{ on } \partial\Omega.$$

If we set $F(x, \alpha_j) = G(x, \alpha_j, d_j) - \log |x - \alpha_j|$, then we can fully express $W(\alpha, d)$ as

$$(1.10) \quad W(\alpha, d) = -\pi \sum_{j \neq k} d_j d_k \log |\alpha_j - \alpha_k| + \pi \sum_{j,k} d_j F(\alpha_j, \alpha_k);$$

see [4] and the appendix of [40].

1.5. Discussion. Rigorous results on vortex dynamics for Gross–Pitaevskii were established when $p(x) \equiv 1$ by Colliander and Jerrard [9] and also Lin and Xin [28], which showed that vortices satisfy the Kirchoff–Onsager ODE for Euler point vortices. When $p(x) = O(1)$ there is a recent result of Jerrard and Smets [16] that showed that vortices travel along level sets of the Thomas–Fermi profile. In the parabolic setting there were rigorous results by Lin [27] and Jerrard and Soner [18] when $p(x) \equiv 1$ and by Jian and Song [22] when $p = O(1)$. Mixed dynamics with $p = O(1)$ and further forcing terms were discussed by Serfaty and Tice [39]. A variational proof of the parabolic dynamics for $p \equiv 1$ was given by Sandier and Serfaty [35].

1.6. Results. We wish to describe the dynamics of vortices in (1.4). If the vortices are located at positions $\alpha_j(t)$ with degree $d_j \in \{-1, 1\}$, then the vortices move via the ODE,

$$(1.11) \quad \pi d_k \dot{\alpha}_k(t) = \nabla_{\alpha_k}^\perp H_0(\alpha, Q_0),$$

where $\nabla^\perp = (\partial_2, -\partial_1)^T$ and

$$(1.12) \quad H_0(\alpha, Q_0) = W(\alpha, d) + \pi \sum_{j=1}^n Q_0(\alpha_j).$$

²The authors of [4] study the case of Dirichlet boundary conditions. See the discussion in the appendix of [40] for the derivation of $W(\alpha, d)$ with Neumann boundary conditions.

Given the ODE generated by (1.11), we can define the collision time,

$$T_{col} := \sup_{t \geq 0} \{r_{\alpha(s)} > 0 \text{ for all } 0 \leq s \leq t\}.$$

Then for any $0 \leq T < T_{col}$ we can define

$$(1.13) \quad r_{min} := \min_{t \in [0, T]} r_{\alpha(t)},$$

which is the minimum vortex-vortex or vortex-boundary distance until time T . Clearly $r_{min} > 0$.

Our main result is the following, which captures both vortex-vortex and vortex-potential interactions. An important hypothesis will be that the initial data has control of the excess energy

$$(1.14) \quad D_{\varepsilon}(w, \alpha) \equiv E_{\varepsilon}^{\eta_{\varepsilon}}(w) - H_{\varepsilon}(\alpha, Q_0).$$

We will shorten the notation to D_{ε} when the w and α are readily apparent.

THEOREM 1.2. *Let $p_{\varepsilon}^2 = 1 + \frac{\rho_{\varepsilon}(x)}{|\log \varepsilon|}$, where the ρ_{ε} satisfies hypotheses of Proposition 1.6 with $k \geq 4$. Let $\{\alpha_j^0, d_j\}$ be a configuration of vortices such that $r_{\alpha_0} > 0$, $d_j \in \{-1, 1\}$ and suppose $w_{\varepsilon}^0, 0 < \varepsilon < 1$, satisfies the well-preparedness hypotheses*

$$(1.15) \quad \|J(w_{\varepsilon}^0) - \pi \sum d_j \delta_{\alpha_j^0}\|_{\dot{W}^{-1,1}} = o_{\varepsilon}(1) \text{ and } D_{\varepsilon}(w_{\varepsilon}^0, \alpha^0) = o_{\varepsilon}(1).$$

If $w_{\varepsilon}(t)$ is a solution to (1.1) with initial data w_{ε}^0 , then there exists a time $T > 0$ such that for all $t \in [0, T]$,

$$\left\| J(w_{\varepsilon}(\cdot, t)) - \pi \sum d_j \delta_{\alpha_j(t)} \right\|_{\dot{W}^{-1,1}(\Omega)} \rightarrow 0$$

as $\varepsilon \rightarrow 0$, where the $\alpha_j(t)$ are defined by (1.11), and T is independent of ε .

Remark 1.3. An important function satisfying the conditions of Proposition 1.6 below is simply $\rho_{\varepsilon} = \rho_0 \in H^{k+1}(\Omega)$, where ρ_0 is any fixed function such that $\nabla \rho_0$ has compact support inside Ω .

One important example that arises from Theorem 1.2 is dipoles that are scattered by inhomogeneities; see Remark 1.5 below and section 5 for some illustrative numerical simulations.

Under slightly less regularity assumptions, we have a similar result for the gradient flow

$$(1.16) \quad \frac{1}{|\log \varepsilon|} \partial_t u_{\varepsilon} = \Delta u_{\varepsilon} + \frac{1}{\varepsilon^2} (p_{\varepsilon}^2 - |u_{\varepsilon}|^2) u_{\varepsilon}$$

with the limit equation

$$(1.17) \quad \pi \dot{\alpha}_k(t) = -\nabla_{\alpha_k} H_0(\alpha, Q_0),$$

THEOREM 1.4. *Let $p_{\varepsilon}^2 = 1 + \frac{\rho_{\varepsilon}(x)}{|\log \varepsilon|}$, where the ρ_{ε} satisfies hypotheses of Proposition 1.6 with $k \geq 3$. If $\{\alpha_j^0, d_j\}$ is a configuration of vortices such that $d_j \in \{-1, 1\}$*

and if $w_\varepsilon^0, 0 < \varepsilon < 1$ satisfies (1.16) and $w_\varepsilon = \frac{u_\varepsilon}{\eta_\varepsilon}$ satisfies the well-preparedness hypotheses

$$(1.18) \quad \|J(w_\varepsilon^0) - \pi \sum d_j \delta_{\alpha_j^0}\|_{\dot{W}^{-1,1}} = o_\varepsilon(1) \text{ and } D_\varepsilon(w_\varepsilon^0, \alpha^0) = o_\varepsilon(1) \text{ and } r_{\alpha_0} > 0,$$

then there exists a time $T > 0$ such that for all $t \in [0, T]$,

$$\left\| J(w_\varepsilon(\cdot, t)) - \pi \sum d_j \delta_{\alpha_j(t)} \right\|_{\dot{W}^{-1,1}(\Omega)} \rightarrow 0$$

as $\varepsilon \rightarrow 0$, where the $\alpha_j(t)$ are defined by (1.17).

Remark 1.5. There are straightforward adaptations of Theorem 1.2 to other contexts, including the following:

- The Gross–Pitaevskii equation (1.1) on $\Omega \equiv \mathbf{R}^2$. This can be achieved by combining the methods here with the arguments in [5, 21]. In this case vortices move according to the same ODE (1.11); however, the renormalized energy $W(a, d)$ is the classical Coulomb potential,

$$W(\alpha, d) = -\pi \sum_{j \neq k} d_j d_k \log |\alpha_j - \alpha_k|.$$

- Mixed Ginzburg–Landau equations of the form

$$\left(\frac{1}{|\log \varepsilon|} \partial_t + i \partial_t \right) u_\varepsilon = \Delta u_\varepsilon + \frac{1}{\varepsilon^2} (p_\varepsilon^2(x) - |u_\varepsilon|^2) u_\varepsilon$$

as studied in [23, 31] which results in a modified ODE

$$\pi \dot{\alpha}_k(t) - \pi d_k (e_3 \times \dot{\alpha}_k(t)) = -\nabla_{\alpha_k} H_0(\alpha, Q_0).$$

In section 4 we present first and second order Gamma-convergence results for the energy $E_\varepsilon^{\eta_\varepsilon}(w_\varepsilon)$; see Proposition 4.1 and Proposition 4.3, respectively. These results are similar to those found in [1], albeit ported to the case with critical inhomogeneities.

1.7. Properties of $\eta_\varepsilon = \eta_\varepsilon(p_\varepsilon)$. We collect here some information required on the behavior of η_ε , as related to the background fluctuations p_ε . The discussion of the proof will be given in Appendix A and is based on standard elliptic theory estimates, so we simply state the results that we require for the vortex dynamics here.

We write

$$(1.19) \quad p_\varepsilon^2 = 1 + \frac{\rho_\varepsilon}{|\log \varepsilon|}, \quad \eta_\varepsilon^2 = 1 + \frac{Q_\varepsilon}{|\log \varepsilon|}.$$

We will assume that $\rho_\varepsilon \rightarrow \rho_0(x)$ in $H^k(\Omega)$ with k made precise below. Using results such as [10] for the Dirichlet problem, we expect that p_ε and η_ε should be quite close. We make precise the nature in which that is true in the following proposition, which will be proved in the appendix.

We remark here that the elliptic analysis required for the convergence that we will implement in the appendix is somewhat nontrivial, as the overall ellipticity of the underlying nonlinear elliptic problem is going to 0 in the limit as $\varepsilon \rightarrow 0$. Hence, estimates must be done with some care, especially to understand regularity up to the boundary of our domain.

PROPOSITION 1.6. *Let ρ_ε be such that $\rho_\varepsilon \rightarrow \rho_0(x)$ in $H^k(\Omega)$, $\nabla \rho_\varepsilon$ is compactly supported strictly on the interior of Ω for all ε and $\varepsilon^2 \rho_\varepsilon \rightarrow 0$ in H^{k+1} . Then, for Q_ε as defined in (1.3), (1.19) satisfying Neumann boundary conditions on $\partial\Omega$, we have $Q_\varepsilon \rightarrow \rho_0$ in $H^k(\Omega)$. In particular, we have $Q_\varepsilon \rightarrow \rho_0$ in $C^{2,\delta}(\overline{\Omega})$ with $\delta > 0$ for integer $k \geq 4$ as required for the Schrödinger dynamics below and $Q_\varepsilon \rightarrow \rho_0$ in $C^{1,\delta}(\overline{\Omega})$, $\delta > 0$, for integer $k \geq 3$ as required for the gradient flow dynamics.³*

Remark 1.7. In the appendix, we will actually decompose $\eta_\varepsilon = 1 + \frac{\tilde{Q}_\varepsilon}{|\log \varepsilon|}$, in which case

$$Q_0 = \lim_{\varepsilon \rightarrow 0} \left[2\tilde{Q}_\varepsilon + \frac{\tilde{Q}_\varepsilon^2}{|\log \varepsilon|} \right];$$

this convention slightly simplifies the resulting elliptic analysis.

Remark 1.8. Similar results hold assuming Dirichlet boundary conditions, but we work here with Neumann as they are the most physically relevant.

2. Excess energy control. In this section we present an excess energy identity similar to [9, 20]. The influence of the background requires some modifications, so we give a full proof. The form of the estimate will be similar to that found in [5].

Recall from (1.9) the quantity

$$r_\alpha = \frac{1}{8} \min \left\{ \min_{i \neq j} |\alpha_i - \alpha_j|, \min_i \text{dist}(\alpha_i, \partial\Omega) \right\}.$$

Then, we have the next proposition.

PROPOSITION 2.1. *Let $w \in H^1(\Omega; \mathbb{C})$ and assume that there exists α such that $D_\varepsilon(\alpha, w) \leq 1$. Given $0 < r \leq r_\alpha$, there exist constants $\varepsilon_0 > 0$ and $\mu_0 > 0$ such that if $0 < \varepsilon < \varepsilon_0$, and*

$$\mu \equiv \left\| J(w) - \pi \sum d_j \delta_{\alpha_j} \right\|_{\dot{W}^{-1,1}} \leq \mu_0,$$

then

$$(2.1) \quad \int_{\Omega_r(\alpha)} e^{\eta_\varepsilon} (|w|) + \frac{1}{8} \left| \frac{j(w)}{|w|} - j(w_*) \right|^2 dx \leq D_\varepsilon + o_{\varepsilon,\mu}(1),$$

where $o_{\mu,\varepsilon}(1)$ vanishes as $\varepsilon, \mu \rightarrow 0$. We also have

$$(2.2) \quad \int_{\Omega} e_\varepsilon^1(|w|) dx \leq C,$$

$$(2.3) \quad \|j(w)\|_{L^p(\Omega)} \leq C, \quad 1 \leq p < 2.$$

Furthermore, there exists points $\xi_j \in B_{r/2}(\alpha_j)$ such that

$$(2.4) \quad \left\| J(w) - \pi \sum d_j \delta_{\xi_j} \right\|_{\dot{W}^{-1,1}} \leq C\varepsilon |\log \varepsilon|$$

³Note, by interpolation arguments, we can actually run the convergence argument which is $H^{3+}(\Omega)$, where by $\cdot + \nu$ we mean $\cdot + \nu$ for any $\nu > 0$. We chose here to work with integer Sobolev spaces for convenience. In fact, the sharp estimate would include Schauder theory estimates directly on Hölder norms, but that would require uniform convergence in C^1 , which does not seem obvious given that the ∇Q term appears to vanish in the limit, preventing the proof of uniform bounds of higher regularity.

and

$$(2.5) \quad \left\| \frac{(\partial_k w, \partial_\ell w)}{|\log \varepsilon|} - \delta_{k\ell} \sum \pi \delta_{\xi_j} \right\|_{\dot{W}^{-1,1}} \leq C |\log \varepsilon|^{-\frac{1}{2}},$$

where $\delta_{k\ell}$ is the Kronecker delta. Here C is a constant depending on $Q_0, \Omega, r_\alpha, n, p, \mu_0,$ and ε_0 .

Remark 2.2. Note that (2.2) and Rellich–Kondrachov imply

$$(2.6) \quad \| |w| - 1 \|_{L^q(\Omega)} = o_{\varepsilon, \mu}$$

for all $q < +\infty$.

Proof. Although it is possible to get explicit control on the error $o_\varepsilon(1)$ in (2.1) by a careful analysis as in [20, 25], the weaker estimate (2.1) is sufficient to prove the vortex motion law.

1. We first decompose the excess energy into

$$\begin{aligned} D_\varepsilon &= E_\varepsilon^{\eta_\varepsilon}(w) - H_\varepsilon(\alpha, Q_0) \\ &= \int_\Omega e_\varepsilon^{\eta_\varepsilon}(w) dx - \left[\sum_{j=1}^n (\pi |\log \varepsilon| + \gamma_0 + \pi Q_0(\alpha_j)) + W(\alpha) \right] \\ &= \int_{\Omega_\sigma(\alpha)} e_\varepsilon^{\eta_\varepsilon}(w) - e_\varepsilon^{\eta_\varepsilon}(w_*) dx + \int_{\Omega_\sigma(\alpha)} e_\varepsilon^{\eta_\varepsilon}(w_*) dx - W(\alpha) \\ &\quad + \sum_{j=1}^n \left[\int_{B_\sigma(\alpha_j)} e_\varepsilon^{\eta_\varepsilon}(w) dx - (\pi |\log \varepsilon| + \gamma_0 + \pi Q_0(\alpha_j)) \right] \\ &= I + II + III, \end{aligned}$$

where σ is appropriately chosen and $\sigma \leq r \leq r_\alpha$.

We will control terms I – III by variants of estimates found in [20, 25, 16]. We will choose

$$(2.7) \quad \sigma = \max\{|\log \varepsilon|^{-\frac{1}{3}}, \sqrt{\mu}\}$$

in the following.

2. By a simple variation of a standard calculation (see, for example, [20]),

$$(2.8) \quad \frac{\eta_\varepsilon^2}{2} \left| \frac{j(w)}{|w|} - j(w_*) \right|^2 + e_\varepsilon^{\eta_\varepsilon}(|w|) = e_\varepsilon^{\eta_\varepsilon}(w) - e_\varepsilon^{\eta_\varepsilon}(w_*) + \eta_\varepsilon^2 j(w_*) \cdot \left(j(w_*) - \frac{j(w)}{|w|} \right),$$

and so our primary concern will be with estimating the integral of (2.8) on a suitable subdomain $\Omega_\sigma(\alpha)$. This last term can be rewritten as

$$\begin{aligned} \eta_\varepsilon^2 j(w_*) \cdot \left(j(w_*) - \frac{j(w)}{|w|} \right) &= (\eta_\varepsilon^2 - 1) j(w_*) \cdot \left(j(w_*) - \frac{j(w)}{|w|} \right) \\ &\quad + j(w_*) \cdot \left(j(w_*) - \frac{j(w)}{|w|} \right). \end{aligned}$$

We now fix our μ_0 and ε_0 in order to be able to cite some estimates from [17, 20]. The tool we wish to use is in the proof of Lemma 4 in [20], which yields an

estimate given some assumptions on the Jacobian and ε . Let K_1 be the constant in the assumptions of Lemma 4 from [20], which depends only on Ω .

We follow an argument in the proof of Theorem 6.1 of [5] for fixing these constants; however, there are several new constraints, including the fact that the vortex ball radius, σ , cannot be too small due to the need to control terms like $\|\eta_\varepsilon^2 - 1\|_{L^\infty} \|j(w_*)\|_{L^\infty(\Omega_\sigma)}$.

We first choose μ_0 . Set μ_1 such that

$$(2.9) \quad 4\mu \leq \sqrt{\mu} \leq \min \left\{ r, \frac{r_\alpha}{nK_1} \right\}$$

for all $0 \leq \mu \leq \mu_1$, i.e., $\mu_1 = \min\{\frac{1}{16}, r^2, \frac{r_\alpha^2}{n^2K_1^2}\}$. We further restrict $\mu_0 = \min\{\mu_1, \frac{r_\alpha}{8K_2n^5}\}$, where K_2 comes from the assumptions in Theorem 3 of [20], needed for the localization result (2.4).

We now choose an ε_0 to allow us to use the necessary array of estimates. First, we choose ε_2 so that for all $\varepsilon < \varepsilon_2$,

$$(2.10) \quad |\log \varepsilon|^{-\frac{1}{3}} \leq \min \left\{ r, \frac{r_\alpha}{nK_1} \right\}.$$

We then set $\varepsilon_1 \leq \varepsilon_2$ so that $\varepsilon E_\varepsilon^1(w) \leq \sqrt{\varepsilon}$ for all $\varepsilon < \varepsilon_1$. Indeed since $\eta_\varepsilon^2 \geq \frac{1}{2}$, then $E_\varepsilon^1(w) \leq \frac{1}{\min \eta_\varepsilon^4} E_\varepsilon^{\eta_\varepsilon}(w) \leq 4C|\log \varepsilon|$, so there exists an ε_1 such that $\varepsilon E_\varepsilon^1(w) \leq 4C\varepsilon|\log \varepsilon| \leq \sqrt{\varepsilon}$ for all $\varepsilon \leq \varepsilon_1$. We next fix $\varepsilon_0 \leq \varepsilon_1$ such that

$$(2.11) \quad \varepsilon \log \frac{r_\alpha}{\varepsilon} \leq |\log \varepsilon|^{-\frac{2}{3}}$$

for all $0 \leq \varepsilon \leq \varepsilon_0$.

If $\mu \leq |\log \varepsilon|^{-\frac{2}{3}}$, then we use $s_\varepsilon = |\log \varepsilon|^{-\frac{2}{3}}$ and $\sigma = |\log \varepsilon|^{-\frac{1}{3}}$ in Lemma 4 of [20]. In particular (2.10) and (2.11) imply that the assumptions in Lemma 4 hold, and we find

$$(2.12) \quad \begin{aligned} & \left| \int_{\Omega_\sigma(\alpha)} j(w_*) \cdot \left(j(w_*) - \frac{j(w)}{|w|} \right) dx \right| \\ & \leq \frac{1}{4} \int_{\Omega_\sigma(\alpha)} \left| \frac{j(w)}{|w|} - j(w_*) \right|^2 dx + C|\log \varepsilon|^{\frac{1}{3}} \left(|\log \varepsilon|^{-\frac{2}{3}} + \sqrt{\varepsilon} \right) + \frac{C}{r_\alpha} |\log \varepsilon|^{-\frac{1}{3}} \\ & \leq \frac{1}{4} \int_{\Omega_\sigma(\alpha)} \left| \frac{j(w)}{|w|} - j(w_*) \right|^2 dx + o_{\varepsilon, \mu}(1). \end{aligned}$$

If $|\log \varepsilon|^{-\frac{2}{3}} \leq \mu$, then again the assumptions hold for Lemma 4 of [20], and choose $s_\varepsilon = \mu$ and $\sigma = \sqrt{\mu}$ so

$$(2.13) \quad \begin{aligned} & \left| \int_{\Omega_\sigma(\alpha)} j(w_*) \cdot \left(j(w_*) - \frac{j(w)}{|w|} \right) dx \right| \\ & \leq \frac{1}{4} \int_{\Omega_\sigma(\alpha)} \left| \frac{j(w)}{|w|} - j(w_*) \right|^2 dx + \frac{C}{\sqrt{\mu}} (\mu + \sqrt{\varepsilon}) + C \frac{\mu}{r_\alpha} \\ & \leq \frac{1}{4} \int_{\Omega_\sigma(\alpha)} \left| \frac{j(w)}{|w|} - j(w_*) \right|^2 dx + o_{\varepsilon, \mu}(1). \end{aligned}$$

Taking the sum over the two errors yields sufficient control on $\int_{\Omega_\sigma} j(w_*) \cdot \left(j(w_*) - \frac{j(w)}{|w|}\right) dx$.

Finally, we note

$$\begin{aligned} & \left| \int_{\Omega_\sigma(\alpha)} (\eta_\varepsilon^2 - 1) j(w_*) \cdot \left(j(w_*) - \frac{j(w)}{|w|}\right) dx \right| \\ & \leq C \|1 - \eta_\varepsilon^2\|_{L^\infty} \|j(w_*)\|_{L^\infty(\Omega_\sigma(\alpha))} \left(1 + \int_{\Omega_\sigma(\alpha)} \left|j(w_*) - \frac{j(w)}{|w|}\right|^2 dx\right) \\ & \leq C \frac{\min\{|\log \varepsilon|^{\frac{1}{3}}, \sqrt{\frac{1}{\mu}}\}}{|\log \varepsilon|} \left(1 + \int_{\Omega_\sigma(\alpha)} \left|j(w_*) - \frac{j(w)}{|w|}\right|^2 dx\right) \\ & \leq o_{\varepsilon, \mu}(1) \left(1 + \int_{\Omega_\sigma(\alpha)} \left|j(w_*) - \frac{j(w)}{|w|}\right|^2 dx\right). \end{aligned}$$

Combining these estimates together controls term I .

3. We control II using the explicit control in Lemma 12 [20],

$$\begin{aligned} & \left| \int_{\Omega_\sigma(\alpha)} e^{\eta_\varepsilon} (w_*) dx - \left(W(\alpha) + n\pi \log \frac{1}{\sigma}\right) \right| \\ & = \left| \frac{1}{2} \int_{\Omega_\sigma(\alpha)} \eta_\varepsilon^2 |\nabla w_*|^2 dx - \left(W(\alpha) + n\pi \log \frac{1}{\sigma}\right) \right| \\ & \leq \left| \frac{1}{2} \int_{\Omega_\sigma(\alpha)} |\nabla w_*|^2 dx - \left(W(\alpha) + n\pi \log \frac{1}{\sigma}\right) \right| + C \|1 - \eta_\varepsilon^2\|_{L^\infty} \|j(w_*)\|_{L^\infty(\Omega_\sigma)}^2 \\ & \leq C \frac{\sigma^2}{r_\alpha^2} + C \frac{1}{\sigma^2 |\log \varepsilon|} \leq C \frac{\max\{|\log \varepsilon|^{-\frac{2}{3}}, \mu\}}{r_\alpha^2} + C \frac{\min\{|\log \varepsilon|^{\frac{2}{3}}, \frac{1}{\mu}\}}{|\log \varepsilon|} = o_{\varepsilon, \mu}(1). \end{aligned}$$

4. We now consider term III. Recall that $Q_\varepsilon(x) = |\log \varepsilon| (\eta_\varepsilon^2(x) - 1)$. We set $q_\varepsilon(r, x_0) = \inf_{x \in B_r(x_0)} Q_\varepsilon(x)$ and $\tilde{\varepsilon}_r = \frac{\varepsilon}{1 + q_\varepsilon(r, x_0)}$. Recall from (2.10) that

$$\|J(w) - \pi d_j \delta_{\alpha_j}\|_{\dot{W}^{-1,1}(B_\sigma(\alpha_j))} \leq \|J(w) - \pi \sum d_k \delta_{\alpha_k}\|_{\dot{W}^{-1,1}(\Omega)} \leq \mu \leq \frac{\sqrt{\mu}}{4} \leq \frac{\sigma}{4};$$

therefore, we can invoke Theorem 1.3 and Lemma 6.8 of [17]:

$$\begin{aligned} \int_{B_\sigma(\alpha_j)} e^{\frac{1}{\tilde{\varepsilon}_\sigma}} (w) dx & \geq \pi \log \frac{\sigma}{\tilde{\varepsilon}_\sigma} + \gamma_0 - C \frac{\tilde{\varepsilon}_\sigma}{\sigma} \sqrt{\log \frac{\sigma}{\tilde{\varepsilon}_\sigma}} - C \frac{\mu}{\sigma} \\ & \geq \pi \log \frac{\sigma}{\varepsilon} + \gamma_0 + \pi \log \left(1 + \frac{Q_\varepsilon(\sigma, \alpha_j)}{|\log \varepsilon|}\right) - C\varepsilon |\log \varepsilon| - C\sqrt{\mu}, \end{aligned}$$

so

$$(2.14) \quad \int_{B_\sigma(\alpha_j)} e^{\frac{1}{\tilde{\varepsilon}_\sigma}} (w) dx \geq \pi \log \frac{\sigma}{\varepsilon} + \gamma_0 - o_{\varepsilon, \mu}(1).$$

Therefore,

$$\begin{aligned} & \int_{B_\sigma(\alpha_j)} e^{\eta_\varepsilon}(w) dx - \left(\pi \log \frac{\sigma}{\varepsilon} + \gamma_0 + \pi Q_0(\alpha) \right) \\ & \geq \left(1 + \frac{Q_\varepsilon(\sigma, \alpha_j)}{|\log \varepsilon|} \right) \left(\pi \log \frac{\sigma}{\varepsilon} + \gamma_0 - o_{\varepsilon, \eta}(1) \right) - \left(\pi \log \frac{\sigma}{\varepsilon} + \gamma_0 + \pi Q_0(\alpha) \right) \\ & \geq -\pi |q_\varepsilon(\sigma, \alpha_j) - Q_0(\alpha_j)| - C \frac{\log \sigma + 1}{|\log \varepsilon|} \\ & \geq -\pi |q_\varepsilon(\sigma, \alpha_j) - Q_0(\alpha_j)| - C \frac{\log |\log \varepsilon|}{|\log \varepsilon|} = -o_{\varepsilon, \mu}(1), \end{aligned}$$

and so (2.1) follows.

5. We next turn to the proof of (2.2) and (2.3). We first choose vortex balls $B_{r_\alpha}(\alpha_j)$; by (2.9),

$$(2.15) \quad \|J(w) - \pi d_j \delta_{\alpha_j}\|_{W^{-1,1}(B_r(\alpha_j))} \leq \|J(w) - \pi \sum d_k \delta_{\alpha_j}\|_{W^{-1,1}(\Omega)} \leq \mu \leq \frac{r_\alpha}{4},$$

and arguing as in step 4,

$$\int_{B_{r_\alpha}(\alpha_j)} e^{\eta_\varepsilon}(w) dx \geq \pi |\log \varepsilon| - C(r_\alpha, Q_0).$$

Therefore,

$$(2.16) \quad \begin{aligned} \int_{\Omega_{r_\alpha}(\alpha)} e^{\eta_\varepsilon}(w) dx &= E_\varepsilon^{\eta_\varepsilon}(w) - \sum_{j=1}^n \int_{B_{r_\alpha}(\alpha_j)} e^{\eta_\varepsilon}(w) dx \\ &\leq D_\varepsilon + n\gamma_0 + \pi \sum_{j=1}^n Q_0(\alpha_j) + W(\alpha) + C(n, r_\alpha, Q_0), \end{aligned}$$

and so,

$$(2.17) \quad \begin{aligned} \int_{B_{r_\alpha}(\alpha_k)} e^{\eta_\varepsilon}(w) dx &= E_\varepsilon^{\eta_\varepsilon}(w) - \int_{\Omega_{r_\alpha}(\alpha)} e^{\eta_\varepsilon}(w) dx - \sum_{j \neq k} \int_{B_{r_\alpha}(\alpha_j)} e^{\eta_\varepsilon}(w) dx \\ &\leq \pi |\log \varepsilon| + C(n, r_\alpha, Q_0). \end{aligned}$$

Finally, $\eta_\varepsilon^2 \geq \frac{1}{2}$, (2.17), and (2.15) allow us to use (4.27) in the proof of Proposition 4.2 in [17], which in turn implies

$$(2.18) \quad \int_{B_r(\alpha_j)} e_\varepsilon^1(|w|) dx \leq C(n, r_\alpha, Q_0).$$

Since $r \leq r_\alpha$, we can combine (2.1), $\eta_\varepsilon^2 \geq \frac{1}{2}$, and (2.18) to get bound (2.2).

To prove (2.3) we use an L^p bound on each vortex ball $B_{r_\alpha}(\alpha_j)$ in Theorem 3.2.1 of [9]. Outside the vortex balls, we use the decomposition argument; in particular, since $\eta_\varepsilon^2(x) \geq \frac{1}{2}$ then $\frac{1}{4} \int_{\Omega_{r_\alpha}(\alpha)} \frac{|j(w)|^2}{|w|} dx \leq \int_{\Omega_{r_\alpha}(\alpha)} e^{\eta_\varepsilon}(w) dx \leq C(D_\varepsilon, r_\alpha)$ from (2.16). From Remark 2.2 and the above bound, one finds

$$\begin{aligned} \|j(w)\|_{L^p(\Omega_{r_\alpha}(\alpha))} &\leq \left\| \frac{j(w)}{|w|} \right\|_{L^p(\Omega_{r_\alpha}(\alpha))} + \left\| \frac{j(w)}{|w|} (1 - |w|) \right\|_{L^p(\Omega_{r_\alpha}(\alpha))} \\ &\leq \left\| \frac{j(w)}{|w|} \right\|_{L^2(\Omega_{r_\alpha}(\alpha))} \left(C + \|1 - |w|\|_{L^{\frac{2p}{2-p}}(\Omega_{r_\alpha}(\alpha))} \right) \\ &\leq C(r_\alpha, Q_0, p), \end{aligned}$$

due to Remark 2.2. Combining the vortex ball bound with the bound in the excised domain yields (2.3).

5. The restriction on μ_0 allows us to use Theorem 3 of [20], and estimate (2.4) follows directly. To establish (2.5) we can follow the proof of Theorem 11 in [25], along with the equipartitioning result of [24], which lets us localize the stress-energy tensor $\nabla w \otimes \nabla w$ up to an error of order $O(\sqrt{|\log \varepsilon|})$. \square

3. Proof of Theorem 1.2. The proof of the vortex motion law entails first establishing the smooth evolution of the vortex paths $\xi_j(t)$ via the compactness theorems in section 2. The second step will compare the vortex paths with the ODE (1.11).

3.1. Lipschitz continuity of the vortex paths.

PROPOSITION 3.1. *Let $\{a_j^0, d_j\}$ be a configuration of vortices such that $d_j \in \{-1, 1\}$. Let $w_\varepsilon, 0 < \varepsilon < 1$, satisfy (1.4) under well-preparedness condition (1.15). There exist a time $T > 0$, a sequence $\varepsilon_k \rightarrow 0$, and $\xi_j \in Lip([0, T_0], \Omega)$ such that $\xi_j(0) = a_j^0$ and*

$$(3.1) \quad \sup_k \left\| J(w_{\varepsilon_k}(\cdot, t)) - \pi \sum d_j \delta_{\xi_j(t)} \right\|_{\dot{W}^{-1,1}(\Omega)} \rightarrow 0$$

as $k \rightarrow \infty$ and $t \in [0, T_0]$.

Proof.

1. We first claim that for any $T > 0$, there exists an ε such that $J(w_\varepsilon)$ is a continuous function on $[0, T]$ into $L^1(\Omega)$. Since $J(u) = -\nabla^\perp \mathcal{R}u \cdot \nabla \mathcal{I}u$ then

$$\|J(w_\varepsilon(t)) - J(w_\varepsilon(s))\|_{L^1} \leq C \|\nabla w_\varepsilon(t) - \nabla w_\varepsilon(s)\|_{L^2} \|\nabla w_\varepsilon(t) + \nabla w_\varepsilon(s)\|_{L^2} \leq C\varepsilon o_{|t-s|}(1),$$

since the solution map is a continuous function into $H^1(\Omega)$.

2. We use (1.8) along with the results in Proposition 2.1. Set

$$(3.2) \quad \varphi_j(x) = x\chi\left(\frac{|x - a_j^0|}{r a^0}\right),$$

where

$$\chi(s) = \begin{cases} 1 & \text{for } s \leq 1, \\ 0 & \text{for } s > 2. \end{cases}$$

Choosing $0 \leq r \leq s$ and using (1.8) and (1.15) we generate the bound

$$\begin{aligned} \left| \int_r^s \int_\Omega \varphi_j(x) \partial_t J(w_\varepsilon(\cdot, t)) dx dt \right| &\leq C \|\nabla^2 \varphi_j\|_{L^\infty} \int_r^s \int_{\Omega_{r a^0}(r)} |\nabla w_\varepsilon(t)|^2 dx \\ &\quad + C \frac{\|\nabla \varphi_j\|_{L^\infty}}{|\log \varepsilon|} \int_r^s \int_\Omega e^{\eta_\varepsilon}(w_\varepsilon) dx ds \\ &\leq C|s - r|, \end{aligned}$$

and so

$$\begin{aligned} \pi |\xi_j(s) - \xi_j(r)| &= \pi \left| \int \varphi_j(x) (\delta_{\xi_j(s)} - \delta_{\xi_j(r)}) dx \right| \\ &\leq C \|\nabla \varphi_j\|_{L^\infty} \sup_{t \in [r,s]} \left\| J(w_\varepsilon(\cdot, t)) - \pi \sum d_j \delta_{\xi_j(t)} \right\|_{\dot{W}^{-1,1}} \\ &\quad + \left| \int_r^s \int_\Omega \varphi_j(x) \partial_t J(w_\varepsilon(\cdot, t)) dx dt \right| \\ &\stackrel{(2.4)}{\leq} C|s - r| + o_\varepsilon(1). \end{aligned}$$

We can now employ a diagonalization argument. Taking subsequence $\varepsilon_k \subset \varepsilon$, we generate a dense, countable collection of times $t \in [0, T]$, in which T depends solely on r_α , such that the $\xi_j(t)$ satisfy $\xi_j(0) = a_j^0$ with (3.1) as $k \rightarrow \infty$. Since the collection is dense, we can take the limit as $k \rightarrow \infty$. The Lipschitz extension of the $\xi_j(t)$'s on $[0, T]$ satisfy the same conditions. \square

3.2. Vortex dynamics. In this subsection we complete the proof of Theorem 1.2. We use the Lipschitz paths $\xi_j(t)$ generated along the subsequence ε_k from Proposition 3.1. We will examine the differences between vortex paths and the ODE, which will lead eventually to a Gronwall argument.

The proof requires two technical calculations. The first describes the gradient of $W(\alpha)$ in terms of the canonical harmonic map w_* .

LEMMA 3.2. *Let $\alpha \in \Omega$; then $w_* = w_*(\cdot, \alpha)$ and the renormalized energy $W(\alpha)$ satisfy*

$$(3.3) \quad \int \mathbb{J}_{k\ell} \partial_k \partial_m \varphi j(w_*(\cdot, \alpha))_m j(w_*(\cdot, \alpha))_\ell dx = - \sum d_j \mathbb{J}_{\ell k} \partial_k \varphi(\alpha_j) (\nabla_{\alpha_j} W(\alpha))_\ell,$$

where $\varphi \in C^2(\Omega)$ and $\nabla^2 \varphi$ has support in a neighborhood of the α_j 's.

Proof. See, for example, [9, 28] for a proof. \square

The second technical result proves weak compactness of the supercurrent away from the vortex cores. This will be used to bound certain cross terms in the Gronwall argument. The argument is along the lines of similar proofs in [9, 28], among other places, and we sketch most of the argument.

LEMMA 3.3. *Let w_ε be a solution to (1.4) under the hypotheses of Proposition 3.1 and suppose $X \in C_0^\infty(\Omega \times [0, T])$ is a smooth vector field with support away from the vortex paths. For any $0 < t_0 < T$ and $\tau > 0$ with $t_0 + \tau < T$, then*

$$(3.4) \quad \int_{t_0}^{t_0+\tau} \int_\Omega X \cdot \left(\frac{j(w_\varepsilon(t))}{|w_\varepsilon(t)|} - j(w_*(\xi(t))) \right) dx dt \rightarrow 0$$

as $\varepsilon \rightarrow 0$, where $\xi(t) = (\xi_1(t), \dots, \xi_n(t))$ are the Lipschitz vortex paths generated in Proposition 3.1.

Proof. We sketch the argument in [23] for simplicity. Since

$$\frac{j(w_\varepsilon(t))}{|w_\varepsilon(t)|} - j(w_*(\xi(t))) = \frac{j(w_\varepsilon(t))}{|w_\varepsilon(t)|} (1 - |w_\varepsilon(t)|) + j(w_\varepsilon(t)) - j(w_*(\xi(t))),$$

then we can control

$$\begin{aligned} & \int_{t_0}^{t_0+\tau} \int_\Omega X \cdot \left(\frac{j(w_\varepsilon(t))}{|w_\varepsilon(t)|} - j(w_*(\xi(t))) \right) dx dt \\ & \leq \int_{t_0}^{t_0+\tau} \int_\Omega X \cdot \frac{j(w_\varepsilon(t))}{|w_\varepsilon(t)|} (1 - |w_\varepsilon(t)|) dx dt \\ & \quad + \int_{t_0}^{t_0+\tau} \int_\Omega X \cdot (j(w_\varepsilon(t)) - j(w_*(\xi(t)))) dx dt \\ & \leq C\varepsilon |\log \varepsilon| \|X\|_{L^2(\Omega \times [0, T])} + \int_{t_0}^{t_0+\tau} \int_\Omega X \cdot (j(w_\varepsilon(t)) - j(w_*(\xi(t)))) dx dt. \end{aligned}$$

We now focus on the last integral.

By (2.3) in Proposition 2.1 the $j(w_\varepsilon)$ are uniformly bounded in L^p for all $1 \leq p < 2$; furthermore, the $j(w_*(\xi(t)))$ are also in L^p for all $1 \leq p < 2$. The uniform bounds allow us to generate a Hodge decomposition, $j(w_\varepsilon) - j(w_*(\xi(t))) = \nabla h_1^\varepsilon(t) + \mathbb{J}\nabla h_2^\varepsilon(t)$, where the $h_j^\varepsilon(t) \in W^{1,p}(\Omega)$ and

$$\begin{aligned} \Delta h_1^\varepsilon(t) &= \operatorname{div} j(w_\varepsilon(t)) \text{ in } \Omega & \text{and} & \quad \partial_\nu h_1^\varepsilon(t) = 0 \text{ on } \partial\Omega, \\ \Delta h_2^\varepsilon(t) &= 2(J(w_\varepsilon(t)) - J(w_*(\xi(t)))) \text{ in } \Omega & \text{and} & \quad h_2^\varepsilon(t) = 0 \text{ on } \partial\Omega. \end{aligned}$$

We first claim that

$$(3.5) \quad \left| \int_{t_0}^{t_0+\tau} \int_{\Omega} X \cdot \nabla h_1^\varepsilon dx dt \right| \rightarrow 0 \text{ as } \varepsilon \rightarrow 0.$$

Since the vector $X(t) \in C_0^\infty(\Omega)$ at each time, we can define the unique Hodge decomposition $X = \nabla\phi + \mathbb{J}\nabla\psi$ such that

$$\begin{aligned} \Delta\phi(t) &= \operatorname{div} X \text{ in } \Omega & \text{and} & \quad \partial_\nu\phi(t) = 0 \text{ on } \partial\Omega, \\ \Delta\psi(t) &= \operatorname{curl} X \text{ in } \Omega & \text{and} & \quad \psi(t) = 0 \text{ on } \partial\Omega. \end{aligned}$$

We set $\bar{\phi}$ to be the average of ϕ over Ω , and consequently from the Neumann boundary conditions on w_ε ,

$$\begin{aligned} \left| \int_0^T \int_{\Omega} X \cdot \nabla h_1^\varepsilon dx dt \right| &= \left| \int_0^T \int_{\Omega} (\nabla\phi + \mathbb{J}\nabla\psi) \cdot \nabla h_1^\varepsilon dx dt \right| \\ &= \left| \int_0^T \int_{\Omega} (\phi - \bar{\phi}) \operatorname{div} j(w_\varepsilon(t)) dx dt \right| \\ &\leq \left| \int_0^T \int_{\Omega} (\phi - \bar{\phi}) \frac{\nabla\eta_\varepsilon^2 \cdot j(w_\varepsilon)}{\eta_\varepsilon^2} dx dt \right| + \left| \int_0^T \int_{\Omega} \frac{|\partial_t(\phi - \bar{\phi})|}{2\eta_\varepsilon^2} |1 - |w_\varepsilon|^2| dx dt \right| \\ &\leq \frac{C}{|\log \varepsilon|} \|X\|_{L^4(\Omega \times [0,T])} + C\varepsilon \|\partial_t X\|_{L^2(\Omega \times [0,T])} = o_\varepsilon(1), \end{aligned}$$

where we used a Hodge decomposition of $\partial_t X$ that arises from differentiating ϕ and ψ in time. In order to control the term with h_2^ε , we use elliptic estimates and bound

$$\|\nabla h_2^\varepsilon(t)\|_{L^p} \leq C \|J(w_\varepsilon(t)) - J(w_*(t))\|_{W^{-1,p}} \leq C \|J(w_\varepsilon(t)) - J(w_*(t))\|_{W^{-1,1}} = o_\varepsilon(1).$$

This implies $|\int_{t_0}^{t_0+\tau} \int_{\Omega} X \cdot \mathbb{J}\nabla h_2^\varepsilon dx dt| = o_\varepsilon(1)$, and the three bounds combine together to yield (3.4). \square

With Lemmas 3.2 and 3.3 in hand, we can now turn to the proof of the main result.

Proof of Theorem 1.2. Define

$$\mu(t) \equiv \sum_j |\alpha_j(t) - \xi_j(t)| = \sum_j \mu_j(t).$$

We will estimate the difference between the vortex paths ξ_j , generated by (1.4), and the positions a_j , extracted by the ODE (1.11). We also let $D_\varepsilon(w_\varepsilon(t), b(t)) = E_\varepsilon^{\eta_\varepsilon}(w_\varepsilon(t)) - W_\varepsilon(b(t))$ denote the time-varying excess energy.

Recall from (1.11) that we will estimate the differences between vortex paths and the ODE positions:

$$(3.6) \quad \begin{aligned} \pi \dot{\alpha}_j &= -\nabla_{\alpha_j}^\perp W(\alpha) + \pi \nabla^\perp Q_0(\alpha_j) \\ &= -\nabla_{\xi_j}^\perp W(\xi) + \pi \nabla^\perp Q_0(\xi_j) + \nabla_{\xi_j}^\perp W(\xi) - \nabla_{\alpha_j}^\perp W(\alpha) + \pi \nabla^\perp Q_0(\alpha_j) - \pi \nabla^\perp Q_0(\xi_j). \end{aligned}$$

Choosing a time interval $0 \leq a < b \leq T$ and using (1.8), (3.2), (3.3), (3.6),

$$(3.7) \quad \begin{aligned} &\pi (\mu_j(b) - \mu_j(a)) \\ &\leq \int_a^b |\nabla_{\xi_j} W(\xi(t)) - \nabla_{\alpha_j} W(\alpha(t))| dt + \pi \int_a^b |\nabla Q_0(\xi_j(t)) - \nabla Q_0(\alpha_j(t))| dt \\ &+ \limsup_{k \rightarrow \infty} \left| \int_a^b \int_{B_r(\alpha_j)} \mathbb{J}_{\ell_p} \partial_p \partial_q \varphi [(\partial_q w_{\varepsilon_k}, \partial_\ell w_{\varepsilon_k}) - (\partial_q w_*(\xi), \partial_\ell w_*(\xi))] dx dt \right| \\ &+ \limsup_{k \rightarrow \infty} \left| \int_a^b \left[\int_{B_r(\alpha_j)} \mathbb{J}_{\ell_p} \partial_p \varphi \frac{\partial_q \eta_{\varepsilon_k}^2}{\eta_{\varepsilon_k}^2} (\partial_\ell w_{\varepsilon_k}, \partial_q w_{\varepsilon_k}) dx - \pi \nabla^\perp Q_0(\xi_j) \right] dt \right| \\ &+ \limsup_{k \rightarrow \infty} \left| \int_a^b \int_{B_r(\alpha_j)} \mathbb{J}_{\ell_p} \partial_p \varphi \partial_\ell \eta_{\varepsilon_k}^2 \frac{(1 - |w_{\varepsilon_k}|^2)^2}{4\varepsilon^2} dx dt \right| \\ (3.7) &= \mathcal{A} + \mathcal{B} + \mathcal{C} + \mathcal{D} + \mathcal{E}. \end{aligned}$$

We can control \mathcal{A} and \mathcal{B} using our Lipschitz bounds; in particular,

$$(3.8) \quad \begin{aligned} \int_a^b |\nabla W(\xi_j(t)) - \nabla W(\alpha_j(t))| dt &\leq \int_a^b \sup_{j: |\xi_j - \alpha_j| \leq r_a/4} |\nabla^2 W(\alpha)| \left(\sum |\xi_j(t) - \alpha_j(t)| \right) dt \\ &\leq C \int_a^b \mu(t) dt, \end{aligned}$$

where we used Lemma 10 of [20] to control the Hessian of $W(a)$, with a bound dependent on r_a . Likewise, we can use Appendix A on the regularity of η_{ε_k} to bound

$$(3.9) \quad \int_a^b |\nabla Q(\xi_j(t)) - \nabla Q(\alpha_j(t))| dt \leq \|Q_0\|_{W^{2,\infty}} \int_a^b \mu(t) dt.$$

We now turn to the control on \mathcal{C} . For shorthand we define components $j_k = \left(\frac{j(w_{\varepsilon_k})}{|w_{\varepsilon_k}|}\right)_k$ and $j_k^* = (j(w_*(\xi)))_k$ of the supercurrents, and since $\partial_k w_{\varepsilon_k} = (\partial_k |w_{\varepsilon_k}| + i j_k) \frac{w_{\varepsilon_k}}{|w_{\varepsilon_k}|}$ then

$$\begin{aligned} \mathcal{C} &\leq \limsup_{k \rightarrow \infty} \left| \int_a^b \int_{\Omega_r(\alpha)} \mathbb{J}_{\ell_p} \partial_p \partial_q \varphi \partial_q |w_{\varepsilon_k}| \partial_\ell |w_{\varepsilon_k}| dx dt \right| \\ &+ \limsup_{k \rightarrow \infty} \left| \int_a^b \int_{\Omega_r(\alpha)} \mathbb{J}_{\ell_p} \partial_p \partial_q \varphi (j_q - j_q^*, j_\ell - j_\ell^*) dx dt \right| \\ &+ \limsup_{k \rightarrow \infty} \left| \int_a^b \int_{\Omega_r(\alpha)} \mathbb{J}_{\ell_p} \partial_p \partial_q \varphi (j_q^* (j_\ell - j_\ell^*)) dx dt \right| \\ &+ \limsup_{k \rightarrow \infty} \left| \int_a^b \int_{\Omega_r(\alpha)} \mathbb{J}_{\ell_p} \partial_p \partial_q \varphi ((j_q - j_q^*) j_\ell^*) dx dt \right| \\ &= \mathcal{C}_1 + \mathcal{C}_2 + \mathcal{C}_3 + \mathcal{C}_4, \end{aligned}$$

and since $D_{\varepsilon_k}(w_{\varepsilon_k}, \xi) = D_{\varepsilon_k}(w_{\varepsilon_k}, \alpha) + W(\xi) - W(\alpha) + \pi Q_0(\xi) - \pi Q_0(\alpha)$, then

$$\begin{aligned} \mathcal{C}_1 + \mathcal{C}_2 &\stackrel{(2.1)}{\lesssim} \int_a^b D_{\varepsilon_k}(w_{\varepsilon_k}, \xi(t)) dt + o_k(1) \\ &\stackrel{(3.8), (3.9)}{\leq} C \int_a^b \mu(t) dt + o_k(1), \end{aligned}$$

since $D_{\varepsilon_k}(w_{\varepsilon_k}(t), \alpha(t)) = D_{\varepsilon_k}(w_{\varepsilon_k}^0, \alpha^0) = o_k(1)$. We set $X = \mathbb{J}_{\ell_p} j_q^* \partial_p \partial_q \varphi$ or $\mathbb{J}_{\ell_p} j_\ell^* \partial_p \partial_q \varphi$, both in $C_0^\infty(\Omega \times [0, T])$, and then from Lemma 3.3, we have that $\lim_{k \rightarrow \infty} \mathcal{C}_3 + \mathcal{C}_4 = 0$; note this is where we needed to consider the time-integral on (a, b) .

Next we control \mathcal{D} .

$$\begin{aligned} \mathcal{D} &\leq \limsup_{k \rightarrow \infty} \left| \int_a^b \int_{B_r(a_j)} \mathbb{J}_{\ell_p} \partial_p \varphi \frac{\partial_q \eta_{\varepsilon_k}^2}{\eta_{\varepsilon_k}^2} \left[(\partial_\ell w_{\varepsilon_k}, \partial_q w_{\varepsilon_k}) - \pi |\log \varepsilon_k| \delta_{\ell q} \sum \delta_{\xi_j(t)} \right] dx dt \right| \\ &\quad + \limsup_{k \rightarrow \infty} \left| \int_a^b \pi |\log \varepsilon_k| \frac{\nabla^\perp \eta_{\varepsilon_k}^2(\xi_j(t))}{\eta_{\varepsilon_k}^2(\xi_j(t))} - \pi \nabla^\perp Q_0(\xi_j(t)) dt \right| \\ &= \mathcal{D}_1 + \mathcal{D}_2. \end{aligned}$$

We can now bound

$$\begin{aligned} \mathcal{D}_1 &\leq T \limsup_{k \rightarrow \infty} \left[\left\| \nabla \left(\partial_p \varphi \frac{\partial_q \eta_{\varepsilon_k}^2}{\eta_{\varepsilon_k}^2} \right) \right\|_{L^\infty} \left\| (\partial_\ell w_{\varepsilon_k}, \partial_q w_{\varepsilon_k}) - \pi |\log \varepsilon_k| \delta_{\ell q} \sum \delta_{\xi_j} \right\|_{\dot{W}^{-1,1}} \right] \\ &\stackrel{(2.5)}{\lesssim} \limsup_{k \rightarrow \infty} \frac{C |\log \varepsilon_k|^{-\frac{1}{2}}}{r_\alpha^2} = o_k(1) \end{aligned}$$

and

$$\mathcal{D}_2 \lesssim \limsup_{k \rightarrow \infty} \int_a^b \left| \frac{\partial_p Q_{\varepsilon_k}(\xi_j)}{\eta_{\varepsilon_k}^2} - \partial_p Q_0(\xi_j) \right| dt = o_k(1).$$

Finally, we bound \mathcal{E} using (2.2):

$$\mathcal{E} \leq \frac{C}{r_a |\log \varepsilon_k|} \int_a^b \int_\Omega \frac{(1 - |w_{\varepsilon_k}(t)|^2)^2}{\varepsilon_k^2} dx dt \leq \frac{CT}{r_a |\log \varepsilon_k|} = o_k(1).$$

Combining together estimates generates a differential inequality,

$$\mu(b) - \mu(a) \leq C \int_a^b \mu(t) dt,$$

and choosing $b = a + \delta$ and dividing by δ implies the classical Gronwall inequality with $\mu(0) = 0$. Therefore, $\mu(t) = 0$ for all $0 \leq t \leq T$. We can then restart the problem at time T and continue the dynamics until $r_a \rightarrow 0$.

So far we have only shown the theorem for a subsequence. However, our argument can be applied to subsequences of any sequence of $\varepsilon_k \rightarrow 0$, and as the limit is independent of the chosen subsequence, we obtain convergence for $\varepsilon \rightarrow 0$ without taking subsequences. \square

4. Second order gamma convergence and gradient flow. In this section we collect and restate some of our results on the static energy in the spirit of Γ -convergence; compare [1] for the corresponding results for $p_\varepsilon \equiv 1$. For dynamics, we show that the approach of [35] can be adapted to treat the gradient flow as opposed to the Schrödinger dynamics studied in the previous sections.

All of our arguments work with either Neumann or Dirichlet boundary conditions, just as those in [1] and [35]. For simplicity, we assume Neumann boundary conditions throughout and sometimes comment on changes necessary for the Dirichlet case.

4.1. Further Γ -convergence results. We discuss Γ -convergence properties for $E_\varepsilon^{\eta_\varepsilon}$. From this, an analysis for the original energy

$$E_\varepsilon(u) = \int_\Omega e_\varepsilon(u) dx = \int_\Omega \frac{1}{2} |\nabla u|^2 + \frac{1}{4\varepsilon^2} (p_\varepsilon^2 - |u|^2)^2 dx$$

can also be deduced by the Lassoued–Mironescu result [26]

$$(4.1) \quad E_\varepsilon(w\eta_\varepsilon) = E_\varepsilon(\eta_\varepsilon) + E_\varepsilon^{\eta_\varepsilon}(w)$$

that follows from the calculation

$$e_\varepsilon(u) = e_\varepsilon(\eta_\varepsilon) + e_\varepsilon^{\eta_\varepsilon}(w) + \frac{1}{2} \operatorname{div}((|w|^2 - 1)\eta_\varepsilon \nabla \eta_\varepsilon).$$

If $|w| = |u| = \eta_\varepsilon = 1$ on $\partial\Omega$ or $\nu \cdot \nabla \eta_\varepsilon = 0$ on $\partial\Omega$, we can deduce (4.1) in the Dirichlet or Neumann case, respectively. However, $E_\varepsilon(\eta_\varepsilon)$ contributes very little to the energy, and all the dynamically interesting information is encoded in $E_\varepsilon^{\eta_\varepsilon}(w)$, so we concentrate on $E_\varepsilon^{\eta_\varepsilon}$ in the following.

PROPOSITION 4.1. *Let $w_\varepsilon \in H^1(\Omega; \mathbb{C})$ be a sequence of functions with*

$$E_\varepsilon^{\eta_\varepsilon}(w_\varepsilon) \leq K_1 |\log \varepsilon|.$$

Then the Jacobians $J(w_\varepsilon)$ are compact in $\dot{W}^{-1,1}(\Omega)$ and for a subsequence,

$$J(w_\varepsilon) \rightarrow J_* = \pi \sum d_i \delta_{a_i},$$

where $a_i \in \Omega$ are distinct points and $d_i \in \mathbb{Z} \setminus \{0\}$. Furthermore,

$$\liminf_{\varepsilon \rightarrow 0} \frac{1}{|\log \varepsilon|} E_\varepsilon^{\eta_\varepsilon}(w_\varepsilon) \geq \pi \sum |d_i| = \|J_*\|_{\mathcal{M}}$$

with $\|J_\|_{\mathcal{M}} \leq K_1$.*

Proof. Comparing with the standard Ginzburg–Landau energy E_ε^1 , we clearly have

$$(4.2) \quad \min_{\Omega} \min(\eta_\varepsilon^2, \eta_\varepsilon^4) E_\varepsilon^1(w_\varepsilon) \leq E_\varepsilon^{\eta_\varepsilon}(w_\varepsilon) \leq \max_{\Omega} \max(\eta_\varepsilon^2, \eta_\varepsilon^4) E_\varepsilon^1(w_\varepsilon).$$

As both the minimum and the maximum can be estimated as $1 + O(\frac{1}{|\log \varepsilon|})$, we obtain that the standard Ginzburg–Landau energy is bounded as

$$E_\varepsilon^1(w_\varepsilon) \leq K_1 |\log \varepsilon| + CK_1.$$

This implies the compactness of $J(w_\varepsilon)$; see [19, 37].

The lower bound for the energy follows from the standard Ginzburg–Landau lower bounds and (4.2). □

The following bound is the heart of the second order Γ -convergence argument.

PROPOSITION 4.2. *Let $B_{r_0}(a) \subset \Omega$. If $J(w_\varepsilon) \rightarrow \pm\pi\delta_\alpha$ in $W^{-1,1}(B_{r_0}(\alpha))$, then*

$$\liminf_{r \rightarrow 0} \liminf_{\varepsilon \rightarrow 0} E_\varepsilon^{\eta_\varepsilon}(w_\varepsilon; B_r(\alpha)) - \pi \log \frac{r}{\varepsilon} \geq \gamma_0 + \pi Q_0(\alpha),$$

where γ_0 is the constant introduced in [4, Lemma IX.1].

Proof. We have shown this in step 4 of the proof of Proposition 2.1.

PROPOSITION 4.3. *Under the assumptions of Proposition 4.1, if we have for some $K_2 > 0$ the sharper upper bound*

$$E_\varepsilon^{\eta_\varepsilon}(w_\varepsilon) \leq \|J_*\|_{\mathcal{M}} |\log \varepsilon| + K_2,$$

then $J_* = \pi \sum_{i=1}^n d_i \delta_{\alpha_i}$ with distinct α_i and $d_i \in \{\pm 1\}$.

Furthermore, (w_ε) are then bounded in $W^{1,p}(\Omega)$ for all $p \in [1, 2)$ and satisfy the energy lower bound

$$\liminf_{\varepsilon \rightarrow 0} (E_\varepsilon^{\eta_\varepsilon}(w_\varepsilon) - \pi n |\log \varepsilon|) \geq n\gamma_0 + W(\alpha_i, d_i) + \pi \sum_{i=1}^n Q_0(\alpha_i),$$

where W is the renormalized energy defined in (1.10).

Also, there is K_3 such that

$$\limsup_{\varepsilon \rightarrow 0} \frac{1}{\varepsilon^2} \int_{\Omega} (1 - |w_\varepsilon|^2)^2 dx \leq K_3.$$

Proof. We clearly have $E_\varepsilon^1(w_\varepsilon) \leq \|J_*\| |\log \varepsilon| + K_4$ with K_4 depending on K_2 and $\|J_*\|$ as well as on Q_0 . Hence we can use Ginzburg–Landau arguments of Colliander and Jerrard [9] or Alicandro and Ponsiglione [1] to obtain the claimed structure of J_* . The L^p bound follows from Proposition 2.1. Let now $r > 0$ be so small that $B_r(a_i)$ are disjoint and do not intersect $\partial\Omega$. Then we apply Proposition 4.2 to obtain

$$\liminf_{r \rightarrow 0} \liminf_{\varepsilon \rightarrow 0} E_\varepsilon^{\eta_\varepsilon} \left(w_\varepsilon; \bigcup B_r(\alpha_i) \right) - \pi n \log \frac{r}{\varepsilon} - n\gamma_0 - \pi \sum Q_0(\alpha_i) \geq 0.$$

In $\Omega_r(\alpha)$, we let w_r be the optimal S^1 valued map as in the definition of the Dirichlet energy W , i.e., the function that minimizes the Dirichlet energy with cut-off radius $r > 0$. Let \hat{w}_* be the subsequential weak limit of w_ε . Then $|\hat{w}_*| = 1$ a.e. and hence by lower semicontinuity of the Dirichlet integral,

$$\int_{\Omega_r(\alpha)} |\nabla w_r|^2 dx \leq \int_{\Omega_r(\alpha)} |\nabla \hat{w}_*|^2 dx \leq \liminf_{\varepsilon \rightarrow 0} \int_{\Omega_r(\alpha)} |\nabla w_\varepsilon|^2 dx.$$

By (4.2), we deduce that also

$$\liminf_{\varepsilon \rightarrow 0} E_\varepsilon^{\eta_\varepsilon}(w_\varepsilon; \Omega_r(\alpha)) \geq \frac{1}{2} \int_{\Omega_r(\alpha)} |\nabla w_r|^2 dx.$$

Using that

$$W(\alpha, d) = \lim_{r \rightarrow 0} \left(\frac{1}{2} \int_{\Omega_r(\alpha)} |\nabla w_r|^2 dx - \pi n \log \frac{1}{r} \right),$$

we obtain the claimed lower bound.

The upper bound for the penalty term has been shown in Proposition 2.1. □

4.2. The gradient flow. Now we study the gradient flow analogue of (1.1),

$$(4.3) \quad \begin{cases} \frac{1}{|\log \varepsilon|} \partial_t u = \Delta u + (p^2(x) - |u|^2)u & \text{in } \Omega, \\ \frac{\partial u}{\partial \nu} = 0 & \text{on } \partial\Omega. \end{cases}$$

Instead of deriving an equation of motion by studying the limits of the energy density, we are going to use the method of Γ -convergence of gradient flows. This has the advantage that we will require slightly less regularity and convergence properties for the potential term $p(x)$ than in the Schrödinger part of this article. We will follow closely the approach of Sandier and Serfaty [35] to obtain the convergence of the gradient flow of $E_\varepsilon^{\eta_\varepsilon}$ to that of F . In fact, the proof goes through with almost no essential changes, and we will therefore assume the reader is familiar with the argument and some of the notation of [35] and will mostly highlight the differences.

Again, we use the approach of dividing by the profile η_ε , leading to an equation for $w_\varepsilon = \frac{u_\varepsilon}{\eta_\varepsilon}$. As we assume Neumann boundary conditions $\partial_\nu \eta_\varepsilon = 0$ for η_ε , then Neumann boundary conditions $\partial_\nu u = 0$ imply Neumann boundary conditions $\partial_\nu w_\varepsilon = 0$. As before, we obtain

$$(4.4) \quad \begin{cases} \frac{1}{|\log \varepsilon|} \eta_\varepsilon^2 \partial_t w_\varepsilon = \operatorname{div}(\eta_\varepsilon^2 \nabla w_\varepsilon) + \frac{\eta_\varepsilon^4}{\varepsilon^2} (1 - |w_\varepsilon|^2) w_\varepsilon & \text{in } \Omega, \\ \frac{\partial w_\varepsilon}{\partial \nu} = 0 & \text{on } \partial\Omega. \end{cases}$$

This is the gradient flow of $E_\varepsilon^{\eta_\varepsilon}$ with respect to the scalar product on $\mathcal{X}_\varepsilon = L^2(\Omega; \mathbb{C})$ given by the quadratic form

$$\|v\|_{\mathcal{X}_\varepsilon}^2 = \frac{1}{|\log \varepsilon|} \int_\Omega \eta_\varepsilon^2 |v|^2 dx.$$

In fact, we have for $\phi \in C^\infty(\Omega; \mathbb{C})$ the Gâteaux derivative

$$dE_\varepsilon^{\eta_\varepsilon}(w_\varepsilon; \phi) = \int_\Omega \eta_\varepsilon^2 \nabla w_\varepsilon \cdot \nabla \phi - \frac{\eta_\varepsilon^4}{\varepsilon^2} (1 - |w_\varepsilon|^2) w_\varepsilon \cdot \phi dx,$$

and on the other hand,

$$\langle \partial_t w_\varepsilon, \phi \rangle_{\mathcal{X}_\varepsilon} = \frac{1}{|\log \varepsilon|} \int_\Omega \eta_\varepsilon^2 \partial_t w_\varepsilon \phi dx,$$

so integrating by parts we see that (4.4) is a gradient flow, more precisely, $\partial_t w_\varepsilon = -\nabla_{\mathcal{X}_\varepsilon} E_\varepsilon^{\eta_\varepsilon}(w)$.

We will show that we can relate this gradient flow to that of $F(\alpha)$, where for $\alpha = (\alpha_1, \dots, \alpha_n) \in \Omega^n$, $\alpha_i \neq \alpha_j$ for $i \neq j$, we set

$$F(\alpha) = W(\alpha, d) + \pi \sum_{j=1}^n Q_0(\alpha_j)$$

and we will use the following metric on $\mathcal{Y} = \mathbf{R}^{2n}$ (the tangential space to Ω^n):

$$\|b\|_{\mathcal{Y}}^2 = \pi \sum_{j=1}^n |b_j|^2.$$

THEOREM 4.4. *Assume that the profiles η_ε solving (1.3) satisfy $Q_\varepsilon = |\log \varepsilon|(\eta_\varepsilon^2 - 1) \rightarrow Q_0$ in $C^1(\bar{\Omega})$.*

Let (w_ε) be a family of solutions of (4.4) with initial data w_ε^0 . Assume that w_ε^0 satisfy $J(w_\varepsilon^0) \rightarrow \pi \sum d_j \delta_{\alpha_j^0}$ in $\dot{W}^{-1,1}$ with $d_j \in \{\pm 1\}$.

Let $\alpha(t) : [0, T^m) \rightarrow \mathbf{R}^{2n}$ be a solution of the system of ODEs

$$(4.5) \quad \pi \dot{\alpha}_j(t) = -\nabla_{\alpha_j} W(\alpha(t), d) - \pi \nabla Q_0(\alpha_j(t)),$$

where $T^m = T(\alpha^0) \in (0, \infty]$ is the maximal time of existence for (4.5), i.e., the smallest time T such that as $t \rightarrow T$, we have $r_{\alpha(t)} \rightarrow 0$.

If the initial data are very well-prepared, i.e., $D_\varepsilon(0) = E_\varepsilon^{\eta_\varepsilon}(w_\varepsilon^0) - H_\varepsilon(\alpha^0) \rightarrow 0$ as $\varepsilon \rightarrow 0$, then for $0 \leq t < T_m$, $J(w_\varepsilon(t)) \rightarrow \pi \sum d_j \delta_{\alpha_j(t)}$ so we have convergence of the gradient flow of $E_\varepsilon^{\eta_\varepsilon}$ to that of F . Furthermore, $D_\varepsilon(t) = E_\varepsilon^{\eta_\varepsilon}(w_\varepsilon(t)) - H_\varepsilon(\alpha(t)) \rightarrow 0$ so the data continue to be well-prepared. Finally, we have for $T < T^m$ and any $B_i(t)$ that are disjoint open balls contained in Ω and centered at $\alpha_i(t)$ that

$$(4.6) \quad \frac{1}{|\log \varepsilon|} \int_0^T \int_\Omega \eta_\varepsilon^2 |\partial_t w_\varepsilon - \chi_{B_i(t)} \dot{\alpha}_i(t) \cdot \nabla w_\varepsilon|^2 dx dt \rightarrow 0.$$

Remark 4.5. See Proposition 1.6 for conditions on the potential p_ε that imply the convergence of Q_ε assumed in the theorem. Note that we require less regularity and convergence than for the Schrödinger case.

Proof of Theorem 4.4. The proof is mostly an exercise in changing E_ε^1 to $E_\varepsilon^{\eta_\varepsilon}$ in the corresponding results in [35]. The main observation, which we will use repeatedly below, is that for any sequence of $f_\varepsilon \in L^2(\Omega)$ we have

$$(4.7) \quad \left(1 + \frac{C}{|\log \varepsilon|}\right)^{-1} \int_\Omega |f_\varepsilon|^2 dx \leq \int_\Omega \eta_\varepsilon^2 |f_\varepsilon|^2 dx \leq \left(1 + \frac{C}{|\log \varepsilon|}\right) \int_\Omega |f_\varepsilon|^2 dx,$$

so in particular if $\int_\Omega \eta_\varepsilon^2 |f_\varepsilon|^2 dx \leq K|\log \varepsilon|$, then $\int_\Omega |f_\varepsilon|^2 dx \leq K|\log \varepsilon| + KC$, and a similar reverse inequality.

We note that any solution of (4.4) satisfies for $t_1 < t_2$

$$E_\varepsilon^{\eta_\varepsilon}(w_\varepsilon(t_2)) = E_\varepsilon^{\eta_\varepsilon}(w_\varepsilon(t_1)) + \frac{1}{|\log \varepsilon|} \int_{t_1}^{t_2} \int_\Omega \eta_\varepsilon^2 |\partial_t w_\varepsilon|^2 dx dt$$

as follows from multiplying (4.4) with $\partial_t w_\varepsilon$ and integrating by parts. In particular, $s \mapsto E_\varepsilon^{\eta_\varepsilon}(w_\varepsilon(s))$ is weakly decreasing and

$$\int_{t_1}^{t_2} \int_\Omega \eta_\varepsilon^2 |\partial_t w_\varepsilon|^2 dx dt \leq C|\log \varepsilon|^2.$$

Essentially verbatim as in [35, Lemma 3.4], we can show the existence of $T_0 > 0$ such that actually

$$\int_0^{T_0} \int_\Omega \eta_\varepsilon^2 |\partial_t w_\varepsilon|^2 dx dt \leq C|\log \varepsilon|,$$

which implies

$$\int_0^{T_0} \int_\Omega |\partial_t w_\varepsilon|^2 dx dt \leq C|\log \varepsilon| + o(|\log \varepsilon|).$$

In addition, we also have for all $t \in [0, T_0)$ with $n = \sum |d_j^0|$ the estimate $E_\varepsilon^{\eta_\varepsilon}(w_\varepsilon(t)) \leq \pi n |\log \varepsilon| + C$ and so $E_\varepsilon^1(w_\varepsilon(t)) \leq \pi n |\log \varepsilon| + o(|\log \varepsilon|)$.

We can now apply Propositions 3.2 and 3.3 of [35] to w_ε and obtain that (for a subsequence) $J(w_\varepsilon(t)) \rightarrow \pi \sum d_j \delta_{b_j(t)}$, where $b_j \in H^1(0, T_0; \mathbf{R}^2)$ with $b_j(0) = \alpha_j^0$, i.e., the d_j are constant. Corollary 7 of [36] now applies to show

$$\liminf_{\varepsilon \rightarrow 0} \frac{1}{|\log \varepsilon|} \int_{t_1}^{t_2} \int_{\Omega} |\partial_t w_\varepsilon|^2 dx dt \geq \pi \sum \int_{t_1}^{t_2} |\dot{b}_i|^2 dt.$$

As $|\eta_\varepsilon^2 - 1| \leq \frac{C}{|\log \varepsilon|}$, this also implies

$$(4.8) \quad \liminf_{\varepsilon \rightarrow 0} \frac{1}{|\log \varepsilon|} \int_{t_1}^{t_2} \int_{\Omega} \eta_\varepsilon^2 |\partial_t w_\varepsilon|^2 dx dt \geq \pi \sum \int_{t_1}^{t_2} |\dot{b}_i|^2 dt.$$

The last equation is the lower bound part needed for the Γ -convergence of gradient flows argument.

The construction also proceeds almost verbatim as in Proposition 3.5 of [35]. Let w_ε satisfy $J(w_\varepsilon) \rightarrow \pi \sum d_j \delta_{\alpha_j}$ and $D_\varepsilon(w_\varepsilon, \alpha) \leq C$ and assume that $\|\nabla_{\mathcal{X}_\varepsilon} E_\varepsilon^{\eta_\varepsilon}(w_\varepsilon)\|_{\mathcal{X}_\varepsilon} \leq C$. This implies that

$$\int_{\Omega} \frac{1}{\eta_\varepsilon^2} \left| \operatorname{div}(\eta_\varepsilon^2 \nabla w_\varepsilon) + \frac{\eta_\varepsilon^4}{\varepsilon^2} (1 - |w_\varepsilon|^2) w_\varepsilon \right|^2 dx \leq \frac{C}{|\log \varepsilon|}.$$

We set

$$f_\varepsilon = \operatorname{div}(\eta_\varepsilon^2 \nabla w_\varepsilon) + \frac{\eta_\varepsilon^4}{\varepsilon^2} (1 - |w_\varepsilon|^2) w_\varepsilon$$

and note that $f_\varepsilon \rightarrow 0$ in L^2 and $(i w_\varepsilon, f_\varepsilon) = (i w_\varepsilon, \operatorname{div}(\eta_\varepsilon^2 \nabla w_\varepsilon))$. Furthermore

$$\operatorname{div} j(w_\varepsilon) = (i w_\varepsilon, \operatorname{div} \nabla w_\varepsilon) = (i w_\varepsilon, \operatorname{div}(\eta_\varepsilon^2 \nabla w_\varepsilon)) - \nabla(\eta_\varepsilon^2) \cdot j(w_\varepsilon).$$

By (2.2) we see $|w_\varepsilon| \rightarrow 1$. From (2.3) together with $|\nabla(\eta_\varepsilon^2)| \rightarrow 0$ we deduce $\operatorname{div} j(w_\varepsilon) \rightarrow 0$ in $L^2(\Omega)$. From this point we can continue as in [35], as now both $\operatorname{curl} j(w_\varepsilon)$ and $\operatorname{div} j(w_\varepsilon)$ have the same properties as the analogous quantities in [35].

Now let $V \in (\mathbf{R}^2)^n$ and let $b(t)$ be a curve satisfying $b(0) = \alpha$ and $\partial_t b(0) = V$. We claim that there exists a path $v_\varepsilon(t)$ with $v_\varepsilon(0) = w_\varepsilon$ and the properties

$$(4.9) \quad \|\partial_t v_\varepsilon(0)\|_{\mathcal{X}_\varepsilon}^2 = \|\partial_t b(0)\|_{\mathcal{Y}}^2 + o(1),$$

$$(4.10) \quad \lim_{\varepsilon \rightarrow 0} \frac{d}{dt} \Big|_{t=0} E_\varepsilon^{\eta_\varepsilon}(v_\varepsilon(t)) = \frac{d}{dt} \Big|_{t=0} F(b(t)) + g(\alpha) D_\varepsilon$$

for some function g that is locally bounded in $\{a \in \Omega^n : r_\alpha > 0\}$.

To construct this path in function space, we use the same “push” map as in [35]. Let $B_i = B_\rho(\alpha_i)$ with $\rho < r_\alpha$ be pairwise disjoint balls and define $\chi_t : \Omega \rightarrow \Omega$ to be a one-parameter family of diffeomorphisms that satisfies

$$\chi_t(x) = x + tV_i \quad \text{in } B_i.$$

With the phase corrector function ψ_t defined as in (3.24) of [35], set

$$v_\varepsilon(\chi_t(x), t) = w_\varepsilon(x) e^{i\psi_t(x)}.$$

The claim (4.9) follows as in [35], keeping in mind our observation (4.7).

To calculate the energy change along this path, we change variables $y = \chi_t(x)$ and obtain with $\text{Jac}(\chi_t)$ denoting the Jacobian determinant,

$$\begin{aligned} E_\varepsilon^{\eta_\varepsilon}(v_\varepsilon) &= \int_\Omega \frac{1}{2} \eta_\varepsilon^2(y) |\nabla v_\varepsilon(y)|^2 + \frac{\eta_\varepsilon^4(y)}{4\varepsilon^2} (1 - |v_\varepsilon(y)|^2)^2 dy \\ &= \int_\Omega \left(\frac{1}{2} (\eta_\varepsilon \circ \chi_t(x))^2 |D\chi_t^{-1} \nabla(w_\varepsilon e^{i\psi_t})|^2 \right. \\ &\quad \left. + \frac{(\eta_\varepsilon \circ \chi_t)^4}{4\varepsilon^2} (1 - |w_\varepsilon(x)|^2)^2 |\text{Jac}(\chi_t)| \right) dx. \end{aligned}$$

We now differentiate in time and set $t = 0$. When the derivative does not hit $\eta_\varepsilon \circ \chi_t$, the resulting terms can be dealt with exactly as in [35] by our observation (4.7). The remaining terms are

$$\mathcal{A}_1 = \int_\Omega \frac{1}{2} \frac{d}{dt} \Big|_{t=0} (\eta_\varepsilon \circ \chi_t)^2 |\nabla w_\varepsilon|^2 dx$$

and

$$\mathcal{A}_2 = \int_\Omega \frac{1}{\varepsilon^2} \frac{d}{dt} \Big|_{t=0} (\eta_\varepsilon \circ \chi_t)^4 (1 - |w_\varepsilon|^2)^2 dx.$$

We note that in B_i , $\frac{d}{dt} \Big|_{t=0} (\eta_\varepsilon \circ \chi_t)^2 = V_i \cdot \nabla(\eta_\varepsilon^2)$. Using the uniform convergence $\frac{1}{|\log \varepsilon|} \nabla(\eta_\varepsilon^2) \rightarrow \nabla Q_0$ and the fact that the logarithmic part of the energy is concentrated in the B_i (recall Proposition 2.1 or (3.17) of [35]), we deduce that $\mathcal{A}_1 \rightarrow \pi \sum V_i \cdot \nabla Q_0(\alpha_i)$ as $\varepsilon \rightarrow 0$. To show that $\mathcal{A}_2 \rightarrow 0$, we recall (2.2) and note that $|\nabla(\eta_\varepsilon^4)| \rightarrow 0$ uniformly in Ω . Together with the results of [35] for the terms not involving derivatives of η_ε , we deduce (4.10).

On the interval $[0, T_0)$, we can now apply Theorem 1.4 of [35] and obtain the claim of Theorem 4.4 up to the time T_0 . The global statement up to T^m follows as in section 3.3 of [35]. \square

5. Numerical simulations of the vortex dynamics. Armed with the dynamical equation (1.11), we can simulate the flow of vortices and observe the impact of the background potential at the scales we have studied in Theorem 1.2. For our simulations, we have used the renormalized energy

$$W(\alpha, d) = -\pi \sum_{j \neq k} d_j d_k \log |\alpha_j - \alpha_k|,$$

which is the correct expression for $\Omega = \mathbf{R}^2$; see Remark 1.5. This is an approximation for the renormalized energy for $\Omega = B_R(0)$ for $R \gg 1$ sufficiently large compared to $|\alpha|$. We will focus here on the case of the dipole (a pair of vortices of opposite charge) interacting with potentials $Q_0 = V_i$ of the form

(5.1) $V_1 = e^{-|\bar{x}|^2}$ single Gaussian,

(5.2) $V_2 = .225 \tanh(x)$ step function to different material background,

(5.3) $V_3 = e^{-|\bar{x}-(1,0)|^2} + e^{-|\bar{x}+(1,0)|^2}$ double Gaussian,

(5.4) $V_4 = \sum_{j,k=-15}^{15} e^{-|\bar{x}-(j,k)|^2}$ lattice of Gaussians.




FIG. 1. Dipole dynamics for $d_1 = 1$, $d_2 = -1$ plotted over contours of Q_0 for the cases of 1. The single Gaussian (5.1) with $\vec{\alpha}_1(0) = (-5.0, 2.0)$, $\vec{\alpha}_2(0) = (-4.99, 1.99)$ (top left). 2. A smooth transition function (5.2) with $\vec{\alpha}_1(0) = (-5.0, -2.0)$, $\vec{\alpha}_2(0) = (-4.99, -1.99)$ (top right). 3. A symmetric double-well Gaussian potential (5.3) with $\vec{\alpha}_1(0) = (-0.01, 1.0)$, $\vec{\alpha}_2(0) = (0.01, 1.0)$ (bottom left). 4. A square lattice array of Gaussians (5.4) with $\vec{\alpha}_1(0) = (-5.0, -1.99)$, $\vec{\alpha}_2(0) = (-4.99, -2.0)$.

While these potentials are not compactly supported, we can apply Theorem 1.2 by truncating the potentials outside a suitably large domain without affecting the dynamics we are plotting.

Recall that in the absence of the background potential, dipole dynamics simply move in a straight line perpendicular to that connecting the vortex centers at a speed correlated to the vortex spacing.

Equations (1.11) for each choice of background are then plugged into the *ode15s* ODE solver in MATLAB and integrated over time scales long enough to observe the impact of the background potential on the dipole dynamics. The results are recorded graphically in Figure 1.

Appendix A. Estimates on η_ε . We provide here the details required to prove Proposition 1.6. We seek to understand bounds on the function η_ε defined by

$$(A.1) \quad \Delta\eta_\varepsilon = -\frac{1}{\varepsilon^2}(p_\varepsilon^2 - \eta_\varepsilon^2)\eta_\varepsilon,$$

with Neumann boundary conditions, where p_ε represents the IGL background on a bounded domain $\Omega \subset \mathbf{R}^2$.

Proof of Proposition 1.6. 1. We first claim there exists an H^{k+2} solution for $k \in \mathbb{N}$.

We use a slightly different ansatz for p_ε and η_ε :

$$p_\varepsilon = 1 + \frac{\tilde{\rho}_\varepsilon}{|\log \varepsilon|}, \quad \eta_\varepsilon = 1 + \frac{\tilde{Q}_\varepsilon}{|\log \varepsilon|}.$$

Recall that we will assume that $p_\varepsilon \in H^k$ for $k \geq 2$ with $\varepsilon^2 \tilde{\rho}_\varepsilon \rightarrow 0$ in H^{k+1} as $\varepsilon \rightarrow 0$ and ∇p_0 is compactly supported strictly on the interior of Ω . The last assumption in

particular will be used for simplicity to allow convergence upto the boundary, though this likely can be relaxed. We first claim that η_ε is the unique positive solution of the following minimization problem:

$$(A.2) \quad \eta_\varepsilon = \arg \min_{\eta \in H^1(\Omega; \mathbf{R}^1)} \int_\Omega \frac{1}{2} |\nabla \eta|^2 + \frac{1}{4\varepsilon^2} (p_\varepsilon^2 - \eta^2)^2 dx.$$

Since we are working on a bounded set Ω , such a nontrivial η_ε exists in H^1 for ε sufficiently small and p_ε sufficiently bounded in H^1 using a direct method and Rellich–Kondrachov. We can improve the regularity away from the boundary. In particular if $\tilde{\rho}_\varepsilon \in H^k$, then $\eta_\varepsilon \in H^{k+2}$, even though the H^{k+2} norm blows up as $\varepsilon \rightarrow 0$. This follows from standard elliptic theory results involving nonlinear Sobolev embeddings to bootstrap regularity; see, for instance, [8, section 2.4]. The existence of Euler–Lagrange equations, strong solutions, and classical solutions follows.

As noted in (1.19) this is a slight shift of notation, as the Q_0 defined in the main text is not quite the limit of this family of functions; rather it takes the form

$$Q_0 = \lim_{\varepsilon \rightarrow 0} \left[2\tilde{Q}_\varepsilon + \frac{\tilde{Q}_\varepsilon^2}{|\log \varepsilon|} \right].$$

The Euler–Lagrange perturbation equation is written as

$$(A.3) \quad 0 = \varepsilon^2 \Delta \tilde{Q}_\varepsilon + |\log \varepsilon| \left(1 + \frac{\tilde{Q}_\varepsilon}{|\log \varepsilon|} \right) \left(\frac{2\tilde{\rho}_\varepsilon}{|\log \varepsilon|} + \frac{\tilde{\rho}_\varepsilon^2}{|\log \varepsilon|^2} - \frac{2\tilde{Q}_\varepsilon}{|\log \varepsilon|} - \frac{\tilde{Q}_\varepsilon^2}{|\log \varepsilon|^2} \right),$$

and so

$$-\frac{\varepsilon^2}{2} \Delta \tilde{Q}_\varepsilon = (\tilde{\rho}_\varepsilon - \tilde{Q}_\varepsilon) \left[\left(1 + \frac{\tilde{Q}_\varepsilon}{|\log \varepsilon|} \right) \left(1 + \frac{\tilde{\rho}_\varepsilon}{2|\log \varepsilon|} + \frac{\tilde{Q}_\varepsilon}{2|\log \varepsilon|} \right) \right].$$

Then we can write this as

$$(A.4) \quad \left(1 - \frac{1}{|\log \varepsilon|} \left(\tilde{\rho}_\varepsilon + \frac{\tilde{\rho}_\varepsilon^2}{2|\log \varepsilon|} \right) - \frac{\varepsilon^2}{2} \Delta \right) \tilde{Q}_\varepsilon = \tilde{\rho}_\varepsilon + \frac{\tilde{\rho}_\varepsilon^2}{2|\log \varepsilon|} - \frac{\tilde{Q}_\varepsilon^2}{2|\log \varepsilon|} + \frac{\tilde{Q}_\varepsilon^3}{|\log \varepsilon|^2}.$$

2. We next establish ε -dependent estimates on a model problem.

The required estimates follow by, for instance, carefully modifying [13, Theorem 7.32], which draws upon ideas originally put forth by Nirenberg [33]. The proof relies upon an analysis of regularity up to the boundary by flattening locally to the half-plane and using the Green’s function there, application of difference operators to gain regularity, and induction on regularity estimates for elliptic problems with Neumann boundary problems. Essentially, the argument boils down to the fact that the operator is coercive for each ε , however. The equation also has a unique set of solutions by the same property.⁴

⁴Perhaps the closest argument of this type for Neumann boundary conditions can be found in [43, Chapter 5.7, Propositions 7.4 and 7.5], where it states that for a smooth enough domain $\Omega \subset \mathbb{R}^2$, there exists a unique solution v to the elliptic equation $(-\Delta + 1)v = f$ in Ω , $\frac{\partial v}{\partial \nu} = 0$ on $\partial\Omega$, and for all $k = 0, 1, 2, \dots$, given $f \in H^k$, we have $\|v\|_{H^{k+2}(\Omega)}^2 \leq C \|\Delta v\|_{H^k(\Omega)}^2 + C \|v\|_{H^{k+1}(\Omega)}^2$. Since we need precise control on the constants of our estimate with respect to ε for a perturbation of this equation, which are not available by rescaling, we have included a proof for completeness.

To establish the bounds we need more precisely, we follow [13, Theorems 7.32 and 7.29], where the author studies Sobolev estimates on coercive elliptic equations. In our setting, these equations take the form

$$L_\varepsilon w := \left(1 + \frac{1}{|\log \varepsilon|} A - \frac{\varepsilon^2}{2} \Delta\right) w = f, \quad \partial_\nu w = 0,$$

on Ω smooth enough, for A a bounded function of the same regularity as p .

We will first consider this equation on half-balls of radius s with flat boundary, say, $N(s)$, where y parametrizes the boundary and x parametrizes the interior. We claim we have an estimate of the form

$$(A.5) \quad \varepsilon^2 \|w\|_{H^{k+2}} + \|w\|_{H^k} \leq C \|f\|_{H^k}.$$

To see this, we first claim that

$$(A.6) \quad \varepsilon^2 \|\partial_x^\gamma w\|_{H^1(N(\tilde{r}))} \leq \|f\|_{H^k(N(r))}$$

with $\tilde{r} < r$, with $\gamma \leq k + 1$. For $\gamma = 0$, this is exactly the coercivity estimate. Otherwise, we use an inductive argument constructed via difference operators inside the Dirichlet form $D(\Delta_h^\gamma \zeta w, \Delta_h^\gamma \zeta w)$ for ζ a cut-off to N that vanishes on the curved part of the boundary of N . Commuting with the cut-off function and integrating by parts when necessary, we observe

$$\begin{aligned} \varepsilon^2 \|\Delta_h^\gamma \zeta w\|_{H^1(N(\tilde{r}))} + \left(1 - \frac{\max |A|}{|\log \varepsilon|}\right) \|\Delta_h^\gamma \zeta w\|_{L^2(N(\tilde{r}))} &\leq D(\Delta_h^\gamma \zeta w, \Delta_h^\gamma \zeta w) \\ &\leq C \varepsilon^2 \|\Delta_h^\gamma \zeta w\|_{H^1(N(\tilde{r}))} (\|f\|_{H^k(N(r))} + \|\partial^{\gamma-1} w\|_{L^2}), \end{aligned}$$

where then by induction we have

$$\begin{aligned} \varepsilon^2 \|\Delta_h^\gamma \zeta w\|_{H^1(N(\tilde{r}))} + \left(1 - \frac{\max |A|}{|\log \varepsilon|}\right) \|\Delta_h^\gamma \zeta w\|_{L^2(N(\tilde{r}))} &\leq D(\Delta_h^\gamma \zeta w, \Delta_h^\gamma \zeta w) \\ &\leq C \|f\|_{H^k(N(r))}. \end{aligned}$$

Similarly, we claim that

$$(A.7) \quad \|\partial_x^\gamma w\|_{L^2(N(\tilde{r}))} \leq \|f\|_{H^k(N(r))}$$

for $0 \leq \gamma < k$. This follows by instead putting all the derivatives onto f and using L^2 instead of H^1 norms for the f term on the right-hand side.

At the boundary, we can then establish (A.5), again proven using induction. To see this, we recognize that if γ is a multi-index and $\gamma_2 = 0$ or 1, then the estimate follows by (A.6) and (A.7). We will use the fact that

$$\partial_y^2 w = -\frac{2}{\varepsilon^2} \left(f + \frac{\varepsilon^2}{2} \partial_x^2 w - w \right).$$

If the number of derivatives on y is 0 or 1, we use the above estimate. For more than that, we pull off the first two derivatives in y , make the substitution, then use the inductive hypothesis.

Then, once such estimates are established on half-balls, we can create a sequence of cut-off functions $U_0 = B(0, R), \dots, U_k = V_0, \bar{U}_{j+1} \subset U_j$, where $V_0 \cup V_1 \cup \dots \cup V_M = B(0, R)$ and V_j can be mapped to a half-ball for $j > 0$ and W_0 is an interior region.

Letting ζ_j be a cut-off to U_j , on the interior we have

$$\begin{aligned} \varepsilon^2 \|w\|_{H^{j+2}(U_{j+1})} + \left(1 - \frac{\max |A|}{|\log \varepsilon|}\right) \|w\|_{H^j(U_{j+1})} &\leq \varepsilon^2 \|\zeta_j w\|_{H^{j+2}(U_{j+1})} + \|\zeta_j w\|_{H^j(U_{j+1})} \\ &\leq C \|L\zeta_j w\|_{H^j} \\ &\leq C \|\zeta_j f\|_{H^j} + \varepsilon^2 \|w\|_{H^{j+1}(U_j)}, \end{aligned}$$

from which the result follows via induction. On regions identified with a half-ball, we apply (A.5).

3. We can use the linear estimates (A.5) to generate a posteriori estimates on the sequence. From the energetic formulation, we have using that $\eta = p_\varepsilon$ provides a natural set of bounds the observation

$$\|p_\varepsilon^2 - \eta_\varepsilon^2\|_{L^2}^2 \leq 2\varepsilon^2 \|\nabla p_\varepsilon\|_{L^2}^2.$$

This gives

$$\|\tilde{\rho}_\varepsilon - \tilde{Q}_\varepsilon\|_{L^2} \leq \sqrt{2}\varepsilon \|\nabla \tilde{\rho}_\varepsilon\|_{L^2}$$

or

$$\|\tilde{Q}_\varepsilon\|_{L^2} \leq C \|\tilde{\rho}_\varepsilon\|_{H^1}.$$

By a similar line of reasoning, we know that

$$\|\nabla \tilde{Q}_\varepsilon\|_{L^2} \leq \|\tilde{\rho}_\varepsilon\|_{H^1},$$

and hence

$$\|\tilde{Q}_\varepsilon\|_{H^1} \leq C \|\tilde{\rho}_\varepsilon\|_{H^1}.$$

We observe directly from (A.4) that

$$\|\tilde{Q}_\varepsilon\|_{H^k} \leq C \left(\|\tilde{\rho}_\varepsilon + \frac{\tilde{\rho}_\varepsilon^2}{2|\log \varepsilon|}\|_{H^k} + \frac{1}{|\log \varepsilon|} \|\tilde{Q}_\varepsilon\|_{H^k}^2 + \frac{1}{|\log \varepsilon|^2} \|\tilde{Q}_\varepsilon\|_{H^k}^3 \right),$$

which gives a uniform control on \tilde{Q}_ε in H^k via a boot-strapping argument. Similarly, we can rearrange (A.4) to observe

$$(A.8) \quad \left(1 - \frac{\varepsilon^2}{2}\Delta\right) (\tilde{Q}_\varepsilon - \tilde{\rho}_\varepsilon) = \frac{\varepsilon^2}{2}\Delta\tilde{\rho}_\varepsilon + \frac{(\tilde{\rho}_\varepsilon - \tilde{Q}_\varepsilon)}{|\log \varepsilon|} \mathcal{P}(\tilde{Q}_\varepsilon, \tilde{\rho}_\varepsilon),$$

where

$$(A.9) \quad \mathcal{P}(\tilde{Q}_\varepsilon, \tilde{\rho}_\varepsilon) := \frac{\tilde{\rho}_\varepsilon}{2} + \frac{3\tilde{Q}_\varepsilon}{2} + \frac{(\tilde{Q}_\varepsilon(\tilde{\rho}_\varepsilon + \tilde{Q}_\varepsilon))}{2|\log \varepsilon|}.$$

Hence, applying the linear estimates, we easily observe

$$\|\tilde{Q}_\varepsilon - \tilde{\rho}_\varepsilon\|_{H^{k-1}} \leq C\varepsilon^2 \|\tilde{\rho}_\varepsilon\|_{H^{k+1}},$$

which converges to 0 as $\varepsilon \rightarrow 0$ for $\tilde{\rho}_\varepsilon$ sufficiently regular. Note that we have used here that $\nabla \rho_\varepsilon$ has compact support in order to integrate by parts. To remove this condition otherwise would require further work controlling the error terms relating to boundary condition of ρ_ε and potentially restrict us to local convergence estimates.

4. To finish the a posteriori convergence, we need to get to H^k convergence. Recall the equation

$$(1 - \varepsilon^2 \Delta)q = g, \quad \partial_\nu q = 0,$$

which we rewrite as

$$(A.10) \quad (q - g) - \varepsilon^2 \Delta q = 0, \quad \partial_\nu q = 0.$$

We will replace q with \tilde{Q}_ε and g by the right-hand side of our elliptic equation.

For higher regularity, we have the formal calculation,

$$\begin{aligned} \int |\partial^k(q - g)|^2 dx &= \varepsilon^2 \int \partial^k(q - g) \partial^k \Delta q dx \\ &= -\varepsilon^2 \int |\nabla \partial^k q|^2 + \frac{\varepsilon^2}{2} \|\partial^k q\|_{L^2}^2 + \frac{\varepsilon^2}{2} \|\partial^k g\|_{L^2}^2 \\ &\quad + \varepsilon^2 \int_{\partial\Omega} \partial_\nu \partial^k q (\partial^k q - \partial^k g) ds, \end{aligned}$$

where the boundary terms can be controlled by careful use of the Neumann boundary condition and the equations.

To establish the result rigorously up to the boundary, we must repeat the argument as in step 2, but acting Δ_h^γ on both sides of (A.10) and then multiplying both sides by $\Delta_h^\gamma(q - g)$ and integrating. Specifically, we can consider the coercive Dirichlet form

$$D(\Delta_h^k \zeta(q - g), \Delta_h^k \zeta(q - g)) + \varepsilon^2 D(\nabla_h \Delta_h^k \zeta q, \nabla_h \Delta_h^k \zeta q)$$

both in neighborhoods of the boundary as well as in the interior to get elliptic estimates as in step 2.

5. To see the uniqueness, let $\eta_j = 1 + \frac{\tilde{Q}_j}{|\log \varepsilon|}$, $j = \{1, 2\}$, be two solutions to (A.1) with Neumann boundary conditions for the same $p_\varepsilon = 1 + \frac{\tilde{\rho}_\varepsilon}{|\log \varepsilon|}$ with $\tilde{\rho}_\varepsilon \in H^k$ for $k > 2$. Then set $w = \eta_1 - \eta_2$; w solves

$$\begin{cases} -\Delta w = \frac{1}{\varepsilon^2} (wp_\varepsilon^2 - (\eta_1^3 - \eta_2^3)) & \text{in } \Omega, \\ \partial_\nu w = 0 & \text{on } \partial\Omega. \end{cases}$$

Note that $\eta_1^3 - \eta_2^3 = w(\eta_1^2 + \eta_1\eta_2 + \eta_2^2) = w(3 + \frac{3}{|\log \varepsilon|}(\tilde{Q}_1 + \tilde{Q}_2) + \frac{1}{|\log \varepsilon|^2}(Q_1^2 + Q_1Q_2 + Q_2^2))$. Multiply by w and integrate over Ω , using the boundary condition

$$\begin{aligned} &\int |\nabla w|^2 + \frac{2}{\varepsilon^2} w^2 dx \\ &= \frac{1}{|\log \varepsilon|^2} \int w^2 \left[2\rho_\varepsilon - 3(Q_1 + Q_2) + \frac{1}{|\log \varepsilon|^2} (\rho_\varepsilon^2 - \tilde{Q}_1^2 - \tilde{Q}_1\tilde{Q}_2 - \tilde{Q}_2^2) \right] dx \\ &\leq \frac{C}{|\log \varepsilon|} \left(1 + \|\rho_\varepsilon\|_{H^k}^2 \right) \left(1 + \|\tilde{Q}_1\|_{H^k}^2 \right) \left(1 + \|\tilde{Q}_2\|_{H^k}^2 \right) \frac{1}{\varepsilon^2} \int w^2 dx \\ &\leq \frac{1.5}{\varepsilon^2} \int w^2 dx \end{aligned}$$

for $\varepsilon \leq \varepsilon_0$ depending on the norms of $\tilde{\rho}_\varepsilon, \tilde{Q}_j$, and so $\|w\|_{H^1} = 0$. □

Acknowledgments. The authors wish to thank Sylvia Serfaty and Rafe Mazzeo for helpful conversations throughout the development of this work and wish to thank the anonymous referees for their comments that helped to improve the paper.

REFERENCES

- [1] R. ALICANDRO AND M. PONSIGLIONE, *Ginzburg-Landau functionals and renormalized energy: A revised Γ -convergence approach*, J. Funct. Anal., 266 (2014), pp. 4890–4907.
- [2] F. ARECCHI, *Space-time complexity in nonlinear optics*, Phys. D, 51 (1991), pp. 450–464.
- [3] F. ARECCHI, G. GIACOMELLI, P. RAMAZZA, AND S. RESIDORI, *Vortices and defect statistics in two-dimensional optical chaos*, Phys. Rev. Lett., 67 (1991), 3749.
- [4] F. BETHUEL, H. BREZIS, AND F. HÉLEIN, *Ginzburg-Landau Vortices*, Progr. Nonlinear Differential Equations Appl. 13. Birkhäuser Boston, Boston, MA, 1994.
- [5] F. BETHUEL, R. L. JERRARD, AND D. SMETS, *On the NLS dynamics for infinite energy vortex configurations on the plane*, Rev. Mat. Iberoam., 24 (2008), pp. 671–702. ‘
- [6] H. BREZIS, J.-M. CORON, AND E. H. LIEB, *Harmonic maps with defects*, Comm. Math. Phys., 107 (1986), pp. 649–705.
- [7] H. BRÉZIS AND T. GALLOUET, *Nonlinear Schrödinger evolution equations*, Nonlinear Anal., 4 (1980), pp. 677–681.
- [8] H. CHRISTIANSON, J. MARZUOLA, J. METCALFE, AND M. TAYLOR, *Nonlinear bound states on weakly homogeneous spaces*, Commun. Partial Differential Equations, 39 (2014), pp. 34–97.
- [9] J. E. COLLIANDER AND R. L. JERRARD, *Vortex dynamics for the Ginzburg-Landau-Schrödinger equation*, Internat. Math. Res. Notices, 1998 (1998), pp. 333–358.
- [10] M. DOS SANTOS, *The Ginzburg-Landau functional with a discontinuous and rapidly oscillating pinning term. Part II: The non-zero degree case*, Indiana Univ. Math. J., 62 (2013), pp. 551–641.
- [11] A. L. FETTER, *Vortices in an imperfect Bose gas. IV. Translational velocity*, Phys. Rev., 151 Nov (1966), pp. 100–104.
- [12] A. L. FETTER AND A. A. SVIDZINSKY, *Vortices in a trapped dilute Bose-Einstein condensate*, J. Phys. Condensed Matter, 13 (2001), R135.
- [13] G. B. FOLLAND, *Introduction to Partial Differential Equations*, Princeton University Press, Princeton, NJ, 1995.
- [14] D. FREILICH, D. BIANCHI, A. KAUFMAN, T. LANGIN, AND D. HALL, *Real-time dynamics of single vortex lines and vortex dipoles in a Bose-Einstein condensate*, Science, 329 (2010), pp. 1182–1185.
- [15] L. GIL, K. EMILSSON, AND G.-L. OPPO, *Dynamics of spiral waves in a spatially inhomogeneous Hopf bifurcation*, Phys. Rev. A, 45 (1992), R567.
- [16] R. JERRARD AND D. SMETS, *Vortex dynamics for the two dimensional non homogeneous Gross-Pitaevskii equation*, Ann. Sc. Norm. Super. Pisa, 14 (2002), pp. 729–766.
- [17] R. JERRARD AND D. SPIRN, *Refined Jacobian estimates for Ginzburg-Landau functionals*, Indiana Univ. Math. J., 56 (2007), pp. 135–186.
- [18] R. L. JERRARD AND H. M. SONER, *Dynamics of Ginzburg-Landau vortices*, Arch. Ration. Mech. Anal., 142 (1998), pp. 99–125.
- [19] R. L. JERRARD AND H. M. SONER, *The Jacobian and the Ginzburg-Landau Energy*, Calc. Var. Partial Differential Equations, 14 (2002), pp. 151–191.
- [20] R. L. JERRARD AND D. SPIRN, *Refined Jacobian estimates and Gross-Pitaevsky vortex dynamics*, Arch. Ration. Mech. Anal., 190 (2008), pp. 425–475.
- [21] R. L. JERRARD AND D. SPIRN, *Hydrodynamic limit of the Gross-Pitaevskii equation*, Commun. Partial Differential Equations, 40 (2015), pp. 135–190.
- [22] H.-Y. JIAN AND B.-H. SONG, *Vortex dynamics of Ginzburg-Landau equations in inhomogeneous superconductors*, J. Differential Equations, 170 (2001), pp. 123–141.
- [23] M. KURZKE, C. MELCHER, R. MOSER, AND D. SPIRN, *Dynamics for Ginzburg-Landau vortices under a mixed flow*, Indiana Univ. Math. J., 58 (2009) pp. 2597–2621.
- [24] M. KURZKE AND D. SPIRN, *Quantitative equipartition of the Ginzburg-Landau energy with applications*, Indiana Univ. Math. J., 59 (2010), pp. 2077–2092.
- [25] M. KURZKE AND D. SPIRN, *Vortex liquids and the Ginzburg-Landau equation*, Forum of Mathematics, Sigma, 2:e11, 2014.
- [26] L. LASSOUÉ AND P. MIRONESCU, *Ginzburg-Landau type energy with discontinuous constraint*, J. Anal. Math., 77 (1999), pp. 1–26.
- [27] F. H. LIN, *Some dynamical properties of Ginzburg-Landau vortices*, Comm. Pure Appl. Math., 49 (1996), pp. 323–359.

- [28] F.-H. LIN AND J. X. XIN, *On the incompressible fluid limit and the vortex motion law of the nonlinear Schrödinger equation*, Comm. Math. Phys., 200 (1999), pp. 249–274.
- [29] S. MIDDELKAMP, P. G. KEVREKIDIS, D. J. FRANTZESKAKIS, R. CARRETERO-GONZÁLEZ, AND P. SCHMELCHER, *Bifurcations, stability, and dynamics of multiple matter-wave vortex states*, Phys. Rev. A, 82 (2010), 013646.
- [30] S. MIDDELKAMP, P. J. TORRES, P. G. KEVREKIDIS, D. J. FRANTZESKAKIS, R. CARRETERO-GONZÁLEZ, P. SCHMELCHER, D. V. FREILICH, AND D. S. HALL, *Guiding-center dynamics of vortex dipoles in Bose-Einstein condensates*, Phys. Rev. A, 84 (2011), 011605.
- [31] E. MIOT, *Dynamics of vortices for the complex Ginzburg-Landau equation*, Anal. PDE, 2 (2009), pp. 159–186.
- [32] T. NEELY, E. SAMSON, A. BRADLEY, M. DAVIS, AND B. ANDERSON, *Observation of vortex dipoles in an oblate Bose-Einstein condensate*, Phys. Rev. Lett., 104 (2010), 160401.
- [33] L. NIRENBERG, *Remarks on strongly elliptic partial differential equations*, Comm. Pure Applied Math., 8 (1955), pp. 648–674.
- [34] B. Y. RUBINSTEIN AND L. M. PISMEN, *Vortex motion in the spatially inhomogeneous conservative Ginzburg-Landau model*, Phys. D, 78 (1994), pp. 1–10.
- [35] E. SANDIER AND S. SERFATY, *Gamma-convergence of gradient flows with applications to Ginzburg-Landau*, Comm. Pure Appl. Math., 57 (2004), pp. 1627–1672.
- [36] E. SANDIER AND S. SERFATY, *A product-estimate for Ginzburg-Landau and corollaries*, J. Funct. Anal., 211 (2004), pp. 219–244.
- [37] E. SANDIER AND S. SERFATY, *Vortices in the Magnetic Ginzburg-Landau Model*, Progr. Nonlinear Differential Equations Appl. 70. Birkhäuser Boston, Boston, MA, 2007.
- [38] K. SCHWARZ, *Three-dimensional vortex dynamics in superfluid He 4: Line-line and line-boundary interactions*, Phys. Rev. B, 31 (1985), 5782.
- [39] S. SERFATY AND I. TICE, *Ginzburg-Landau vortex dynamics with pinning and strong applied currents*, Arch. Ration. Mech. Anal., 201 (2011), pp. 413–464.
- [40] D. SPIRN, *Vortex motion law for the Schrödinger-Ginzburg-Landau equations*, SIAM J. Math. Anal., 34 (2003), pp. 1435–1476.
- [41] J. STOCKHOFE, P. G. KEVREKIDIS, AND P. SCHMELCHER, *Existence, Stability and Nonlinear Dynamics of Vortices and Vortex Clusters in Anisotropic Bose-Einstein Condensates*, arXiv:1203.4762, 2013.
- [42] J. STOCKHOFE, S. MIDDELKAMP, P. G. KEVREKIDIS, AND P. SCHMELCHER, *Impact of anisotropy on vortex clusters and their dynamics*, Europhys. Lett., 93 (2011), 20008.
- [43] M. TAYLOR, *Partial Differential Equations I: Basic Theory*, Appl. Math. Sci. 115, Springer, New York, 2013.
- [44] P. TORRES, P. KEVREKIDIS, D. FRANTZESKAKIS, R. CARRETERO-GONZÁLEZ, P. SCHMELCHER, AND D. HALL, *Dynamics of vortex dipoles in confined Bose-Einstein condensates*, Phys. Lett. A, 375 (2011), pp. 3044–3050.
- [45] M. TSUBOTA AND S. MAEKAWA, *Pinning and depinning of two quantized vortices in superfluid He 4*, Phys. Rev. B, 47 (1993), 12040.
- [46] E. YARMCHUK AND R. PACKARD, *Photographic studies of quantized vortex lines*, J. Low Temperature Phys., 46 (1982), pp. 479–515.