

## Article

# Effect of high and low frequency exercise therapy in patients after coronary artery bypass graft surgery

Periyasami, Muruganandam, Rao, Pasupuleti Visweswara and Soon, Jan Mei

Available at <http://clock.uclan.ac.uk/17213/>

*Periyasami, Muruganandam, Rao, Pasupuleti Visweswara and Soon, Jan Mei (2017) Effect of high and low frequency exercise therapy in patients after coronary artery bypass graft surgery. The Lancet, 389 (S79). ISSN 0140-6736*

It is advisable to refer to the publisher's version if you intend to cite from the work.  
[http://dx.doi.org/10.1016/S0140-6736\(17\)30475-0](http://dx.doi.org/10.1016/S0140-6736(17)30475-0)

For more information about UCLan's research in this area go to <http://www.uclan.ac.uk/researchgroups/> and search for <name of research Group>.

For information about Research generally at UCLan please go to <http://www.uclan.ac.uk/research/>

All outputs in CLoK are protected by Intellectual Property Rights law, including Copyright law. Copyright, IPR and Moral Rights for the works on this site are retained by the individual authors and/or other copyright owners. Terms and conditions for use of this material are defined in the <http://clock.uclan.ac.uk/policies/>

## **Effect of high and low frequency exercise therapy in patients after coronary artery bypass graft surgery**

Muruganandam Periyasami, Pasupuleti Visweswara Rao, Jan Mei Soon

Lovely Professional University, Phagwara, India (M Periyasami MSc); Universiti Malaysia Kelantan, Kelantan, Malaysia (P V Rao PhD); and University of Central Lancashire, Preston, UK (J M Soon PhD)

### **Correspondence to:**

Dr Jan Mei Soon, School of Sport and Wellbeing, University of Central Lancashire, Preston, Lancashire, PR1 2HE, UK

[jmsoon@uclan.ac.uk](mailto:jmsoon@uclan.ac.uk)

### **Abstract**

**Background** Coronary artery bypass graft (CABG) surgery is one of the major surgeries requiring long-term stay in hospital. This generally leads to the detrimental effects of bed-rest, including dependency in self-care, transfer, and locomotion. Our aim was to compare the effect of high-frequency and low-frequency exercise therapy in patients who had undergone CABG.

**Methods** Patients who had undergone CABG were recruited from PSG Medical College and Hospital, Coimbatore, India, between Jan 1 and March 31, 2006. Functional Independence Measure (FIM) and modified Borg Rating of Perceived Exertion (RPE) were used to assess functional outcome. In a quasi-experimental design, patients received either high-frequency exercise therapy (exercise three times a day for 10 days, group 1), or low-frequency exercise therapy (once a day for 10 days, group 2). Data were analysed with paired *t* tests.

**Findings** 30 patients were recruited (15 in each group). Mean FIM was 75 (SD 1.77) in group 1 and 64 (1.65) in group 2. There was a significant difference between the pretest and post-test FIM values in group 1 patients (49.07 [2.43] vs 124.07 [1.75],  $p < 0.0001$ ) but not in group 2 patients. The RPE in group 1 and group 2 was 6.3 (0.62) and 4.2 (0.7), respectively.

**Interpretation** Patients given high-frequency exercise therapy had a significant improvement in their physical activity, but low-frequency exercise did not lead to significantly improved changes. In conclusion, the high-frequency exercise therapy improves the functional ability of patients with CABG.

**Funding** None.

### **Contributors**

MP designed the study, and collected and analysed the data. PVR contributed to the planning, design, and analysis of the study, and to the writing of the abstract. JMS analysed data and wrote the abstract.

**Declaration of interests**

We declare no competing interests.