

Ovarian follicular fluid reflects the clinical condition
and oocyte cumulus homeostasis

Tawiwani Pantasri

1206585

Discipline of Obstetrics and Gynaecology,
The School of Paediatrics and Reproductive Health,
Robinson Institute
The University of Adelaide

Supervisors:

Professor Robert Norman

Doctor Rebecca Robker

A thesis submitted to the University of Adelaide
for the degree of Master in Medical Science November 2012

Table of Contents

Declaration.....	i
Acknowledgements	ii
Abstract.....	vi
List of Figures.....	ix
List of Tables	x
Chapter 1 Literature review	1
1.1 The effects of being overweight and obese on reproduction.....	2
1.1.1 Overweight and obesity.....	2
1.1.2 Glucose and lipid in overweight-obese patients.....	10
1.2. The ovarian follicle.....	15
1.2.1 The oocyte	15
1.2.2 Follicular glucose and lipid.....	26
1.2.3 Follicular cell stress and apoptosis	30
1.3 Hypothesis and objectives.....	35
1.3.1 Hypothesis:.....	35
1.3.2 Objectives:.....	35
Chapter 2 General methods and demographic data	36
2.1 Methods	36

2.1.2 Patients	36
2.1.3 Blood samples	37
2.1.4 The follicular fluid	42
2.1.5 The cumulus cells	43
2.1.6 The unfertilised oocyte	43
2.1.6 Terms and definition	43
2.1.7 Sample size calculation	45
2.1.8 Statistical analysis.....	46
2.2 Demographic data	46
Chapter3 Effect of BMI, body fat distribution and metabolic disturbance on follicular metabolism	48
3.1 Introduction.....	48
3.2 Methods	49
3.3 Results	49
3.4 Summary of the results	57
3.5 Discussion	57
Chapter 4 Relationship between serum and follicular glucose and lipids.....	61
4.1 Introduction.....	61
4.2 Assay methods	62
4.3 Results	62

4.4 Summary of the results	76
4.5 Discussion	77
Chapter 5 Oocyte lipid, cumulus cell ER stress and embryo and pregnancy outcome	82
5.1 Introduction.....	82
5.2 Methods	83
5.3 Results	88
5.4 Summary of the results	105
5.5 Discussion	106
Chapter 6 Overall discussion and conclusion	111
6.1 Introduction.....	111
6.2 Effect of systemic metabolic disturbance on follicular metabolism.....	111
6.3 Relationship between serum and follicular glucose and lipids	113
6.4 Endoplasmic stress marker and the oocyte study.....	115
6.5 Study limitation and future research	117
6.6 Conclusion	117
References	119
Appendix1	134
Information and consent forms	134
Appendix 2.....	140
The serum and follicular fatty acid-BMI relationship.....	140

Appendix 3.....	148
The number of women who had undetectable levels of particular fatty acids among overall 84 women	148
.....	148

Declaration

This work contains no material which has been accepted for the award of any other degree or diploma in any university or other tertiary institution and, to the best of my knowledge and belief, contains no material previously published or written by another person, except where due reference has been made in the text.

I give consent to this copy of my thesis, when deposited in the University Library, being made available for loan and photocopying, subject to the provisions of the Copyright Act 1968.

I also give permission for the digital version of my thesis to be made available on the web, via the University's digital research repository, the Library catalogue, the Australasian Digital Theses Program (ADTP) and also through web search engines, unless permission has been granted by the university to restrict access for a period of time.

Signature:

Date:

Acknowledgements

I would like firstly to acknowledge Chiang Mai University for the staff development scholarship for my financial support. Acknowledgement also to my supervisors, Professor Robert Norman and Dr. Rebecca Robker who have supported this project.

It is a pleasure to thank the helpful teams of FertilitySA, St Andrews Hospital and Flinders University helping me collecting the samples for this project. This includes, the doctors, nurses, operative staff, embryologists and administrative staff. I also wish to thank all the infertile women participating in this study.

I would like to thank Adelaide Microscopy, Associate Professor David Kennaway and his team, Dr. Linda Wu, Dr. Anne Mcpherson and Mr. David Apps for laboratory assessment. Acknowledgement also go to Mr. Thomas Sullivan for his statistical support.

Special thank for the ovarian biology lab staff for their suggestions in my experiments, presentations and sample transfer. The thanks also go to my PhD candidate friends and the administrative staff of the school of Paediatrics and Reproductive health for their general help and guidance in Medical school.

It is the pleasure to thank the IT service, Dr. Sakrapee Paisitkriangkrai and IBP staff for their wonderful IT support and IBP course.

Finally, I would like to thank my family and friends for their support and encouragement.

Abstract

Infertility is a worldwide problem that is often overlooked. Although assisted reproductive technology has been developed over decades, many women still suffer from infertility. More knowledge is needed to understand ovarian homeostasis to optimise fertility treatment. This study aimed to explore the relationship of lipids and glucose levels in blood and follicular fluid, and compare these substrates among women with normal or abnormal metabolic condition. It also sought to measure lipid content within human oocytes as well as the expression of endoplasmic reticulum stress marker genes in cumulus cells, and their relationship with metabolic substances, obesity and IVF outcome.

The blood, follicular fluid, cumulus cells and unfertilised oocytes from 88 women, who underwent IVF in FertilitySA from February 2011 to August 2011, were collected and analysed for glucose, lipids and endoplasmic reticulum stress markers.

Follicular glucose, insulin, high density lipoprotein cholesterol (HDL-C) and majority of polyunsaturated fatty acid (PUFA) and monounsaturated fatty acid (MUFA) levels correlated with the serum levels ($r= 0.16-0.23$). Insulin was associated with the BMI, waist circumference, metabolic syndrome and many fatty acids, but not follicular glucose. However, the immaturity rate of the retrieved oocytes correlated with the follicular glucose and total fatty

acids ($r = 0.19-0.26$, $p < 0.04$). Variability of the unfertilised oocyte morphology correlated with follicular glucose, and the immaturity rate also differed among the metabolic syndrome group.

An increase of follicular 18:3 n-3 (alpha-linolenic) and decrease of 20:3 n-3 (eicosatrienoic acid; ETA) existed in women with a waist circumference of more than 80 cm. The follicular 20:5 n-3 (eicosapentaenoic acid; EPA) percentage correlated with fertilisation and cleavage rate ($r = -0.32$, $p = 0.003$ and $r = -0.35$, $p = 0.001$). The follicular low density lipoprotein cholesterol (LDL-C) and HDL-C related to follicular fatty acids. Higher levels of serum LDL-C (2.31 ± 0.65 and 1.98 ± 0.61 mmol/L, $p 0.02$) and follicular fatty acid (0.26 ± 0.09 and 0.22 ± 0.05 mmol/L, $p = 0.03$) were found in non-pregnant women.

There was a wide variability of ER stress expression in cumulus cells among women in this study. There was no obvious correlation between ER stress markers and other measurements. The unfertilised oocyte BODIPY fluorescence intensity had high variability among women and individuals. However, the unfertilised oocyte lipid content correlated with the serum LDL-C level.

Substances with a good follicular-serum relationship may be transported directly from blood to the follicle. The disconnection might be affected by intrafollicular metabolism. Insulin may be involved in follicular lipid metabolism because it correlated with many follicular fatty acids and cholesterols. The equilibrium between follicular fatty acids involving insulin modulation may affect oocyte quality.

Overall, this study found there were correlations between serum and follicular lipids, follicular insulin and cholesterol with follicular fatty acids and the importance of serum LDL-C and follicular omega-3 fatty acids. Serum insulin and LDL-C screening might be another tool for predicting the follicular lipid dysequilibrium and poor IVF outcome. The unfertilised oocyte may be a useful tool for a study on oocyte quality. A larger study is needed to recruit more women of different ages and BMI for a stronger correlation between follicular insulin, glucose and lipid metabolism, and ER stress markers.

List of Figures

Figure 3.1 The follicular eicosatrienoic acid correlation with BMI.	52
Figure 4.1 Correlation between serum and follicular substances among 84 women.	63
Figure 4.2 Correlation between serum and follicular fatty acids	65
Figure 5.1 The boxed area determining the average of the fluorescence pixel dot.	84
Figure 5.2 The morphology of oocytes from the same patient	87
Figure 5.3 The distribution of ER stress marker gene expression in cumulus cells.....	89
Figure 5.4 The correlation between ATF4 and GRP78 expression among 45 women.	90
Figure 5.5 Picture of individual BODIPY staining of a good normal oocyte from the same patient:	98
Figure 5.6 Fluorescence intensity of individual mature oocytes from 55 women according to the BMI.	99
Figure 5.7 Mean fluorescence intensity of mature and immature oocytes according to the BMI.	100
Figure 5.8 Mean fluorescence intensity of mature and immature oocytes from women with both.....	101
Figure 5.9 The mean fluorescence intensity of the mature and immature oocytes in each BMI subgroup.	102

List of Tables

Table 1.1 Phenotypes of PCOS patients according to different criteria.....	9
Table 1.2 Abnormal morphology of oocytes related to poor IVF outcome.....	21
Table 1.3 Summary of data about substances and cell stress markers in serum and follicular fluid of PCOS and obese patients.....	33
Table 2.1 Fatty acid equivalent names in the study	41
Table 3.1 Comparison between normal weight , overweight and obese subgroups	50
Table 3.2 Comparison between women with a normal waist circumference (n=18), 80-88 cm (n=20), and the group with more than 88 cm (n=44)	50
Table 3.3 Serum analytes and significant correlation with BMI and waist circumference among 84 women	51
Table 3.4 Significant correlation of follicular substances with BMI and waist circumference among 84 women	51
Table 3.5 Significant comparison between women with a HOMA score < 3.8 (n=63) and those who had more than 3.8 (n=21).....	54
Table 3.6 Significant differences of follicular substances between women with normal serum HDL-C level (n=25) and those with less than 1.29 mmol/L (n=59)	55
Table 3.7 Significant difference of fatty acid percentage between women with normal serum LDL-C level (n=67) and those with more than 2.56 mmol/L (n=17)	55
Table 3.8 Significant comparison between women with none of the NCEP metabolic syndrome criteria (n=29), 1 criterion (n=25), 2 criteria (n=20) and 3-4 criteria (n=10) group.....	56
Table 3.9 Significant difference of substances between women with none or some of the IDF metabolic syndrome criteria (n=15), 1 IDF criterion (n=35), 2 criteria (n=22) and 3-4 criteria (n=12) group.....	56
Table 4.1 Mean concentrations of metabolic substances in serum and follicular fluid of 84 women	62

Table 4.2 Mean percentage of each total fatty acid class in serum and follicular fluid of 84 women	64
Table 4.3 Mean percentage of fatty acid in serum and follicular fluid	67
Table 4.4 Correlation between serum and follicular fatty acids	69
Table 4.5 Significant correlation of follicular glucose with other substances.....	70
Table 4.6 Significant correlation of follicular insulin with other substances	71
Table 4.7 Significant correlation of follicular insulin with follicular fatty acids.....	71
Table 4.8 Significant correlation of follicular triglyceride with other substances	71
Table 4.9 Significant correlation of follicular LDL-C with other substances	72
Table 4.10 Significant correlation of follicular LDL-C with other substances.....	72
Table 4.11 Significant differences between women with and without detectable follicular LDL-C	74
Table 4.12 Significant correlation of follicular HDL-C with other substances	75
Table 4.13 Significant correlation of follicular HDL-C with fatty acids.....	75
Table 4.15 Significant correlation of serum LDL-C with other substances	76
Table 5.1 Significant comparisons between women with less and more than the 80th percentile of GRP78 expression	91
Table 5.2 Significant comparisons between women with less and more than the 80th percentile of ATF6 expression	92
Table 5.3 ER stress markers in PCOS and non-PCOS women	93
Table 5.4 Significant correlations of the degree of morphologic variability in unfertilised oocytes with other measurements	94
Table 5.5 Significant comparisons between women with all good unfertilised oocytes and those with at least one abnormally morphologic oocyte.....	95
Table 5.6 Significant comparisons between women with good polar body in every one of their unfertilised oocytes and those who had abnormal polar body in some of their oocytes.....	96

Table 5.7 Comparison between women with all good unfertilised oocytes and those who had some abnormally morphologic unfertilised oocytes104

Table 5.8 Comparison between biochemically pregnant women and women who were non-pregnant from the fresh cycle104