

ACCEPTED VERSION

Hanru Wang, Mark S Geier and Gordon S Howarth

Prebiotics: a potential treatment strategy for the chemotherapy-damaged gut?

Critical Reviews in Food Science and Nutrition, 2016; 56(6):946-956

Copyright © Taylor and Francis Group, LLC

"This is an Accepted Manuscript of an article published by Taylor & Francis in Critical Reviews in Food Science and Nutrition on 27 Aug 2014 available online:

<http://dx.doi.org/10.1080/10408398.2012.741082>

PERMISSIONS

<http://authorservices.taylorandfrancis.com/sharing-your-work/>

Accepted Manuscript (AM)

As a Taylor & Francis author, you can post your Accepted Manuscript (AM) on your personal website at any point after publication of your article (this includes posting to Facebook, Google groups, and LinkedIn, and linking from Twitter). To encourage citation of your work we recommend that you insert a link from your posted AM to the published article on [Taylor & Francis Online](#) with the following text:

"This is an Accepted Manuscript of an article published by Taylor & Francis in [JOURNAL TITLE] on [date of publication], available online: [http://www.tandfonline.com/\[Article DOI\]](http://www.tandfonline.com/[Article DOI])."

For example: *"This is an Accepted Manuscript of an article published by Taylor & Francis Group in Africa Review on 17/04/2014, available online: <http://www.tandfonline.com/10.1080/12345678.1234.123456>."*

N.B. Using a real DOI will form a link to the Version of Record on [Taylor & Francis Online](#).

The AM is defined by the [National Information Standards Organization](#) as:
"The version of a journal article that has been accepted for publication in a journal."

This means the version that has been through peer review and been accepted by a journal editor. When you receive the acceptance email from the Editorial Office we recommend that you retain this article for future posting.

Embargoes apply if you are posting the AM to an institutional or subject repository, or to academic social networks such as Mendeley, ResearchGate, or Academia.edu.

Embargo

[Critical Reviews in Food Science and Nutrition](#) 12 months

1 June 2017

<http://hdl.handle.net/2440/96358>

Prebiotics: a potential treatment strategy for the chemotherapy-damaged gut?

Hanru Wang¹, Mark S Geier^{1,2} and Gordon S Howarth^{1,3}

¹School of Animal and Veterinary Sciences, University of Adelaide, Roseworthy Campus, South Australia

²South Australian Research and Development Institute, Pig and Poultry Production Institute, Nutrition Research Laboratory, Roseworthy, South Australia

³Centre for Paediatric and Adolescent Gastroenterology, Children, Youth and Women's Health Service, North Adelaide, South Australia

Corresponding author:

Professor Gordon S. Howarth, School of Animal and Veterinary Sciences, University of Adelaide, Roseworthy Campus, South Australia

Tel: +61 8 8313 7885

Fax: +61 8 8313 7972

Email: gordon.howarth@adelaide.edu.au

Short title: Prebiotics and Mucositis

Key words: Animal models, Mucositis, Prebiotics, Chemotherapy

Abstract

Mucositis, characterized by ulcerative lesions along the alimentary tract, is a common consequence of many chemotherapy regimens. Chemotherapy negatively disrupts the intestinal microbiota, resulting in increased numbers of potentially pathogenic bacteria, such as Clostridia and Enterobacteriaceae, and decreased numbers of “beneficial” bacteria, such as Lactobacilli and Bifidobacteria. Agents capable of restoring homeostasis in the bowel microbiota could therefore be applicable to mucositis. Prebiotics are indigestible compounds, commonly oligosaccharides, which seek to reverse chemotherapy-induced intestinal dysbiosis through selective colonization of the intestinal microbiota by probiotic bacteria. In addition, evidence is emerging that certain prebiotics contribute to nutrient digestibility and absorption, modulate intestinal barrier function through effects on mucin expression, and also modify mucosal immune responses, possibly via inflammasome-mediated processes. This review examines the known mechanisms of prebiotic action and explores their potential to reduce the severity of chemotherapy-induced mucositis in the intestine.

MUCOSITIS

The treatment of cancer by chemotherapy, sometimes in combination with radiotherapy, is associated with an array of acute side-effects, including alopecia, neutropenia and nausea (Logan et al., 2007). Furthermore, these treatments can also induce inflammation and deterioration of the mucosal membranes of the alimentary tract, a condition referred to as mucositis. Mucositis can occur throughout the entire gastrointestinal tract (Gibson et al., 2005); however, it most commonly affects the mucosa of the mouth (oral mucositis) and small intestine (gastrointestinal mucositis), with symptoms including pain, vomiting, bloating, and diarrhoea. Severe cases are often associated with secondary complications such as malnutrition and septicemia, resulting in poor patient outcomes. Indeed, the pathobiology of mucositis is influenced by the nature of the causative agent (Logan et al., 2009), with temporal variations in pathogenesis, in addition to the location and severity of damage (Logan et al., 2009). Consequently, the multi-factorial nature of mucositis makes it a difficult condition to understand, and subsequently treat.

Despite mucositis being one of the most common side-effects of cancer treatment (Choi et al., 2007) its incidence and severity is highly variable, and is dependent on the treatment regimen and individual variation in terms of tolerability (Lalla et al., 2006). Approximately 40% of all patients undergoing chemotherapy develop mucositis (Gibson et al., 2002; Lalla et al., 2006) with this number rising to between 60% and 100% in patients receiving high-dose chemotherapy regimens for conditions such as the haematological malignancies (Gibson et al., 2002; Lalla et al., 2006).

To date, effective treatments for intestinal mucositis have been elusive. Potential treatment options are limited by the requirement for treatments to protect the mucosa and promote the repair process, without compromising cytotoxic effects on the neoplasm. Certain growth factors such as epidermal growth factor, transforming growth factor- α and betacellulin (Choi et al., 2008; Dahlhoff et al., 2008; Sukhotnik et al., 2008a; Sukhotnik et al., 2008b; Boukhettala et al., 2010; Kim et al., 2010) have the potential to promote differentiation and proliferation of epithelial cells, and ‘prime’ the intestine prior to chemotherapy, in addition to accelerating the repair process. Previously, keratinocyte growth factor-1 (KGF-1) has shown efficacy in the oral form of the condition (Blijlevens and Sonis, 2007), and decreased diarrhoea in a rat model of irinotecan-induced mucositis (Gibson et al., 2005), however, it is yet to be applied extensively in the treatment of intestinal mucositis in humans. Other growth factors, such as whey growth factor extract (WGFE) (Howarth et al., 1996; Clarke et al., 2002; Tran et al., 2003), insulin-like growth factor-I (IGF-I) (Howarth et al., 1998; Howarth, 2003; Cool et al., 2005) and fibroblast growth factor-20 (FGF-20) (Gibson et al., 2007) have also been investigated, demonstrating variable efficacy. Although in their infancy, nutraceutical agents have recently been tested in experimental models for efficacy against mucositis. These include plant extracts such as grape seed extract (Cheah et al., 2009) and Iberogast (Wright et al., 2009), and animal-sourced oils such as Lyprinol (Torres et al., 2008) and Emu Oil (Lindsay et al., 2010).

Development of Mucositis

Mechanisms underlying the development of mucositis are complex, and still not fully understood; however, it is hypothesized that the condition progresses through five stages. The

initiation of mucositis occurs directly after chemotherapy administration and is characterised by the production of reactive oxygen species (ROS) (Sonis et al., 2004; Lalla and Peterson, 2006; Logan et al., 2007). These then activate nuclear factor- κ B, (NF- κ B) (Logan et al., 2009), a transcription factor implicated in promotion of the inflammatory response in conditions including arthritis and the inflammatory bowel diseases (Mankan et al., 2009). NF- κ B is thought to be directly responsible for the up-regulation of pro-inflammatory cytokines such as tumor necrosis factor- α (TNF- α), interleukin-1 β (IL-1 β) and IL-6 (Logan et al., 2009). These cytokines further amplify damage to the mucosal cells (Sonis et al., 2004). Mucositis first becomes clinically evident during the fourth phase, termed 'ulceration and inflammation' (Sonis, 2004a). It is during this stage that the epithelial layer increases in permeability, allowing the translocation of bacteria into the blood stream. This is accompanied by severe pain and discomfort, and sometimes secondary complications such as malnutrition and septicaemia (Logan et al., 2007). Further changes occur during the ulcerative stage, including crypt ablation and villus atrophy, leading to a decrease in the overall absorptive area of the intestine, a decrease in digestive enzymes, and an increase in intestinal permeability, with subsequent increases in bacterial translocation and alterations in the gut microbiota (Stringer et al., 2009b). Following the discontinuation of chemotherapy or radiotherapy, symptoms resolve and mucosal repair begins, characterised by proliferation and differentiation of tissues and cells (Sonis et al., 2004; Lalla and Peterson, 2006). Currently, our understanding of the mechanisms underlying these changes is incomplete, and a role for apoptosis as an early event has been proposed (Sonis, 2004b). Through a better understanding of the role of the microbiota in the progression and resolution of intestinal

mucositis, and the mechanisms by which these interactions can be influenced, it is possible that the local microbiota could be altered in such a way as to lessen the severity of the condition.

The Bowel Microbiota and Development of Intestinal Mucositis

The role of the gut microbiota in the development of intestinal mucositis awaits clear definition; although evidence is emerging that the symbiotic relationship between the microbiota and its host is negatively disrupted by chemotherapy (Stringer et al., 2009a; Stringer et al., 2009b). This may have important implications for both the severity of mucositis, and the overall health of the host. The microbiota may influence the development and severity of mucositis through at least five different mechanisms. These include influencing the inflammatory response, intestinal permeability, the composition of the mucin layer, epithelial repair, resistance to harmful stimuli, and the gut immune system (van Vliet et al., 2010). In mucositis, there is a shift in composition of the microbiota away from commensals and ‘beneficial’ bacterial species, such as *Bifidobacteria* spp. and *Lactobacillus* spp., towards more ‘pathogenic’ microbes such as Clostridia and Enterobacteriaceae (Stringer et al., 2009a). Recent studies have also demonstrated decreases in goblet cell numbers and mucin secretion associated with chemotherapy administration (Stringer et al., 2009b). Maintaining a healthy microbiota and mucin layer during chemotherapy treatment could minimise the likelihood of bacterial translocation and secondary infection arising from mucositis. Probiotic microorganisms hold potential in this context. Probiotics are defined as living microorganisms which when administered in adequate amounts, exert desirable health benefits on the host (Reid, 2005). A recent review by Prisciandaro et al. (2011) reported that certain probiotics such as *Lactobacillus*, *Bifidobacterium* genera,

Enterococcus and Streptococcus species may reduce the severity of chemotherapy-induced mucositis. Modalities capable of stimulating colonisation of the bowel by probiotic bacteria more globally could be a potential strategy to restore intestinal homeostasis. Prebiotics could represent such a strategy.

PREBIOTICS

The concept of prebiotics, and targeted modulation of the gut microbiota was introduced in 1994-1995 (Delzenne and Roberfroid, 1994; Gibson and Roberfroid, 1995), with the definition of prebiotics as compounds that ‘selectively stimulate the growth and/or activity of microbial species in the gut microbiota and confer health benefits to the host’ (Roberfroid et al., 2010). In order to be classified as a prebiotic, a given substance must be neither hydrolysed nor absorbed in the proximal gastrointestinal tract, must be selectively fermented by one or a limited number of beneficial bacteria in the intestine and be able to alter the colonic microbiota toward a healthier composition (Gibson and Roberfroid, 1995; Looijer-van Langen and Dieleman, 2009; Roberfroid, 2007). The fermentation of prebiotics in the colon produces short-chain fatty acids (SCFA), which support proliferation of the colonic microbiota (Yanahira et al., 1995), Oligosaccharides (OS) tend to predominate as compounds that meet the definition of a prebiotic (Roberfroid, 2007).

A wide variety of compounds have the potential to meet the requirements for definition as a prebiotic, although at this stage, not all have been studied sufficiently to classify them as such. The most intensively researched prebiotic compounds to date include inulin (Abrams et al.,

2005), a plant derived fructan, lactulose (Bovee-Oudenhoven et al., 2003), galacto-oligosaccharide (GOS) (Shoaf et al., 2006; Alliet et al., 2007), derived from milk and fructo-oligosaccharide (FOS) (Cherbut et al., 2003), also having plant origins. Other compounds studied for their prebiotic potential include mannan-oligosaccharides (MOS) (Yang et al., 2008b), β -1,4-mannobiose (MNB) (Ibuki et al., 2010), sialyl-oligosaccharides (SOS) (Sinclair et al., 2008), xylo-oligosaccharides (XOS) (Xu et al., 2009), germinated barley foodstuff (GBF) (Kanauchi et al., 2008), epilactose (Nishimukai et al., 2008; Watanabe et al., 2008), arabinoxylan-oligosaccharides (AXOS) (Maki et al., 2012) and chito-oligosaccharides (COS) (Liu et al., 2008).

Different prebiotics may exert an array of biological effects, including alteration of gut morphology (Leforestier et al., 2009), stimulation of mucosal and systemic immune responses (Eiwegger et al., 2010; Gopalakrishnan et al., 2012), alteration of local gut microbial communities, such as by enhancing the proliferation of probiotics (*Lactobacillus plantarum* L12 and *Bifidobacterium pseudocatenulatum* B7003) *in vitro* (Marotti et al., 2012), modulation of inflammation (Pilzner et al., 2012), changes in mucin production (Leforestier et al., 2009) and a decreased occurrence of infection in the first six months of life (Arslanoglu et al., 2007). Prebiotics have also been reported to alter the uptake of various nutrients and minerals, such as calcium, from the intestine (Abrams et al., 2005; Bakker-Zierikzee et al., 2005). Considered together, certain prebiotics could have the potential to combat specific aspects of mucositis pathogenesis.

Prebiotics and Intestinal Dysfunction

One of the earliest studies to investigate the influence of indigestible carbohydrates on gut dysfunction examined the effect of FOS ingestion in immunoglobulin A (IgA) deficient dogs with small bowel bacterial overgrowth (Willard et al., 1994). This study concluded that FOS ingestion altered the microbial communities in the small intestine towards fewer aerobic/facultative anaerobic species, although the exact mechanism and implications of these findings were not determined. The following year, Gibson and Roberfroid (1995) introduced the concept of a ‘prebiotic’ as it is defined today, and in later studies, introduced two other major postulates. Firstly, the concept of targeted modulation of the intestinal microbiota (Gibson, 1998) and secondly, the use of prebiotics and probiotics concurrently, which has come to be known as ‘synbiotics’ (Roberfroid, 1998).

This early work, and the establishment of set criteria for a ‘prebiotic’ resulted in a number of studies examining the effect of oligosaccharides on the microbiota (Yanahira et al., 1995) and digestibility studies on substances suspected of containing indigestible oligosaccharides (Brand-Miller et al., 1998). Recently, Bogusławska-Tryk et al. (2012) reported that inulin and FOS type fructans increased probiotic bacteria, while decreasing certain pathogens. Additionally, these investigators reported that inulin and FOS increased the length of the small and large intestines in chickens and this contributed to an increase in intestinal enzymatic activity, which in turn led to an improvement in nutrient digestibility and absorption (Bogusławska-Tryk et al., 2012).

De Souza Oliveira et al. (2012) suggested that one of the beneficial effects of the prebiotic inulin, during the fermentation process of skim milk by *Streptococcus thermophilus* and *Lactobacillus rhamnosus*, was due to the release of fructose, which could act as a carbon and energy source for the host. In the setting of clinical cancer treatment, this carbon and energy source could potentially be used to enhance the energy levels available to chemotherapy patients struggling with nutrient malabsorption due to mucositis.

These studies were followed by investigations into disease states characterised by an imbalance in the commensal microbiota, particularly in the colon. Certain prebiotics have revealed therapeutic promise in the treatment of inflammatory bowel disease (Kumar et al., 2012) and researchers are now investigating prebiotics for their potential to treat other gut disorders which include acute infantile diarrhoea (Agustina et al., 2007) and irritable bowel syndrome (Fedorak and Madsen, 2004). As mucositis is characterised by a disrupted intestinal microbiota it could be clinically rewarding to test the capacity for prebiotics to normalize the composition of the gut flora.

Prebiotics and Cancer

Some studies have indicated that prebiotics such as FOS may play an important role in cancer prevention (Daddaoua et al., 2007; Sung and Choi, 2008). Clark et al. (2012) recently reviewed the effect of prebiotics on biomarkers of colorectal cancer (CRC) and concluded that certain prebiotics had the potential to prevent colonic cancer. In their review, potential biomarkers of prebiotic efficacy on CRC were identified (Clark et al., 2012). These endpoints included

neoplasia occurrence, cell proliferation, gene expression, and DNA methylation, which were likely more indicative relating to CRC risk (Clark et al., 2012). Furthermore, it has been suggested that secondary markers, such as toxic compounds in the colon or rectum, protein degradation, and bacterial enzyme activity could also be considered in CRC prevention (Clark et al., 2012). Indeed, recurrence rate of adenomatous polyps was decreased after administration of non-digestible lactulose in humans (Roncucci et al., 1993; Clark et al., 2012). These endpoints could be used in the future study of prebiotics on mucositis. It has become evident that prebiotic-type compounds present in milk, such as GOS, play an important role in the development of the neonatal gut, and immune system (Fanaro et al., 2005; Arslanoglu et al., 2007). Modulating cancer-related biomarkers of proliferation, apoptosis and immune modulation via prebiotic administration could be equally applied in mucositis pathogenesis, although we await appropriately controlled studies in this context.

Prebiotics and the Microbiota

The mucosal barrier on the epithelial surface of the intestine consists, in part, of colonized bacterial communities, which may contribute to the defensive gut barrier and guard against other pathogen colonization and subsequently decrease the likelihood of bacterial translocation (Gori et al., 2011). However, chemotherapy administration can induce changes in bacterial communities to a more pathogenic composition (Stringer et al., 2009b). Microbiota composition in the small and large intestine were significantly altered by 5-FU injection in rats. Specifically, levels of *Clostridium* spp., *Lactobacillus* spp. and *Streptococcus* spp. were decreased and

Escherichia spp. increased in the jejunum, whereas, levels of *Enterococcus* spp., *Lactobacillus* spp. and *Streptococcus* spp. were decreased in the colon.

The ability for prebiotics to alter the commensal microbiota towards a beneficial composition has been demonstrated in several studies. For example, a prebiotic oligosaccharide mixture administered to humans improved gut microbiota composition, as measured by an increase in Bifidobacteria and a reduction in the levels of pathogenic Clostridia-related species (Gori et al., 2011). In broiler chickens, MOS and FOS affected the microbiota composition of the small intestine. Yang et al. (2008a) reported that MOS ingestion decreased mucosa-associated *Coliform* spp. in the jejunum in an *Escherichia coli* challenge model in broiler chickens (Yang et al., 2008a), and the same authors also demonstrated that FOS increased *Lactobacillus* spp. levels in the jejunum, and lactic acid levels in the ileum in both the *Escherichia coli* challenged and non-challenged model. Several other studies have shown similar results, with an increase in *Bifidobacteria* spp. and *Lactobacillus* spp. in both rat large intestine and human fecal material following administration of FOS (Lara-Villoslada et al., 2006; Ten Bruggencate et al., 2006). Moreover, when FOS was administered in conjunction with *L. paracasei* to gnotobiotic piglets, the increase in *Bifidobacteria* spp. and *Lactobacillus* spp. was prolonged and enhanced, compared to piglets administered the probiotic alone (Bomba et al., 2002). AXOS has also been reported to alter lactic acid bacteria and *Clostridium* sp. composition in sturgeon fish species (Geraylou et al., 2012). Recently, it has been reported that the administration of a specific prebiotic mixture (short-chain galacto-oligosaccharides and long-chain fructo-oligosaccharides, ratio 9:1) to newborn human babies resulted in higher levels of fecal bifidobacteria and

lactobacilli than the placebo group (Salvini et al., 2011). Together, these studies demonstrated that prebiotics have the potential to consolidate the beneficial bacterial composition of the mucosal barrier in the small and large intestine as well as altering the profile of the microbiota in fecal material.

The altered microbiota may in turn compete with pathogenic bacteria for adhesion sites in the gut. However, evidence is still lacking to confirm the capacity for prebiotics to normalise the composition of the microbiota in specific areas of the small and large intestine in mucositis. In addition, prebiotics are able to decrease gut pH and modulate the SCFA pattern, resulting in an inhospitable environment for pathogens (Eiwegger et al., 2010; Gori et al., 2011). Collectively, the stimulation of favourable strains of bacteria in the gut, and a reduction in gut pH, may also contribute to enhanced immune defences in the mucosal barrier. This could represent a fertile area for future research.

Nevertheless, not all prebiotic studies have resulted in clinical improvements. For example, the prebiotic, MOS, failed to improve growth performance in pigeons, nor improve the immune response against the Newcastle disease virus (Khan et al., 2012). This indicates that our understanding of the potential applications for prebiotics is limited and suggests that a range of factors, particularly the amount and type of prebiotic supplement used, the species and health status of the subject, as well as a range of other factors, all play an important role.

Further evidence is emerging that administration of prebiotics can alter gene expression in the gut through their fermentation products, in particular, butyrate (Sauer et al., 2007; Rodenburg et al., 2008). Butyrate has been identified as a potential anti-neoplastic agent in the colon (Scharlau et al., 2009), has been shown to induce apoptosis in gastric cancer cells (Matthews et al., 2007) and also plays an essential role in the regeneration of the mucosa (Sauer et al., 2007) and in inhibition of pro-inflammatory cytokine production in rat mesenteric lymph nodes (Looijer-van Langen and Dieleman, 2009).

Prebiotics and Mucins

The mucin layer is an integral component of barrier function in the intestine (Stringer et al., 2009b). Mucins form gels and generate a protective mucus blanket overlying the epithelial surface protecting the mucosa from bacterial overgrowth and/or penetration (Specian and Oliver, 1991). Mucins also produce a physical and chemical barrier that protects the epithelium from luminal agents such as enteric bacterial toxins (Specian and Oliver, 1991). In addition, mucins create an enormous repertoire of potential binding sites for microorganisms, such as commensal and pathogenic bacteria (Robbe et al., 2004).

It has been proposed that inhibition of bacterial adhesion occurs as a result of terminal sugars on prebiotics interfering with receptors on the bacteria that facilitate adhesion to the intestinal mucin layer (Forchielli and Walker, 2005). Sinclair et al. (2008) demonstrated that this interference was not unique to bacterial cells, with a study using sialyloligosaccharide shown to inhibit binding of the cholera toxin to its receptor on the cell. FOS and GOS have been linked to

increased mucin production (Ten Bruggencate et al., 2006; Leforestier et al., 2009) resulting in reduced bacterial translocation and consequently, a lower incidence of pathogenic infection (Looijer-van Langen and Dieleman, 2009). In contrast, chemotherapy agents significantly decreased epithelial mucin levels in the jejunum in rats treated with 5-FU (Stringer et al., 2009b) although, mucus secretion in the large intestine was increased in irinotecan treated rats (Gibson et al., 2003). Therefore, it is clear that chemotherapy can induce mucin changes in rats. However, there is still insufficient evidence to definitively predict mucin changes induced by different chemotherapy agents.

Prebiotics, on the other hand, could have the ability to prevent chemotherapy-induced damage to the gut by maintaining mucosal integrity through mucin modification. However, the capacity for prebiotics to alter mucin levels in the different intestinal regions in the mucositis setting requires future research.

Prebiotics and the Gut Immune System

The gut mucosal lining contains organized lymphoid tissues that support the initiation of anti-microbial immune responses (Rancez et al., 2012). Lymphoid cells can be orchestrated by chemokines, trafficking and positioning into different mucosa-associated lymphoid tissues, contributing to immune defence (Rancez et al., 2012). Prebiotics therefore have the potential to modulate lymphoid cells and stimulate the gut immune system (Forchielli and Walker, 2005; Daddaoua et al., 2007; Geraylou et al., 2012). For example, administration of FOS directly stimulated the mucosal immune system in mice and humans, in the form of increased IgA

production (Looijer-van Langen and Dieleman, 2009; Yasuda et al., 2012). IgA is the dominant antibody associated with mucosal immunity, and plays an important role in inhibition of pathogen adhesion to the intestinal wall and the activation of macrophages (Looijer-van Langen and Dieleman, 2009). Recently, Yasuda et al. (2012) reported that FOS decreased allergic peritoneal inflammation in mice through increased local synthesis of IgA. However, prebiotics such as FOS and inulin did not affect IgA levels in dogs (Patra, 2011) or in humans treated with AXOS (Walton et al., 2012).

In addition to the expression of IgA, a mixture of the prebiotics FOS and GOS enhanced delayed type hypersensitivity in an influenza vaccine model as a marker of T helper cell (T_H1) immunity (Vos et al., 2006) and induced an immunoglobulin profile considered beneficial in infants at high risk for allergy (van Hoffen et al., 2009). FOS has also been demonstrated to increase circulating levels of IgM and IgG in plasma, suggesting that the influence of prebiotics is not limited to the gut (Janardhana et al., 2009). Furthermore, GOS has recently shown the ability to reduce the severity of colitis in mice through increasing the activity of natural killer (NK) cells in the spleen and mesenteric lymph nodes (MsLN) (Gopalakrishnan et al., 2012). GOS also stimulated NK expression of the chemokine receptor CCR9, which is involved in lymphocyte trafficking to the gut, pre-infection in the blood, spleen and MsLN (Gopalakrishnan et al., 2012). GOS mixture has also been reported to improve NK cell activity in highly active antiretroviral therapy (HAART)-naïve HIV-infected human individuals (Gori et al., 2011). In addition, prebiotics such as GOS could produce SCFA during fermentation (Lamsal, 2012), which have been reported to induce NK cytotoxic activity in rats (Pratt et al., 1996).

FOS facilitated the proliferation of Lactobacilli, Bifidobacteria, both of which are involved in the induction of regulatory T cells (Treg), which play an important role in modulation of the immune system in C57BL/6J mice in response to Flu-vaccination (van't Land et al., 2010). FOS is also involved in the direct induction of Treg (van't Land et al., 2010). Therefore, FOS could modulate the chemotherapy-affected immune response through activation of suppressive T cell populations including Treg. To this end, bacteria, such as *Bifidobacterium infantis* 35624 induced to proliferate by certain prebiotics, may enhance the activity of indoleamin 2,3-dioxygenase which in turn may contribute to Treg production and T cell tolerance (Konieczna et al., 2012; Yasuda et al., 2012).

In addition, β -1,4-mannobiose has been reported to exert prebiotic activity by stimulating both systemic and innate immune responses in the small intestine in one-day-old chicks (Ibuki et al., 2010). MNB up-regulated the peptide transporter associated with antigen processing (TAP)-2 expression (Ibuki et al., 2010), an essential factor for expression of the major histocompatibility complex (MHC) molecule on cell surfaces (Panter et al., 2012). Moreover, MNB up-regulated TNFSF15 gene expression in healthy chicks, related to the differentiation, proliferation and apoptosis of immune cells (Zhang and Li, 2012). Likewise, MNB enhanced healthy chick immunity through up-regulated expression of BLB-1, BF-2, IRF-1, IRF-7 and TLR3 genes (Ibuki et al., 2010).

Furthermore, it has been shown that AXOS up-regulated the immune response in piglets by influencing the inflammatory and antibacterial protein pancreatitis associated protein (PAP) and down regulated *Escherichia coli* induced responses (Niewold et al., 2012). Interestingly, the higher degree of polymerization of AXOS seems to have a greater beneficial impact on health, as measured by enhanced phagocytic activity of fish macrophages as well as improved alternative haemolytic complement activity and total serum peroxidase content (Geraylou et al., 2012). As chemotherapy-induced mucositis is associated with a compromised immune system, prebiotics may have the potential to beneficially modify the immune response.

Prebiotics and Inflammation

The inflammatory process associated with mucositis is complex, and is characterised by a number of different mechanisms, including the production of pro-inflammatory cytokines, such as TNF- α , IL-6, IL-1 β , nitric oxide (NO) and ROS (Abdelouhab et al., 2012; Liu et al., 2012). Administration of prebiotics such as inulin, FOS and germinated foodstuff have demonstrated efficacy in models of inflammatory intestinal disorders, including dextran sulphate sodium (DSS)-induced colitis, dimethylhydrazine-induced colon cancer, and chemotherapy (5-FU) induced rats mucositis (Araki et al., 2000; Kanauchi et al., 2003; Lara-Villoslada et al., 2006; Winkler et al., 2007; Smith et al., 2008; Abdelouhab et al., 2012).

Initially, the beneficial effects demonstrated by prebiotics were believed to be due solely to their interaction with the commensal intestinal microbiota. However, recent indications suggest that prebiotic administration may have independent effects. A study in obese mice revealed that

oligofructose ingestion stimulated secretion of glucagon-like peptide-2 (Cani et al., 2009), a potent intestinal growth factor, in a rat model of DSS-induced colitis (Brubaker et al., 1997). Sung and Choi (2008) found that cyclooxygenase-2 expression in rats treated with dimethylhydrazine, was decreased following FOS administration, and it was proposed that this could represent a potential mechanism of action for certain prebiotics.

Recently, inulin has been shown to significantly reduce colonic lesions in DSS-induced colitis in rats (Abdelouhab et al., 2012). Moreover, FOS ameliorated allergic peritoneal inflammation induced by ovalbumin (OVA), partially through the suppression of interleukin-5 (IL-5), and chemokine eotaxin in mice (Hosono et al., 2003; Yasuda et al., 2012). The IL-5 gene cluster and eotaxin are concomitant with the infiltration of eosinophils, neutrophils, mast cells and T lymphocytes into the airways of mice contributing to inflammation, particularly in Th2 type chronic inflammatory disease (Pilzner et al., 2012). Thus, it is likely that FOS influences cell populations such as T cells, basophils, mast cells, and epithelial cells (Hosono et al., 2003; Yasuda et al., 2012).

GOS has demonstrated efficacy in inhibiting the adherence of pathogenic bacteria to Caco-2 and HEp-2 cells in culture (Shoaf et al., 2006) and *in vitro* tests also demonstrated that GOS, in the absence of bacteria, stimulated sucrase activity in a Caco-2 cell line (Leforestier et al., 2009). More recently, it has been reported that the prebiotic, inulin has the ability to normalise increased NO levels, in peritoneal macrophage cultures from Swiss mice (Abdelouhab et al., 2012). In addition, prebiotics could mediate the secretion of pro-inflammatory cytokines such as TNF- α

via modification of Toll-like receptor (TLR) gene expression. Trevisi et al. (2008) found that FOS in combination with *Bifidobacterium animalis* increased expression of the TLR-2 encoding gene in the lymph nodes of piglets. The authors further described that TLR-2 and TLR-4 encoding gene expressions were significantly associated with TNF- α encoding gene expression (Trevisi et al., 2008). On the other hand, Fukuda et al. (2011) revealed that germinated barley food stuff (containing prebiotics) decreased TLR-4 expression by increasing butyrate production and decreasing cyclooxygenase 2 mRNA expression in azoxymethane treated F344 rats. Furthermore, the consumption of XOS in combination with inulin decreased blood lipopolysaccharide (LPS) concentrations and attenuated LPS-induced increases in gene expression following IL-1 β and LPS-induced decreases in gene expression of IL-13 in the blood of healthy humans (Lecerf et al., 2012).

LPS can be recognized by TLR-4 with the MD-2, CD14 and LPS-binding protein (LBP). Initiation of this TLR signaling has also been involved in proteins of MyD88, Tirap, Trif, and Tram (Yamamoto et al., 2002; Yamamoto et al., 2003; Shigeoka et al., 2007). Ultimately, this LPS-TLR signaling leads to the induction or suppression of genes that orchestrate the inflammatory response through both MyD88 dependent and independent expression (Yamamoto et al., 2002; Yamamoto et al., 2003; Shigeoka et al., 2007). However, it is still not clear if prebiotics increase the immune response of intestinal epithelial cells by modifying TLR-4 in the MyD88 dependent, independent or other pathways. Future studies should focus on the effects of prebiotics on TLR-4 modification and its signaling pathways compared to LPS, as well as the key proteins involved.

In a recent commentary, Howarth (2012) proposed that prebiotics have the potential to influence inflammasomes and modulate intestinal inflammation and function. Inflammasomes are a group of protein complexes built around several proteins including NLRP3, NLRC4, AIM2 and NLRP6 (Strowig et al., 2012). These inflammasomes can recognize a wide range of microbial, stress and damage signals and following the activation of caspase-1, subsequently induce the secretion of pro-inflammatory cytokines such as IL-1 and IL-18 (Strowig et al., 2012). Inflammasomes can also induce pyroptosis, a form of cell death (Strowig et al., 2012). Furthermore, inflammasome-mediated processes play an important role in microbial infections, regulating metabolic processes, and mucosal immune responses in human diseases (Strowig et al., 2012). Indeed, a recent study by Qu et al. (2012) found that inflammasomes NLRC4 phosphorylation (NLRC4 phospho-Ser 533) successfully inhibited the activation of caspase-1 and pyroptosis in response to *Salmonella typhimurium*. This indicated that NLRC4 phosphorylation could be a decisive point for NLRC4 inflammasome activation and host innate immunity (Qu et al., 2012). Future studies could investigate the potential for prebiotics to modify inflammasomes through different signaling pathways. This would further our understanding of prebiotics and their direct and indirect effects on inflammation in the chemotherapy-damaged intestine (Howarth, 2012).

The influence of prebiotics on inflammation in mucositis requires further investigation. Future studies should not only focus on histological or inflammatory cytokine related responses, but also on molecular responses, such as gene expression and DNA methylation (Clark et al., 2012).

The journey to identify the key genes related to chemotherapy-induced inflammation, and whether prebiotics could control or alter this inflammation at the gene or protein level, promises to be an exciting and worthwhile focus for the future.

Prebiotics and Synbiotics as Potential Treatment Strategies for Mucositis

To date, there have been few rigorous scientific investigations into the interactions of prebiotics in the healthy and diseased small intestine. In particular, the area of intestinal mucositis remains largely unexplored, with only one published study investigating FOS in a rat model of 5-Fluorouracil (5-FU) induced mucositis (Smith et al., 2008). This study demonstrated that FOS was able to marginally reduce myeloperoxidase activity, an indicator of neutrophil infiltration, in the proximal small intestine of chemotherapy-treated rats. However, no other beneficial effects were evident in the other parameters measured. Similarly, there is only a single report of a synbiotic investigation in mucositis, whereby FOS and the new probiotic *Lactobacillus fermentum* BR11, was administered to rats (Smith et al., 2008). Nevertheless, their combination did not seem to exert a further improvement in the severity of mucositis than administered *L. fermentum* BR11 to rats alone (Smith et al., 2008). More recently, the combination of GOS and Bifidobacterium strain (*B. longum* subsp. *longum* PCB133) significantly inhibited pathogenic *Campylobacter jejuni* levels, a pathogen responsible for food-borne gastroenteritis in humans (Baffoni et al., 2012). Considered together, investigations of prebiotics and synbiotics provide a fertile area for further research in disorders of the small bowel, such as intestinal mucositis.

When used in combination, prebiotics and probiotics may complement each other to achieve a synergistic or additive effect in an attempt to ameliorate mucositis (Geraylou et al., 2012).

Probiotics have been studied extensively in relation to gut health (Parnell and Reimer, 2012; Shimizu et al., 2012). However, only recently have prebiotics been examined specifically in relation to chemotherapy-induced mucositis (Smith et al., 2008). Although it remains unknown whether they can be considered as new agents for the treatment of chemotherapy-induced mucositis, prebiotics and synbiotics require investigation in relation to factors such as composition, dose, molecular mechanisms and gene expression, both *in vitro* and *in vivo*, specifically as they relate to the different mechanisms underlying mucositis pathogenesis.

In conclusion, prebiotics are demonstrating efficacy as growth promoters, anti-inflammatory agents and microbial modulators making them promising candidates for the potential treatment of chemotherapy-induced mucositis. The challenge for the future will be the development of predictive *in vitro*, animal model and clinical trial systems to specifically identify the most efficacious probiotics, prebiotics and synbiotic combinations for this debilitating condition.

Acknowledgements

Professor Gordon S. Howarth is supported by a South Australian Health and Medical Research Institute Cancer Council South Australia Senior Research Fellowship. There are no other conflicts of interest or financial agreements in place.

REFERENCES

- Abdelouhab, K., Rafa, H., Toumi, R., Bouaziz, S., Medjeber, O., and Touil-Boukoffa, C. (2012). Mucosal intestinal alteration in experimental colitis correlates with nitric oxide production by peritoneal macrophages: Effect of probiotics and prebiotics. *Immunopharmacol Immunotoxicol.* **34**: 590-597.
- Abrams, S.A., Griffin, I.J., Hawthorne, K.M., Liang, L., Gunn, S.K., Darlington, G., and Ellis, K.J. (2005). A combination of prebiotic short- and long-chain inulin-type fructans enhances calcium absorption and bone mineralization in young adolescents. *Am J Clin Nutr.* **82**: 471-476.
- Agustina, R., Lukito, W., Firmansyah, A., Suhardjo, H.N., Murniati, D., and Bindels, J. (2007). The effect of early nutritional supplementation with a mixture of probiotic, prebiotic, fiber and micronutrients in infants with acute diarrhea in Indonesia. *Asia Pac J Clin Nutr.* **16**: 435-442.
- Alliet, P., Scholtens, P., Raes, M., Hensen, K., Jongen, H., Rummens, J.L., Boehm, G., and Vandenplas, Y. (2007). Effect of prebiotic galacto-oligosaccharide, long-chain fructo-oligosaccharide infant formula on serum cholesterol and triacylglycerol levels. *Nutrition.* **23**: 719-723.
- Araki, Y., Fujiyama, Y., Andoh, A., Koyama, S., Kanauchi, O., and Bamba, T. (2000). The Dietary Combination of Germinated Barley Foodstuff plus *Clostridium butyricum* Suppresses the Dextran Sulfate Sodium-induced Experimental Colitis in Rats. *Scand J Gastroenterol.* **35**: 1060-1067.

- Arslanoglu, S., Moro, G.E., and Boehm, G. (2007). Early supplementation of prebiotic oligosaccharides protects formula-fed infants against infections during the first 6 months of life. *J Nutr.* **137**: 2420-2424.
- Baffoni, L., Gaggìa, F., Di Gioia, D., Santini, C., Mogna, L., and Biavati, B. (2012). A Bifidobacterium-based synbiotic product to reduce the transmission of *C. jejuni* along the poultry food chain. *Int J Food Microbiol.* **157**: 156-161.
- Bakker-Zierikzee, A.M., Alles, M.S., Knol, J., Kok, F.J., Tolboom, J.J., and Bindels, J.G. (2005). Effects of infant formula containing a mixture of galacto- and fructo-oligosaccharides or viable Bifidobacterium animalis on the intestinal microflora during the first 4 months of life. *Br J Nutr.* **94**: 783-790.
- Blijlevens, N., and Sonis, S. (2007). Palifermin (recombinant keratinocyte growth factor-1): a pleiotropic growth factor with multiple biological activities in preventing chemotherapy- and radiotherapy-induced mucositis. *Ann Oncol.* **18**: 817-826.
- Bogusławska-Tryk, M., Piotrowska, A., and Burlikowska, K. (2012). Dietary fructans and their potential beneficial influence on health and performance parameters in broiler chickens. *J Cent Eur Agric.* **13**: 272-291.
- Bomba, A., Nemcova, R., Gancarc̣ı́kova, S., Herich, R., Guba, P., and Mudroṇova, D. (2002). Improvement of the probiotic effect of micro-organisms by their combination with maltodextrins, fructo-oligosaccharides and polyunsaturated fatty acids. *Br J Nutr.* **88**: S95-S99.
- Boukhettala, N., Ibrahim, A., Aziz, M., Vuichoud, J., Saudan, K.Y., Blum, S., Dechelotte, P., Breuille, D., and Coeffier, M. (2010). A diet containing whey protein, free glutamine, and

transforming growth factor-beta ameliorates nutritional outcome and intestinal mucositis during repeated chemotherapeutic challenges in rats. *J Nutr.* **140**: 799-805.

Bovee-Oudenhoven, I.M., ten Bruggencate, S.J., Lettink-Wissink, M.L., and van der Meer, R. (2003). Dietary fructo-oligosaccharides and lactulose inhibit intestinal colonisation but stimulate translocation of salmonella in rats. *Gut.* **52**: 1572-1578.

Brand-Miller, J.C., McVeagh, P., McNeil, Y., and Messer, M. (1998). Digestion of human milk oligosaccharides by healthy infants evaluated by the lactulose hydrogen breath test. *J Pediatr.* **133**: 95-98.

Brubaker, P.L., Izzo, A., Hill, M., and Drucker, D.J. (1997). Intestinal function in mice with small bowel growth induced by glucagon-like peptide-2. *Am J Physiol.* **272**: E1050-1058.

Cani, P.D., Possemiers, S., Van de Wiele, T., Guiot, Y., Everard, A., Rottier, O., Geurts, L., Naslain, D., Neyrinck, A., Lambert, D.M., Muccioli, G.G., and Delzenne, N.M. (2009). Changes in gut microbiota control inflammation in obese mice through a mechanism involving GLP-2-driven improvement of gut permeability. *Gut.* **58**: 1091-1103.

Cheah, K.Y., Howarth, G.S., Yazbeck, R., Wright, T.H., Whitford, E.J., Payne, C., Butler, R.N., and Bastian, S.E. (2009). Grape seed extract protects IEC-6 cells from chemotherapy-induced cytotoxicity and improves parameters of small intestinal mucositis in rats with experimentally-induced mucositis. *Cancer Biol Ther.* **8**: 382-390.

Cherbut, C., Michel, C., and Lecannu, G. (2003). The prebiotic characteristics of fructooligosaccharides are necessary for reduction of TNBS-induced colitis in rats. *J Nutr.* **133**: 21-27.

- Choi, G.H., Park, H.S., Kim, K.R., Choi, H.N., Jang, K.Y., Chung, M.J., Kang, M.J., Lee, D.G., and Moon, W.S. (2008). Increased expression of epidermal growth factor receptor and betacellulin during the early stage of gastric ulcer healing. *Mol Med Report*. **1**: 505-510.
- Choi, K., Lee, S.S., Oh, S.J., Lim, S.Y., Jeon, W.K., Oh, T.Y., and Kim, J.W. (2007). The effect of oral glutamine on 5-fluorouracil/leucovorin-induced mucositis/stomatitis assessed by intestinal permeability test. *Clin Nutr*. **26**: 57-62.
- Clark, M.J., Robien, K., and Slavin, J.L. (2012). Effect of prebiotics on biomarkers of colorectal cancer in humans: a systematic review. *Nutr Rev*. **70**: 436-443.
- Clarke, J., Butler, R., Howarth, G., Read, L., and Regester, G. (2002). Exposure of oral mucosa to bioactive milk factors reduces severity of chemotherapy-induced mucositis in the hamster. *Oral Oncol*. **38**: 478-485.
- Cool, J.C., Dyer, J.L., Xian, C.J., Butler, R.N., Geier, M.S., and Howarth, G.S. (2005). Pre-treatment with insulin-like growth factor-I partially ameliorates 5-fluorouracil-induced intestinal mucositis in rats. *Growth Horm IGF Res*. **15**: 72-82.
- Daddaoua, A., Martinez-Plata, E., Lopez-Posadas, R., Vieites, J.M., Gonzalez, M., Requena, P., Zarzuelo, A., Suarez, M.D., de Medina, F.S., and Martinez-Augustin, O. (2007). Active hexose correlated compound acts as a prebiotic and is antiinflammatory in rats with hapten-induced colitis. *J Nutr*. **137**: 1222-1228.
- Dahlhoff, M., Horst, D., Gerhard, M., Kolligs, F.T., Wolf, E., and Schneider, M.R. (2008). Betacellulin stimulates growth of the mouse intestinal epithelium and increases adenoma multiplicity in Apc+/Min mice. *FEBS Lett*. **582**: 2911-2915.

- de Souza Oliveira, R.P., Perego, P., de Oliveira, M.N., and Converti, A. (2012). Effect of inulin on the growth and metabolism of a probiotic strain of *Lactobacillus rhamnosus* in co-culture with *Streptococcus thermophilus*. *LWT - Food Sci Technol.* **47**: 358-363.
- Delzenne, N.M., and Roberfroid, M.R. (1994). Physiological effects of non-digestible oligosaccharides. *Lebensm.-Wiss.u. Technol.* **27**: 1-6.
- Eiwegger, T., Stahl, B., Haidl, P., Schmitt, J., Boehm, G., Dehlink, E., Urbanek, R., and Szepefalusi, Z. (2010). Prebiotic oligosaccharides: in vitro evidence for gastrointestinal epithelial transfer and immunomodulatory properties. *Pediatr Allergy Immunol.* **21**: 1179-1188.
- Fanaro, S., Boehm, G., Garssen, J., Knol, J., Mosca, F., Stahl, B., and Vigi, V. (2005). Galacto-oligosaccharides and long-chain fructo-oligosaccharides as prebiotics in infant formulas: a review. *Acta Paediatr Suppl.* **94**: 22-26.
- Fedorak, R.N., and Madsen, K.L. (2004). Probiotics and prebiotics in gastrointestinal disorders. *Curr Opin Gastroenterol.* **20**: 146-155.
- Forchielli, M.L., and Walker, W.A. (2005). The role of gut-associated lymphoid tissues and mucosal defence. *Br J Nutr.* **93**(Suppl 1): S41-48.
- Fukuda, M., Komiyama, Y., Mitsuyama, K., Andoh, A., Aoyama, T., Matsumoto, Y., and Kanauchi, O. (2011). Prebiotic treatment reduced preneoplastic lesions through the downregulation of toll like receptor 4 in a chemo-induced carcinogenic model. *J Clin Biochem Nutr.* **49**: 57-61.
- Geraylou, Z., Souffreau, C., Rurangwa, E., D'Hondt, S., Callewaert, L., Courtin, C.M., Delcour, J.A., Buyse, J., and Ollevier, F. (2012). Effects of arabinoxylan-oligosaccharides (AXOS) on

juvenile Siberian sturgeon (*Acipenser baerii*) performance, immune responses and gastrointestinal microbial community. *Fish Shellfish Immunol.* (In Press)

Gibson, G.R. (1998). Dietary modulation of the human gut microflora using prebiotics. *Br J Nutr.* **80**: S209-212.

Gibson, G.R., and Roberfroid, M.B. (1995). Dietary modulation of the human colonic microbiota: introducing the concept of prebiotics. *J Nutr.* **125**: 1401-1412.

Gibson, R.J., Bowen, J.M., and Keefe, D.M. (2005). Palifermin reduces diarrhea and increases survival following irinotecan treatment in tumor-bearing DA rats. *Int J Cancer.* **116**: 464-470.

Gibson, R.J., Bowen, J.M., Inglis, M.R., Cummins, A.G., and Keefe, D.M. (2003). Irinotecan causes severe small intestinal damage, as well as colonic damage, in the rat with implanted breast cancer. *J Gastroenterol Hepatol.* **18**: 1095-1100.

Gibson, R.J., Keefe, D.M., Clarke, J.M., Register, G.O., Thompson, F.M., Goland, G.J., Edwards, B.G., and Cummins, A.G. (2002). The effect of keratinocyte growth factor on tumour growth and small intestinal mucositis after chemotherapy in the rat with breast cancer. *Cancer Chemother Pharmacol.* **50**: 53-58.

Gibson, R.J., Stringer, A.M., Bowen, J.M., Logan, R.M., Yeoh, A.S., Burns, J., Alvarez, E., and Keefe, D.M. (2007). Velafermin Improves Gastrointestinal Mucositis Following Irinotecan Treatment in Tumor-Bearing DA Rats. *Cancer Biol Ther.* **6**: 541-547.

Gopalakrishnan, A., Clinthorne, J.F., Rondini, E.A., McCaskey, S.J., Gurzell, E.A., Langohr, I.M., Gardner, E.M., and Fenton, J.I. (2012). Supplementation with galacto-oligosaccharides increases the percentage of NK cells and reduces colitis severity in smad3-deficient mice. *J Nutr.* **142**: 1336-1342.

- Gori, A., Rizzardini, G., Van't Land, B., Amor, K.B., van Schaik, J., Torti, C., Quirino, T., Tincati, C., Bandera, A., Knol, J., Benlhassan-Chahour, K., Trabattoni, D., Bray, D., Vriesema, A., Welling, G., Garssen, J., and Clerici, M. (2011). Specific prebiotics modulate gut microbiota and immune activation in HAART-naive HIV-infected adults: results of the "COPA" pilot randomized trial. *Mucosal Immunol.* **4**: 554-563.
- Hosono, A., Ozawa, A., Kato, R., Ohnishi, Y., Nakanishi, Y., Kimura, T., and Nakamura, R. (2003). Dietary fructooligosaccharides induce immunoregulation of intestinal IgA secretion by murine Peyer's patch cells. *Biosci Biotechnol Biochem.* **67**: 758-764.
- Howarth, G.S. (2003). Insulin-like growth factor-I and the gastrointestinal system: therapeutic indications and safety implications. *J Nutr.* **133**: 2109-2112.
- Howarth, G.S. (2012). Commentary on prebiotic utility in colitis: will inflammasomics hold the key? *J Nutr.* **142**: 1189-1190.
- Howarth, G.S., Cool, J.C., Bourne, A.J., Ballard, F.J., and Read, L.C. (1998). Insulin-like growth factor-I (IGF-I) stimulates regrowth of the damaged intestine in rats, when administered following, but not concurrent with, methotrexate. *Growth Factors.* **15**: 279-292.
- Howarth, G.S., Francis, G.L., Cool, J.C., Xu, X., Byard, R.W., and Read, L.C. (1996). Milk growth factors enriched from cheese whey ameliorate intestinal damage by methotrexate when administered orally to rats. *J Nutr.* **126**: 2519-2530.
- Ibuki, M., Kovacs-Nolan, J., Fukui, K., Kanatani, H., and Mine, Y. (2010). Analysis of gut immune-modulating activity of beta-1,4-mannobiose using microarray and real-time reverse transcription polymerase chain reaction. *Poult Sci.* **89**: 1894-1904.

- Janardhana, V., Broadway, M.M., Bruce, M.P., Lowenthal, J.W., Geier, M.S., Hughes, R.J., and Bean, A.G. (2009). Prebiotics modulate immune responses in the gut-associated lymphoid tissue of chickens. *J Nutr.* **139**: 1404-1409.
- Kanauchi, O., Oshima, T., Andoh, A., Shioya, M., and Mitsuyama, K. (2008). Germinated barley foodstuff ameliorates inflammation in mice with colitis through modulation of mucosal immune system. *Scand J Gastroenterol.* **43**: 1346-1352.
- Kanauchi, O., Serizawa, I., Araki, Y., Suzuki, A., Andoh, A., Fujiyama, Y., Mitsuyama, K., Takaki, K., Toyonaga, A., Sata, M., and Bamba, T. (2003). Germinated barley foodstuff, a prebiotic product, ameliorates inflammation of colitis through modulation of the enteric environment. *J Gastroenterol.* **38**: 134-141.
- Khan, A.R., Yousaf, M.S., Rehman, H., Zaneb, H., Pasha, T.N., Fatima, N., Khushal, A., Ijaz, A., Ashraf, K., and Afzal, M. (2012). Response of maternally isolated rock pigeons (*Columba livia domestica*) to different dietary concentrations of mannan-oligosaccharide. *Poult Sci.* **91**: 1598-1603.
- Kim, J.K., Kim, C.S., Ahn, H.J., Bang, S.R., Kim, S.J., Yoo, H.Y., and Jeong, H.S. (2010). Early recombinant human epidermal growth factor treatment recovers the irradiation-induced decrease of Na⁺ absorption prior to the definite histological mucositis. *Biomed Pharmacother.* **64**: 594-599.
- Konieczna, P., Groeger, D., Ziegler, M., Frei, R., Ferstl, R., Shanahan, F., Quigley, E.M., Kiely, B., Akdis, C.A., and O'Mahony, L. (2012). Bifidobacterium infantis 35624 administration induces Foxp3 T regulatory cells in human peripheral blood: potential role for myeloid and plasmacytoid dendritic cells. *Gut.* **61**: 354-366.

- Kumar, V., Sinha, A.K., Makkar, H.P., de Boeck, G., and Becker, K. (2012). Dietary roles of non-starch polysachharides in human nutrition: a review. *Crit Rev Food Sci Nutr.* **52**: 899-935.
- Lalla, R.V., and Peterson, D.E. (2006). Treatment of mucositis, including new medications. *Cancer J.* **12**: 348-354.
- Lalla, R.V., Schubert, M.M., Bensadoun, R.J., and Keefe, D. (2006). Anti-inflammatory agents in the management of alimentary mucositis. *Support Care Cancer.* **14**: 558-565.
- Lamsal, B.P. (2012). Production, health aspects and potential food uses of dairy prebiotic galactooligosaccharides. *J Sci Food Agric.* **92**: 2020-2028.
- Lara-Villoslada, F., de Haro, O., Camuesco, D., Comalada, M., Velasco, J., Zarzuelo, A., Xaus, J., and Galvez, J. (2006). Short-chain fructooligosaccharides, in spite of being fermented in the upper part of the large intestine, have anti-inflammatory activity in the TNBS model of colitis. *Eur J Nutr.* **45**: 418-425.
- Lecerf, J.M., Depeint, F., Clerc, E., Dugenet, Y., Niamba, C.N., Rhazi, L., Cayzeele, A., Abdelnour, G., Jaruga, A., Younes, H., Jacobs, H., Lambrey, G., Abdelnour, A.M., and Pouillart, P.R. (2012). Xylo-oligosaccharide (XOS) in combination with inulin modulates both the intestinal environment and immune status in healthy subjects, while XOS alone only shows prebiotic properties. *Br J Nutr.* (In Press)
- Leforestier, G., Blais, A., Blachier, F., Marsset-Baglieri, A., Davila-Gay, A.M., Perrin, E., and Tome, D. (2009). Effects of galacto-oligosaccharide ingestion on the mucosa-associated mucins and sucrase activity in the small intestine of mice. *Eur J Nutr.* **48**: 457-464.

- Lindsay, R.J., Geier, M.S., Yazbeck, R., Butler, R.N., and Howarth, G.S. (2010). Orally administered emu oil decreases acute inflammation and alters selected small intestinal parameters in a rat model of mucositis. *Br J Nutr.* **104**: 513-519.
- Liu, P., Piao, X.S., Kim, S.W., Wang, L., Shen, Y.B., Lee, H.S., and Li, S.Y. (2008). Effects of chito-oligosaccharide supplementation on the growth performance, nutrient digestibility, intestinal morphology, and fecal shedding of *Escherichia coli* and *Lactobacillus* in weaning pigs. *J Anim Sci.* **86**: 2609-2618.
- Liu, W., Li, X., Wong, Y.S., Zheng, W., Zhang, Y., Cao, W., and Chen, T. (2012). Selenium Nanoparticles as a Carrier of 5-Fluorouracil to Achieve Anticancer Synergism. *ACS Nano.* (In Press)
- Logan, R.M., Stringer, A.M., Bowen, J.M., Gibson, R.J., Sonis, S.T., and Keefe, D.M. (2009). Is the pathobiology of chemotherapy-induced alimentary tract mucositis influenced by the type of mucotoxic drug administered? *Cancer Chemother Pharmacol.* **63**: 239-251.
- Logan, R.M., Stringer, A.M., Bowen, J.M., Yeoh, A.S., Gibson, R.J., Sonis, S.T., and Keefe, D.M. (2007). The role of pro-inflammatory cytokines in cancer treatment-induced alimentary tract mucositis: Pathobiology, animal models and cytotoxic drugs. *Cancer Treat Rev.* **33**: 448-460.
- Looijer-van Langen, M.A., and Dieleman, L.A. (2009). Probiotics in chronic intestinal inflammation. *Inflamm Bowel Dis.* **15**: 454-462.
- Maki, K.C., Gibson, G.R., Dickmann, R.S., Kendall, C.W., Oliver Chen, C.Y., Costabile, A., Comelli, E.M., McKay, D.L., Almeida, N.G., Jenkins, D., Zello, G.A., and Blumberg, J.B.

- (2012). Digestive and physiologic effects of a wheat bran extract, arabino-xylan-oligosaccharide, in breakfast cereal. *Nutrition*. (In Press)
- Mankan, A.K., Lawless, M.W., Gray, S.G., Kelleher, D., and McManus, R. (2009). NF-kappaB regulation: the nuclear response. *J Cell Mol Med*. **13**: 631-643.
- Marotti, I., Bregola, V., Aloisio, I., Di Gioia, D., Bosi, S., Di Silvestro, R., Quinn, R., and Dinelli, G. (2012). Prebiotic effect of soluble fibres from modern and old durum-type wheat varieties on Lactobacillus and Bifidobacterium strains. *J Sci Food Agric*. **92**: 2133-2140.
- Matthews, G.M., Howarth, G.S., and Butler, R.N. (2007). Short-chain fatty acid modulation of apoptosis in the Kato III human gastric carcinoma cell line. *Cancer Biol Ther*. **6**: 1051-1057.
- Niewold, T.A., Schroyen, M., Geens, M.M., Verhelst, R.S.B., and Courtin, C.M. (2012). Dietary inclusion of arabinoxylan oligosaccharides (AXOS) down regulates mucosal responses to a bacterial challenge in a piglet model. *J Funct Foods*. **4**: 626-635.
- Nishimukai, M., Watanabe, J., Taguchi, H., Senoura, T., Hamada, S., Matsui, H., Yamamoto, T., Wasaki, J., Hara, H., and Ito, S. (2008). Effects of epilactose on calcium absorption and serum lipid metabolism in rats. *J Agric Food Chem*. **56**: 10340-10345.
- Panter, M.S., Jain, A., Leonhardt, R.M., Ha, T., and Cresswell, P. (2012). Dynamics of major histocompatibility complex class I association with the human peptide-loading complex. *J Biol Chem*. (In Press)
- Parnell, J.A., and Reimer, R.A. (2012). Prebiotic fiber modulation of the gut microbiota improves risk factors for obesity and the metabolic syndrome. *Gut Microbes*. **3**: 29-34.

- Patra, A.K. (2011). Responses of feeding prebiotics on nutrient digestibility, faecal microbiota composition and short-chain fatty acid concentrations in dogs: a meta-analysis. *Animal*. **5**: 1743-1750.
- Pilzner, C., Buhling, F., Reinheckel, T., Chwieralski, C., Rathinasamy, A., Lauenstein, H.D., Wex, T., Welte, T., Braun, A., and Groneberg, D.A. (2012). Allergic Airway Inflammation in Mice Deficient for the Antigen-Processing Protease Cathepsin E. *Int Arch Allergy Immunol*. **159**: 367-383.
- Pratt, V.C., Tappenden, K.A., McBurney, M.I., and Field, C.J. (1996). Short-chain fatty acid-supplemented total parenteral nutrition improves nonspecific immunity after intestinal resection in rats. *JPEN J Parenter Enteral Nutr*. **20**: 264-271.
- Prisciandaro, L.D., Geier, M.S., Butler, R.N., Cummins, A.G., and Howarth, G.S. (2011). Evidence supporting the use of probiotics for the prevention and treatment of chemotherapy-induced intestinal mucositis. *Crit Rev Food Sci Nutr*. **51**: 239-247.
- Qu, Y., Misaghi, S., Izrael-Tomasevic, A., Newton, K., Gilmour, L.L., Lamkanfi, M., Louie, S., Kayagaki, N., Liu, J., Komuves, L., Cupp, J.E., Arnott, D., Monack, D., and Dixit, V.M. (2012). Phosphorylation of NLRC4 is critical for inflammasome activation. *Nature*. (In Press)
- Rancez, M., Couedel-Courteille, A., and Cheynier, R. (2012). Chemokines at mucosal barriers and their impact on HIV infection. *Cytokine Growth Factor Rev*. **23**: 233-243.
- Reid, G. (2005). The importance of guidelines in the development and application of probiotics. *Curr Pharm Des*. **11**: 11-16.

- Robbe, C., Capon, C., Coddeville, B., and Michalski, J.C. (2004). Structural diversity and specific distribution of O-glycans in normal human mucins along the intestinal tract. *Biochem J.* **384**: 307-316.
- Roberfroid, M. (2007). Prebiotics: the concept revisited. *J Nutr.* **137**: 830S-837S.
- Roberfroid, M., Gibson, G.R., Hoyles, L., McCartney, A.L., Rastall, R., Rowland, I., Wolvers, D., Watzl, B., Szajewska, H., Stahl, B., Guarner, F., Respondek, F., Whelan, K., Coxam, V., Davicco, M.J., Leotoing, L., Wittrant, Y., Delzenne, N.M., Cani, P.D., Neyrinck, A.M., and Meheust, A. (2010). Prebiotic effects: metabolic and health benefits. *Br J Nutr.* **104**(Suppl 2): S1-63.
- Roberfroid, M.B. (1998). Prebiotics and synbiotics: concepts and nutritional properties. *Br J Nutr.* **80**: S197-202.
- Rodenburg, W., Keijer, J., Kramer, E., Vink, C., van der Meer, R., and Bovee-Oudenhoven, I.M. (2008). Impaired barrier function by dietary fructo-oligosaccharides (FOS) in rats is accompanied by increased colonic mitochondrial gene expression. *BMC Genomics.* **9**: 144.
- Roncucci, L., Di Donato, P., Carati, L., Ferrari, A., Perini, M., Bertoni, G., Bedogni, G., Paris, B., Svanoni, F., Girola, M., and et al. (1993). Antioxidant vitamins or lactulose for the prevention of the recurrence of colorectal adenomas. Colorectal Cancer Study Group of the University of Modena and the Health Care District 16. *Dis Colon Rectum.* **36**: 227-234.
- Salvini, F., Riva, E., Salvatici, E., Boehm, G., Jelinek, J., Banderali, G., and Giovannini, M. (2011). A specific prebiotic mixture added to starting infant formula has long-lasting bifidogenic effects. *J Nutr.* **141**: 1335-1339.

- Sauer, J., Richter, K.K., and Pool-Zobel, B.L. (2007). Products formed during fermentation of the prebiotic inulin with human gut flora enhance expression of biotransformation genes in human primary colon cells. *Br J Nutr.* **97**: 928-937.
- Scharlau, D., Borowicki, A., Habermann, N., Hofmann, T., Klenow, S., Miene, C., Munjal, U., Stein, K., and Gleib, M. (2009). Mechanisms of primary cancer prevention by butyrate and other products formed during gut flora-mediated fermentation of dietary fibre. *Mutat Res.* **682**: 39-53.
- Shigeoka, A.A., Holscher, T.D., King, A.J., Hall, F.W., Kiosses, W.B., Tobias, P.S., Mackman, N., and McKay, D.B. (2007). TLR2 is constitutively expressed within the kidney and participates in ischemic renal injury through both MyD88-dependent and -independent pathways. *J Immunol.* **178**: 6252-6258.
- Shimizu, K., Ogura, H., Asahara, T., Nomoto, K., Morotomi, M., Tasaki, O., Matsushima, A., Kuwagata, Y., Shimazu, T., and Sugimoto, H. (2012). Probiotic/Synbiotic Therapy for Treating Critically Ill Patients from a Gut Microbiota Perspective. *Dig Dis Sci.* (In Press)
- Shoaf, K., Mulvey, G.L., Armstrong, G.D., and Hutkins, R.W. (2006). Prebiotic galactooligosaccharides reduce adherence of enteropathogenic *Escherichia coli* to tissue culture cells. *Infect Immun.* **74**: 6920-6928.
- Sinclair, H.R., Smejkal, C.W., Glister, C., Kemp, F., van den Heuvel, E., de Slegte, J., Gibson, G.R., and Rastall, R.A. (2008). Sialyloligosaccharides inhibit cholera toxin binding to the GM1 receptor. *Carbohydr Res.* **343**: 2589-2594.

- Smith, C.L., Geier, M.S., Yazbeck, R., Torres, D.M., Butler, R.N., and Howarth, G.S. (2008). *Lactobacillus fermentum* BR11 and fructo-oligosaccharide partially reduce jejunal inflammation in a model of intestinal mucositis in rats. *Nutr Cancer*. **60**: 757-767.
- Sonis, S.T. (2004a). Pathobiology of mucositis. *Semin Oncol Nurs*. **20**: 11-15.
- Sonis, S.T. (2004b). The pathobiology of mucositis. *Nature Reviews Cancer*. **4**: 277-284.
- Sonis, S.T., Elting, L.S., Keefe, D., Peterson, D.E., Schubert, M., Hauer-Jensen, M., Bekele, B.N., Raber-Durlacher, J., Donnelly, J.P., and Rubenstein, E.B. (2004). Perspectives on cancer therapy-induced mucosal injury: pathogenesis, measurement, epidemiology, and consequences for patients. *Cancer*. **100**: 1995-2025.
- Specian, R.D., and Oliver, M.G. (1991). Functional biology of intestinal goblet cells. *Am J Physiol*. **260**: C183-193.
- Stringer, A.M., Gibson, R.J., Bowen, J.M., and Keefe, D.M. (2009a). Chemotherapy-induced modifications to gastrointestinal microflora: evidence and implications of change. *Curr Drug Metab*. **10**: 79-83.
- Stringer, A.M., Gibson, R.J., Logan, R.M., Bowen, J.M., Yeoh, A.S., Hamilton, J., and Keefe, D.M. (2009b). Gastrointestinal microflora and mucins may play a critical role in the development of 5-Fluorouracil-induced gastrointestinal mucositis. *Exp Biol Med (Maywood)*. **234**: 430-441.
- Strowig, T., Henao-Mejia, J., Elinav, E., and Flavell, R. (2012). Inflammasomes in health and disease. *Nature*. **481**: 278-286.
- Sukhotnik, I., Shteinberg, D., Ben Lulu, S., Bashenko, Y., Mogilner, J.G., Ure, B.M., Shaoul, R., and Coran, A.G. (2008a). Effect of transforming growth factor-alpha on enterocyte apoptosis is

correlated with EGF receptor expression along the villus-crypt axis during methotrexate-induced intestinal mucositis in a rat. *Apoptosis*. **13**: 1344-1355.

Sukhotnik, I., Shteinberg, D., Ben Lulu, S., Bashenko, Y., Mogilner, J.G., Ure, B.M., Shaoul, R., Shamian, B., and Coran, A.G. (2008b). Transforming growth factor-alpha stimulates enterocyte proliferation and accelerates intestinal recovery following methotrexate-induced intestinal mucositis in a rat and a cell culture model. *Pediatr Surg Int*. **24**: 1303-1311.

Sung, H.Y., and Choi, Y.S. (2008). Fructooligosaccharide and soy isoflavone suppress colonic aberrant crypt foci and cyclooxygenase-2 expression in dimethylhydrazine-treated rats. *J Med Food*. **11**: 78-85.

Ten Bruggencate, S.J., Bovee-Oudenhoven, I.M., Lettink-Wissink, M.L., Katan, M.B., and van der Meer, R. (2006). Dietary fructooligosaccharides affect intestinal barrier function in healthy men. *J Nutr*. **136**: 70-74.

Torres, D.M., Tooley, K.L., Butler, R.N., Smith, C.L., Geier, M.S., and Howarth, G.S. (2008). Lyprinol only partially improves indicators of small intestinal integrity in a rat model of 5-fluorouracil-induced mucositis. *Cancer Biol Ther*. **7**: 295-302.

Tran, C.D., Howarth, G.S., Coyle, P., Philcox, J.C., Rofe, A.M., and Butler, R.N. (2003). Dietary supplementation with zinc and a growth factor extract derived from bovine cheese whey improves methotrexate-damaged rat intestine. *Am J Clin Nutr*. **77**: 1296-1303.

Trevisi, P., De Filippi, S., Minieri, L., Mazzoni, M., Modesto, M., Biavati, B., and Bosi, P. (2008). Effect of fructo-oligosaccharides and different doses of *Bifidobacterium animalis* in a weaning diet on bacterial translocation and Toll-like receptor gene expression in pigs. *Nutrition*. **24**: 1023-1029.

- van Hoffen, E., Ruiter, B., Faber, J., M'Rabet, L., Knol, E.F., Stahl, B., Arslanoglu, S., Moro, G., Boehm, G., and Garssen, J. (2009). A specific mixture of short-chain galacto-oligosaccharides and long-chain fructo-oligosaccharides induces a beneficial immunoglobulin profile in infants at high risk for allergy. *Allergy*. **64**: 484-487.
- van Vliet, M.J., Harmsen, H.J., de Bont, E.S., and Tissing, W.J. (2010). The role of intestinal microbiota in the development and severity of chemotherapy-induced mucositis. *PLoS Pathog.* **6**: e1000879.
- van't Land, B., Schijf, M., van Esch, B.C., van Bergenhenegouwen, J., Bastiaans, J., Schouten, B., Boon, L., and Garssen, J. (2010). Regulatory T-cells have a prominent role in the immune modulated vaccine response by specific oligosaccharides. *Vaccine*. **28**: 5711-5717.
- Vos, A.P., Haarman, M., Bucu, A., Govers, M., Knol, J., Garssen, J., Stahl, B., Boehm, G., and M'Rabet, L. (2006). A specific prebiotic oligosaccharide mixture stimulates delayed-type hypersensitivity in a murine influenza vaccination model. *Int Immunopharmacol.* **6**: 1277-1286.
- Walton, G.E., Lu, C., Trogh, I., Arnaut, F., and Gibson, G.R. (2012). A randomised, double-blind, placebo controlled cross-over study to determine the gastrointestinal effects of consumption of arabinoxylnoligosaccharides enriched bread in healthy volunteers. *Nutr J.* **11**: 36.
- Watanabe, J., Nishimukai, M., Taguchi, H., Senoura, T., Hamada, S., Matsui, H., Yamamoto, T., Wasaki, J., Hara, H., and Ito, S. (2008). Prebiotic properties of epilactose. *J Dairy Sci.* **91**: 4518-4526.

- Willard, M.D., Simpson, R.B., Delles, E.K., Cohen, N.D., Fossum, T.W., Kolp, D., and Reinhart, G. (1994). Effects of dietary supplementation of fructo-oligosaccharides on small intestinal bacterial overgrowth in dogs. *Am J Vet Res.* **55**: 654-659.
- Winkler, J., Butler, R., and Symonds, E. (2007). Fructo-oligosaccharide reduces inflammation in a dextran sodium sulphate mouse model of colitis. *Dig Dis Sci.* **52**: 52-58.
- Wright, T.H., Yazbeck, R., Lymn, K.A., Whitford, E.J., Cheah, K.Y., Butler, R.N., Feinle-Bisset, C., Pilichiewicz, A.N., Mashtoub, S., and Howarth, G.S. (2009). The herbal extract, Iberogast, improves jejunal integrity in rats with 5-Fluorouracil (5-FU)-induced mucositis. *Cancer Biol Ther.* **8**: 923-929.
- Xu, B., Wang, Y., Li, J., and Lin, Q. (2009). Effect of prebiotic xylooligosaccharides on growth performances and digestive enzyme activities of allogynogenetic crucian carp (*Carassius auratus gibelio*). *Fish Physiol Biochem.* **35**: 351-357.
- Yamamoto, M., Sato, S., Hemmi, H., Sanjo, H., Uematsu, S., Kaisho, T., Hoshino, K., Takeuchi, O., Kobayashi, M., Fujita, T., Takeda, K., and Akira, S. (2002). Essential role for TIRAP in activation of the signalling cascade shared by TLR2 and TLR4. *Nature.* **420**: 324-329.
- Yamamoto, M., Sato, S., Hemmi, H., Uematsu, S., Hoshino, K., Kaisho, T., Takeuchi, O., Takeda, K., and Akira, S. (2003). TRAM is specifically involved in the Toll-like receptor 4-mediated MyD88-independent signaling pathway. *Nat Immunol.* **4**: 1144-1150.
- Yanahira, S., Morita, M., Aoe, S., Suguri, T., Nakajima, I., and Deya, E. (1995). Effects of lactitol-oligosaccharides on the intestinal microflora in rats. *J Nutr Sci Vitaminol (Tokyo).* **41**: 83-94.

- Yang, Y., Iji, P.A., Kocher, A., Mikkelsen, L.L., and Choct, M. (2008a). Effects of mannanoligosaccharide and fructooligosaccharide on the response of broilers to pathogenic *Escherichia coli* challenge. *Br Poult Sci.* **49**: 550-559.
- Yang, Y., Iji, P.A., Kocher, A., Thomson, E., Mikkelsen, L.L., and Choct, M. (2008b). Effects of mannanoligosaccharide in broiler chicken diets on growth performance, energy utilisation, nutrient digestibility and intestinal microflora. *Br Poult Sci.* **49**: 186-194.
- Yasuda, A., Inoue, K., Sanbongi, C., Yanagisawa, R., Ichinose, T., Tanaka, M., Yoshikawa, T., and Takano, H. (2012). Dietary supplementation with fructooligosaccharides attenuates allergic peritonitis in mice. *Biochem Biophys Res Commun.* **422**: 546-550.
- Zhang, Z., and Li, L.Y. (2012). TNFSF15 Modulates Neovascularization and Inflammation. *Cancer Microenviron.* (In Press)