

## MA... MÄNNISTE

Physiological ecology of greenfinches:  
information content of feathers in relation  
to immune function and behavior





## MARJU MÄNNISTE

Physiological ecology of greenfinches:  
information content of feathers in relation  
to immune function and behavior



Department of Zoology, Institute of Ecology and Earth Sciences, Faculty of Science and Technology, University of Tartu

Dissertation was accepted for the commencement of the degree of *Doctor philosophiae* in animal ecology at the University of Tartu on May 19, 2014 by the Scientific Council of the Institute of Ecology and Earth Sciences, University of Tartu.

Supervisor: Prof. Peeter Hõrak, University of Tartu, Estonia

Opponent: Prof. Anders Pape Møller, Université Paris-Sud, France

Commencement: Room 301, 46 Vanemuise Street, Tartu, on September 19, 10.15 a.m., 2014

Publication of this thesis is granted by the Institute of Ecology and Earth Sciences, University of Tartu, and by the Doctoral School of Earth Sciences and Ecology created under the auspices of European Social Fund.



European Union  
European Social Fund



Investing in your future

ISSN 1024-6479

ISBN 978-9949-32-643-3 (print)

ISBN 978-9949-32-644-0 (pdf)

Copyright: Marju Männiste, 2014

University of Tartu Press

[www.tyk.ee](http://www.tyk.ee)

# CONTENTS

LIST OF ORIGINAL PAPERS .....	6
LIST OF ABBREVIATIONS .....	7
1. INTRODUCTION.....	8
1.1. Physiological ecology and neuro-endocrine-immune communication .....	8
1.2. Information content of feathers .....	11
1.2.1. Feather size.....	11
1.2.2. Feather corticosterone content .....	12
1.2.3. Eumelanotic coloration .....	13
1.3. Aims of the thesis .....	14
2. METHODS .....	17
3. RESULTS AND DISCUSSION .....	19
3.1. How do immune activation and glucocorticoid administration affect feather growth?.....	19
3.2. Synthetic glucocorticoid inhibits CORT deposition into feathers....	21
3.3. High feather CORT indicates better resistance to experimental coccidian infection.....	23
3.4. Is feather CORT indicative of avian personalities?.....	25
3.5. Emerging infectious disease selects for darker plumage coloration.....	27
SUMMARY .....	30
SUMMARY IN ESTONIAN .....	35
ACKNOWLEDGEMENTS .....	40
REFERENCES.....	42
PUBLICATIONS.....	51
CURRICULUM VITAE .....	103

## LIST OF ORIGINAL PAPERS

- I. Männiste M., Hõrak P. (2011). Effects of Immune Activation and Glucocorticoid Administration on Feather Growth in Greenfinches. *Journal of Experimental Zoology Part A: Ecological Genetics and Physiology*, 315A (9), 527–535.
- II. Hõrak P., Männiste M., Meitern R., Sild E., Saks L., Sepp T. (2013). Dexamethasone inhibits corticosterone deposition in feathers of greenfinches. *General and Comparative Endocrinology*, 191, 210–214.
- III. Sild E., Meitern R., Männiste M., Karu U., Hõrak P. (2014). High feather corticosterone indicates better coccidian infection resistance in greenfinches. *General and comparative Endocrinology*, 204, 203–210.
- IV. Sepp T., Männiste M., Kaasik A., Hõrak P. (2014). Multidimensionality of fear in captive greenfinches (*Carduelis chloris*). *Behavioral Ecology and Sociobiology*, 68, 1173–1181.
- V. Männiste M., Hõrak P., (2014). Emerging infectious disease selects for darker plumage coloration in greenfinches. *Frontiers in Ecology and Evolution: Behavioral and Evolutionary Ecology*, 2:4, doi: 10.3389/fevo.2014.00004.

Published papers are reproduced with the permission of the publishers.

Author's contribution to the papers

	I	II	III	IV	V
Original idea	*				*
Study design	*			*	*
Data collection	*	*	*	*	*
Data analysis	*	*			*
Manuscript preparation	*	*	*	*	*

## **LIST OF ABBREVIATIONS**

- HPA-axis** – hypothalamus-pituitary-adrenal-axis  
**CORT** – corticosterone  
**MC1R** – melanocortin 1 receptor  
**DEX** – dexamethasone  
**PHA** – phytohaemagglutinin  
**SEM** – structural equation modeling  
**H/L** – heterophil/lymphocyte

# I. INTRODUCTION

## I.1. Physiological ecology and neuro-endocrine-immune communication

Physiological ecology is a hybrid field integrating disciplines of physiology and ecology. Research in this field has two major goals: to understand how the interaction between an organism and its environment determine characteristics of individuals that are relevant to ecology, and to understand how these individual characteristics affect population and interspecific dynamics (Kingsolver 1989). In response to interaction with the environment organisms have developed highly complex nervous, endocrine and immunological systems to elicit appropriate responses to various biotic and abiotic stimuli. Up until recently, however, these systems were regarded as separate entities, especially in the case of immunity. Realization that nervous, endocrine and immunological systems are closely interconnected and are involved in a bidirectional communication gave rise to the field of immunoeology (ecoimmunology) in the 1990s (Martin *et al.* 2011) and psychoneuroimmunology in the early 1970s (Ader 2001).

Immunoeology is a newly emergent, interdisciplinary research field that examines interactions among host physiology and disease ecology in a wide range of environmentally relevant contexts (Demas & Nelson 2011). Immunoeological studies give great attention to extrinsic and intrinsic factors that explain why organisms vary in their immune responses to pathogens in different environments and time points. To ensure ultimate protection, it would make sense to use one's immune system maximally all the time. Yet organisms exhibit remarkable variability and seasonality in their immune function (Nelson & Demas 1996; Martin *et al.* 2008) and in capacity of defence against pathogens and parasites (Graham *et al.* 2011). Immunoeology proposes that susceptibility persists because immune defense is just one element of a context-dependent, integrated, whole-organism response to parasitism (Graham *et al.* 2011; Martin *et al.* 2011). Immunoeological research also assumes trade-offs between investment of resources into life-history traits (Ricklefs & Wikelski 2002) and various physiological processes (Moore & Hopkins 2009) that integrate the organism to form a whole. In order to understand these trade-offs, it is imperative to quantify the costs and benefits that accompany immune defenses and other vital physiological functions of an organism.

Among the many factors that can affect disease outcome is organism's stress response. In response to an acute stressor, the stress response prepares the nervous, cardiovascular and neuroendocrine systems for fight or flight response; it may also prepare the immune system for wounding or infection which may be imposed by the stressor (Dhabhar 2002). As the link between stress and immune function is bidirectional, the activation of immune system can also cause a rise in stress hormone levels (Adamo 2010). Although the general principles of



interaction between the stress response and immune system are known, the adaptive explanations are often incomplete (Adamo 2012). For instance, the question about why chronic stress in general appears immunosuppressive lacks definite answers (Miller *et al.* 2002; Segerstrom 2007; Martin 2009). E.g., assuming that the CORT response serves to “turn off” heightened immune responses that could result in autoimmune damage if sustained over long periods (Sapolsky *et al.* 2000; Martin 2009), still leaves the question about whether the benefits of avoidance of immunopathology would normally outweigh the costs of infection in immunocompromised organisms under natural conditions (e.g. Graham *et al.* 2005).

Brain-immune communication necessitates integrating behavior into the study of stress and immunity. Psychoneuroimmunology is a research field that combines the studies of the relationships among behavioral, neural, endocrine and immune processes (Ader 2001). Recently, researchers from a broad array of ecological sub-disciplines (e.g. population biology, ecophysiology, behavioral ecology) have begun to consider between-individual variation in behavior as an important ecological and evolutionary characteristic of wild populations (Careau *et al.* 2008). Consistent between-individual differences in behavior have been found in many animal species (Van Oers *et al.* 2004). Correlated repeatable and heritable behavioral traits form individually consistent behavior that have been referred to as temperament, personality, behavioral syndrome or coping style. The notion of behavioral syndrome or coping style suggests that behavior is not completely flexible and different behavioral traits do not vary independently of each other (Carere *et al.* 2010). For instance, aggressive individuals tend to be aggressive across different contexts, i.e. they cannot display optimal behavior in all situations. Aggressive behavior should be advantageous in contexts like territory defense or maintaining dominance in social hierarchy, but not in parental care. Studying correlations between different behavioral traits is a key to fully comprehend complex behavioral responses (Sih *et al.* 2004).

The extremes in individually consistent behavior and physiological characteristics in coping with stressful situations can be described as reactive versus proactive coping styles (Koolhaas *et al.* 1999). Individuals with proactive coping style are aggressive, bold and relatively inflexible in their behavioral responses, on the other hand, individuals with reactive coping style can be described as shy, precautious and behaviorally flexible. As there are individually consistent differences in behavior there are also respective consistent differences in stress responsiveness (Cockrem 2007; Bonier *et al.* 2009). These differences are related to differentially activated hypothalamus-pituitary-adrenal-axis (HPA-axis) and sympathetic nervous system, which could link stress responsiveness with personality (Koolhaas *et al.* 1999; Carere *et al.* 2010). Typically proactive, cue-independent, risk-prone individuals try to prevent or manipulate stressors and react to stressful situations with activation of the sympathetic nervous system with the increase of noradrenergic

stimulation. Reactive individuals on the contrary react with activation of the HPA-axis with the consequent rise in blood glucocorticoid (GC) levels (Carere *et al.* 2010). However little is known about the exact nature of causal relationship between differential behavioral and neuroendocrine characteristics (Koolhaas *et al.* 1999).

An excellent example of a mechanism how different physiological systems can be interconnected is provided by pleiotropic effects of genes that bind together physiological, morphological and behavioral traits. For example, genes regulating the synthesis of the main animal coloration pigment – melanin – are also involved in modulation of immune system, stress responses and behavior (Ducrest *et al.* 2008; Gangoso *et al.* 2011). Different animal color morphs have been associated with exploitation of alternative ecological niches, differential responses to immune activation, distinct wing, tail, tarsus, bill length and alternative mating strategies (reviewed by Roulin 2004). The information content of melanotic signals in birds is very diverse. Melanotic color signals are involved in sexual-selection, indicative of social rank or involved in inconspicuous appearance (e.g. camouflage) (Jawor & Breitwisch 2003). Already in 1982, William Hamilton and Marlene Zuk proposed that individual's appearance could reveal cues about organism's ability to successfully deal with infections (Hamilton & Zuk 1982). Now the evidence has started to emerge that melanin-based plumage color could be associated with organism's immune defenses (Ducrest *et al.* 2008; Gangoso *et al.* 2011). Differences in immune system responses could be reflected in melanotic plumage coloration which would thus eventually indicate disease susceptibility (Keyser & Siefferman 2005). Considering the enormous importance of parasites and pathogens as selective agents in the wild, one would also expect that melanotic pigmentation would provide cues predicting morbidity and mortality due to infections.

The second most widespread class of pigments – carotenoids – is fundamentally different from melanins as carotenoids cannot be synthesized by animals and must be acquired from food. As carotenoid pigments are thought to play important role not only in integument coloration but also in other vital physiological functions of the body, the carotenoid pigments have received a lot more attention in immunological and medical research. Therefore the number of studies documenting direct associations between melanotic plumage coloration and infection-induced mortality or morbidity in birds is far more limited (Keyser & Siefferman 2005) than there is for carotenoid-based plumage coloration (Nolan *et al.* 1998; Lindström & Lundström 2000; Hill & Farmer 2005; Van Oort & Dawson 2005; Svensson & Wong 2011; Simons *et al.* 2012).

In addition to pigmentation, other feather properties, such as size, integrity and hormonal content, are potentially valuable sources of information about physiological processes experienced by the organism during feather growth. Avian plumage could thus appear to be a powerful tool for studying the nexus between immune and neuroendocrine systems. Birds are regularly used for studying trade-offs between costly physiological systems, e.g. in the context of

survival, reproduction and development (Bennett & Owens 2002; De La Hera *et al.* 2009). However, lots of information that is decoded in the external morphology of birds appears still underused (Telleria *et al.* 2013) and researches have just started to realize the full potential of information gained from feathers and incorporating it to the main research areas in physiological ecology.

## **1.2. Information content of feathers**

Feathers are the most conspicuous, complex, and varied (both in structure and color pattern) integumentary derivatives of birds (Stettenheim 2000; Prum & Williamson 2002). Essentially, a feather is a tubular integumentary appendage that grows at its base from a feather follicle (Prum & Williamson 2002). A developing feather is sensitive to energetic, nutritional and hormonal milieu (Murphy 1996; Stettenheim 2000; Hill & McGraw 2006). Therefore, various feather properties can reflect intrinsic and extrinsic conditions experienced by an organism during feather growth (Hill & McGraw 2006). The sensitivity to various internal and external stimuli is one of many qualities that makes feathers a unique and exceptional research material (Bortolotti 2010).

### **1.2.1. Feather size**

Feathers are imperative for bird's survival, therefore all factors affecting molt have direct impact on bird's fitness. Reduction in plumage quality can influence survival by decreasing flight performance and increasing thermoregulatory costs (Nilsson & Svensson 1996; Swaddle *et al.* 1996; Dawson *et al.* 2000; DesRochers *et al.* 2009). Indirectly, plumage quality can impinge on fitness by mate choice (e.g. Pryke & Andersson 2005). The functional relationships between different measures of feather size are diverse. Feather mass has been shown to correlate with the number of barbs, and hence the tightness of feather surface (Butler *et al.* 2008). Feather tightness enhances heat absorption, sustained flight (Butler *et al.* 2008) and resistance to abrasion (Dawson *et al.* 2000). Feather length affects take-off speed and maneuverability (Swaddle *et al.* 1996; Matyjasiak *et al.* 2004), and hence predator escape performance. Rachis diameter affects rigidity of feathers (Dawson *et al.* 2000). Fault bars are translucent areas across feathers that can decrease aerial performance due to brakeage of feathers at these sites (Møller *et al.* 2009). Fault bars form when feathers grow under stressful conditions. The incidence of fault bars have been associated with increased mortality in different bird species (Bortolotti *et al.* 2002; Møller *et al.* 2009). In captive birds, an interesting plumage characteristic that conveys information about individual behavior is a tendency to accrue damage to tail feathers in cages. It has been proposed that this trait could be indicative of the ability of bird to cope with captive conditions and is associated

with different immune parameters (Sild *et al.* 2011) and stress responsiveness (Krams *et al.* 2013). All these feather parameters could be potentially affected by physical, physiological or psychological stressors. Feather parameters could thus implicate its bearer's stress responsiveness, amount of encountered stressors or the ability to avoid stressful situations altogether. However, the information about the effects of pathogens and parasites on feather growth has only recently started to emerge (Pap *et al.* 2013), so it would be important to find out how stress- and immunity-related factors affect feather growth or quality and how it associates with fitness.

### **1.2.2. Feather corticosterone content**

In vertebrates, the brain and immune system are linked by two primary pathways: the sympatho-adrenal system and the HPA-axis with the subsequent release of glucocorticoids (e.g. cortisol, corticosterone (CORT)) from the adrenal cortex (Demas *et al.* 2011). Glucocorticoid hormones are the end-product, primary effectors and principal negative regulators of the HPA-axis (Sapolsky *et al.* 2000). The main glucocorticoid to be released in response to stressful stimuli in birds is CORT. It has been proposed that CORT could be the most important hormone in the mediation of environmental effects on physiology and behavior (Demas & Nelson 2011). CORT has many different physiological and behavioral effects aimed to promote fitness at times of stressful events that include suppression of reproductive behavior, regulation of the immune system, increased gluconeogenesis and proteolysis, energy reallocation from fat stores and promotion of escape behavior (Sapolsky *et al.* 2000; Wingfield & Kitaysky 2002).

Due to its sensitivity to environmental perturbations, increases in CORT levels are often used to evaluate individual stress levels (Romero 2004) in ecological, physiological and conservation studies of wild animals. Stress is defined here as a physiological and behavioral state engaged to endure, avoid or recover from an aversive stimulus or condition (Martin 2009). The most common practice is to measure CORT from blood, either to evaluate baseline CORT levels or rise in CORT levels in response to stressor. Measuring CORT from blood though proves to be problematic due to rapid rise of CORT levels caused by investigator-induced stress of capture and handling (Romero & Reed 2005). Measuring CORT from urine or feces has also its own obstacles: problematic sample collection, species specificity and representation of relatively short time frame (Bortolotti *et al.* 2008).

A new method – measuring CORT from feathers – is expected to overcome all above-listed problems. Feather CORT measurement does not necessarily require capture of individuals, it is adjustable to all avian species and most of all, represents longer time periods as CORT is incorporated into feathers during feather growth (from days to weeks) (Bortolotti *et al.* 2008). Due to cumulative nature of CORT deposition into feathers, the feather CORT measurement has

the potential to integrate different aspects of HPA activity, including variation in baseline levels, magnitude and duration of the stimulated stress response and the number of stressors experienced (Bortolotti *et al.* 2008; Lattin *et al.* 2011). The most important advantage in assessing CORT levels from feathers is long-term retrospective measure of stress (Bortolotti *et al.* 2008), which has the potential to add new insights to avian ecology and stress physiology research. For example, feather CORT measurement can be used to tease apart long-term costs on feather growth relating to immune activation or suppression, a question that has been previously overlooked. Feather CORT measurement can also be implemented in conservational biology, avian ecology studies or for example in long-term monitoring of populations or historical trends in CORT using museum specimens, as CORT is stable over long time and in different environmental conditions (Bortolotti *et al.* 2009).

However, several basic aspects of information content of feather CORT measurement are still unknown. For example, it is not known exactly how and when CORT is deposited into feathers and to what extent the variation in feather CORT represents variation found in blood baseline vs induced CORT levels (Bortolotti *et al.* 2008; Lattin *et al.* 2011). Furthermore, virtually nothing is known of the impact of immune system activation on feather CORT or how it relates to other physiological and behavioral measures of stress. Addressing these issues is important as it would facilitate understanding and interpretation of the information content of feather CORT levels.

### **1.2.3. Eumelanotic coloration**

Melanin is the most abundant and widespread pigment in birds and other animals (Hill & McGraw 2006). Melanins consist of two main pigment types—eumelanin and pheomelanin. Eumelanin forms grey and black colors whereas pheomelanin forms brown, red or buff colors (McGraw 2006). Melanin-based coloration is frequently implicated in social communication (Hill & McGraw 2006a) and covaries with many other physiological, morphological and behavioral traits (Roulin 2004). Ducrest and coauthors (2008) suggest that pleiotropic effects of key regulators of melanogenesis might be responsible for the widespread association between melanin-based coloration and other phenotypic traits in wild vertebrates (Emaresi *et al.* 2013). Melanocortins are posttranslational products of the proopiomelanocortin gene. In addition to binding to melanocortin 1 receptor (MC1R) that triggers production of melanin pigments, melanocortins also bind to four other melanocortin receptors (MC2-5R) which have very different functions. These functions include, among others, anti-inflammatory activity, HPA stress response, energy expenditure, sexual activity and aggressiveness (Ducrest *et al.* 2008). By binding to MC4R, melanocortins increase organism's resistance to stressors. By binding to MC1R and MC3-5R, melanocortins reduce acute, allergic and systemic inflammation, septic shock and increase antipyretic activity (Gangoso *et al.* 2011). Based on

this, Ducrest and coauthors (2008) proposed that darker eumelanotic individuals should be less sensitive to stressful factors and have better anti-inflammatory, antipyretic and anti-oxidative responses than lighter individuals. These predictions need to be tested with special attention directed to ecological, social and physiological factors that influence and maintain inter-individual variation in melanin-based coloration and associated phenotypic traits.

### I.3. Aims of the thesis

Some feather characteristics have been used in ecological research for a long time, e.g. ptilochronology (the study of growth rates of feathers by the measuring of growth bars (Grubb 1989)) has been used in the field of nutritional ecology for several decades. Interest in feather coloration dates back even further to Darwin and Wallace, with the initial focus on evolution and function of feather coloration (Hill & McGraw 2006). Although thoroughly studied, morphological and color characteristics can still provide novel information about organism's endogenous and exogenous processes in conjunction with the advances in methods used in ecological physiology and with emergence of new fields, such as immunoecology and psychoneuroimmunology. One of such new methods is measuring glucocorticoid hormones from feathers which I will employ in my thesis in collaboration with feather size, integrity and color parameters.

Three papers of this thesis address the question about effects of experimental manipulation of physiology of greenfinches on different parameters of feathers. In the first study, I asked whether a single administration of synthetic glucocorticoid, dexamethasone (DEX) and induction of inflammatory response by administration of plant toxin (phytohaemagglutinin, PHA) affects feather growth in adult birds (**Paper I**). Elevation of glucocorticoid hormone levels is an integral part of stress response (as well as its termination) and immunomodulation (Sapolsky *et al.* 2000; Perretti & D'Acquisto 2009). These hormones are also responsible for mobilizing energy stores by stimulation of gluconeogenesis and inhibition of protein synthesis (Ramage-Healey & Romero 2000). Elevation of glucocorticoids is thus incompatible with other protein-demanding processes like molt. Previous studies have shown that chronic elevation of GC hormones suppresses feather growth (Romero *et al.* 2005; Strohlic & Romero 2008; DesRochers *et al.* 2009). Here I asked whether similar effect would also occur in the case of acute GC elevation and induction of an inflammatory response by foreign antigen and whether the potential somatic cost of elevated CORT levels is of comparable magnitude with the one induced by the immune system activation. The latter question is interesting because both immune and stress responses deplete somatic resources; however, the question about which of those is more costly has never been directly assessed. I measured three different parameters of feather size in order to find

out which of these are most sensitive to treatments, and hence most indicative of physiological insults experienced during feather growth. This information is of methodological interest for studies aiming to quantify somatic impacts of physiological manipulations in birds.

To add further ecological dimension to the study, I asked whether supplementation of dietary carotenoids affects the expenditures imposed by immune activation and elevated CORT levels. Carotenoids are dietary pigments with proposed immunomodulatory and antioxidative properties (Chew & Park 2004). Few previous studies have shown that carotenoid supplementation can attenuate body mass loss imposed by immune challenge (Hörak *et al.* 2006; Pap *et al.* 2009) and that CORT administration can interfere with carotenoid sequestration to integument (Loiseau *et al.* 2008; Cote *et al.* 2010, but see Costantini *et al.* 2008). Therefore, I tested whether carotenoid supplementation will positively affect feather growth of birds receiving immune activation and GC challenges.

In the second study, I tested the effects of immune activation and suppression on feather CORT content (**Paper II**). Until recently (Meitern *et al.* 2013), there was no information about whether and how activation or suppression of immune system affects feather CORT levels and how feather CORT relates to other physiological measures of stress, such as differential leukocyte counts and an individual's ability to mount immune responses. Explaining these issues is important for interpretation and understanding the information content of feather CORT levels in studies using this marker for characterizing stress history of individuals and populations. For instance, both the acute phase response and avian stress response share some of the same neuroendocrine control mechanisms such as the HPA-axis (Owen-Ashley & Wingfield 2007) and inflammatory agents can be considered as 'immunological stressors' (e.g. Laugero & Moberg 2000; Xie *et al.* 2000). Because inflammatory responses are usually accompanied by the rise in blood CORT levels (Sapolsky *et al.* 2000), I was interested in testing whether induction of inflammation will affect deposition of CORT in feathers.

In the **Paper III** I continued with measuring the effects of immune system manipulation on feather CORT by asking whether experimental infection with a natural pathogen (coccidian intestinal parasites) affects feather CORT levels. Additionally I asked whether and how are feather CORT levels of greenfinches related to resistance to experimental infection and the measure of captivity-intolerance, i.e., the amount of damage accrued to tail feathers in captivity. Extent of tail damage is expected to reflect captivity-intolerance in greenfinches, because this trait correlates with the frequency of flapping bouts against cage walls (Sild *et al.* 2011). I therefore predicted that the extent of tail damage accrued in captivity correlates positively with feather CORT content as indicator of susceptibility to stress.

In the **Paper IV** I tested whether feather CORT levels are indicative of personalities of greenfinches assuming the existence of a "fearfulness

syndrome”, based on expected covariation of behavioral responses to different fear-eliciting situations. For this purpose I recorded individual variation and consistency of five behavioral traits that were presumably indicative of responses to fear-eliciting and stressful situations. Because individual differences in behavioral responses to stressful situations are at least partly mediated by differences in neuroendocrine activity (Koolhaas *et al.* 1999; Carere *et al.* 2003; Kralj-Fiser *et al.* 2007), I expected to discover a covariation between the amount of CORT deposited into feathers grown in captivity and at least some of the behavioral measures of fearfulness.

In the **Paper V** I studied whether feather size and color parameters predict survival in greenfinches facing epidemic of emerging infectious disease, finch trichomonosis. Plumage characteristics have been long thought to reveal individual’s ability to deal successfully with infections (Hamilton & Zuk 1982; Lozano 1994; Moreno & Møller 2006; Simons *et al.* 2012). However, despite the relatively well-documented evidence about covariation between feather characteristics and survival (see Jennions *et al.* 2001; Roulin & Altwegg 2007; Saino *et al.* 2013), the number of studies documenting direct associations between plumage characteristics and infection-induced mortality or morbidity has remained low. Here I report about associations between plumage characteristics and trichomonosis-induced mortality in wild-caught greenfinches in captivity. Trichomonosis is caused by the flagellate protozoan parasite *Trichomonas gallinae*, which is well known historically in pigeons and raptors and can cause high mortality. The parasite inhabits the upper digestive tract of the bird and spreads through saliva. Spread of trichomonosis since 2005 has coincided with extensive reductions in breeding passerine (especially greenfinch) populations in Great Britain and Northern Europe. In 2013, 11 % of female wild-caught greenfinches brought into aviary died of trichomonosis, which provided a unique opportunity to measure survival selection in relation to feather parameters. I first asked whether the variation in parameters of feather size that were sensitive to glucocorticoid administration and immune system activation predict survival of epidemics. In addition, I tested whether survival could be predicted on the basis of the occurrence of fault bars and eumelanin-based coloration of black feather tips.



## 2. METHODS

Using wild-caught captive greenfinches (*Carduelis chloris*) in research of physiological ecology has many advantages. First, they tolerate captivity stress well (Sepp *et al.* 2010) and are easy to maintain in laboratory conditions. Second, their biology is well known as greenfinches have been widely used as a model species in ecological studies (Lilliendahl 2000; Lindström *et al.* 2001; Peters *et al.* 2008; Samas *et al.* 2012; Minias & Iciek 2013). Third, they are naturally infected with coccidian parasites that are relatively easy to trace and manipulate.

Greenfinches are medium-sized (ca. 28 g) sexually dichromatic gregarious seed-eating passerines native to the western Palearctic region. Males are larger and more colorful than females, with older males developing olive-green plumage on the back, bright or greenish yellow color on the breast, and striking bright yellow markings on the primaries, primary coverts and sides of the tail feathers (Cramp & Perrins 1994). Females are more olive-brown and yellowish-buff, having faint brown streaks on back and lacking full yellow tints in their plumage.

All the birds (N = 45–93) participating in our studies were captured in mist-nets at bird feeders either in the Sõrve Birds Station in Saaremaa or in a garden in the city of Tartu in December or January and released in early March. The birds were housed indoors in individual cages (27 × 51 × 55 cm) with sand-covered floors. The birds were supplied *ad libitum* with sunflower seeds and tap water. Birds were held on the natural day-length cycle on artificial lighting by luminophore tubes.

Feathers of birds have the distinct advantage of being periodically replaced in a predictable fashion and time scale that allows for repeated assessment of developmental history of birds (Bortolotti *et al.* 2002). Another useful property of feathers is that they provide information about organism's physiology in a non-invasive way, for example as compared to blood sampling or killing of birds. In addition, feather size, integrity and color are easy to measure. I used feather mass, length and rachis diameter as a proxy to feather quality (**Papers I, V**). Direct measures of feather strength, durability or aerodynamic properties are more difficult to obtain and may be less reliably measured (than using high-resolution scale, simple ruler or spessimeter) (De La Hera *et al.* 2009). Several studies have also shown that rachis width and feather mass explain a large part of variation in mechanical performance of feathers within populations (Dawson *et al.* 2000; De La Hera *et al.* 2009). Feather integrity (**Papers III, IV**) and fault bars (**Paper V**) are directly observable and assessable; the color of feathers can be measured spectrophotometrically (**Paper V**).

Mortality by infectious diseases enables to observe natural selection in action, as well as consequences of these selection pressures on the appearance of birds. Differential survival of infected birds is also potentially informative about how coloration and immune processes are interconnected and how feather

coloration has evolved (**Paper V**). Many infectious diseases, however, are not highly virulent and hosts are able to tolerate chronic infections. One of such infections is coccidiosis, which is caused by obligate intracellular parasite of the genus *Isospora* that infect a number of songbird species (Greiner 2008). All greenfinches that have been caught and examined in our laboratory have been naturally chronically infected with coccidians. Coccidiosis of wild birds is becoming a popular model of parasite-mediated selection that enables to avoid assessment of immune function on the basis of responses to artificial antigens that may not appear optimal predictors of infection resistance or disease outcome (Adamo 2004; Saks *et al.* 2006). High prevalence of coccidiosis in our research model enabled us to test whether chronic infections relate to feather CORT content and how feather CORT levels are related to resistance to experimental infection (**Paper III**).

In order to find out more about information content of feather CORT levels, I tested the hypothesis that feather CORT levels reflect variation in avian personalities (**Paper IV**). This hypothesis relies on the assumption that feather CORT is indicative of consistent between-individual differences in responsiveness to fear-eliciting situations. To test this hypothesis I had first to quantify behavioral responses to different stimuli and test whether between-individual differences in such responses are repeatable in time. The variables that included behavioral responses to fear-eliciting stimuli were propensity to give distress calls at handling, latency to feed at the presence of a predator image, and changes in locomotor activity in response to distress calls of conspecifics. Additionally, I measured the frequency of flapping flight movements against cage bars and the amount of damage accrued to tail feathers in captivity as a measure of captivity-intolerance. The next step was to test whether and to what extent the measured behavioral traits and feather CORT covary and if this covariation reveals the existence of general “fearfulness syndrome”. This was analysed by structural equation modelling (SEM).

## 3. RESULTS AND DISCUSSION

### 3.1. How do immune activation and glucocorticoid administration affect feather growth?

Immune defenses are considered extremely complicated and resource demanding organismal functions (Lochmiller & Deerenberg 2000; Demas 2004; Schulenburg *et al.* 2009). Activated immune responses impose costs, which eventually can lead to physiological trade-offs in allocation of resources between immunity and other vital functions. Glucocorticoid hormones are important down-regulators and terminators of immune (Perretti & D'Acquisto 2009) and also stress responses (Sapolsky *et al.* 2000). Rise in baseline, as well as in stress-induced levels of CORT has been shown to suppress feather growth (Strochlic & Romero 2008; Müller *et al.* 2009). This occurs via inhibition of protein synthesis and mobilization of endogenous glucose stores (Remage-Healey & Romero 2000). Quality of feathers can in turn affect thermoregulation, flight (Butler *et al.* 2008), resistance to abrasion, rigidity of feathers (Dawson *et al.* 2000), take-off speed and maneuverability (Swaddle *et al.* 1996; Matyjasiak *et al.* 2004) with potential effects on fitness. In the **Paper I**, I asked whether a single, short-term elevation of CORT levels can also affect feather growth in adult birds. To put this information into wider ecophysiological context, I tested whether such potential somatic cost is of comparable magnitude with the one induced by immune system activation. Both immune and stress responses deplete somatic resources; however, the question about which of those is more expensive has never been directly assessed. To add further ecological dimension to the study, I also asked whether supplementation of dietary carotenoids affects the possible somatic cost imposed by immune activation and immune suppression. Carotenoid-based signals have been extensively studied by animal ecologists since recognition of their possible role in health maintenance and signaling. Carotenoids can modulate immune function (Chew & Park 2004) and may also act as antioxidants (Krinsky 1989, but see Costantini & Møller 2008). So far, few studies have shown that carotenoid supplementation can attenuate body mass loss imposed by immune challenge (Hörak *et al.* 2006; Pap *et al.* 2009; Sepp *et al.* 2011) and that CORT administration can interfere with carotenoid sequestration to integument (Loiseau *et al.* 2008; Cote *et al.* 2010).

93 male greenfinches were assigned to groups of immune activation and glucocorticoid administration in a factorial manner. The experiment involved induction of immune response by subcutaneous injection of phytohaemagglutinin (PHA), a plant lectin known to induce localized swelling response which is used as an indicator of strength of immune response (Tella *et al.* 2008). Glucocorticoid administration consisted of a single injection of synthetic glucocorticoid dexamethasone (DEX). DEX is a CORT agonist which, binding to the same receptors, decrease endogenous CORT production via

negative feedback on the hypothalamic-pituitary cascade (reviewed by Holberton *et al.* 2007). DEX injections are routinely used for immune suppression (Huff *et al.* 1999) and manipulation of CORT levels (Rich & Romero 2005). Carotenoid supplementation consisted of birds receiving high and low doses of carotenoids or water. Upon arrival to the laboratory one tail feather (rectrix) from all the birds was removed. Feather mass, length and rachis diameter of wild-grown as well as replacement feathers were recorded.

This study clearly showed that a single boost of glucocorticoid hormone is sufficient to restrict feather growth of wild birds, despite the benign conditions and *ad libitum* access to food in captivity. This raises the possibility that experiencing an acute stress or termination of immune responses by elevation of glucocorticoid hormones may induce long-term somatic costs with a potential impact on fitness. DEX injection significantly reduced all three measures of feather size. The most probable way how DEX could affect feather growth is through its property of suppressing protein synthesis (Sapolsky *et al.* 2000). This indicates that feather size could contain information about the ability of its bearer to avoid stressful situations or block the catabolic effects of CORT during stress. As DEX injection suppressed feather growth, it has practical implications for studies using DEX to study stress physiology, where possible somatic costs, accompanying the physiological effects of DEX, need to be considered. In further studies it would be most interesting to evaluate whether a natural downregulation of HPA-axis by negative feedback mechanism can also induce similar somatic costs to an exogenous glucocorticoid administration.

Immunostimulation by PHA reduced feather length, but did not affect feather mass or rachis diameter. This implies, that in face of resource limitation, investment into feather mass (which increases feather strength) is prioritized over investment into feather length. This study adds another piece of evidence to what we know about somatic costs of induced immune responses in terms of feather growth. Importantly, this study also exemplifies the value of measuring several parameters of feather size. For example, I would not have detected the impact of immune activation by measuring only feather mass.

Previous studies in greenfinches have shown that carotenoid supplementation can incline birds to fattening and alleviate body mass loss induced by immune challenge with sheep erythrocytes (Hörak *et al.* 2006) or PHA (Sepp *et al.* 2011). Current experiment showed that dietary carotenoids had no effect on feather growth, suggesting that any anabolic effects of carotenoids could involve lipid rather than protein metabolism. Altogether these results demonstrate the sensitivity of feather growth to manipulations of immune and adrenal functions. These findings reinforce the potential utility of feather parameters for assessment of somatic impacts of physiological manipulations in birds.

### 3.2. Synthetic glucocorticoid inhibits CORT deposition into feathers

Assessment of individual well-being on the basis of glucocorticoid hormone levels has become a standard tool in ecological, physiological and conservation studies of wild animals. In birds, CORT is the primary glucocorticoid hormone that has been linked to the promotion of an emergency life-history stage (Wingfield & Sapolsky 2003), and which stimulates behaviors and physiological processes aimed at recovering from disruption of homeostasis caused by external stressors (reviewed by Johnstone *et al.* 2012). Feather CORT content is thus a potent source of information relating to activation of the HPA-axis during feather growth which can be used for assessment of well-being and stress history of individuals. However, despite increasing popularity of feather CORT measurement in avian studies, the research aimed at understanding the processes underlying CORT deposition into feathers and their interconnections to other physiological processes is still only in its infancy.

My objective in the **Paper II** was to study different factors affecting CORT deposition into feathers and how feather CORT covaries with other markers of stress imposed upon individuals during feather growth. I used the same dataset as in the **Paper I**, consisting of 93 male greenfinches that were assigned to treatment groups in a factorial manner. To mimic acute stress half of the birds were injected with DEX, half with PHA to induce inflammation. Upon arrival to the laboratory one tail feather from all the birds was removed. CORT levels of wild-grown as well as replacement feathers grown in the captivity were recorded. Birds were blood sampled three times during the period of feather growth, which enabled to test whether leukocytic markers of stress (concentrations of total leukocytes, heterophils, lymphocytes and heterophil/lymphocyte (H/L) ratio) correlate with feather CORT levels. Heterophils and lymphocytes comprise the majority of circulating immune cells in birds and H/L ratios are specifically sensitive to either natural stressors or administration of stress hormones, so that relatively high heterophil counts in relation to lymphocytes reliably indicate high glucocorticoid levels (Ots *et al.* 1998; Davis *et al.* 2008). Assessment of swelling response to PHA enabled to test whether feather CORT levels relate to the magnitude of cutaneous inflammatory response to the antigen.

From the two experimental treatments applied, only DEX injection had significant effect on feather CORT levels. DEX-injected birds had on average 37% lower level of CORT in their feathers than saline-injected birds. Because the physiological effects of DEX are mediated by glucocorticoid receptors and DEX, like endogenous glucocorticoids, ultimately reduces CORT concentrations by forming the negative feedback loop that controls CORT secretion (Bennett *et al.* 2013), the DEX-induced down-regulation of circulating CORT levels provides the most parsimonious explanation for the observed pattern. This finding provides novel evidence that feather CORT levels are

sensitive to manipulation of hormonal balance of birds, thereby supporting the diagnostic value of feather CORT measurements. This result compares favorably with those of Lattin and coauthors (2011) who used CORT implants to increase feather CORT levels and with a recent study by Fairhurst and coauthors (2013a) showing that feather CORT correlated with circulating CORT levels in some (but not all) stages of feather growth.

Absence of the effect of PHA injection on feather CORT is noteworthy as inflammatory responses generally increase circulating CORT levels (Sapolsky *et al.* 2000; Owen-Ashley & Wingfield 2007). Therefore, I expected that PHA-injected birds deposit more CORT into their feathers than saline-injected birds. One possible explanation is that PHA injection induced negative feedback loop that would down-regulate inflammation-induced surge of CORT, so that the eventual amount of CORT secreted during feather growth would equal that of control group. Alternative explanation would be that although PHA injection caused mild stress, increase in chronic stress level and CORT concentration was not severe enough to increase feather CORT. Cutaneous swelling response to PHA and feather CORT levels did not correlate either. This suggests that measuring feather CORT would give information about more severely increased chronic stress level and would appear a relatively “noise-free” method, not affected by smaller stressors.

Surprisingly there were no significant correlations between feather CORT levels and circulating leukocyte concentrations, measured at three stages of feather growth. Assessment of stress on the basis of H/L ratio is considered as an indicator of stress over longer period of time than on the basis of glucocorticoid hormone levels in the blood (Davis *et al.* 2008; Müller *et al.* 2010). Therefore I expected that birds with high heterophil haemoconcentrations and H/L ratios deposit more CORT into feathers grown in captivity, especially given that PHA injection in the current experiment resulted in elevated heterophil haemoconcentration and H/L ratios for at least 30 days (Sarv & Hõrak 2009). Absence of such correlations indicates that circulating leukocyte counts and feather CORT levels convey differential information about HPA-axis activation in birds. Hence, these markers cannot substitute each other, but appear rather complimentary indicators of stress status of birds (Davis *et al.* 2008; Müller *et al.* 2010). In conclusion, this study suggest that assessment of health state and stress history of birds on the basis of feather CORT levels appears more complicated than previously thought and calls for further research on information content of feather CORT.

### 3.3. High feather CORT indicates better resistance to experimental coccidian infection

Even when exposed to the same parasites and pathogens, some individuals manage to cope with their infections better than others. The questions about the causes and consequences of these processes have remained largely unclear, particularly as regards wild animals and their natural pathogens. Differential exposure or sensitivity to stressors may have substantial effects on the variation in immune responsiveness of animals. Furthermore, there is a reason to expect that interrelationships between stress and immune responses relate to animal personalities or coping styles (Seegerstrom 2007; Koolhaas 2008). Proactive coping styles tend to be coupled with efforts to prevent or manipulate stressors, high sympathetic nervous system activity and a Th1 biased (inflammatory) immune system. Reactive coping styles entail high HPA-axis reactivity and a Th2 biased (humoral) immune responses (Koolhaas 2008; Martin 2009, but see Martins *et al.* 2007). One might thus expect that individual differences in disease resistance relate to behavioral phenotypes. In the **Paper III**, I ask whether infection with a natural pathogen affects feather CORT levels and how a potential marker of stress responsiveness, the feather CORT content, reflects the resistance to an experimental infection with natural coccidian parasites. To study the link with behavior, I measured the captivity-intolerance (as a potential marker of individual susceptibility to stress on the basis of the extent of tail damage accrued in captivity) of birds and predicted that it will correlate positively with feather CORT content.

All 56 male greenfinches that were involved in the study appeared naturally infected with isosporan coccidians at capture. The birds were subsequently divided to control group and experimental infection group. At the beginning of the experiment all birds were subjected to anticoccidian medication with Toltrazuril in order to standardize their infection status. Toltrazuril specifically reacts to apicoplast of coccidians, which is unique to the Apicomplexan parasites (see Sepp *et al.* 2012). Subsequently, infection group was infected orally with sporulated heterologous coccidian oocysts. At the same time control group received another round of medication in order to increase differences in infection intensity between infected and control birds. In order to monitor the course of infection, fecal samples were collected at different time points of the study. Upon arrival to the laboratory one tail feather from all the birds was removed and CORT levels of wild-grown as well as replacement captivity-grown feathers were recorded. Before the birds were released their tail feathers were photographed under standard conditions in order to assess tail damage.

As expected, I found that extent of tail damage accrued in captivity correlated positively with captivity-grown feather CORT content. The most likely explanation for this pattern would be that some greenfinches, i.e., those with the most damaged tails, possessed more stress-prone personalities than others and that these differences were mirrored in feather CORT content. This

explanation is consistent with finding of Sild and coauthors (2011), showing that the propensity to perform flapping flight bouts against cage walls (which correlated with tail damage scores) was individually consistent over 57 days; qualifying thus as a personality trait according to Reale and others (2007).

This study demonstrated that coccidian infection (or its suppression) is associated with CORT secretion in wild bird species. Birds subjected to experimental infection had on average 23% higher CORT content in their captivity-grown tail feathers than sham-inoculated individuals. Unexpectedly, the pre-infection medication had long-term effects, so that post-infection levels of oocyst shedding among experimental birds never reached pre-experimental levels. Thus, I cannot distinguish whether differences in feather CORT between infected and un-infected birds were primarily caused by enhanced CORT secretion due to novel infection or due to reduction of CORT levels among control birds that were relieved from chronic infection.

The most surprising result of this study was positive correlation between feather CORT content and measures of resistance to an experimental infection. Experimental birds with lowest feather CORT levels developed highest infection intensities at peak infection phase as well as highest increase in average infection levels. The most likely explanation for such pattern is that birds that invest more into fighting novel coccidian infections have higher stress responsiveness or that birds that experience captivity as more stressful have higher resistance to novel infections. On the basis of current data these explanations are difficult to discern. Interpretation of these findings is further complicated by conjunctions between chronic and acute stress responses and CORT secretion. Captivity can cause both acute and chronic stress; in turn, chronic stress can suppress baseline and acute-stress-induced CORT levels in birds (Rich & Romero 2005; Cyr & Romero 2008; Cyr *et al.* 2009). Hence, it would be difficult to distinguish whether elevated CORT levels in more resistant birds reflect unhealthy chronic stress or a healthy coping response (Cyr & Romero 2007). Assuming that feather CORT is affected more by stress-induced stimulation than baseline levels of CORT (Bortolotti *et al.* 2008), our results would support an idea that sustaining higher stress responsiveness in captivity is beneficial. Under this scenario, birds that were experiencing less stressful events, or perceiving their environment least stressful, appeared most susceptible to novel infection with heterologous parasite strain. Finally, I cannot exclude the possibility that high feather CORT levels could primarily reflect the importance of this hormone as a mediator of energy-allocation processes, rather than stress responsiveness *per se*. Thus, high feather CORT among the most resistant individuals could have reflected high energetic demand/expenditure associated with efficient immune responses within the normal stress reactive scope (see (Romero *et al.* 2009).

Coccidians are the most prevalent avian parasites (Greiner 2008) with well documented pathological effects (reviewed by Pap *et al.* 2011; Sepp *et al.* 2012), so the benefits of increased resistance in the wild is obvious. Findings of



this study suggests that further research on CORT levels in relation to resistance to coccidiosis might increase our understanding about the mechanisms linking individual variation in CORT levels to fitness (Breuner *et al.* 2008; Angelier *et al.* 2009).

### 3.4. Is feather CORT indicative of avian personalities?

Animal behavior is not completely flexible and different behavioral traits do not vary independently of each other (reviewed by Sih *et al.* 2004; Carere *et al.* 2010). Research on consistent between-individual differences in suites of correlated traits, known as behavioral syndromes or personalities is becoming increasingly popular in behavioral ecology (Sih *et al.* 2004; Dingemanse & Réale 2005; Careau *et al.* 2008).

Individual differences in behavior are particularly apparent when individuals face mildly stressful situations (Koolhaas *et al.* 1999). Challenging conditions, for example the presence of a predator in the environment, seem to induce correlations between behavioral traits (Bell & Sih 2007; Dingemanse *et al.* 2007). Fear is prompted by imminent and real danger, and galvanizes active defensive responses (Davis *et al.* 2010). Fearfulness can thus be defined as a basic psychological characteristic of the individual that predisposes it to perceive and react in a similar manner to a wide range of potentially frightening events (Boissy & Bouissou 1995).

Individual differences in behavioral responses to stressful situations are at least partly mediated by differences in neuroendocrine activity (Koolhaas *et al.* 1999; Carere *et al.* 2003; Kralj-Fiser *et al.* 2007). Individuals with proactive, risk-prone, aggressive and bold personalities respond to stress with a strong sympathetic activation and increase in noradrenergic stimulation. In contrast, individuals with a passive, risk-averse, non-aggressive, shy personality respond to stress with a strong HPA stimulation and a consequent increase in circulating glucocorticoids (Carere *et al.* 2010, but see Kralj-Fiser *et al.* 2010 and Koolhaas *et al.* 2011 for opposite results). Such patterns indicate that individual variation in stress responsiveness is linked to behavioral syndromes (Carere *et al.* 2010; Garamszegi *et al.* 2012). In this context, the covariation between stress hormone secretion and behavioral fear responses is of particular interest. Aim of the **Paper IV** was (1) to test for the presence of “fearfulness syndrome” in captive greenfinches and (2) to determine whether stress responsiveness (or susceptibility) of individuals, assessed on the basis of feather CORT is a part of this syndrome.

In 45 male greenfinches, altogether five behavioral traits were recorded that were expected to contribute to the general “fearfulness syndrome” on the basis of previous observations in greenfinches (Sild *et al.* 2011) and great tits (*Parus major*) (Krams *et al.* 2013). CORT levels of wild-grown and replacement tail feathers grown in captivity during the study were measured. Measured

behavioral variables included latency to feed at the presence of a predator image, the propensity to give distress calls at handling, changes in locomotor activity in response to distress calls of conspecifics, frequency of flapping flight movements against cage bars and tendency to damage tail feathers in captivity. Associations between these traits and repeatability in time were assessed in correlation analyses. The presence of a behavioral syndrome was tested by structural equation modelling (SEM).

All behavioral traits were repeatable in time. In an unprovoked situation, without any outside disturbance, flapping frequency against cage bars correlated with damage accrued to tail feathers. This confirms the findings of Sild and coauthors (2011) and indicates that in this model system, damage to tail feathers describes the behavior of birds in captivity and can be used as a proxy of captivity tolerance.

None of the behavioral parameters correlated with feather CORT levels. This result was unexpected but compares favorably to a study of collared flycatchers (*Ficedula hypoleuca*), where individuals with higher levels of fecal CORT metabolites did not show consistently higher or lower behavioral scores along the shy/bold spectrum, and where environmental challenges owing to the presence of a novel object, territorial intruder and a potential predator caused no elevation in the level of fecal CORT metabolites (Garamszegi *et al.* 2012). The authors of this study emphasise that the relationship between CORT levels and behavior, if it exists at all, is likely to be of small magnitude, a conclusion that is also supported by our data. Generality of this conclusion, however, awaits for further confirmation because feather CORT levels correlated with problem-solving performance in a recent study of house sparrows (*Passer domesticus*) (Bókony *et al.* 2014). It is also possible that feather CORT levels did not correlate with behavioral measures of fear because the behaviors are taking place on a different time scale than the growth of captivity-grown feathers. For instance, Fairhurst and coauthors (2013a) recently showed that feather CORT correlated with blood hormone levels only in some days of feather growth in nestling tree swallows (*Tachycineta bicolor*).

I was particularly surprised about the absence of correlation between CORT deposited to captivity-grown feathers and flapping frequency. This finding contradicts that of the previous study (**Paper III**) where a positive association between feather CORT and tail damage was detected. This contradiction between the findings of two studies in the identical model system is difficult to explain. One might argue that exposure of birds to different potentially fear-eliciting stimuli in the current study could have obscured the relationship between HPA-axis activity and tail damage that appeared in the previous study, where such stimuli were not present. On the other hand, we also failed to find any evidence that behavioral responses to fear-eliciting stimuli had correlated with feather CORT or tail damage scores. A conclusion from this part of the study is that tail damage does not appear a universal correlate of stress responsiveness in captive greenfinches.

None of the measured behavioral parameters correlated with each other. This is an intriguing result, considering that all of these traits (latency to feed in a presence of a predator/novel object, propensity to give distress calls, response to stress calls of conspecifics, flapping frequency in cages) can be intuitively associated with fearfulness or anxiety. However, no “fearfulness syndrome” emerged from our data, as shown also in the SEM analysis. None of the a priori composed hypothesis about causal relationships between measured traits proved to be better than the null model assuming independence of traits. This indicates that responses to different fear eliciting situations might appear more uncoupled than expected and could be condition- and context-dependent. At the same time, all the behavioral variables showed individual consistency in time, which hints that these traits might reflect some aspect of personality (Réale *et al.* 2007; Dingemanse & Wolf 2010).

According to Boissy and Boissou (1995), consistency of individual responses to different fear-eliciting situations reflects the existence of an underlying psychological profile, currently labelled fearfulness. However, the question of fear as a single variable has been challenged by the recognition of its multidimensionality: very diverse events may generate fear responses, and it is likely that these events are not perceived equally by animals in diverse internal states (Boissy 1995; Saint-Dizier *et al.* 2008). The findings of this study challenge the concept of a single internal variable responsible for fearfulness and support the proposed multidimensional nature of fear responses.

### **3.5. Emerging infectious disease selects for darker plumage coloration**

Parasites exert persistent selection pressure on all organisms. Hosts differ in their vulnerability to infections and these differences could be mirrored in individual’s appearance as immune system is integrated with almost any physiological function of an organism. Color and morphology of feathers provide ample cues about the physiological condition of individual birds during feather growth. For instance, most common feather pigments – melanins and carotenoids – can be indicative of the individual’s developmental and hormonal milieu, their ability to find and process specific micronutrients and of immunological and oxidative challenges experienced during feather growth (reviewed by Hill & McGraw 2006). Additional information about the individual’s state during feather growth is provided by the physical properties of the feathers such as different parameters of size and structure (e.g. DesRochers *et al.* 2009, **Paper I**). All of these parameters have a potential to affect organism’s fitness. However, despite the relatively well-documented evidence about covariation between feather characteristics and survival (Jennions *et al.* 2001; Roulin & Altwegg 2007), the number of studies documenting direct associations between plumage characteristics and infection-induced mortality or

morbidity is limited. The spread of trichomonosis epidemics from Great Britain over Fennoscandia to Estonia provided an excellent opportunity to study associations between plumage characteristics and trichomonosis-induced mortality. The protozoan parasite *Trichomonas gallinae* causes necrotic inflammation of the oropharynx that eventually inhibits swallowing and leads to starvation and dehydration (Atkinson *et al.* 2008). Given the extreme pathogenicity and recent emergence of trichomonosis in greenfinches, it has been suggested that infection may have been recently spilled over from another host species, leaving little or no time for host adaptations to evolve (Neimanis *et al.* 2010).

The aim of the **Paper V** was to establish a connection between plumage characteristics of greenfinches and mortality caused by trichomonosis. Therefore melanin-based black plumage patches and three parameters of feather size (mass, length and rachis diameter) of greenfinch's tail feathers were measured. Additionally I was interested if the occurrence of fault bars can predict mortality as fault bars have been associated with increased mortality in different bird species (Bortolotti *et al.* 2002; Møller *et al.* 2009).

Out of 46 wild female greenfinches, five birds died during the first 11 days in captivity. Three of them displayed clear symptoms of trichomonosis (ingluvitis), while the remaining two exhibited similar pattern of total depletion of bodily reserves. Feathers of survivors were on average 22 % darker than feathers of dead birds. There are two alternative explanations why birds with higher plumage reflectance were selected against.

First explanation involves covariation between melanin-based coloration and some relevant aspect of immune function. Accumulating evidence indicates that melanin-based coloration is often correlated with other physiological, morphological and behavioral traits as an outcome of pleiotropic effects of key regulators of melanogenesis (Ducrest *et al.* 2008). Melanocortin peptide hormones, that trigger the production of melanin pigments (by binding to MC1R), also bind to four other melanocortin receptors, causing the reduction of acute, allergic and systemic inflammation and septic shock (Catania *et al.* ; Ducrest *et al.* 2008). In addition, activation of MC1R causes a collective reduction in the pro-inflammatory cytokines and prostaglandins and an increase in the anti-inflammatory mediators involved in the inflammatory process (Gangoso *et al.* 2011). Under such mechanism, one would predict enhanced inflammatory responses in paler, less-eumelanotic individuals (which has been established in at least one avian species (Gangoso *et al.* 2011, but see Vinkler *et al.* 2014). Inflammation, leading to terminal septicemia is a typical hallmark of *Trichomonas gallinae* infection (Neimanis *et al.* 2010), so one might hypothesize that paler greenfinches (i.e. the ones with suppressed expression of genes involved in eumelanin production) were more likely to die due to their over-responsive inflammatory defenses.

Another possible mechanism relates to pleiotropic covariation between melanin production and stress susceptibility. Association between melanin-

based pigmentation and behavioral-physiological aspects of animal personality seems to be widespread in vertebrates, with more distinctly pigmented individuals typically being more proactive, stress-resistant, and socially dominant (Kittilsen *et al.* 2009; Kittilsen *et al.* 2012). It is possible that birds with paler eumelanotic tail coloration appear to be more susceptible to stress and consequently stress-induced immune suppression would turn pale individuals more susceptible to trichomonosis. Higher stress-susceptibility of pale greenfinches is consistent with significantly paler plumage coloration among birds with fault bars.

Prevalence of fault bars was five times higher among the feathers of dead birds than among survivors. In addition, feathers which contained fault bars were significantly paler. Fault bars are developmental irregularities in feathers that result from a variety of stressors that occur during feather growth (Bortolotti *et al.* 2002; Møller *et al.* 2009). High prevalence of fault bars in dead birds thus suggests that those individuals either experienced more stressors or appeared more susceptible to stress already in the preceding autumn when those feathers were growing. Therefore, it seems unlikely that trichomonosis caused fault bars in finches, but instead, birds that were more stressed during preceding autumn molt appear to be more susceptible to trichomonosis in winter. The finding about associations between occurrence of fault bars and mortality is consistent with those of two previous studies. In female American kestrels (*Falco sparverius*), high number of fault bars was associated with lower survival probabilities (Bortolotti *et al.* 2002). More specifically, Møller and colleagues (2009) demonstrated that prey items of the goshawk (*Accipiter gentilis*) had nearly three times higher occurrence of fault bars than the general population of different prey species. The suggested mechanism involves decrease of aerial performance of prey due to breakage of feathers at fault bars. My results corroborate the diagnostic value of fault bars in predicting mortality as found in two above-cited studies. However, they also show that reduction of aerial performance is not the single mechanism connecting occurrence of fault bars with mortality. This study also shows that fault bars and feather size parameters convey different information about the stressful experiences during feather growth as feather size parameters did not differ between dead birds and survivors.

Current study is the first one to detect trichomonosis-induced viability selection on plumage characteristics. Detection of extremely strong selection for lower plumage reflectance is particularly interesting. Assuming non-zero heritability for tail feather coloration, one would predict that the recent outbreak of trichomonosis leads to evolution of darker melanin-based coloration in greenfinches.

## SUMMARY

There are many characteristics that make birds perfect model organisms for evolutionary ecology research. Among the most notable of these are feathers that are considered the most complex integumentary structures of vertebrates (Prum & Williamson 2002). The interplay of feather growth and pigmentation produces tremendous diversity of within- and across-taxa coloration patterns, from elaborated sexual ornaments to exceptional camouflage matching (Landeén & Badyaev 2012). Not less importantly for ecologists, coloration, size, integrity and hormone content of feathers provide information about specific endogenous or exogenous processes and events of interest in the life of individuals or ecology of species, as well as the environment from which they came (Bortolotti 2010). The full potential of this approach has been realized only recently.

In this thesis, I relied on experiments and correlational data on wild-caught captive greenfinches to explore the applicability of different feather parameters for immunoecological and behavioral research. The first three papers are experimental. They aim at explaining whether and how artificial and natural immune challenge and manipulation of hormonal milieu by administration of synthetic stress hormone dexamethasone affect feather growth and corticosterone (CORT) content. However, in addition to specifying diagnostic value of feather parameters, these experiments also provided valuable information about somatic costs and correlates of up- and down-regulation of immunity and stress responses, which belong to the central research questions of physiological ecology. Remaining two papers are correlational. One of these asks whether feather CORT content provides any information about fear-related personality traits. The final paper takes an advantage of an outbreak of emerging infectious disease – trichomonosis – that enabled me to study associations between feather parameters and infection-induced mortality under controlled laboratory conditions.

One of the main aims of my thesis was to test the suitability of newly developed feather CORT measurement method for studying the nexus between neural, endocrine and immunological functions. Despite increasing popularity of integrated measures of glucocorticoid hormones, the potential for feather CORT measurements to advance avian ecology has been hindered by a lack of experimental research into what CORT values represent physiologically (Fairhurst *et al.* 2013b). For example, we still do not exactly know what impact frequently encountered physiological (immune activation or rise in blood CORT levels) or psychological (encounters with predators) stressors have on feather CORT content. Therefore, I tested the effects of immune activation and glucocorticoid administration on feather CORT content. In addition, I recorded the effects of these manipulations on feather mass, length and rachis diameter. These morphological characteristics are responsible for the mechanical performance of feathers and can therefore directly affect fitness of birds.

Glucocorticoid administration significantly decreased feather CORT levels. This evidence that feather CORT levels are sensitive to hormonal manipulations supports the suitability of feather CORT for characterization of stress history of birds. Surprisingly, feather CORT levels did not correlate with other well-established measures of stress such as leukocyte differentials. Neither did I find that immune activation with artificial antigen had affected feather CORT levels. This finding, along with those of Meitern (2013) and Fairhurst (2013a) suggests that assessment of health state and stress history of birds on the basis of feather CORT levels is not as straightforward as previously thought and calls for further research on information content of feather CORT.

In addition to reducing CORT content of feathers, glucocorticoid administration also reduced all three measures of feather size. This points to the possibility that experiencing an acute stress or termination of immune responses by rising the concentration of circulating glucocorticoid hormones may induce long-term somatic costs with potential impact on fitness. Immunostimulation experiment showed that in the face of resource limitation investment into feather mass (which increases feather strength) is prioritized over investment into feather length. These results add another piece of evidence about somatic costs of induced immune responses in terms of feather growth.

Taking a step further from artificial antigens, I asked whether infection with a natural pathogen affects feather CORT level and how feather CORT content, a potential marker of stress responsiveness, reflects the resistance to an experimental infection with natural coccidian parasites. I also asked whether individual differences in disease resistance relate to behavioral phenotypes, as can be expected on the basis of previous work on animal personalities and endocrine activity (Martins *et al.* 2007; Koolhaas 2008). To study the link with behaviour, I assessed the extent of damage to tail feathers accrued in captivity. Extent of tail damage was suggested to reflect the captivity-intolerance in greenfinches (Sild *et al.* 2011). The same study also showed that greenfinches that inflicted greater extent of damage to their tail feathers exhibited compromised immune responsiveness in terms of reduced capability for producing antibody response to a novel *Brucella abortus* antigen and reduced oxidative burst capability of phagocytes, which suggests a possibility of CORT-induced immune suppression in birds with most damaged tail feathers.

Coccidian infection (or its suppression) indeed was associated with CORT secretion as birds subjected to experimental infection had on average 23% higher feather CORT content than control birds. In addition we found that experimental birds with lowest feather CORT levels developed highest infection intensities at the peak infection phase as well as highest increase in average infection levels. The most likely explanation for such pattern is that birds that invest more into fighting coccidiosis have higher stress responsiveness or that birds that experience captivity as more stressful have higher coccidian resistance. The benefits of increased resistance to coccidian parasites in the wild are obvious as coccidians are the most prevalent avian parasites (Greiner 2008)

with well documented pathological effects (Pap *et al.* 2011; Sepp *et al.* 2012). As predicted, I found that extent of tail damage accrued in captivity correlated positively with feather CORT content. The most likely explanation for this pattern would be that some greenfinches possessed more stress-prone personalities than others, consequently depositing more CORT in their feathers. These findings suggest that further research on CORT levels in relation to resistance of coccidiosis might increase our understanding about the mechanisms linking individual variation in CORT levels to fitness (see Breuner *et al.* 2008; Angelier *et al.* 2009).

Confirming that feather CORT content and morphological characteristics of feathers are sensitive to manipulations of organism's hormonal status and immune function, I proceeded with the second aim of my study, namely, with the investigation of whether and how feather parameters relate to behavioral traits. Between-individual variation in behavior is an important ecological and evolutionary characteristic of wild populations (Careau *et al.* 2008). Ample evidence suggests that behavior is not completely flexible and different behavioral traits do not vary independently of each other, thus giving rise to behavioral syndromes or personalities (Carere *et al.* 2010). I recorded five behavioral traits that are associated with different immune parameters (Sild *et al.* 2011) and stress responsiveness (Krams *et al.* 2013), and asked whether and which of these traits covary and manifest in general "fearfulness syndrome". The measured traits included latency to feed at the presence of a predator, the propensity to give distress calls at handling, changes in locomotor activity in response to distress calls of conspecifics, frequency of flapping flight bouts against cage bars and tendency to damage tail feathers in captivity. All of these traits show a response to an acute stressor and can therefore indicate different sides of a fear response. To study the associations between endocrine activity and behavior I recorded CORT levels of feathers. To my surprise, none of the behavioral parameters correlated with each other. These findings challenge the concept of a single internal variable responsible of fearfulness and support the proposed multidimensional nature of fear responses. Similarly, none of the behavioral parameters correlated with feather CORT levels. Furthermore, tail damage accrued in captivity did not correlate with the amount of CORT deposited into feathers grown in captivity. This finding contradicts my previous study where I detected a positive association between feather CORT and tail damage. This contradiction between the findings of two studies in the identical model system is difficult to explain and it calls for caution of generalizing the interpretations of feather CORT measurements obtained in different settings. Importantly, this study also showed that tail damage does not appear a universal correlate of stress responsiveness in captive greenfinches.

In the last part of my thesis I asked whether feather size and color parameters predict survival in greenfinches facing epidemics of emerging infectious disease, finch trichomonosis. Animals differ in their vulnerability to infections and these differences could be mirrored in their appearance as



immune system is integrated with almost any physiological function of an organism. The spread of trichomonosis epidemics to Estonia provided an excellent opportunity to study associations between plumage characteristics and trichomonosis-induced mortality. I measured the reflectance of black eumelanin-based tips of tail feathers because I expected that black coloration may relate to disease resistance. This expectation was based on the information that eumelanotic coloration is linked to expression of pleiotropic pro-opiomelanocortin genes that affect inflammatory and stress responses (Ducrest *et al.* 2008). Additionally, I was interested if the occurrence of fault bars can predict mortality as fault bars have been associated with increased mortality in different bird species (Bortolotti *et al.* 2002; Møller *et al.* 2009).

Out of 46 female greenfinches five birds died in captivity. Three of them displayed symptoms of trichomonosis, and the remaining two died most likely in the same reasons. Feathers of survivors were 22 % darker than feathers of dead birds. One possible explanation would be covariation between melanin-based coloration and some relevant aspect of immune function. Another possible explanation involves pleiotropic covariation between melanin production and stress susceptibility. Feathers of dead birds had significantly higher prevalence of fault bars. In addition, feathers with fault bars were significantly paler. As fault bars result from experiencing stress during feather growth (Bortolotti *et al.* 2002; Møller *et al.* 2009), high prevalence of fault bars in dead birds suggests that those individuals either experienced more stressors or appeared more susceptible to stress already in the preceding autumn when those feathers were growing. My results thus corroborate the findings of Bortolotti (2002) and Møller (2009), showing that fault bars can be used as a diagnostic cue in predicting mortality. Notably, this study also showed that fault bars and feather size parameters convey different information about stressful experiences during feather growth as feather size parameters did not differ between dead and alive birds. This study is the first to detect trichomonosis-induced viability selection on plumage characteristics. It enables to predict that the recent outbreak of trichomonosis leads to evolution of darker melanin-based coloration in greenfinches. Importantly, this study also suggests that melanin-based coloration can evolve via mechanisms that are independent of visual information provided by the pigment. This idea is relevant for understanding the mechanisms of evolution of melanin-based coloration in general. So far, the research in this area has been mainly concentrating on finding the costs of melanin-based display (McGraw, 2006). I suggest that the possibility that melanin-based integument coloration can evolve as a by-product of selection on pleiotropically linked physiological traits (see also Bize *et al.* 2006) deserves serious consideration in future studies.

In conclusion, my study showed that morphological characteristics, as well as CORT content of feathers are sensitive to manipulations of hormonal milieu and immune function, which makes these parameters potentially valuable indicators about the events that occurred in the period of feather growth. In

particular, feather mass, length and rachis diameter can be used to assess somatic costs of different stressors or trade-offs between feather growth and other demanding organismal functions. Feather wear in captivity can reflect behavior of birds and possibly be linked to stress susceptibility. Fault bars can be used as a diagnostic cue in predicting mortality. My study also indirectly suggests that eumelanotic color of feathers is linked to immune and/or stress responses, which helps us to understand the mechanisms of evolution of melanin-based coloration in general. Detected associations between feather CORT content and resistance to coccidiosis are particularly interesting, as they suggest a novel route of selection for higher stress responsiveness.

## SUMMARY IN ESTONIAN

### Rohevintide füsioloogiline ökoloogia: sulgedes sisalduv informatsioon immuunfunktsiooni ja käitumise kontekstis

Füsioloogiline ökoloogia uurib organismide füsioloogiliste funktsioonide ja keskkonna vahelisi suhteid. Selgroogsete närvi-, endokriin- ning immuunsüsteemid kuuluvad tõenäoliselt kõige keerukamate evolutsiooniliste kohastumuste hulka. Alles hiljuti on jõutud arusaamisele, et nimetatud süsteemid ei toimi üksteisest sõltumatult, vaid on omavahel ühendatud ühtseks tervikuks. Sellest tõdemusest kasvas välja kaks uut teadusharu: immuunökoloogia ja psühhoneuroimmunoloogia. Immuunökoloogia otsib vastuseid küsimustele, miks organismid erinevad vastuvõtlikkuses haigustele ja mis takistab immuunsüsteemi paremini toimimast. Immuunfunktsiooni isenditevahelise varieeruvuse üheks põhjuseks võib olla immuunsüsteemi regulatsioonimehhanismide keerukus ja hävitusemehhanismide ohtlikkus, mis muudab immuunsüsteemi kasutamise kulukaks. Kui organismi mingi funktsioon on kulukas, siis peab see paratamatult teiste funktsioonidega ühise ressursi pärast konkureerima; tulemuseks on erinevate funktsioonide vahelised lõivsuhted (*trade-off*). Närvi-, endokriin- ning immuunsüsteemi tööga kaasnevate lõivuhete mõistmiseks on oluline nende toimimisega kaasnevate kulude ja tulude kvantifitseerimine.

Haiguse kulgu võivad mõjutada paljud erinevad faktorid; üheks nendest on organismi stressivastus. Vastuseks kogetud stressitekitajale vabanevad hüpotaalamuse-ajuripatsi-neerupealise kaskaadi kaudu stressihormoonid (lindude puhul eelkõige kortikosteron (KORT)), mis võivad läbi füsioloogiliste ja käitumuslike mehhanismide mõjutada näiteks immuun- ja sigimissüsteemi ning ainevahetust. Kuigi immuun- ning stressivastuse vahelised seosed on üldjoontes teada, ei osata kõiki nähtusi adaptiivselt seletada. Näiteks pole teada, miks pikaajaline stress pärsib immuunsüsteemi funktsionaalsust.

Psühhoneuroimmunoloogia ühendab immuun- ning stressivastuse uuringud käitumise uurimisega. Ajas püsivaid isenditevahelisi erinevusi käitumises on leitud paljudel loomaliikidel. Päritavad, ajas ning erinevates olukordades püsivad ja omavahel korreleeruvad käitumistunnused moodustavad isendi isiksuse või iseloomu. Erineva isiksusega loomad erinevad üksteisest ka füsioloogia poolest, seega on põhjust eeldada, et käitumine ja immuun- ning stressivastused on omavahel seotud.

Üheks füsioloogilisi, morfoloogilisi ja käitumuslikke tunnuseid siduvaks mehhanismiks on geenide pleiotroopne mõju (ühe geeni mõju mitmele tunnusele). Näiteks peamise loomse pigmendi melaniini sünteesi eest vastutavad geenid osalevad nii stressi- ja immuunvastuse kui ka käitumise kujunemisel. Nii on erineva värvusega loomi seostatud alternatiivsete ökoloogiliste nišside, erinevate immuunvastuste, iseloomulike morfoloogiliste tunnuste või sigimisstrateegiatega.

Füsioloogilise ökoloogia peamiseks töömeetodiks on biokeemiliste markerite mõõtmine uurimisobjekti erinevatest kudedest. Lindude suled pakuvad selles kontekstis mitmeid unikaalseid võimalusi. Sulgede kasutamine uurimistöös on väheinvasiivne meetod, mille abil saab uurida looduslikke populatsioone ja sealhulgas ka kaitsealuseid liike. Sulgedes leidub rohkelt teavet organismisiseste ja organismiväliste protsesside, kogetud sündmuste või keskkonnamuutuste kohta. Informatsiooni võivad anda sulgede värv, suurus, terviklikkus ning erinevate hormoonide sisaldus. Selle informatsiooni potentsiaali on füsioloogilise ökoloogia uuringutes hakatud mõistma alles hiljuti.

Käesolevas töös kasutasin mudelorganismina rohevinte (*Carduelis chloris*), et välja selgitada erinevate suleparameetrite rakendatavus immuunökoloogilistes ja käitumisuuringutes. Esimesed kolm artiklit baseeruvad eksperimentaalsetel uuringutel. Nende eesmärgiks oli uurida immuunvastuse ning stressihormooni kunstliku analoogi deksametasooni manustamise mõju sule kasvule ja sule KORTi sisaldusele. Lisaks suleparameetrite diagnostilise väärtuse täpsustamisele võimaldasid need eksperimendid saada väärtuslikku informatsiooni aktiveeritud või alla surutud immuun- ja stressivastuste somaatiliste kulude ning korreleeruvate füsioloogiliste ja käitumuslike parameetrite kohta. Nende küsimuste uurimine on füsioloogilise ökoloogia keskmeks. Järgnevad kaks uurimust toetuvad korrelatiivsetele andmetele. Esimeses neist uurin, kas sule KORTi sisalduse kaudu on võimalik saada uut teavet hirmuga seostuvate käitumistunnuste kohta. Teine uurimus põhineb rohevintide uuel nakkushaigusel trihomonooosil (põhjustaja algloom *Trichomonas gallinae*), mis võimaldab laboritingimustes uurida suleparameetrite ning nakkushaigusest põhjustatud suremuse seoseid.

Üheks minu töö peamiseks eesmärgiks oli testida, kas uus meetod, KORTi mõõtmine sulgedest, sobib neuro-, endokriin- ning immuunsüsteemi omavahealise seoste uurimiseks. Kuigi see meetod on viimastel aastatel kiiresti populaarsust kogunud, puuduvad eksperimentaalsed uuringud sulgede KORTi sisalduse füsioloogilise tähenduse kohta. Näiteks pole ikka veel täpselt teada, milline on erinevate füsioloogiliste (nt immuunsüsteemi aktiveerumine või vere KORTi sisalduse tõus) ja psühholoogiliste (nt kiskja kohtamine) tegurite mõju sulgede KORTi sisaldusele. Seepärast testisin immuunsüsteemi aktiveerimise ja stressihormooni manustamise mõju sulgede KORTi sisaldusele. Lisaks uurisin, kuidas sellised manipulatsioonid mõjutavad sule massi, pikkust ning suleroo diameetrit. Need sule morfoloogilised tunnused vastutavad sulgede mehhaanilise kvaliteedi eest ja võivad seega otseselt mõjutada lindude kohasust.

Minu töö tulemuse kohaselt vähendas deksametasooni manustamine oluliselt sulgede KORTi sisaldust (artikkel II). Kunstliku stressihormooni lisandumine vereringesse annab organismile signaali vähendada kehaomase stressihormooni tootmist. Sulgede KORTi sisalduse tundlikkus hormoonide taset mõjutavale manipulatsioonile kinnitab selle parameetri sobivust lindude stressivastuse kirjeldamiseks. Huvitav on avastus, et sulgede KORTi sisaldus ei korreleerunud teise linnu-uuringutes laialt kasutusel oleva stressiindikaatoriga, heterofiilide ja

lümfotsüütide suhtega veres. Lisaks ei mõjutanud sulgede KORTi sisaldust immuunsüsteemi aktiveerimine kunstliku antigeeniga (fütohemaglutiniin, PHA) (artikkel I). Need tulemused koos mõnede teiste uurimustega lubavad oletada, et lindude tervisliku seisundi ja stressivastuse hindamine sule KORTi sisalduse põhjal ei ole nii lihtne kui arvatakse.

Lisaks sulgede KORTi sisalduse vähendamisele vähendas kunstliku stressihormooni manustamine kõiki sulgede suuruse parameetreid (artikkel I). See viitab võimalusele, et akuutne stress või vere KORTi sisalduse kunstliku tõstmise tulemusena inhibeeritud stressivastus (negatiivne tagaside) võib esile kutsuda pikaajalisi somaatilisi kulusid, mis võivad mõjutada isendi kohasust. Kui KORT mõjutab nii sulgede pikkust, massi kui roo diameetrit, siis immuunsüsteemi aktiveerimine PHA-ga vähendas vaid sule pikkust, mis näitab seda, et ressursside limiteerituse korral panustatakse rohkem sule massi (mis suurendab sule tugevust). Need tulemused täiendavad meie teadmisi aktiveeritud immuunsüsteemi ja stressivastuse somaatiliste kulude kohta.

Kolmandas artiklis uurisin, kuidas mõjutab sule KORTi sisaldust nakatamine koktsiididega (rohevinte looduslikult nakatavad rakusisesed sooleparasiidid perekonnast *Isospora*) (artikkel III). Lisaks huvitas mind, kas sule KORTi sisaldus (kui potentsiaalne stressivastuse marker) peegeldab resistentsust eksperimentaalsele koktsiidinakkusele. Katse võimaldas välja selgitada, kas erinevused haigusresistentsuses seostuvad lindude isiksusega, nagu on viidanud mitmed varasemad füsioloogia- ja käitumisalased uuringud. Käitumise kvantifitseerimiseks hindasin rohevintidel vangistuses tekkinud sabakahjustusi. Eelnevad uuringud on näidanud, et erinevused sabakahjustuste ulatuses võivad peegeldada lindude erinevat võimet taluda vangistust.

Koktsiidinakkus tõstis sule KORTi sisaldust: eksperimentaalselt nakatatud lindude suled sisaldasid 23% rohkem KORTi kui kontrollgrupi lindude suled. Lisaks leidsin, et eksperimentaalselt nakatatud lindude hulgas oli nakkus kõige intensiivsem nendel, kel oli sulgedes kõige vähem KORTi. Sellel tulemusel on kaks võimalikku seletust. Esiteks võib lindudel, kes panustasid kõige rohkem nakkusega võitlemisele, olla keskmisest kõrgem stressivastus. Teiseks võivad linnud, kellel on tugevam vangistusstress, olla nakkusele resistentsamad. Resistentsus koktsiidinakkusele on looduses ilmselgelt kasulik, sest koktsiidid on levinuimad lindude patogeensed parasiidid. Ootuspäraselt korreleerus sabasulgedele tekkinud kahjustuste määr sule KORTi sisaldusega. Võib arvata, et mõned linnud olid madalama stressitaluvusega kui teised, mille tagajärjel sisaldasid nende lindude suled rohkem KORTi ja nad tekitasid sabasulgedele suuremas ulatuses kahjustusi. Need tulemused aitavad selgitada, kuidas võib sule KORTi sisaldus olla seotud isendi kohasusega.

Töö teises pooles uurisin, kas ja kuidas on suleparameetrid seotud lindude käitumisega (artikkel IV). Isendite erinev käitumine (erinevad isiksused ehk iseloomud) on looduslike populatsioonide tähtis evolutsiooniline ja ökoloogiline omadus. Eelnevad uuringud on näidanud käitumise piiratud paindlikkust, mis tuleneb erinevate käitumistunnuste omavahelistest seostest. Neljandas artiklis

mõõtsin viit käitumistunnust, mis on seotud vastusega akuutsele stressitekitajale. Uurisin, kas need käitumistunnused omavahel korreleeruvad ja moodustavad „hirmu sündroomi“. Mõõdetavad parameetrid olid toitumislattentsus kiskja juuresolekul, kalduvus käsitsemisel hirmukisa kuuldavale tuua, aktiivsuse muutus vastuseks liigikaaslaste hirmukisale, vastu puuriseina raplemise sagedus ning vangistuses tekkinud sabakahjustuste ulatus.

Vastupidiselt oodatule mõõdetud käitumistunnused omavahel ei korreleerunud. See tulemus on vastuolus arusaamaga, et hirmukäitumine on ühedimensiooniline isiksuseomadus, ning näitab, et lindude hirmureaktsioonid on konteksti- ja seisundispetsiifilised.

Ükski käitumistunnus ei korreleerunud sule KORTi sisaldusega. Sule KORTi sisaldus ei korreleerunud ka vangistuses sabasulgede tekitatud kahjustustega. See tulemus on vastuolus varasema tulemusega (artikkel III), kus sule KORTi sisaldus korreleerus positiivselt sabasulgede tekitatud kahjustustega. Kahe sarnase katse vastukäivad tulemused seavad kahtluse alla üksikutel sule KORT-i käsitlevatel uurimustel põhinevate järelduste üldistusvõime.

Oma töö viimases osas (artikkel V) uurisin, kas rohevintide sule suurus ja värvus ennustab vastupanuvõimet uudsele ning epideemilisele trihhomonoosinakkusele. Organismide vastuvõtlikkus haigustele on erinev ning need erinevused võivad peegelduda välimuses, sest immuunsüsteem on seotud paljude teiste füsioloogiliste funktsioonidega. Trihhomonoosi hiljutine levik Eestisse andis võimaluse uurida trihhomonoosist põhjustatud suremuse seoseid sule parameetritega. Mõõtsin sulgede musta (eumelaniinse) suletipu heledust, sest eeldasin sule musta värvuse seotust haigusresistentsusega. See eeldus põhineb varasematel uuringutel, kus näidati, et eumelaniini süntees on seotud pro-opiomelanokortiini geeni aktiivsusega, mis omakorda mõjutab ka põletikulisi protsesse ning stressivastust. Lisaks uurisin, kas sulgedel esinevad kasvulüngad ennustavad vintide suremust, kuna seda tunnust on erinevatel linnuliikidel varem suremusega seostatud.

Loodusest püütud 46 rohevindist suri vangistuses viis lindu. Kolmel neist esinesid selged trihhomonoosi sümptomid ning kaks ülejäänut surid tõenäoliselt samal põhjusel. Ellujäänud lindude suled olid võrreldes surnud lindude sulgedega 22% tumedamad. Üheks võimalikuks seletuseks on melaniinil põhineva värvuse füsioloogiline seos immuunsüsteemiga. Teiseks võimalikuks seletuseks on seos melaniini sünteesi ja stressivastuse vahel. Surnud lindude suled sisaldasid rohkem kasvulünki, lisaks olid kasvulünkadega suled ka oluliselt heledamad. Kuna kasvulüngad tekivad sule kasvamise ajal kogetud stressi tulemusena, viitab rohkemate kasvulünkade esinemine surnud lindude sulgedes sellele, et need isendid olid stressile vastuvõtlikumad või puutusid juba eelneval sügisel, sulgede kasvu ajal, kokku tõsisemate stressitekitajatega. See kinnitab varasemaid tulemusi, mille kohaselt kasvulünki on võimalik kasutada suremuse ennustamisel. Märkimist väärib ka järeldus, et kasvulüngad ja sule suuruse parameetrid annavad sule kasvu ajal kogetud stressitekitajate kohta erinevat informatsiooni, kuna sule suuruse parameetrid surnud ja ellujäänud

lindudel ei erinenud. Käesolev uurimus on esimene, mis näitab trihhomonoosi valikusurvet sule parameetritele. Selle põhjal võib ennustada, et hiljutine trihhomonoosipuhang viib rohevintidel tumedama sulevärvuse evolutsioneerumiseni. Uurimus viitab ka sellele, et eumelaniinil põhinev sulevärvus võib evolutsioneeruda sõltumatult värvipigmenti poolt edastatud visuaalsest informatsioonist. See avastus aitab mõista melaniinil põhineva värvuse üldisi evolutsioonimehhanisme. Siiani on peamiselt keskendunud melaniinsete ornamentide kulude kvantifitseerimisele. Võimalus, et melaniinse värvuse kujunemisel on oluline roll värvuse tootmisega pleiotroopselt seotud füsioloogilistele omadustele avalduva valikusurvega, väärib tulevastel uuringutes tõsist tähelepanu.

Kokkuvõttes võib minu töö põhjal järeldada, et suled sisaldavad külluslikult erinevat informatsiooni. Sule erinevad parameetrid on tundlikud organismi hormonaalse sisekeskkonna ja immuunsüsteemi manipulatsioonidele, andes seega väärtuslikku informatsiooni sulgede kasvu ajal kogetud sündmuste kohta. Sule massi, pikkust ja suleroo diameetrit on võimalik kasutada erinevate stressitekitajate somaatiliste kulude hindamiseks või sulekasvu ja organismi teiste oluliste füsioloogiliste protsesside vaheliste lõivuhete kirjeldamiseks. Vangistuses tekkinud sulekahjustused peegeldavad lindude käitumist ning võivad iseloomustada vastuvõtlikkust stressile. Kasvulünki on võimalik kasutada suremuse ennustamisel. Minu uurimistöö viitab võimalusele, et sulgede melaniinil põhinev värvus on seotud immuun- ja stressivastusega, mis aitab mõista melaniinil põhineva värvuse üldisi evolutsioonimehhanisme. Sule KORTi ja koktsidiosisiresistentsuse vaheline seos on eriti huvitav, sest see võib aidata selgitada, miks looduslik valik võib soosida kõrgeenenud stressivastust.

## ACKNOWLEDGEMENTS

My first and foremost gratitude goes to my supervisor Peeter Hõrak. Without his infectious enthusiasm and profound knowledge in the field of ecology I would not have achieved my bachelor's, master's and not to mention a PhD degree. He assembled a high functioning and supportive workgroup that I have really enjoyed working in. Richard Meitern saved me hours and days worth of work by providing me computer codes that I was not able to come up with on my own. Ulvi Karu kindly taught me how to do spectrophotometric measurements and introduced the life of coccidians. From Tuul Sepp I have learned how to be patient, work hard and how to write scientific text. Especially, I am grateful to Ulvi and Tuul who helped with the thesis manuscript preparation. Many thanks to Elin Sild for adjusting the corticosterone assay for greenfinch feathers and for being always ready to elaborate on biochemical analyses. Diana Osuna, Jolanta Vrublevska, Pirko Jalakas, Mari-Ann Lind, Janek Urvik, Grete Lüütsepp, Liis Rebase and many others have helped with bird maintenance and sample collection. Thanks to Stefaan Van Dyck (Kemin Agrifoods Europe) we and our greenfinches were continuously supplied with carotenoids. Ingmar Tulva repeatedly lent us the color spectrometer. Jaanis Lodjak provided some excellent pictures of greenfinches that illustrated my conference posters and presentations. Christine R. Lattin consulted us in the determination of feather corticosterone and Margus Leppik and Aivar Liiv helped with the radioimmunoassay. Ants Kaasik and Lauri Saks were the co-authors of some studies. I would like to thank all the reviewers for providing constructive criticism on the manuscripts. Mati Martinson was of great help with catching the greenfinches in Sõrve.

I would like to thank Raivo Mänd and all other members of the faculty for the knowledge, support and good advice I frequently needed. Thank you also for the interesting discussions in the animal ecology seminars. Margret Sisask, Triin Lett and Eve Möls were always ready to help with administrative questions. Indrikis Krams from Latvia was always ready to discuss the newest and most interesting ecological problems. I would like to thank my fellow PhD students for the motivation and support in my studies.

I would especially like to thank my friends and family for the patience and support throughout my studies. Special thanks to Madis Prink for tremendous support and motivation.

Thank you all.



The study was performed at the Department of Zoology, University of Tartu, at the Institute of Ecology and Earth Sciences.

The study was financially supported by the Estonian Ministry of Education and Science (target-financing project no. 0180004s09), the European Union through the European Regional Development Fund (Centre of Excellence Frontiers In Biodiversity Research), the Estonian Science Foundation grant no. 7737 (to Peeter Hõrak), the Estonian Doctoral School of Ecology and Environmental Sciences and the scholarship of Lydia and Felix Krabi.

## REFERENCES

- Adamo S.A. (2004). How should behavioural ecologists interpret measurements of immunity? *Animal Behaviour*, 68, 1443–1449.
- Adamo S.A. (2010). Why should an immune response activate the stress response? Insights from the insects (the cricket *Gryllus texensis*). *Brain, Behavior, and Immunity*, 24, 194–200.
- Adamo S.A. (2012). The effects of the stress response on immune function in invertebrates: an evolutionary perspective on an ancient connection. *Hormones and Behavior*, 62, 324–330.
- Ader R. (2001). Psychoneuroimmunology. *Current Directions in Psychological Science*, 10, 94–98.
- Angelier F., Holberton R.L. & Marra P.P. (2009). Does stress response predict return rate in a migratory bird species? A study of American redstarts and their non-breeding habitat. *Proceedings of the Royal Society B: Biological Sciences*, 276, 3545–3551.
- Atkinson C.T., Thomas N.J. & Hunter D.B. (2008). Parasitic diseases of wild birds. *Wiley Online Library*.
- Bell A.M. & Sih A. (2007). Exposure to predation generates personality in threespined sticklebacks (*Gasterosteus aculeatus*). *Ecology Letters*, 10, 828–834.
- Bennett K.A., Fedak M.A., Moss S.E.W., Pomeroy P.P., Speakman J.R. & Hall A.J. (2013). The role of glucocorticoids in naturally fasting grey seal (*Halichoerus grypus*) pups: dexamethasone stimulates mass loss and protein utilization, but not departure from the colony. *The Journal of Experimental Biology*, 216, 984–991.
- Bennett P.M. & Owens I.P.F. (2002). *Evolutionary ecology of birds: life histories, mating systems and extinction*. Oxford University Press. New York.
- Bize P., Gasparini J., Klopfenstein A., Altwegg R. & Roulin A. (2006). Melanin-based coloration is a nondirectionally selected sex-specific signal of offspring development in the alpine swift. *Evolution*, 60, 2370–2380.
- Boissy A. (1995). Fear and fearfulness in animals. *Quarterly Review of Biology*, 70, 165–191.
- Boissy A. & Bouissou M.F. (1995). Assessment of individual differences in behavioural reactions of heifers exposed to various fear-eliciting situations. *Applied Animal Behaviour Science*, 46, 17–31.
- Bókony V., Lendvai Á.Z., Vágási C.I., Pátraş L., Pap P.L., Németh J., Vincze E., Papp S., Preiszner B. & Seress G. (2014). Necessity or capacity? Physiological state predicts problem-solving performance in house sparrows. *Behavioral Ecology*, 25, 124–135.
- Bonier F., Martin P.R., Moore I.T. & Wingfield J.C. (2009). Do baseline glucocorticoids predict fitness? *Trends in Ecology & Evolution*, 24, 634–642.
- Bortolotti G.R. (2010). Flaws and pitfalls in the chemical analysis of feathers: bad news-good news for avian chemoecology and toxicology. *Ecological Applications*, 20, 1766–1774.
- Bortolotti G.R., Dawson R.D. & Murza G.L. (2002). Stress during feather development predicts fitness potential. *Journal of Animal Ecology*, 71, 333–342.
- Bortolotti G.R., Marchant T., Blas J. & Cabezas S. (2009). Tracking stress: localisation, deposition and stability of corticosterone in feathers. *Journal of Experimental Biology*, 212, 1477–1482.

- Bortolotti G.R., Marchant T.A., Blas J. & German T. (2008). Corticosterone in feathers is a long-term, integrated measure of avian stress physiology. *Functional Ecology*, 22, 494–500.
- Breuner C.W., Patterson S.H. & Hahn T.P. (2008). In search of relationships between the acute adrenocortical response and fitness. *General and Comparative Endocrinology*, 157, 288–295.
- Butler L.K., Rohwer S. & Speidel M.G. (2008). Quantifying structural variation in contour feathers to address functional variation and life history trade-offs. *Journal of Avian Biology*, 39, 629–639.
- Careau V., Thomas D., Humphries M.M. & Réale D. (2008). Energy metabolism and animal personality. *Oikos*, 117, 641–653.
- Carere C., Caramaschi D. & Fawcett T.W. (2010). Covariation between personalities and individual differences in coping with stress: converging evidence and hypotheses. *Current Zoology*, 56, 728–740.
- Carere C., Groothuis T.G.G., Mostl E., Daan S. & Koolhaas J.M. (2003). Fecal corticosteroids in a territorial bird selected for different personalities: daily rhythm and the response to social stress. *Hormones and Behavior*, 43, 540–548.
- Catania A., Lonati C., Sordi A., Carlin A., Leonardi P. & Gatti S. The Melanocortin System in Control of Inflammation. *TheScientificWorldJournal*, 10, 1840–1853.
- Chew B.P. & Park J.S. (2004). Carotenoid Action on the Immune Response. *Journal of Nutrition*, 134, S257–S261.
- Cockrem J. (2007). Stress, corticosterone responses and avian personalities. *Journal of Ornithology*, 148, S169–S178.
- Costantini D., Fanfani A. & Dell’Omo G. (2008). Effects of corticosteroids on oxidative damage and circulating carotenoids in captive adult kestrels (*Falco tinnunculus*). *Journal of Comparative Physiology B: Biochemical, Systemic, and Environmental Physiology*, 178, 829–835.
- Costantini D. & Møller A.P. (2008). Carotenoids are minor antioxidants for birds. *Functional Ecology*, 22, 367–370.
- Cote J., Meylan S., Clobert J. & Voituron Y. (2010). Carotenoid-based coloration, oxidative stress and corticosterone in common lizards. *Journal of Experimental Biology*, 213, 2116–2124.
- Cramp P. & Perrins C.M. (1994). *The Birds of Western Palearctic*. Oxford University Press, Oxford.
- Cyr N.E., Dickens M.J. & Romero L.M. (2009). Heart Rate and Heart-Rate Variability Responses to Acute and Chronic Stress in a Wild-Caught Passerine Bird. *Physiological and Biochemical Zoology*, 82, 332–344.
- Cyr N.E. & Michael Romero L. (2007). Chronic stress in free-living European starlings reduces corticosterone concentrations and reproductive success. *General and Comparative Endocrinology*, 151, 82–89.
- Cyr N.E. & Romero L.M. (2008). Fecal glucocorticoid metabolites of experimentally stressed captive and free-living starlings: implications for conservation research. *General and Comparative Endocrinology*, 158, 20–28.
- Davis A.K., Maney A.K. & Maerz J.C. (2008). The use of leukocyte profiles to measure stress in vertebrates: a review for ecologists. *Functional Ecology*, 760–772.
- Davis M., Walker D.L., Miles L. & Grillon C. (2010). Phasic vs Sustained Fear in Rats and Humans: role of the Extended Amygdala in Fear vs Anxiety. *Neuropsychopharmacology*, 35, 105–135.

- Dawson A., Hinsley S.A., Ferns P.N., Bonser R.H.C. & Eccleston L. (2000). Rate of moult affects feather quality: a mechanism linking current reproductive effort to future survival. *Proceedings of the Royal Society of London Series B-Biological Sciences*, 267, 2093–2098.
- De La Hera I., Perez-Tris J. & Telleria J.L. (2009). Migratory behaviour affects the trade-off between feather growth rate and feather quality in a passerine bird. *Biological Journal of the Linnean Society*, 97, 98–105.
- Demas G.E. (2004). The energetics of immunity: a neuroendocrine link between energy balance and immune function. *Hormones and Behavior*, 45, 173–180.
- Demas G.E., Adamo S.A. & French S.S. (2011). Neuroendocrine-immune crosstalk in vertebrates and invertebrates: implications for host defence. *Functional Ecology*, 25, 29–39.
- Demas G.E. & Nelson R.J. (2011). *Ecoimmunology*. Oxford University Press, New York.
- DesRochers D.W., Reed J.M., Awerman J., Kluge J.A., Wilkinson J., van Griethuijsen L.I., Aman J. & Romero L.M. (2009). Exogenous and endogenous corticosterone alter feather quality. *Comparative Biochemistry and Physiology – A Molecular and Integrative Physiology*, 152, 46–52.
- Dhabhar F.S. (2002). A hassle a day may keep the doctor away: stress and the augmentation of immune function. *Integrative and Comparative Biology*, 42, 556–564.
- Dingemanse N.J. & Réale D. (2005). Natural selection and animal personality. *Behaviour*, 142, 9, 1159–1184.
- Dingemanse N.J. & Wolf M. (2010). Recent models for adaptive personality differences: a review. *Philosophical Transactions of the Royal Society B-Biological Sciences*, 365, 3947–3958.
- Dingemanse N.J., Wright J., Kazem A.J.N., Thomas D.K., Hickling R. & Dawnay N. (2007). Behavioural syndromes differ predictably between 12 populations of three spined stickleback. *Journal of Animal Ecology*, 76, 1128–1138.
- Ducrest A.L., Keller L. & Roulin A. (2008). Pleiotropy in the melanocortin system, coloration and behavioural syndromes. *Trends in Ecology & Evolution*, 23, 502–510.
- Emaresi G., Ducrest A.-L., Bize P., Richter H., Simon C. & Roulin A. (2013). Pleiotropy in the melanocortin system: expression levels of this system are associated with melanogenesis and pigmentation in the tawny owl (*Strix aluco*). *Molecular Ecology*, 22, 4915–4930.
- Fairhurst G.D., Marchant T.A., Soos C., Machin K.L. & Clark R.G. (2013a). Experimental relationships between levels of corticosterone in plasma and feathers in a free-living bird. *The Journal of Experimental Biology*, 216, 4071–4081.
- Fairhurst G.D., Vögeli M., Serrano D., Delgado A., Tella J.L. & Bortolotti G.R. (2013b). Can synchronizing feather-based measures of corticosterone and stable isotopes help us better understand habitat–physiology relationships? *Oecologia*, 1–13.
- Gangoso L., Grande J.M., Ducrest A.L., Figuerola J., Bortolotti G.R., Andres J.A. & Roulin A. (2011). MC1R-dependent, melanin-based colour polymorphism is associated with cell-mediated response in the Eleonora's falcon. *Journal of Evolutionary Biology*, 24, 2055–2063.
- Garamszegi L.Z., Rosivall B., Rettenbacher S., Marko G., Zsebok S., Szollosi E., Eens M., Potti J. & Torok J. (2012). Corticosterone, avoidance of novelty, risk-taking and

- aggression in a wild bird: no evidence for pleiotropic effects. *Ethology*, 118, 621–635.
- Graham A.L., Allen J.E. & Read A.F. (2005). Evolutionary causes and consequences of immunopathology. *Annual Review of Ecology Evolution and Systematics*, 36, 373–397.
- Graham A.L., Shuker D.M., Pollitt L.C., Auld S.K.J.R., Wilson A.J. & Little T.J. (2011). Fitness consequences of immune responses: strengthening the empirical framework for ecoimmunology. *Functional Ecology*, 25, 5–17.
- Greiner E.C. (2008). *Parasitic Diseases of Wild Birds*. Wiley-Blackwell, New York.
- Grubb T.C. (1989). Ptilochronology- feather growth bars as indicators of nutritional status. *Auk*, 106, 314–320.
- Hamilton W.D. & Zuk M. (1982). Heritable true fitness and bright birds: a role of parasites. *Science*, 218, 384–387.
- Hill G.E. & Farmer K.L. (2005). Carotenoid-based plumage coloration predicts resistance to a novel parasite in the house finch. *Naturwissenschaften*, 92, 30–34.
- Hill G.E. & McGraw K.J. (2006). *Bird Coloration. Volume I. Mechanisms and Measurements*. Harvard University Press, Cambridge.
- Hill G.E. & McGraw K.J. (2006a). *Bird Coloration. Volume II. Function and Evolution*. Harvard University Press, Cambridge.
- Holberton R.L., Wilson C.M., Hunter M.J., Cash W.B. & Sims C.G. (2007). The role of corticosterone in supporting migratory lipogenesis in the dark-eyed Junco (*Junco hyemalis*): a model for central and peripheral regulation. *Physiol Biochem Zool*, 80, 125–37.
- Huff G.R., Huff W.E., Balog J.M. & Rath N.C. (1999). The effect of a second dexamethasone treatment on turkeys previously challenged in an experimental *Escherichia coli* respiratory model of turkey osteomyelitis complex. *Poultry Science*, 78, 1116–1125.
- Hörak P., Zilmer M., Saks L., Ots I., Karu U. & Zilmer K. (2006). Antioxidant protection, carotenoids, and the costs of immune challenge in greenfinches. *Journal of Experimental Biology*, 209, 4329–4338.
- Jawor J.M. & Breitwisch R. (2003). Melanin ornaments, honesty, and sexual selection. *Auk*, 120, 249–265.
- Jennions M.D., Møller A.P. & Petrie M. (2001). Sexually selected traits and adult survival: a meta-analysis. *Quarterly Review of Biology*, 76, 3–36.
- Johnstone C.P., Reina R.D. & Lill A. (2012). Interpreting indices of physiological stress in free-living vertebrates. *Journal of Comparative Physiology B: Biochemical, Systemic, and Environmental Physiology*, 1–19.
- Keyser A.J. & Siefferman L.M. (2005). Viability selection against highly ornamented males. *Evolutionary Ecology Research*, 7, 595–606.
- Kingsolver J.G. (1989). Weather and the population dynamics of insects- integrating physiological and population ecology. *Physiological Zoology*, 62, 314–334.
- Kittilsen S., Johansen I.B., Braastad B.O. & Overli O. (2012). Pigments, parasites and personality: towards a unifying role for steroid hormones? *PLoS ONE*, 7.
- Kittilsen S., Schjolden J., Beitnes-Johansen I., Shaw J.C., Pottinger T.G., Sørensen C., Braastad B.O., Bakken M. & Overli O. (2009). Melanin-based skin spots reflect stress responsiveness in salmonid fish. *Hormones and Behavior*, 56, 292–298.
- Koolhaas J., Bartolomucci A., Buwalda B., De Boer S., Flüggé G., Korte S., Meerlo P., Murison R., Olivier B. & Palanza P. (2011). Stress revisited: a critical evaluation of the stress concept. *Neuroscience & Biobehavioral Reviews*, 35, 1291–1301.

- Koolhaas J.M. (2008). Coping style and immunity in animals: making sense of individual variation. *Brain Behavior and Immunity*, 22, 662–667.
- Koolhaas J.M., Korte S.M., De Boer S.F., Van Der Vegt B.J., Van Reenen C.G., Hopster H., De Jong I.C., Ruis M.A.W. & Blokhuis H.J. (1999). Coping styles in animals: current status in behavior and stress-physiology. *Neuroscience and Biobehavioral Reviews*, 23, 925–935.
- Kralj-Fiser S., Scheiber I.B.R., Blejec A., Mostl E. & Kotrschal K. (2007). Individualities in a flock of free-roaming greylag geese: behavioral and physiological consistency over time and across situations. *Hormones and Behavior*, 51, 239–248.
- Kralj-Fiser S., Weiss B.M. & Kotrschal K. (2010). Behavioural and physiological correlates of personality in greylag geese (*Anser anser*). *Journal of Ethology*, 28, 363–370.
- Krams I., Vrublevska J., Cīrule D., Kivleniece I., Krama T., Rantala M.J., Kaasik A., Hōrak P. & Sepp T. (2013). Stress, behaviour and immunity in wild-caught wintering great tits (*Parus major*). *Ethology*, 119, 397–406.
- Krinsky N.I. (1989). Antioxidant functions of carotenoids. *Free Radical Biology and Medicine*, 7, 617–635.
- Landeau E.A. & Badyaev A.V. (2012). Developmental integration of feather growth and pigmentation and its implications for the evolution of diet-derived coloration. *Journal of Experimental Zoology Part B-Molecular and Developmental Evolution*, 318B, 59–70.
- Lattin C.R., Reed J.M., DesRochers D.W. & Romero L.M. (2011). Elevated corticosterone in feathers correlates with corticosterone-induced decreased feather quality: a validation study. *Journal of Avian Biology*, 42, 247–252.
- Laugero K.D. & Moberg G.P. (2000). Summation of behavioral and immunological stress: metabolic consequences to the growing mouse. *American Journal of Physiology-Endocrinology and Metabolism*, 279, E44–E49.
- Lilliendahl K. (2000). Daily accumulation of body reserves under increased predation risk in captive greenfinches (*Carduelis chloris*). *Ibis*, 142, 587–595.
- Lindström K., Krakower D., Lundström J.O. & Silverin B. (2001). The effects of testosterone on a viral infection in greenfinches (*Carduelis chloris*): an experimental test of the immunocompetence-handicap hypothesis. *Proceedings of the Royal Society of London, Series B: Biological Sciences*, 268, 207–211.
- Lindström K. & Lundström J. (2000). Male greenfinches (*Carduelis chloris*) with brighter ornaments have higher virus infection clearance rate. *Behavioral Ecology and Sociobiology*, 48, 44–51.
- Lochmiller R.L. & Deerenberg C. (2000). Trade-offs in evolutionary immunology: just what is the cost of immunity? *Oikos*, 88, 87–98.
- Loiseau C., Fellous S., Haussy C., Chastel O. & Sorci G. (2008). Condition-dependent effects of corticosterone on a carotenoid-based begging signal in house sparrows. *Hormones and Behavior*, 53, 266–273.
- Lozano G.A. (1994). Carotenoids, parasites and sexual selection. *Oikos*, 70, 309–311.
- Martin L.B. (2009). Stress and immunity in wild vertebrates: timing is everything. *General and Comparative Endocrinology*, 163, 70–76.
- Martin L.B., Hawley D.M. & Ardia D.R. (2011). An introduction to ecological immunology. *Functional Ecology*, 25, 1–4.
- Martin L.B., Weil Z.M. & Nelson R.J. (2008). Seasonal changes in vertebrate immune activity: mediation by physiological trade-offs. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 363, 321–339.

- Martins T.L.F., Roberts M.L., Giblin I., Huxham R. & Evans M.R. (2007). Speed of exploration and risk-taking behavior are linked to corticosterone titres in zebra finches. *Hormones and Behavior*, 52, 445–453.
- Matyjasiak P., Matyjasiak J., de Lope F. & Møller A.P. (2004). Vane emargination of outer tail feathers improves flight maneuverability in streamerless hirundines, *Hirundinidae*. *Proceedings of the Royal Society of London Series B-Biological Sciences*, 271, 1831–1838.
- Meitern R., Sild E., Lind M.-A., Männiste M., Sepp T., Karu U. & Hõrak P. (2013). Effects of endotoxin and psychological stress on redox physiology, immunity and feather corticosterone in greenfinches. *PLOS ONE*, 8, e67545.
- Miller G.E., Cohen S. & Ritchey A.K. (2002). Chronic psychological stress and the regulation of pro-inflammatory cytokines: a glucocorticoid-resistance model. *Health Psychology*, 21, 531–541.
- Minias P. & Iciek T. (2013). Extent and symmetry of post-juvenile moult as predictors of future performance in greenfinch (*Carduelis chloris*). *Journal of Ornithology*, 154, 465–468.
- Møller A.P., Erritzoe J. & Nielsen J.T. (2009). Frequency of fault bars in feathers of birds and susceptibility to predation. *Biological Journal of the Linnean Society*, 97, 334–345.
- Møller A.P., Erritzoe J. & Nielsen J.T. (2009). Frequency of fault bars in feathers of birds and susceptibility to predation. *Biological Journal of the Linnean Society*, 97, 334–345.
- Moore I.T. & Hopkins W.A. (2009). Interactions and trade-offs among physiological determinants of performance and reproductive success. *Integrative and Comparative Biology*, 49, 441–451.
- Moreno J. & Møller A. (2006). Are melanin ornaments signals of antioxidant and immune capacity in birds. *Acta Zoologica Sinica*, 52, 202–208.
- Murphy M.E. (1996). Energetics and nutrition of molt. *Avian Energetics and Nutritional Ecology*, 158–198.
- Müller C., Jenni-Eiermann S. & Jenni L. (2009). Effects of a short period of elevated circulating corticosterone on postnatal growth in free-living Eurasian kestrels (*Falco tinnunculus*). *Journal of Experimental Biology*, 212, 1405–1412.
- Müller C., Jenni-Eiermann S. & Jenni L. (2010). Heterophils/Lymphocytes-ratio and circulating corticosterone do not indicate the same stress imposed on Eurasian kestrel nestlings. *Functional Ecology*, 25, 566–576.
- Neimanis A.S., Handeland K., Isomursu M., Agren E., Mattsson R., Hammes I.S., Bergsjø B. & Hirvela-Koski V. (2010). First report of epizootic trichomoniasis in wild finches (Family *Fringillidae*) in southern fennoscandia. *Avian diseases*, 54, 136–141.
- Nelson R.J. & Demas G.E. (1996). Seasonal changes in immune function. *Quarterly Review of Biology*, 71, 511–549.
- Nilsson J. & Svensson E. (1996). The cost of reproduction: a new link between current reproductive effort and future reproductive success. *Proceedings of the Royal Society of London, Series B: Biological Sciences*, 263, 711–714.
- Nolan P.M., Hill G.E. & Stoehr A.M. (1998). Sex, size and plumage redness predict house finch survival in an epidemic. *Proceedings of the Royal Society of London, Series B: Biological Sciences*, 265, 961–965.
- Ots I., Murumägi A. & Hõrak P. (1998). Haematological health state indices of reproducing great tits: methodology and sources of natural variation. *Functional Ecology*, 12, 700–707.

- Owen-Ashley N. & Wingfield J. (2007). Acute phase responses of passerine birds: characterization and seasonal variation. *Journal of Ornithology*, 148, 583–591.
- Pap P.L., Vagasi C.I., Barbos L. & Marton A. (2013). Chronic coccidian infestation compromises flight feather quality in house sparrows (*Passer domesticus*). *Biological Journal of the Linnean Society*, 108, 414–428.
- Pap P.L., Vagasi C.I., Czirjak G.A., Titilincu A., Pintea A. & Barta Z. (2009). Carotenoids modulate the effect of coccidian infection on the condition and immune response in moulting house sparrows. *Journal of Experimental Biology*, 212, 3228–3235.
- Pap P.L., Vagasi C.I., Czirjak G.A., Titilincu A., Pintea A., Osvath G., Fulop A. & Barta Z. (2011). The effect of coccidians on the condition and immune profile of molting house sparrows (*Passer domesticus*). *Auk*, 128, 330–339.
- Perretti M. & D'Acquisto F. (2009). Annexin A1 and glucocorticoids as effectors of the resolution of inflammation. *Nature Reviews Immunology*, 9, 62–70.
- Peters A., Delhey K., Andersson S., van Noordwijk H. & Forschler M.I. (2008). Condition-dependence of multiple carotenoid-based plumage traits: an experimental study. *Functional Ecology*, 22, 831–839.
- Prum R.O. & Williamson S. (2002). Reaction-diffusion models of within-feather pigmentation patterning. *Proceedings of the Royal Society B-Biological Sciences*, 269, 781–792.
- Pryke S.R. & Andersson S. (2005). Experimental evidence for female choice and energetic costs of male tail elongation in red-collared widowbirds. *Biological Journal of the Linnean Society*, 86, 35–43.
- Réale D., Reader S.M., Sol D., McDougall P.T. & Dingemanse N.J. (2007). Integrating animal temperament within ecology and evolution. *Biological Reviews*, 82, 291–318.
- Remage-Healey L. & Romero L.M. (2000). Daily and Seasonal Variation in Response to Stress in Captive Starlings (*Sturnus Vulgaris*): glucose. *General and Comparative Endocrinology*, 119, 60–68.
- Rich E.L. & Romero L.M. (2005). Exposure to chronic stress downregulates corticosterone responses to acute stressors. *American Journal of Physiology-Regulatory Integrative and Comparative Physiology*, 288, R1628–R1636.
- Ricklefs R.E. & Wikelski M. (2002). The physiology/life-history nexus. *Trends in Ecology & Evolution*, 17, 462–468.
- Romero L.M. (2004). Physiological stress in ecology: lessons from biomedical research. *Trends in Ecology & Evolution*, 19, 249–255.
- Romero L.M., Dickens M.J. & Cyr N.E. (2009). The reactive scope model – a new model integrating homeostasis, allostasis, and stress. *Hormones and Behavior*, 55, 375–389.
- Romero L.M. & Reed J.M. (2005). Collecting baseline corticosterone samples in the field: is under 3 min good enough? *Comparative Biochemistry and Physiology-Part A: Molecular & Integrative Physiology*, 140, 73–79.
- Romero L.M., Strohlic D. & Wingfield J.C. (2005). Corticosterone inhibits feather growth: potential mechanism explaining seasonal down regulation of corticosterone during molt. *Comparative Biochemistry and Physiology – A Molecular and Integrative Physiology*, 142, 65–73.
- Roulin A. (2004). The evolution, maintenance and adaptive function of genetic colour polymorphism in birds. *Biological Reviews*, 79, 815–848.



- Roulin A. & Altwegg R. (2007). Breeding rate is associated with pheomelanism in male and with eumelanism in female barn owls. *Behavioral Ecology*, 18, 563–570.
- Saino N., Romano M., Rubolini D., Ambrosini R., Caprioli M., Milzani A., Costanzo A., Colombo G., Canova L. & Wakamatsu K. (2013). Viability is associated with melanin-based coloration in the barn swallow (*Hirundo rustica*). *PLoS ONE*, 8, e60426.
- Saint-Dizier H., Leterrier C., Levy F. & Richard S. (2008). Selection for tonic immobility duration does not affect the response to novelty in quail. *Applied Animal Behaviour Science*, 112, 297–306.
- Saks L., Karu U., Ots I. & Hõrak P. (2006). Do standard measures of immunocompetence reflect parasite resistance? The case of greenfinch coccidiosis. *Functional Ecology*, 20, 75–82.
- Samas P., Polacikova L., Hauber M.E., Cassey P. & Grim T. (2012). Egg rejection behavior and clutch characteristics of the european greenfinch introduced to New Zealand. *Chinese Birds*, 3, 330–338.
- Sapolsky R.M., Romero L.M. & Munck A.U. (2000). How do glucocorticoids influence stress responses? Integrating permissive, suppressive, stimulatory, and preparative actions. *Endocrine Reviews*, 21, 55–89.
- Sarv T. & Hõrak P. (2009). Phytohaemagglutinin injection has a long-lasting effect on immune cells. *Journal of Avian Biology*, 40, 569–571.
- Schulenburg H., Kurtz J., Moret Y. & Siva-Jothy M.T. (2009). Introduction. Ecological immunology. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 364, 3–14.
- Segerstrom S.C. (2007). Stress, energy, and immunity: An ecological view. *Current Directions in Psychological Science*, 16, 326–330.
- Sepp T., Karu U., Blount J.D., Sild E., Männiste M. & Hõrak P. (2012). Coccidian Infection Causes Oxidative Damage in Greenfinches. *PLoS ONE*, 7, e36495.
- Sepp T., Karu U., Sild E., Männiste M. & Hõrak P. (2011). Effects of carotenoids, immune activation and immune suppression on the intensity of chronic coccidiosis in greenfinches. *Experimental Parasitology*, 127, 651–657.
- Sepp T., Sild E. & Hõrak P. (2010). Hematological condition indexes in greenfinches: effects of captivity and diurnal variation. *Physiological and Biochemical Zoology*, 83, 276–282.
- Sih A., Bell A. & Johnson J.C. (2004). Behavioral syndromes: an ecological and evolutionary overview. *Trends in Ecology & Evolution*, 19, 372–378.
- Sild E., Sepp T. & Hõrak P. (2011). Behavioural trait covaries with immune responsiveness in a wild passerine. *Brain Behavior and Immunity*, 25, 1349–1354.
- Simons M.J.P., Briga M., Koetsier E., Folkertsma R., Wubs M.D., Dijkstra C. & Verhulst S. (2012). Bill redness is positively associated with reproduction and survival in male and female zebra finches. *PLoS ONE*, 7, e40721.
- Stettenheim P.R. (2000). The integumentary morphology of modern birds – an overview. *American Zoologist*, 40, 461–477.
- Strochlic D.E. & Romero L.M. (2008). The effects of chronic psychological and physical stress on feather replacement in European starlings (*Sturnus vulgaris*). *Comparative Biochemistry and Physiology – A Molecular and Integrative Physiology*, 149, 68–79.
- Swaddle J.P., Witter M.S., Cuthill I.C., Budden A. & McCowen P. (1996). Plumage condition affects flight performance in common starlings: implications for

- developmental homeostasis, abrasion and moult. *Journal of Avian Biology*, 27, 103–111.
- Svensson P. & Wong B. (2011). Carotenoid-based signals in behavioural ecology: a review. *Behaviour*, 148, 131–189.
- Tella J.L., Lemus J.A., Carrete M. & Blanco G. (2008). The PHA test reflects acquired T-cell mediated immunocompetence in birds. *PLoS ONE*, 3, e3295.
- Telleria J.L., de la Hera I. & Perez-Tris J. (2013). Morphological variation as a tool for monitoring bird populations: a review. *Ardeola*, 60, 191–224.
- Van Oers K., Drent P.J., De Goede P. & Van Noordwijk A.J. (2004). Realized heritability and repeatability of risk-taking behaviour in relation to avian personalities. *Proceedings of the Royal Society B: Biological Sciences*, 271, 65–73.
- Van Oort H. & Dawson R.D. (2005). Carotenoid ornamentation of adult male common redpolls predicts probability of dying in a salmonellosis outbreak. *Functional Ecology*, 19, 822–827.
- Wingfield J.C. & Kitaysky A.S. (2002). Endocrine responses to unpredictable environmental events: stress or anti-stress hormones? *Integrative and Comparative Biology*, 42, 600–609.
- Wingfield J.C. & Sapolsky R.M. (2003). Reproduction and resistance to stress: when and how. *Journal of Neuroendocrinology*, 15, 711–724.
- Vinkler M., Svobodová J., Gabrielová B., Bainová H. & Bryjová A. (2014). Cytokine expression in phytohaemagglutinin-induced skin inflammation in a galliform bird. *Journal of Avian Biology*, 45, 43–50.
- Xie H., Rath N., Huff G., Huff W. & Balog J. (2000). Effects of *Salmonella typhimurium* lipopolysaccharide on broiler chickens. *Poultry Science*, 79, 33–40.

## CURRICULUM VITAE

**Name:** Marju Männiste  
**Date of birth:** 14.11.1985  
**Citizenship:** Estonian  
**Phone:** (+372) 5122085  
**Contact:** Department of Zoology, Institute of Ecology and Earth Sciences, Vanemuise 46, Tartu, 51014, Estonia  
**E-mail:** marju.manniste@ut.ee

**Education:** Röpina Ühisgümnaasium, 2004  
University of Tartu, bachelor's degree (Biology), 2008  
University of Tartu, master's degree (Zoology and Hydrobiology), 2010

**Language skills:** Estonian (native), English (fluent), German (advanced), Russian (beginner)

### Employment history:

2014 AS BIT, Publishing house Avita, editor in the field of biology

### Honours and Awards:

I prize in Contest for Students' Scientific Research, organised by Estonian Ministry of Education, 2011  
Scholarship of Lydia and Felix Krabi, 2013

**Research interests:** immunoecology, animal coloration, feathers, behavior

### Publications:

Sild E., Meitern R., **Männiste M.**, Karu U., Hõrak P. (2014). High feather corticosterone indicates better coccidian infection resistance in greenfinches. *General and comparative Endocrinology*, 204, 203–210.

Sepp T., **Männiste M.**, Kaasik A., Hõrak P. (2014). Multidimensionality of fear in captive greenfinches (*Carduelis chloris*). *Behavioral Ecology and Sociobiology*, 68, 1173–1181.

**Männiste M.** & Hõrak P. (2014). Emerging infectious disease selects for darker plumage coloration in greenfinches. *Frontiers in Ecology and Evolution – Behavioral and Evolutionary Ecology*, 2:4, doi: 10.3389/fevo.2014.00004.

Hõrak P., **Männiste M.**, Meitern R., Sild E., Saks L. & Sepp T. (2013). Dexamethasone inhibits corticosterone deposition in feathers of greenfinches. *General and Comparative Endocrinology*, 191, 210–214.

Meitern R., Sild E., Lind M.-A., **Männiste M.**, Sepp T., Karu U. & Hõrak P. (2013). Effects of Endotoxin and Psychological Stress on Redox Physiology,

- Immunity and Feather Corticosterone in Greenfinches. *PLOS ONE*, 8, e67545.
- Männiste M.**, Sepp T. & Hõrak P. (2013). Locomotor activity of captive greenfinches involves two different behavioural traits. *Ethology*, 581–591.
- Sepp T., Karu U., Blount J.D., Sild E., **Männiste M.** & Hõrak P. (2012). Coccidian infection causes oxidative damage in greenfinches. *PLoS ONE*, 7, e36495.
- Sepp T., Sild E., Blount J.D., **Männiste M.**, Karu U. & Hõrak P. (2012). Individual consistency and covariation of measures of oxidative status in greenfinches. *Physiological and Biochemical Zoology*, 85, 299–307.
- Männiste M.** & Hõrak P. (2011). Effects of immune activation and glucocorticoid administration on feather growth in greenfinches. *Journal of Experimental Zoology Part A: Ecological Genetics and Physiology*, 315A, 527–535.
- Sepp T., Karu U., Sild E., **Männiste M.** & Hõrak P. (2011). Effects of carotenoids, immune activation and immune suppression on the intensity of chronic coccidiosis in greenfinches. *Experimental Parasitology*, 127, 651–657.
- Sild E., Sepp T., **Männiste M.** & Hõrak P. (2011). Carotenoid intake does not affect immune-stimulated oxidative burst in greenfinches. *The Journal of Experimental Biology*, 214, 3467–3473.

#### **Conference theses:**

- Männiste M., Hõrak P., poster presentation: “Surm heledatele vintidele: trihho-monoos tapab värvi järgi”, XII Ecology conference “Eesti teadus- ja arendustegevuse konkurentsivõime”, 18–19 october, Tartu, Estonia.
- Männiste M., Sepp T., Hõrak P., poster presentation: “The effects of environmental enrichment on feather corticosterone content in captive wild-caught birds”, Conference “Behaviour”, 4–8 August 2013, Newcastle Gateshead, Great Britain.
- Männiste M., Sild E., Karu U., Meitern R., Hõrak P. poster presentation: „Can feather corticosterone predict behavior under stressful conditions?“, Conference on Behavioural Biology to celebrate the 20th anniversary of the Netherlands Society for Behavioural Biology (NVG), 28–30 November 2012, Soesterberg, Netherlands.
- Männiste M., oral presentation: „Behavior and plumage quality in greenfinches“, International Society for Behavioral Ecology conference (ISBE 2012), 12–18 August 2012, Lund, Sweden.
- Männiste M., oral presentation: “Acute glucocorticoid administration suppresses feather growth in greenfinches”, 8th Conference of European Ornithology Union, 27–30 August 2011, Riga, Latvia.
- Männiste M., Hõrak P., poster presentation: “Effects of immune activation and glucocorticoid administration on feather quality in greenfinches”, Science Meeting “Thermadapt”, 28–30 April 2011, Tartu, Estonia.

Männiste M., oral presentation: “The phenotypic plasticity of eumelanic black coloration in greenfinches”, Autumn School “Phenotypic plasticity”, 29–31 October 2011, Otepää, Estonia.

**Review work:**

Reviewed manuscripts for journals *Ornis Fennica*, *Animal Biology*, *Journal of Ornithology*, *The Condor*, *Physiological and Biochemical Zoology*

## ELULOOKIRJELDUS

- Nimi:** Marju Männiste  
**Sünniaeg:** 14.11.1985  
**Kodakondsus:** eesti  
**Telefon:** (+372) 5122085  
**Kontakt:** Zooloogia osakond, Ökoloogia ja Maateaduste Instituut, Vanemuise 46, Tartu, 51014, Eesti  
**E-post:** marju.manniste@ut.ee
- Haridus:** Räpina Ühisgümnaasium, 2004  
Tartu Ülikool, loodusteaduse bakalaureuse kraad, (bioloogia), 2008  
Tartu Ülikool, loodusteaduse magistri kraad (zooloogia ja hüdrobioloogia), 2010
- Keelteoskus:** eesti keel (emakeel), inglise keel (väga hea), saksa keel (kesktase), vene keel (algtase)
- Töökogemus:**  
2014 AS BIT, Kirjastus Avita, Bioloogia-alase õppekirjanduse toimetaja

### Teaduspreemiad ja -tunnustused:

I preemia Eesti Vabariigi Haridus ja Teadusministeeriumi üliõpilaste teadustööde riiklikul konkursil, 2011  
Lydia ja Felix Krabi stipendium, 2013

**Peamised uurimisvaldkonnad:** immuunökoloogia, loomade värvus, suled, käitumine

### Publikatsioonide loetelu:

- Sild E., Meitern R., **Männiste M.**, Karu U., Hõrak P. (2014). High feather corticosterone indicates better coccidian infection resistance in greenfinches. *General and comparative Endocrinology*, 204, 203–210.
- Sepp T., **Männiste M.**, Kaasik A., Hõrak P. (2014). Multidimensionality of fear in captive greenfinches (*Carduelis chloris*). *Behavioral Ecology and Sociobiology*, 68, 1173–1181.
- Männiste M.** & Hõrak P. (2014). Emerging infectious disease selects for darker plumage coloration in greenfinches. *Frontiers in Ecology and Evolution – Behavioral and Evolutionary Ecology*, 2:4, doi: 10.3389/fevo.2014.00004.
- Hõrak P., **Männiste M.**, Meitern R., Sild E., Saks L. & Sepp T. (2013). Dexamethasone inhibits corticosterone deposition in feathers of greenfinches. *General and Comparative Endocrinology*, 191, 210–214.

- Meitern R., Sild E., Lind M.-A., **Männiste M.**, Sepp T., Karu U. & Hõrak P. (2013). Effects of Endotoxin and Psychological Stress on Redox Physiology, Immunity and Feather Corticosterone in Greenfinches. *PLOS ONE*, 8, e67545.
- Männiste M.**, Sepp T. & Hõrak P. (2013). Locomotor activity of captive greenfinches involves two different behavioural traits. *Ethology*, 581–591.
- Sepp T., Karu U., Blount J.D., Sild E., **Männiste M.** & Hõrak P. (2012). Coccidian infection causes oxidative damage in greenfinches. *PLoS ONE*, 7, e36495.
- Sepp T., Sild E., Blount J.D., **Männiste M.**, Karu U. & Hõrak P. (2012). Individual consistency and covariation of measures of oxidative status in greenfinches. *Physiological and Biochemical Zoology*, 85, 299–307.
- Männiste M.** & Hõrak P. (2011). Effects of immune activation and glucocorticoid administration on feather growth in greenfinches. *Journal of Experimental Zoology Part A: Ecological Genetics and Physiology*, 315A, 527–535.
- Sepp T., Karu U., Sild E., **Männiste M.** & Hõrak P. (2011). Effects of carotenoids, immune activation and immune suppression on the intensity of chronic coccidiosis in greenfinches. *Experimental Parasitology*, 127, 651–657.
- Sild E., Sepp T., **Männiste M.** & Hõrak P. (2011). Carotenoid intake does not affect immune-stimulated oxidative burst in greenfinches. *The Journal of Experimental Biology*, 214, 3467–3473.

#### **Konverentside ettekanded:**

- Männiste M., Hõrak P., posterettekannet: „Surm heledatele vintidele: trihhomonoos tapab värvi järgi”, XII Ökoloogiakonverents “Eesti teadus- ja arendustegevuse konkurentsivõime”, 18–19 oktoober, Tartu, Eesti.
- Männiste M., Sepp T., Hõrak P., posterettekannet: “The effects of environmental enrichment on feather corticosterone content in captive wild-caught birds”, konverents “Behaviour”, 4–8 august 2013, Newcastle Gateshead, Suurbritannia.
- Männiste M., Sild E., Karu U., Meitern R., Hõrak P. posterettekannet: „Can feather corticosterone predict behavior under stressful conditions?”, Käitumisbioloogia konverents (NVG), 28–30 november 2012, Soesterberg, Holland.
- Männiste M., suuline ettekannet: „Behavior and plumage quality in greenfinches”, Rahvusvaheline käitumisökoloogia konverents (ISBE), 12–18 august 2012, Lund, Rootsi.
- Männiste M., suuline ettekannet: “Acute glucocorticoid administration suppresses feather growth in greenfinches”, 8. Euroopa Ornitoloogide ühingu konverents, 27–30 august 2011, Riia, Läti.

Männiste M., Hõrak P., posterettekannet: "Effects of immune activation and glucocorticoid administration on feather quality in greenfinches", Teaduskonverents "Thermadapt", 28–30 aprill 2011, Tartu, Eesti.

Männiste M., suuline ettekanne: "The phenotypic plasticity of eumelanic black coloration in greenfinches", Sügiskool "Phenotypic plasticity", 29–31 oktoober 2011, Otepää, Eesti.

**Muu teaduslik tegevus:**

Olen retsenseerinud teadusartikleid rahvusvahelistele teadusajakirjadele *Ornis Fennica*, *Animal Biology*, *Journal of Ornithology*, *The Condor*, *Physiological and Biochemical Zoology*



## DISSERTATIONES BIOLOGICAE UNIVERSITATIS TARTUENSIS

1. **Toivo Maimets.** Studies of human oncoprotein p53. Tartu, 1991, 96 p.
2. **Enn K. Seppet.** Thyroid state control over energy metabolism, ion transport and contractile functions in rat heart. Tartu, 1991, 135 p.
3. **Kristjan Zobel.** Epifüütsete makrosamblike väärtus õhu saastuse indikaatoritena Hamar-Dobani boreaalsetes mägimetsades. Tartu, 1992, 131 lk.
4. **Andres Mäe.** Conjugal mobilization of catabolic plasmids by transposable elements in helper plasmids. Tartu, 1992, 91 p.
5. **Maia Kivisaar.** Studies on phenol degradation genes of *Pseudomonas* sp. strain EST 1001. Tartu, 1992, 61 p.
6. **Allan Nurk.** Nucleotide sequences of phenol degradative genes from *Pseudomonas* sp. strain EST 1001 and their transcriptional activation in *Pseudomonas putida*. Tartu, 1992, 72 p.
7. **Ülo Tamm.** The genus *Populus* L. in Estonia: variation of the species biology and introduction. Tartu, 1993, 91 p.
8. **Jaanus Remme.** Studies on the peptidyltransferase centre of the *E.coli* ribosome. Tartu, 1993, 68 p.
9. **Ülo Langel.** Galanin and galanin antagonists. Tartu, 1993, 97 p.
10. **Arvo Käär.** The development of an automatic online dynamic fluorescence-based pH-dependent fiber optic penicillin flowthrough biosensor for the control of the benzylpenicillin hydrolysis. Tartu, 1993, 117 p.
11. **Lilian Järvekülg.** Antigenic analysis and development of sensitive immunoassay for potato viruses. Tartu, 1993, 147 p.
12. **Jaak Palumets.** Analysis of phytomass partition in Norway spruce. Tartu, 1993, 47 p.
13. **Arne Sellin.** Variation in hydraulic architecture of *Picea abies* (L.) Karst. trees grown under different environmental conditions. Tartu, 1994, 119 p.
13. **Mati Reeben.** Regulation of light neurofilament gene expression. Tartu, 1994, 108 p.
14. **Urmas Tartes.** Respiration rhythms in insects. Tartu, 1995, 109 p.
15. **Ülo Puurand.** The complete nucleotide sequence and infections *in vitro* transcripts from cloned cDNA of a potato A potyvirus. Tartu, 1995, 96 p.
16. **Peeter Hõrak.** Pathways of selection in avian reproduction: a functional framework and its application in the population study of the great tit (*Parus major*). Tartu, 1995, 118 p.
17. **Erkki Truve.** Studies on specific and broad spectrum virus resistance in transgenic plants. Tartu, 1996, 158 p.
18. **Illar Pata.** Cloning and characterization of human and mouse ribosomal protein S6-encoding genes. Tartu, 1996, 60 p.
19. **Ülo Niinemets.** Importance of structural features of leaves and canopy in determining species shade-tolerance in temperature deciduous woody taxa. Tartu, 1996, 150 p.

20. **Ants Kurg.** Bovine leukemia virus: molecular studies on the packaging region and DNA diagnostics in cattle. Tartu, 1996, 104 p.
21. **Ene Ustav.** E2 as the modulator of the BPV1 DNA replication. Tartu, 1996, 100 p.
22. **Aksel Soosaar.** Role of helix-loop-helix and nuclear hormone receptor transcription factors in neurogenesis. Tartu, 1996, 109 p.
23. **Maido Remm.** Human papillomavirus type 18: replication, transformation and gene expression. Tartu, 1997, 117 p.
24. **Tiiu Kull.** Population dynamics in *Cypripedium calceolus* L. Tartu, 1997, 124 p.
25. **Kalle Olli.** Evolutionary life-strategies of autotrophic planktonic microorganisms in the Baltic Sea. Tartu, 1997, 180 p.
26. **Meelis Pärtel.** Species diversity and community dynamics in calcareous grassland communities in Western Estonia. Tartu, 1997, 124 p.
27. **Malle Leht.** The Genus *Potentilla* L. in Estonia, Latvia and Lithuania: distribution, morphology and taxonomy. Tartu, 1997, 186 p.
28. **Tanel Tenson.** Ribosomes, peptides and antibiotic resistance. Tartu, 1997, 80 p.
29. **Arvo Tuvikene.** Assessment of inland water pollution using biomarker responses in fish *in vivo* and *in vitro*. Tartu, 1997, 160 p.
30. **Urmas Saarma.** Tuning ribosomal elongation cycle by mutagenesis of 23S rRNA. Tartu, 1997, 134 p.
31. **Henn Ojaveer.** Composition and dynamics of fish stocks in the gulf of Riga ecosystem. Tartu, 1997, 138 p.
32. **Lembi Lõugas.** Post-glacial development of vertebrate fauna in Estonian water bodies. Tartu, 1997, 138 p.
33. **Margus Pooga.** Cell penetrating peptide, transportan, and its predecessors, galanin-based chimeric peptides. Tartu, 1998, 110 p.
34. **Andres Saag.** Evolutionary relationships in some cetrarioid genera (Lichenized Ascomycota). Tartu, 1998, 196 p.
35. **Aivar Liiv.** Ribosomal large subunit assembly *in vivo*. Tartu, 1998, 158 p.
36. **Tatjana Oja.** Isoenzyme diversity and phylogenetic affinities among the eurasian annual bromes (*Bromus* L., Poaceae). Tartu, 1998, 92 p.
37. **Mari Moora.** The influence of arbuscular mycorrhizal (AM) symbiosis on the competition and coexistence of calcareous grassland plant species. Tartu, 1998, 78 p.
38. **Olavi Kurina.** Fungus gnats in Estonia (*Diptera: Bolitophilidae, Keroplattidae, Macroceridae, Ditomyiidae, Diadocidiidae, Mycetophilidae*). Tartu, 1998, 200 p.
39. **Andrus Tasa.** Biological leaching of shales: black shale and oil shale. Tartu, 1998, 98 p.
40. **Arnold Kristjuhan.** Studies on transcriptional activator properties of tumor suppressor protein p53. Tartu, 1998, 86 p.
41. **Sulev Ingerpuu.** Characterization of some human myeloid cell surface and nuclear differentiation antigens. Tartu, 1998, 163 p.

42. **Veljo Kisand.** Responses of planktonic bacteria to the abiotic and biotic factors in the shallow lake Võrtsjärv. Tartu, 1998, 118 p.
43. **Kadri Põldmaa.** Studies in the systematics of hypomyces and allied genera (Hypocreales, Ascomycota). Tartu, 1998, 178 p.
44. **Markus Vetemaa.** Reproduction parameters of fish as indicators in environmental monitoring. Tartu, 1998, 117 p.
45. **Heli Talvik.** Prepatent periods and species composition of different *Oesophagostomum* spp. populations in Estonia and Denmark. Tartu, 1998, 104 p.
46. **Katrin Heinsoo.** Cuticular and stomatal antechamber conductance to water vapour diffusion in *Picea abies* (L.) karst. Tartu, 1999, 133 p.
47. **Tarmo Annilo.** Studies on mammalian ribosomal protein S7. Tartu, 1998, 77 p.
48. **Indrek Ots.** Health state indicies of reproducing great tits (*Parus major*): sources of variation and connections with life-history traits. Tartu, 1999, 117 p.
49. **Juan Jose Cantero.** Plant community diversity and habitat relationships in central Argentina grasslands. Tartu, 1999, 161 p.
50. **Rein Kalamees.** Seed bank, seed rain and community regeneration in Estonian calcareous grasslands. Tartu, 1999, 107 p.
51. **Sulev Kõks.** Cholecystokinin (CCK) — induced anxiety in rats: influence of environmental stimuli and involvement of endopioid mechanisms and erotonin. Tartu, 1999, 123 p.
52. **Ebe Sild.** Impact of increasing concentrations of O<sub>3</sub> and CO<sub>2</sub> on wheat, clover and pasture. Tartu, 1999, 123 p.
53. **Ljudmilla Timofejeva.** Electron microscopical analysis of the synaptone-mal complex formation in cereals. Tartu, 1999, 99 p.
54. **Andres Valkna.** Interactions of galanin receptor with ligands and G-proteins: studies with synthetic peptides. Tartu, 1999, 103 p.
55. **Taavi Virro.** Life cycles of planktonic rotifers in lake Peipsi. Tartu, 1999, 101 p.
56. **Ana Rebane.** Mammalian ribosomal protein S3a genes and intron-encoded small nucleolar RNAs U73 and U82. Tartu, 1999, 85 p.
57. **Tiina Tamm.** Cocksfoot mottle virus: the genome organisation and transla-tional strategies. Tartu, 2000, 101 p.
58. **Reet Kurg.** Structure-function relationship of the bovine papilloma virus E2 protein. Tartu, 2000, 89 p.
59. **Toomas Kivisild.** The origins of Southern and Western Eurasian popula-tions: an mtDNA study. Tartu, 2000, 121 p.
60. **Niilo Kaldalu.** Studies of the TOL plasmid transcription factor XylS. Tartu 2000. 88 p.
61. **Dina Lepik.** Modulation of viral DNA replication by tumor suppressor protein p53. Tartu 2000. 106 p.
62. **Kai Vellak.** Influence of different factors on the diversity of the bryophyte vegetation in forest and wooded meadow communities. Tartu 2000. 122 p.

63. **Jonne Kotta.** Impact of eutrophication and biological invasions on the structure and functions of benthic macrofauna. Tartu 2000. 160 p.
64. **Georg Martin.** Phytobenthic communities of the Gulf of Riga and the inner sea the West-Estonian archipelago. Tartu, 2000. 139 p.
65. **Silvia Sepp.** Morphological and genetical variation of *Alchemilla L.* in Estonia. Tartu, 2000. 124 p.
66. **Jaan Liira.** On the determinants of structure and diversity in herbaceous plant communities. Tartu, 2000. 96 p.
67. **Priit Zingel.** The role of planktonic ciliates in lake ecosystems. Tartu 2001. 111 p.
68. **Tiit Teder.** Direct and indirect effects in Host-parasitoid interactions: ecological and evolutionary consequences. Tartu 2001. 122 p.
69. **Hannes Kollist.** Leaf apoplastic ascorbate as ozone scavenger and its transport across the plasma membrane. Tartu 2001. 80 p.
70. **Reet Marits.** Role of two-component regulator system PehR-PehS and extracellular protease PrtW in virulence of *Erwinia Carotovora* subsp. *Carotovora*. Tartu 2001. 112 p.
71. **Vallo Tilgar.** Effect of calcium supplementation on reproductive performance of the pied flycatcher *Ficedula hypoleuca* and the great tit *Parus major*, breeding in Northern temperate forests. Tartu, 2002. 126 p.
72. **Rita Hõrak.** Regulation of transposition of transposon Tn4652 in *Pseudomonas putida*. Tartu, 2002. 108 p.
73. **Liina Eek-Piirsoo.** The effect of fertilization, mowing and additional illumination on the structure of a species-rich grassland community. Tartu, 2002. 74 p.
74. **Krõõt Aasamaa.** Shoot hydraulic conductance and stomatal conductance of six temperate deciduous tree species. Tartu, 2002. 110 p.
75. **Nele Ingerpuu.** Bryophyte diversity and vascular plants. Tartu, 2002. 112 p.
76. **Neeme Tõnisson.** Mutation detection by primer extension on oligonucleotide microarrays. Tartu, 2002. 124 p.
77. **Margus Pensa.** Variation in needle retention of Scots pine in relation to leaf morphology, nitrogen conservation and tree age. Tartu, 2003. 110 p.
78. **Asko Lõhmus.** Habitat preferences and quality for birds of prey: from principles to applications. Tartu, 2003. 168 p.
79. **Viljar Jaks.** p53 — a switch in cellular circuit. Tartu, 2003. 160 p.
80. **Jaana Männik.** Characterization and genetic studies of four ATP-binding cassette (ABC) transporters. Tartu, 2003. 140 p.
81. **Marek Sammul.** Competition and coexistence of clonal plants in relation to productivity. Tartu, 2003. 159 p.
82. **Ivar Iives.** Virus-cell interactions in the replication cycle of bovine papillomavirus type 1. Tartu, 2003. 89 p.
83. **Andres Männik.** Design and characterization of a novel vector system based on the stable replicator of bovine papillomavirus type 1. Tartu, 2003. 109 p.

84. **Ivika Ostonen.** Fine root structure, dynamics and proportion in net primary production of Norway spruce forest ecosystem in relation to site conditions. Tartu, 2003. 158 p.
85. **Gudrun Veldre.** Somatic status of 12–15-year-old Tartu schoolchildren. Tartu, 2003. 199 p.
86. **Ülo Väli.** The greater spotted eagle *Aquila clanga* and the lesser spotted eagle *A. pomarina*: taxonomy, phylogeography and ecology. Tartu, 2004. 159 p.
87. **Aare Abroi.** The determinants for the native activities of the bovine papillomavirus type 1 E2 protein are separable. Tartu, 2004. 135 p.
88. **Tiina Kahre.** Cystic fibrosis in Estonia. Tartu, 2004. 116 p.
89. **Helen Orav-Kotta.** Habitat choice and feeding activity of benthic suspension feeders and mesograzers in the northern Baltic Sea. Tartu, 2004. 117 p.
90. **Maarja Öpik.** Diversity of arbuscular mycorrhizal fungi in the roots of perennial plants and their effect on plant performance. Tartu, 2004. 175 p.
91. **Kadri Tali.** Species structure of *Neotinea ustulata*. Tartu, 2004. 109 p.
92. **Kristiina Tambets.** Towards the understanding of post-glacial spread of human mitochondrial DNA haplogroups in Europe and beyond: a phylogeographic approach. Tartu, 2004. 163 p.
93. **Arvi Jõers.** Regulation of p53-dependent transcription. Tartu, 2004. 103 p.
94. **Lilian Kadaja.** Studies on modulation of the activity of tumor suppressor protein p53. Tartu, 2004. 103 p.
95. **Jaak Truu.** Oil shale industry wastewater: impact on river microbial community and possibilities for bioremediation. Tartu, 2004. 128 p.
96. **Maire Peters.** Natural horizontal transfer of the *pheBA* operon. Tartu, 2004. 105 p.
97. **Ülo Maiväli.** Studies on the structure-function relationship of the bacterial ribosome. Tartu, 2004. 130 p.
98. **Merit Otsus.** Plant community regeneration and species diversity in dry calcareous grasslands. Tartu, 2004. 103 p.
99. **Mikk Heidema.** Systematic studies on sawflies of the genera *Dolerus*, *Empria*, and *Caliroa* (Hymenoptera: Tenthredinidae). Tartu, 2004. 167 p.
100. **Ilmar Tõnno.** The impact of nitrogen and phosphorus concentration and N/P ratio on cyanobacterial dominance and N<sub>2</sub> fixation in some Estonian lakes. Tartu, 2004. 111 p.
101. **Lauri Saks.** Immune function, parasites, and carotenoid-based ornaments in greenfinches. Tartu, 2004. 144 p.
102. **Siiri Rootsi.** Human Y-chromosomal variation in European populations. Tartu, 2004. 142 p.
103. **Eve Vedler.** Structure of the 2,4-dichloro-phenoxyacetic acid-degradative plasmid pEST4011. Tartu, 2005. 106 p.
104. **Andres Tover.** Regulation of transcription of the phenol degradation *pheBA* operon in *Pseudomonas putida*. Tartu, 2005. 126 p.
105. **Helen Udras.** Hexose kinases and glucose transport in the yeast *Hansenula polymorpha*. Tartu, 2005. 100 p.

106. **Ave Suija.** Lichens and lichenicolous fungi in Estonia: diversity, distribution patterns, taxonomy. Tartu, 2005. 162 p.
107. **Piret Lõhmus.** Forest lichens and their substrata in Estonia. Tartu, 2005. 162 p.
108. **Inga Lips.** Abiotic factors controlling the cyanobacterial bloom occurrence in the Gulf of Finland. Tartu, 2005. 156 p.
109. **Kaasik, Krista.** Circadian clock genes in mammalian clockwork, metabolism and behaviour. Tartu, 2005. 121 p.
110. **Juhan Javoiš.** The effects of experience on host acceptance in ovipositing moths. Tartu, 2005. 112 p.
111. **Tiina Sedman.** Characterization of the yeast *Saccharomyces cerevisiae* mitochondrial DNA helicase Hmi1. Tartu, 2005. 103 p.
112. **Ruth Aguraiuja.** Hawaiian endemic fern lineage *Diellia* (Aspleniaceae): distribution, population structure and ecology. Tartu, 2005. 112 p.
113. **Riho Teras.** Regulation of transcription from the fusion promoters generated by transposition of Tn4652 into the upstream region of *pheBA* operon in *Pseudomonas putida*. Tartu, 2005. 106 p.
114. **Mait Metspalu.** Through the course of prehistory in india: tracing the mtDNA trail. Tartu, 2005. 138 p.
115. **Elin Lõhmussaar.** The comparative patterns of linkage disequilibrium in European populations and its implication for genetic association studies. Tartu, 2006. 124 p.
116. **Priit Kupper.** Hydraulic and environmental limitations to leaf water relations in trees with respect to canopy position. Tartu, 2006. 126 p.
117. **Heili Ilves.** Stress-induced transposition of Tn4652 in *Pseudomonas Putida*. Tartu, 2006. 120 p.
118. **Silja Kuusk.** Biochemical properties of Hmi1p, a DNA helicase from *Saccharomyces cerevisiae* mitochondria. Tartu, 2006. 126 p.
119. **Kersti Püssa.** Forest edges on medium resolution landsat thematic mapper satellite images. Tartu, 2006. 90 p.
120. **Lea Tummeleht.** Physiological condition and immune function in great tits (*Parus major* L.): Sources of variation and trade-offs in relation to growth. Tartu, 2006. 94 p.
121. **Toomas Esperk.** Larval instar as a key element of insect growth schedules. Tartu, 2006. 186 p.
122. **Harri Valdmann.** Lynx (*Lynx lynx*) and wolf (*Canis lupus*) in the Baltic region: Diets, helminth parasites and genetic variation. Tartu, 2006. 102 p.
123. **Priit Jõers.** Studies of the mitochondrial helicase Hmi1p in *Candida albicans* and *Saccharomyces cerevisia*. Tartu, 2006. 113 p.
124. **Kersti Lilleväli.** Gata3 and Gata2 in inner ear development. Tartu, 2007. 123 p.
125. **Kai Rünk.** Comparative ecology of three fern species: *Dryopteris carthusiana* (Vill.) H.P. Fuchs, *D. expansa* (C. Presl) Fraser-Jenkins & Jermy and *D. dilatata* (Hoffm.) A. Gray (Dryopteridaceae). Tartu, 2007. 143 p.

126. **Aveliina Helm.** Formation and persistence of dry grassland diversity: role of human history and landscape structure. Tartu, 2007. 89 p.
127. **Leho Tedersoo.** Ectomycorrhizal fungi: diversity and community structure in Estonia, Seychelles and Australia. Tartu, 2007. 233 p.
128. **Marko Mägi.** The habitat-related variation of reproductive performance of great tits in a deciduous-coniferous forest mosaic: looking for causes and consequences. Tartu, 2007. 135 p.
129. **Valeria Lulla.** Replication strategies and applications of Semliki Forest virus. Tartu, 2007. 109 p.
130. **Ülle Reier.** Estonian threatened vascular plant species: causes of rarity and conservation. Tartu, 2007. 79 p.
131. **Inga Jüriado.** Diversity of lichen species in Estonia: influence of regional and local factors. Tartu, 2007. 171 p.
132. **Tatjana Krama.** Mobbing behaviour in birds: costs and reciprocity based cooperation. Tartu, 2007. 112 p.
133. **Signe Saumaa.** The role of DNA mismatch repair and oxidative DNA damage defense systems in avoidance of stationary phase mutations in *Pseudomonas putida*. Tartu, 2007. 172 p.
134. **Reedik Mägi.** The linkage disequilibrium and the selection of genetic markers for association studies in european populations. Tartu, 2007. 96 p.
135. **Priit Kilgas.** Blood parameters as indicators of physiological condition and skeletal development in great tits (*Parus major*): natural variation and application in the reproductive ecology of birds. Tartu, 2007. 129 p.
136. **Anu Albert.** The role of water salinity in structuring eastern Baltic coastal fish communities. Tartu, 2007. 95 p.
137. **Kärt Padari.** Protein transduction mechanisms of transportans. Tartu, 2008. 128 p.
138. **Siiri-Lii Sandre.** Selective forces on larval colouration in a moth. Tartu, 2008. 125 p.
139. **Ülle Jõgar.** Conservation and restoration of semi-natural floodplain meadows and their rare plant species. Tartu, 2008. 99 p.
140. **Lauri Laanisto.** Macroecological approach in vegetation science: generality of ecological relationships at the global scale. Tartu, 2008. 133 p.
141. **Reidar Andreson.** Methods and software for predicting PCR failure rate in large genomes. Tartu, 2008. 105 p.
142. **Birgot Paavel.** Bio-optical properties of turbid lakes. Tartu, 2008. 175 p.
143. **Kaire Torn.** Distribution and ecology of charophytes in the Baltic Sea. Tartu, 2008, 98 p.
144. **Vladimir Vimberg.** Peptide mediated macrolide resistance. Tartu, 2008, 190 p.
145. **Daima Örd.** Studies on the stress-inducible pseudokinase TRB3, a novel inhibitor of transcription factor ATF4. Tartu, 2008, 108 p.
146. **Lauri Saag.** Taxonomic and ecologic problems in the genus *Lepraria* (*Stereocaulaceae*, lichenised *Ascomycota*). Tartu, 2008, 175 p.

147. **Ulvi Karu.** Antioxidant protection, carotenoids and coccidians in greenfinches – assessment of the costs of immune activation and mechanisms of parasite resistance in a passerine with carotenoid-based ornaments. Tartu, 2008, 124 p.
148. **Jaanus Remm.** Tree-cavities in forests: density, characteristics and occupancy by animals. Tartu, 2008, 128 p.
149. **Epp Moks.** Tapeworm parasites *Echinococcus multilocularis* and *E. granulosus* in Estonia: phylogenetic relationships and occurrence in wild carnivores and ungulates. Tartu, 2008, 82 p.
150. **Eve Eensalu.** Acclimation of stomatal structure and function in tree canopy: effect of light and CO<sub>2</sub> concentration. Tartu, 2008, 108 p.
151. **Janne Pullat.** Design, functionlization and application of an *in situ* synthesized oligonucleotide microarray. Tartu, 2008, 108 p.
152. **Marta Putrinš.** Responses of *Pseudomonas putida* to phenol-induced metabolic and stress signals. Tartu, 2008, 142 p.
153. **Marina Semtsenko.** Plant root behaviour: responses to neighbours and physical obstructions. Tartu, 2008, 106 p.
154. **Marge Starast.** Influence of cultivation techniques on productivity and fruit quality of some *Vaccinium* and *Rubus* taxa. Tartu, 2008, 154 p.
155. **Age Tats.** Sequence motifs influencing the efficiency of translation. Tartu, 2009, 104 p.
156. **Radi Tegova.** The role of specialized DNA polymerases in mutagenesis in *Pseudomonas putida*. Tartu, 2009, 124 p.
157. **Tsipe Aavik.** Plant species richness, composition and functional trait pattern in agricultural landscapes – the role of land use intensity and landscape structure. Tartu, 2009, 112 p.
158. **Kaja Kiiver.** Semliki forest virus based vectors and cell lines for studying the replication and interactions of alphaviruses and hepaciviruses. Tartu, 2009, 104 p.
159. **Meelis Kadaja.** Papillomavirus Replication Machinery Induces Genomic Instability in its Host Cell. Tartu, 2009, 126 p.
160. **Pille Hallast.** Human and chimpanzee Luteinizing hormone/Chorionic Gonadotropin beta (*LHB/CGB*) gene clusters: diversity and divergence of young duplicated genes. Tartu, 2009, 168 p.
161. **Ain Vellak.** Spatial and temporal aspects of plant species conservation. Tartu, 2009, 86 p.
162. **Triinu Remmel.** Body size evolution in insects with different colouration strategies: the role of predation risk. Tartu, 2009, 168 p.
163. **Jaana Salujõe.** Zooplankton as the indicator of ecological quality and fish predation in lake ecosystems. Tartu, 2009, 129 p.
164. **Ele Vahtmäe.** Mapping benthic habitat with remote sensing in optically complex coastal environments. Tartu, 2009, 109 p.
165. **Liisa Metsamaa.** Model-based assessment to improve the use of remote sensing in recognition and quantitative mapping of cyanobacteria. Tartu, 2009, 114 p.



166. **Pille Säälük.** The role of endocytosis in the protein transduction by cell-penetrating peptides. Tartu, 2009, 155 p.
167. **Lauri Peil.** Ribosome assembly factors in *Escherichia coli*. Tartu, 2009, 147 p.
168. **Lea Hallik.** Generality and specificity in light harvesting, carbon gain capacity and shade tolerance among plant functional groups. Tartu, 2009, 99 p.
169. **Mariliis Tark.** Mutagenic potential of DNA damage repair and tolerance mechanisms under starvation stress. Tartu, 2009, 191 p.
170. **Riinu Rannap.** Impacts of habitat loss and restoration on amphibian populations. Tartu, 2009, 117 p.
171. **Maarja Adojaan.** Molecular variation of HIV-1 and the use of this knowledge in vaccine development. Tartu, 2009, 95 p.
172. **Signe Altmäe.** Genomics and transcriptomics of human induced ovarian folliculogenesis. Tartu, 2010, 179 p.
173. **Triin Suvi.** Mycorrhizal fungi of native and introduced trees in the Seychelles Islands. Tartu, 2010, 107 p.
174. **Velda Lauringson.** Role of suspension feeding in a brackish-water coastal sea. Tartu, 2010, 123 p.
175. **Eero Talts.** Photosynthetic cyclic electron transport – measurement and variably proton-coupled mechanism. Tartu, 2010, 121 p.
176. **Mari Nelis.** Genetic structure of the Estonian population and genetic distance from other populations of European descent. Tartu, 2010, 97 p.
177. **Kaarel Krjutškov.** Arrayed Primer Extension-2 as a multiplex PCR-based method for nucleic acid variation analysis: method and applications. Tartu, 2010, 129 p.
178. **Egle Köster.** Morphological and genetical variation within species complexes: *Anthyllis vulneraria* s. l. and *Alchemilla vulgaris* (coll.). Tartu, 2010, 101 p.
179. **Erki Õunap.** Systematic studies on the subfamily Sterrhinae (Lepidoptera: Geometridae). Tartu, 2010, 111 p.
180. **Merike Jõesaar.** Diversity of key catabolic genes at degradation of phenol and *p*-cresol in pseudomonads. Tartu, 2010, 125 p.
181. **Kristjan Herkül.** Effects of physical disturbance and habitat-modifying species on sediment properties and benthic communities in the northern Baltic Sea. Tartu, 2010, 123 p.
182. **Arto Pulk.** Studies on bacterial ribosomes by chemical modification approaches. Tartu, 2010, 161 p.
183. **Maria Põllupüü.** Ecological relations of cladocerans in a brackish-water ecosystem. Tartu, 2010, 126 p.
184. **Toomas Silla.** Study of the segregation mechanism of the Bovine Papillomavirus Type 1. Tartu, 2010, 188 p.
185. **Gyaneshwer Chaubey.** The demographic history of India: A perspective based on genetic evidence. Tartu, 2010, 184 p.
186. **Katrin Kepp.** Genes involved in cardiovascular traits: detection of genetic variation in Estonian and Czech populations. Tartu, 2010, 164 p.

187. **Virve Sõber.** The role of biotic interactions in plant reproductive performance. Tartu, 2010, 92 p.
188. **Kersti Kangro.** The response of phytoplankton community to the changes in nutrient loading. Tartu, 2010, 144 p.
189. **Joachim M. Gerhold.** Replication and Recombination of mitochondrial DNA in Yeast. Tartu, 2010, 120 p.
190. **Helen Tammert.** Ecological role of physiological and phylogenetic diversity in aquatic bacterial communities. Tartu, 2010, 140 p.
191. **Elle Rajandu.** Factors determining plant and lichen species diversity and composition in Estonian *Calamagrostis* and *Hepatica* site type forests. Tartu, 2010, 123 p.
192. **Paula Ann Kivistik.** ColR-ColS signalling system and transposition of Tn4652 in the adaptation of *Pseudomonas putida*. Tartu, 2010, 118 p.
193. **Siim Sõber.** Blood pressure genetics: from candidate genes to genome-wide association studies. Tartu, 2011, 120 p.
194. **Kalle Kipper.** Studies on the role of helix 69 of 23S rRNA in the factor-dependent stages of translation initiation, elongation, and termination. Tartu, 2011, 178 p.
195. **Triinu Siibak.** Effect of antibiotics on ribosome assembly is indirect. Tartu, 2011, 134 p.
196. **Tambet Tõnissoo.** Identification and molecular analysis of the role of guanine nucleotide exchange factor RIC-8 in mouse development and neural function. Tartu, 2011, 110 p.
197. **Helin Räägel.** Multiple faces of cell-penetrating peptides – their intracellular trafficking, stability and endosomal escape during protein transduction. Tartu, 2011, 161 p.
198. **Andres Jaanus.** Phytoplankton in Estonian coastal waters – variability, trends and response to environmental pressures. Tartu, 2011, 157 p.
199. **Tiit Nikopensius.** Genetic predisposition to nonsyndromic orofacial clefts. Tartu, 2011, 152 p.
200. **Signe Värvi.** Studies on the mechanisms of RNA polymerase II-dependent transcription elongation. Tartu, 2011, 108 p.
201. **Kristjan Vällk.** Gene expression profiling and genome-wide association studies of non-small cell lung cancer. Tartu, 2011, 98 p.
202. **Arno Põllumäe.** Spatio-temporal patterns of native and invasive zooplankton species under changing climate and eutrophication conditions. Tartu, 2011, 153 p.
203. **Egle Tammeleht.** Brown bear (*Ursus arctos*) population structure, demographic processes and variations in diet in northern Eurasia. Tartu, 2011, 143 p.
205. **Teele Jairus.** Species composition and host preference among ectomycorrhizal fungi in Australian and African ecosystems. Tartu, 2011, 106 p.
206. **Kessy Abarenkov.** PlutoF – cloud database and computing services supporting biological research. Tartu, 2011, 125 p.

207. **Marina Grigorova.** Fine-scale genetic variation of follicle-stimulating hormone beta-subunit coding gene (*FSHB*) and its association with reproductive health. Tartu, 2011, 184 p.
208. **Anu Tiitsaar.** The effects of predation risk and habitat history on butterfly communities. Tartu, 2011, 97 p.
209. **Elin Sild.** Oxidative defences in immunoeological context: validation and application of assays for nitric oxide production and oxidative burst in a wild passerine. Tartu, 2011, 105 p.
210. **Irja Saar.** The taxonomy and phylogeny of the genera *Cystoderma* and *Cystodermella* (Agaricales, Fungi). Tartu, 2012, 167 p.
211. **Pauli Saag.** Natural variation in plumage bacterial assemblages in two wild breeding passerines. Tartu, 2012, 113 p.
212. **Aleksei Lulla.** Alphaviral nonstructural protease and its polyprotein substrate: arrangements for the perfect marriage. Tartu, 2012, 143 p.
213. **Mari Järve.** Different genetic perspectives on human history in Europe and the Caucasus: the stories told by uniparental and autosomal markers. Tartu, 2012, 119 p.
214. **Ott Scheler.** The application of tmRNA as a marker molecule in bacterial diagnostics using microarray and biosensor technology. Tartu, 2012, 93 p.
215. **Anna Balikova.** Studies on the functions of tumor-associated mucin-like leukosialin (CD43) in human cancer cells. Tartu, 2012, 129 p.
216. **Triinu Kõressaar.** Improvement of PCR primer design for detection of prokaryotic species. Tartu, 2012, 83 p.
217. **Tuul Sepp.** Hematological health state indices of greenfinches: sources of individual variation and responses to immune system manipulation. Tartu, 2012, 117 p.
218. **Rya Ero.** Modifier view of the bacterial ribosome. Tartu, 2012, 146 p.
219. **Mohammad Bahram.** Biogeography of ectomycorrhizal fungi across different spatial scales. Tartu, 2012, 165 p.
220. **Anneli Lorents.** Overcoming the plasma membrane barrier: uptake of amphipathic cell-penetrating peptides induces influx of calcium ions and downstream responses. Tartu, 2012, 113 p.
221. **Katrin Männik.** Exploring the genomics of cognitive impairment: whole-genome SNP genotyping experience in Estonian patients and general population. Tartu, 2012, 171 p.
222. **Marko Prouš.** Taxonomy and phylogeny of the sawfly genus *Empria* (Hymenoptera, Tenthredinidae). Tartu, 2012, 192 p.
223. **Triinu Visnapuu.** Levansucrases encoded in the genome of *Pseudomonas syringae* pv. tomato DC3000: heterologous expression, biochemical characterization, mutational analysis and spectrum of polymerization products. Tartu, 2012, 160 p.
224. **Nele Tamberg.** Studies on Semliki Forest virus replication and pathogenesis. Tartu, 2012, 109 p.

225. **Tõnu Esko.** Novel applications of SNP array data in the analysis of the genetic structure of Europeans and in genetic association studies. Tartu, 2012, 149 p.
226. **Timo Arula.** Ecology of early life-history stages of herring *Clupea harengus membras* in the northeastern Baltic Sea. Tartu, 2012, 143 p.
227. **Inga Hiiesalu.** Belowground plant diversity and coexistence patterns in grassland ecosystems. Tartu, 2012, 130 p.
228. **Kadri Koorem.** The influence of abiotic and biotic factors on small-scale plant community patterns and regeneration in boreonemoral forest. Tartu, 2012, 114 p.
229. **Liis Andresen.** Regulation of virulence in plant-pathogenic pectobacteria. Tartu, 2012, 122 p.
230. **Kaupo Kohv.** The direct and indirect effects of management on boreal forest structure and field layer vegetation. Tartu, 2012, 124 p.
231. **Mart Jüssi.** Living on an edge: landlocked seals in changing climate. Tartu, 2012, 114 p.
232. **Riina Klais.** Phytoplankton trends in the Baltic Sea. Tartu, 2012, 136 p.
233. **Rauno Veeroja.** Effects of winter weather, population density and timing of reproduction on life-history traits and population dynamics of moose (*Alces alces*) in Estonia. Tartu, 2012, 92 p.
234. **Marju Keis.** Brown bear (*Ursus arctos*) phylogeography in northern Eurasia. Tartu, 2013, 142 p.
235. **Sergei Põlme.** Biogeography and ecology of *alnus*- associated ectomycorrhizal fungi – from regional to global scale. Tartu, 2013, 90 p.
236. **Liis Uusküla.** Placental gene expression in normal and complicated pregnancy. Tartu, 2013, 173 p.
237. **Marko Lõoke.** Studies on DNA replication initiation in *Saccharomyces cerevisiae*. Tartu, 2013, 112 p.
238. **Anne Aan.** Light- and nitrogen-use and biomass allocation along productivity gradients in multilayer plant communities. Tartu, 2013, 127 p.
239. **Heidi Tamm.** Comprehending phylogenetic diversity – case studies in three groups of ascomycetes. Tartu, 2013, 136 p.
240. **Liina Kangur.** High-Pressure Spectroscopy Study of Chromophore-Binding Hydrogen Bonds in Light-Harvesting Complexes of Photosynthetic Bacteria. Tartu, 2013, 150 p.
241. **Margus Leppik.** Substrate specificity of the multisite specific pseudouridine synthase RluD. Tartu, 2013, 111 p.
242. **Lauris Kaplinski.** The application of oligonucleotide hybridization model for PCR and microarray optimization. Tartu, 2013, 103 p.
243. **Merli Pärnoja.** Patterns of macrophyte distribution and productivity in coastal ecosystems: effect of abiotic and biotic forcing. Tartu, 2013, 155 p.
244. **Tõnu Margus.** Distribution and phylogeny of the bacterial translational GTPases and the Mqsr/YgiT regulatory system. Tartu, 2013, 126 p.
245. **Pille Mänd.** Light use capacity and carbon and nitrogen budget of plants: remote assessment and physiological determinants. Tartu, 2013, 128 p.

246. **Mario Plaas.** Animal model of Wolfram Syndrome in mice: behavioural, biochemical and psychopharmacological characterization. Tartu, 2013, 144 p.
247. **Georgi Hudjašov.** Maps of mitochondrial DNA, Y-chromosome and tyrosinase variation in Eurasian and Oceanian populations. Tartu, 2013, 115 p.
248. **Mari Lepik.** Plasticity to light in herbaceous plants and its importance for community structure and diversity. Tartu, 2013, 102 p.
249. **Ede Leppik.** Diversity of lichens in semi-natural habitats of Estonia. Tartu, 2013, 151 p.
250. **Ülle Saks.** Arbuscular mycorrhizal fungal diversity patterns in boreo-nemoral forest ecosystems. Tartu, 2013, 151 p.
251. **Eneli Oitmaa.** Development of arrayed primer extension microarray assays for molecular diagnostic applications. Tartu, 2013, 147 p.
252. **Jekaterina Jutkina.** The horizontal gene pool for aromatics degradation: bacterial catabolic plasmids of the Baltic Sea aquatic system. Tartu, 2013, 121 p.
253. **Helen Vellau.** Reaction norms for size and age at maturity in insects: rules and exceptions. Tartu, 2014, 132 p.
254. **Randel Kreitsberg.** Using biomarkers in assessment of environmental contamination in fish – new perspectives. Tartu, 2014, 107 p.
255. **Krista Takkis.** Changes in plant species richness and population performance in response to habitat loss and fragmentation. Tartu, 2014, 141 p.
256. **Liina Nagirnaja.** Global and fine-scale genetic determinants of recurrent pregnancy loss. Tartu, 2014, 211 p.
257. **Triin Triisberg.** Factors influencing the re-vegetation of abandoned extracted peatlands in Estonia. Tartu, 2014, 133 p.
258. **Villu Soon.** A phylogenetic revision of the *Chrysis ignita* species group (Hymenoptera: Chrysididae) with emphasis on the northern European fauna. Tartu, 2014, 211 p.
259. **Andrei Nikonov.** RNA-Dependent RNA Polymerase Activity as a Basis for the Detection of Positive-Strand RNA Viruses by Vertebrate Host Cells. Tartu, 2014, 207 p.
260. **Eele Õunapuu-Pikas.** Spatio-temporal variability of leaf hydraulic conductance in woody plants: ecophysiological consequences. Tartu, 2014, 135 p.