

KAIRI KÄIRO

Biological Quality According
to Macroinvertebrates in Streams
of Estonia (Baltic Ecoregion of Europe):
Effects of Human-Induced
Hydromorphological Changes



DISSERTATIONES BIOLOGICAE UNIVERSITATIS TARTUENSIS

289

KAIRI KÄIRO

Biological Quality According
to Macroinvertebrates in Streams
of Estonia (Baltic Ecoregion of Europe):
Effects of Human-Induced
Hydromorphological Changes



Department of Zoology, Institute of Ecology and Earth Sciences,
Faculty of Science and Technology, University of Tartu, Estonia

Dissertation was accepted for the commencement of the degree of *Doctor philosophiae* in Hydrobiology at the University of Tartu on 21 March, 2016 by the Scientific Council of the Institute of Ecology and Earth Sciences, University of Tartu.

Supervisors: PhD Taavi Virro, University of Tartu, Estonia
PhD Henn Timm, Estonian University of Life Sciences,
Tartu, Estonia

Opponent: PhD Kęstutis Arbačiauskas, Nature Research Centre,
Vilnius, Lithuania

Commencement: Room 301, 46 Vanemuise Street, Tartu, on June 3, 2016 at
10.15 a.m.

The study was carried out at the Estonian University of Life Sciences, Centre for Limnology.

Publication of this thesis is granted by the Institute of Ecology and Earth Sciences, University of Tartu

ISSN 1024-6479
ISBN 978-9949-77-077-9 (print)
ISBN 978-9949-77-078-6 (pdf)

Copyright: Kairi Käiro, 2016

University of Tartu Press
www.tyk.ee

CONTENTS

LIST OF ORIGINAL PUBLICATIONS	6
1. INTRODUCTION.....	7
2. AIMS.....	10
3. MATERIAL AND METHODS	11
3.1. Study area and sampling sites	11
3.2. Sampling	11
3.3. Estimation of biological quality	13
3.4. Data analysis	14
4. RESULTS AND DISCUSSION	16
4.1. Taxonomic composition of macroinvertebrates.....	16
4.1.1. Effects of damming	16
4.1.2. Effects of channelization	16
4.2. Biological, hydromorphological and hydrochemical quality	17
5. CONCLUSIONS.....	21
SUMMARY	22
SUMMARY IN ESTONIAN	23
ACKNOWLEDGEMENTS	25
REFERENCES.....	26
APPENDIX.....	31
PUBLICATIONS.....	33
CURRICULUM VITAE	108
ELULOOKIRJELDUS.....	110

LIST OF ORIGINAL PUBLICATIONS

The thesis is based on the following papers referred to in the text by their Roman numerals:

- I.** Käiro, K., Möls, T., Timm, H., Virro, T. & Järvekülg, R. 2011. The effect of damming on biological quality according to macroinvertebrates in some Estonian streams, Central – Baltic Europe: a pilot study. *River Research and Applications* 27: 895–907.
- II.** Timm, H., Käiro, K., Möls, T. & Virro, T. 2011. An index to assess hydromorphological quality of Estonian surface waters based on macroinvertebrate taxonomic composition. *Limnologica* 41: 398–410.
- III.** Käiro, K., Timm, H., Haldna, M. & Virro, T. 2012. Biological quality on the basis of macroinvertebrates in dammed habitats of some Estonian streams, Central – Baltic Europe. *International Review of Hydrobiology* 97: 497–508.
- IV.** Käiro, K., Timm, H., Haldna, M. & Virro, T. (submitted to *Hydrobiologia*). The effect of channelization on the biological quality of lowland streams according to macroinvertebrates.

The published papers are reproduced with permission from the publishers: John Wiley & Sons, Inc. (papers I and III), Elsevier B.V. (paper II) and Springer Science+Business Media (paper IV).

The participation of the author in preparation of the listed publications is as follows:

	Paper I	Paper II	Paper III	Paper IV
Idea and design	HT, KK, RJ	HT, KK	KK, HT	KK, HT
Data collection	KK, HT	KK, HT	KK, HT	KK, HT
Data analysis	TM	TM	KK, MH	KK, MH
Manuscript preparation	KK, HT, TM, TV	HT, KK, TM, TV	KK, HT, MH, TV	KK, HT, MH, TV

KK – Kairi Käiro, HT – Henn Timm, TV – Taavi Virro, TM – Tõnu Möls, MH – Marina Haldna, RJ – Rein Järvekülg.

1. INTRODUCTION

As defined in the Water Framework Directive (2002), hydromorphology is the physical characteristics of the shape, boundaries and content of a water body. Alteration of hydromorphology by human activities (e.g. damming, channelization, urbanization, drainage) poses a significant threat to habitat quality and in turn to riverine communities (Poff et al., 1997; Bunn & Arthington, 2002; Santucci et al., 2005; Horsák et al., 2009; Navarro-Llácer et al., 2010; Kennedy & Turner, 2011). The most conspicuous human influence modifying flow velocity and hydromorphological conditions in running waters is damming. In this process part of the lotic ecosystem is transformed into lentic one. The direct impacts of damming are reduction in river flow and increase of sedimentation upstream of the dam (Petts, 1984; Poff & Zimmerman, 2010). At the same time, water velocity increases immediately downstream. Dams can also be formed as a result of the action of natural geological forces or the activity of beavers, which will cause significant changes in the stream biota, particularly upstream (Silliman et al., 2007; Arndt & Domdei, 2012; Pliūraite & Kesminas, 2012). The effects of dams on the biota depend on the size, age and construction type of dams. Below-dam water often has different temperature and hydrochemical characteristics, compared to normal lotic conditions. If water is released from the near-bottom layer of the reservoir, it is usually cold and oxygen-depleted but nutrient-rich. On the contrary, the water released from the near-surface layer has usually higher temperature than the water downstream (Petts, 1984; Lessard & Hayes, 2003). A well-known issue is the negative effect of dams on migratory fishes. Impoundments can block the way upstream and thus contribute to the decline and even extinction of species that depend on longitudinal movements (Porto et al., 1999; Liermann et al., 2012). Less is known about how the stream biota is affected by changed water velocity and bottom's character.

Channelization is another worldwide threat to stream biodiversity (Rosenberg et al. 2000; Nakamura & Yamada, 2005; Horsák et al., 2009). It results in a temporary increase of flow velocity and flooding, while reduced discharge occurs most of the time. Natural pool-riffle sequences, which provide shelter for fish and macroinvertebrates, are lost. These streams have little or no stable structure in the channel, have few or no meanders or pools, and have no native riparian vegetation to shade the channel and buffer the stream from adjacent human activities (Maxted et al., 2000; Nakano et al., 2008). The effects of channelization on the stream biota depend on the total length of modified reaches, on the proportion of unmodified reaches and on recovery time. The stream biota may recover more slowly in impounded areas than in channelized areas, due to complete loss of normal flow (Hortle & Lake, 1982; Smiley & Dibble, 2008). Channelization may also have long-term negative effects due to decreased habitat diversity (Horsák et al., 2009).

Excessive **water abstraction** from streams, widespread in areas with a warm and dry climate, has much less importance in cooler regions, compared to

damming and channelization. The possible effects of it were not estimated in this study.

Benthic macroinvertebrates are small bottom-living animals, visible by naked eye. Their taxonomic composition has been widely used as an indicator of running water pollution by organic, acidic or toxic pollutants (Rosenberg & Resh, 1993). In comparison to fishes or macrophytes, macroinvertebrates stand out with high species number (e.g. > 400 bioindicators in Estonian waterbodies; Timm & Vilbaste, 2010) and are easy to collect. They are widely distributed (occur also in canopied areas, underground and upstream of steep barriers). Therefore, the probability to obtain erroneous results because of the incidental absence of some species is relatively low. Macroinvertebrates are by far the most commonly used group for use in assessing freshwater quality (Rosenberg & Resh, 1993; Cranston et al., 1996). They are also considered a necessary measure of biological quality of freshwaters according to the Water Framework Directive (2002). As damming has a significant negative effect on the chemical and physical elements of streams, it consequently leads to changes in the macroinvertebrate community (Pozo et al., 1997; Ahearn et al., 2005; Takao et al., 2008). For example, flow fluctuation caused by power stations disturbs macroinvertebrates which in turn respond with lower biodiversity (Cortes et al., 1998; Bruno et al., 2016). It also disrupts colonization of many macroinvertebrates, causing lower taxa richness (Fisher & LaVoy, 1972; Munn & Brusven, 1991). High level periods have a significant negative effect on the diversity of macroinvertebrates in channelized sections compared to natural ones (Negishi et al., 2002).

Biological quality is a measure of the condition of a waterbody relative to the requirements of one or more biotic groups (such as aquatic flora, macroinvertebrates or fishes). In terms of the Water Framework Directive (2002), these groups are called biological elements. Many countries have developed tools to assess the biological quality of streams on the basis of macroinvertebrates. Some use multivariate assessment approaches, some use multi-metric indices (e.g. Dahl & Johnson, 2004; Ofenböck et al., 2004; Pinto et al., 2004; Vlek et al., 2004). However, most of the calculation methods focus on organic pollution or acidification, only few reflect hydromorphological modifications (Balestrini et al., 2004; Lorenz et al., 2004). During the recent decades, hydrochemical pollution of streams has been significantly reduced in Western Europe. Instead, hydromorphological modification, considered earlier as a second-rate disturbance after hydrochemical impacts, is now recognized as a primary adverse factor for the stream biota, including macroinvertebrates (Feld, 2004; Lorenz et al., 2004). In Europe, there are currently two indication systems to estimate the direct effects of hydromorphological degradation on stream macroinvertebrates: one used in Germany (Schmedtje & Colling, 1996) and the other, in Great Britain (Extence et al., 1999). Similar tools are also available for British Columbia, Canada (Armanini et al., 2011), New Zealand (Greenwood et al., 2016) and South Korea (Kong, 2016).

Estonia (together with Latvia and Lithuania) forms a separate Baltic province ecoregion in terms of the Water Framework Directive. This province is bordered with the Eastern plains in the East, the Fenno-Scandian shield in the north and the Central plains in the west. Estonia seems to be a suitable area to study the effects of different hydromorphological disturbances on macroinvertebrates, for the following reasons: (1) existence of a large number of differently hydromorphologically disturbed waterbodies, (2) lack of appropriate information for this region, as well as for the neighboring areas and (3) location of the Baltic province on the border of three other large ecoregions, which enables to test the results within the surrounding areas with a similar species list.

A multimetric index was developed to estimate the effects of organic pollution and general degradation on stream macroinvertebrate communities in Estonia (Status classes..., 2009; Timm & Vilbaste, 2010). However, it is not clear whether the included metrics (different kinds of taxon richness, Shannon diversity etc.) reflect hydromorphological disturbances appropriately. None of the specific hydromorphological indication systems listed above is applicable to the Baltic conditions: they contain too many missing or redundant species in their indicator lists. Moreover, calculation of the British score on the basis of the local material was not possible because of the incompatibility of sampling procedures.

According to the Estonian Environmental Register, there are more than 1000 dams, including destroyed objects, on Estonian running waters. Most of them are lower than 5 m and are used mainly for recreation and/or irrigation. As the hydroenergetic potential of Estonian streams is low because of the too low gradient of most stream channels, the number of hydroelectric power stations is also small. In 2011, there were only 47 working stations and watermills with a total capacity of only 8.09 MW (Raesaar, 2005).

Purposeful excavation of ditches on the current territory of Estonia started already in the 1820s. Stream channelization works became particularly intensive in the 1950s (Etverk, 1974). Most small streams were straightened and/or dredged (Tuulmets & Aasalo, 1980). In addition, up to 140,000 km of ditches are found nowadays (Lindström & Koff, 2005). This is 4.5 times more than the total length of all Estonian natural streams. However, woodland ponds and ditches in Estonia were considered species rich and diverse habitats. Although drainage reduces their taxonomic richness, ditches in semi-naturally managed forests can serve as replacement habitat for most taxa (Vaikre et al., 2015).

2. AIMS

The aims of this study were the following:

- 1) to assess whether and how much the macroinvertebrate communities were changed in the hydromorphologically modified (dammed or channelized) running waters compared to the streams in near-natural conditions (papers **I, III, IV**). The communities were characterized a) by indicator species, b) by biological quality (in terms of the current national multimetric estimation system);
- 2) to develop a special index, on the basis of macroinvertebrates, for assessment of only hydromorphological disturbances (paper **II**);
- 3) to compare the ability of the old and new indices to reflect relationships between hydromorphological changes and macroinvertebrates (all papers).

3. MATERIAL AND METHODS

3.1. Study area and sampling sites

Estonia is a small country (area 45,200 km²) with a flat landscape (mean altitude 50 m above sea level), situated on the eastern shore of the Baltic Sea, with a prevailing altitude lower than 200 m. It belongs to the area of mixed forests of the temperate zone, bordering the taiga. In comparison with the southern areas, it is characterized by extensive areas of raised bogs and forests (Raukas & Rõuk, 1995). There occur more than 7,300 running waters, with a total length of ca 31,000 km (Loopmann, 1979).

When selecting of the modified sampling areas, the availability of appropriate reference areas, as well as their accessibility for researchers were taken into consideration. As a rule, the reference areas were located not farther than 2 km from the modified areas (to ensure that both were of the same stream size), and were situated upstream (to avoid self-disturbance). In the pairwise study (paper IV), the areas with obvious evidence of channelization were first chosen using maps and their suitability was then checked in the field. The reference sites were located in near-natural areas without hydrochemical pollution and/or hydromorphological (human- of beaver-induced) disturbances.

3.2. Sampling

The macroinvertebrate sampling sites included in the papers I–IV are shown in Fig. 1. The samples were collected from shallow areas (depth < 1 m) in spring after the highwater period (papers I, III, IV) or partly in late autumn (paper I). Summer was excluded to avoid the period when many heterotopic insects have left the water environment. The areas with direct hydrochemical pollution were avoided, to focus on hydromorphological changes only. A standard handnet with an edge length of 25 cm and a mesh size of 500 µm (European..., 1994) was used. Altogether six subsamples were taken from each site: five subsamples collected from the most typical bottom (together 1.25 m²), and a qualitative sample which included different available habitats (Medin et al., 2001). Most of the samples from 1985–2000 (paper II) were qualitative and a significant part of them were collected also in summer. The collected animals were fixed, together with sieving residues with 96% ethanol and transported to the laboratory for analysis. The animals were identified to the species or to the genus level, according to Johnson (1999). Chironomids, oligochaetes, water mites and other organisms, whose identification required higher magnification, were not determined further.

At each site, flow velocity and composition of the bottom substrate were estimated visually on a four-point scale: 0 – no flow and/or muddy bottom, 1 –

slow flow ($< 0.2 \text{ m s}^{-1}$) and/or sandy bottom, 2 – fast flow ($> 0.2 \text{ m s}^{-1}$) and/or sandy-stony bottom, 3 – very fast flow (0.2 m s^{-1}) and/or stony bottom.

For papers **III** and **IV** (pairwise study), water temperature, dissolved oxygen (O_2), oxygen saturation ($\text{O}\%$), pH and conductivity were measured in the field.




Figure 1. Study area. A – paper **I** (16 dams, 32 sites in 2005–2006), B – paper **II** (3,282 sites in 1985–2009; including also the lake littoral areas), C – paper **III** (24 dams, 72 sites in 2009–2010), D1 – paper **IV**, pairwise dataset (24 channelized streams, 48 sites in 2009–2010), D2 – paper **IV**, background dataset (73 channelized streams and 73 unchannelized streams in 2000–2010).

3.3. Estimation of biological quality

To assess the biological quality of running waters, a national multimetric index, based on five components, was used (Status classes..., 2009). These indices were the following: total taxon richness T (indicating total biological diversity); Ephemeroptera, Plecoptera and Trichoptera larvae taxon richness EPT (indicating pollution-sensitive taxon diversity; Lenat, 1988); Shannon diversity H' (indicating how many different species there are in a dataset, and simultaneously taking into account how evenly the individuals are distributed among those species); Average Score Per Taxon ASPT (mean sensitivity of taxon; Armitage et al., 1983); and Danish Stream Fauna Index (DSFI) (indicating the level of organic pollution; Skriver et al., 2000). The H' was calculated on the basis of five area-based subsamples only. For the other metrics, the qualitative subsample was also included. The corresponding table of the biological quality classes (Appendix) was used to attribute quality levels to all five indices. All results of high quality were assigned five points, the values of good quality, four points, the values of moderate quality, two points, and the values of poor and bad quality, zero points. The difference between good level and moderate level was intentionally emphasized in order to underline the principal difference between them in terms of the Water Framework Directive (2002). According to the Directive, the European member states must protect and improve their surface waters to achieve at least good ecological status. Instead, the difference between poor level and bad level was ignored, as they had no actual difference. Multimetric quality (MMQ) was then calculated by adding up the corresponding points. Hence, the sum 23–25 was considered to indicate high quality, 18–22, good, 10–17, moderate, 6–9, poor and < 6, bad quality.

The values of all indices and the estimates of multimetric quality were then divided by their corresponding reference values (5 for each single index and 25 for the multimetric quality, respectively), resulting in observed/expected ratios, called also Ecological Quality Ratios (**EQR**). The EQR-s are universal values that enable to compare different metrics. The acceptable range for the boundary between high quality and good quality in European streams is 0.84 – 0.94 and the range between good quality and moderate quality is 0.64 – 0.75 (Van de Bund, 2009). For Estonian streams, these boundaries were classified as follows: high > 0.9, good 0.7–0.9, moderate 0.4–0.7, poor and bad < 0.4 (Timm & Vilbaste, 2010).

The MESH index (developed with the participation of the author), is based on the affinities of macroinvertebrate taxa to flow velocity and bottom substrate (paper II). The corresponding quality levels and thus the EQR values of MESH are not yet available.

3.4. Data analysis

In paper **I**, the effect of damming on the structure of the macroinvertebrate community and biological quality was studied for two types of reservoirs: 1) with a large amount of accumulated fine sediments, and 2) with hard bottom; and for the corresponding below-dam areas. Generalized linear models implemented in the SAS Genmod procedure (SAS 9.1.3 Help and Documentation, 2004) were used. In calculations, the differences between the sampling sites were partly eliminated by using repeated measures analysis as realized in the Repeated clause of the Genmod procedure. Comparison of the biological indices was performed in a similar way but, instead of the Genmod procedure, the SAS Mixed procedure of mixed-type analysis of variance was used. Following multiple comparisons (183 taxa, 24 indices), significance level α was corrected in the Bonferroni sense using $\alpha = 0.0005$ when comparing the taxa and $\alpha = 0.002$ when comparing the indices.

In paper **II**, a new index was developed to assess the hydromorphological quality of Estonian surface waters, based on the taxonomic composition of macroinvertebrate (**MESH** – **Macroinvertebrates in Estonia: Score of Hydro-morphology**). Both flow velocity and bottom type were estimated visually using a four-level scale. The lowest flow velocity and the softest bottom were coded with “0”, the fastest flow and the hardest bottom, with “3”. Combinations of the values of Flow Score and Bottom Score were named Flow-Bottom Score, because of a highly significant correlation between them. This score was then calculated for 690 macroinvertebrate taxa, adjusting the scores over the occurrence of taxa in different habitats. The taxa were analyzed to calculate informativeness (IG) with respect to the habitats using the Kullback-Liebler difference (Renyi, 1970). The statistical significance of IG was modelled using the SAS IML procedure, repeating the IG calculation 200,000 times. In total, 394 taxa were qualified as indicators. Finally, all indicators were divided into clusters again, with Flow-Bottom Score values of 0 (muddy littoral), 1 (sandy littoral), 2 (slow-flowing stream, or stony littoral), or 3 (fast-flowing stream). The MESH is the average Flow-Bottom Score consisting of the sum of the individual scores of all indicator taxa in a sample divided by the number of the indicator taxa in the sample.

In paper **III**, the impact of damming (above-dam versus below-dam versus undisturbed areas) and in paper **IV**, the effect of channelization (channelized versus undisturbed areas) on macroinvertebrates and stream physicochemical parameters was estimated. In both cases, the data were analyzed using the R (R Development..., 2009) or Statistics 8.0 software (R Development Core Team, Vienna, Austria, 2009). A general linear model was used by comparing the contrasts of three-factor level (above-dam, below-dam, undisturbed). The effect of channelization on the continuous variables of macroinvertebrates and on physicochemical parameters was assessed using one-way analysis of variance (ANOVA). For the categorical variable (Danish Stream Fauna Index), non-

parametric Kruskal-Wallis test or Mann-Whitney U test were employed. For comparison of the metrics, the EQR-s were used. In both papers, the Bonferroni p-value was adjusted according to the number of tests. In addition, Indicator Species Analysis (Dufrene & Legendre, 1997) was used in paper IV, to detect indicator species for the channelized stream sections.

To generalize the effects of damming and channelization on biological and hydromorphological indices, the data from papers I, III and IV were combined and analyzed separately in the Thesis. The residuals of the macroinvertebrates metrics were compared for deviation from a normal distribution using the Shapiro-Wilk's test. In most cases, the one-way analysis of variance (ANOVA) was used. The Danish Stream Fauna Index and MESH were examined by the non-parametric Kruskal-Wallis test, followed by multiple comparison of mean ranks. The critical Bonferroni p-value was 0.007. The Kruskal-Wallis non-parametric test was run in Statistica 8.0 (StatSoft, Inc., 2011) and ANOVA was performed with the software R 3.2.4 (R Development Core Team, Vienna, Austria, 2016).

4. RESULTS AND DISCUSSION

4.1. Taxonomic composition of macroinvertebrates

4.1.1. Effects of damming

In the sediment-rich reservoirs, some insect larvae (Chironomidae, lentic ephemeropteran *Cloeon dipterum*), crustaceans *Asellus aquaticus*, *Gammarus pulex*, and oligochaetes were the most abundant taxa. Except for the highly mobile *G. pulex*, they all usually inhabit slow-flowing or standing waters. The larvae of several other insects (among them the rheophilic ephemeropteran *Baetis rhodani* and Simuliidae) and *G. pulex* dominated in the below-dam reaches of the sediment-rich reservoirs. In the hard-bottomed reservoirs, the riffle beetle *Elmis aenea*, oligochaetes, chironomid larvae and *G. pulex* were the most abundant. In the corresponding below-dam reaches, Chironomidae, larvae of the filtering caddisfly (*Hydropsyche pellucidula*) and another riffle beetle, *Limnius volckmari*, prevailed (paper I). Similarly to this study, Shao et al. (2008) considered chironomids and oligochaetes typical inhabitants of reservoirs. Boles (1981) noted that Simuliidae, *Baetis* and Chironomidae were usually found in below-dam areas, owing to abundant food supply.

In the larger damming-based dataset, the above-dam taxa were similarly usually lentic (paper III). The typical representatives were leeches, snails and water mites, but also some mayfly, damselfly, and caddisfly larvae. The below-dam taxa often included numerous filtrators (such as pea clams, caddisfly and Simuliidae larvae), typical of natural lake outflows. In turn, the taxa of the undisturbed sites included many rheophiles, such as the snail *Theodoxus fluviatilis*, the crustacean *Gammarus pulex*, several mayfly, stonefly and caddisfly larvae; riffle beetles, and the water bug *Aphelocheirus aestivalis*. However, the rare mayfly species *Arthroplea congener* was only found in lentic reservoirs. Changes in the functional feeding groups above and below dams can be primarily explained with differences in food supply (Spence & Hynes, 1971). On the basis of the current study, hydromorphological conditions (such as flow velocity and bottom substrate type) are of not less importance than food.

4.1.2. Effects of channelization

According to Indicator Species Analysis, only two taxa (the caddisfly larvae of *Limnephilus* sp. and the leech *Glossiphonia complanata*) were significantly ($p < 0.05$; uncorrected for multiple comparisons) associated to channelized sites in the small (pairwise) dataset (paper IV). Four taxa (*Elmis aenea*, mayfly *Heptagenia sulphurea* larvae, and two species of caddisfly larvae – *Cheumatopsyche lepida* and *Rhyacophila* sp.) were significantly associated with near-natural (reference) areas. In the large (background) dataset, four taxa (the dipteran family Ceratopogonidae, and three species of caddisfly larvae:

Limnephilus flavicornis, *L. lunatus* and *Ironoquia dubia*) were significantly associated with channelized sites. Nine taxa were significant indicators of the unchannelized sites: seven species of caddisfly larvae (*Hydropsyche pellucidula*, *Sericostoma personatum*, *Lepidostoma hirtum*, *Cheumatopsyche lepida*, *Ithytrichia lamellaris*, *Odontocerum albicorne*, *Hydropsyche siltalai*), a whirligig beetle larva (*Orectochilus villosus*), and a crane fly larva (*Antocha* sp.). The indicators of the channelized sites were mostly typical of slow-flowing waters (according to MESH, Paper II), while the indicators of the reference sites were usually found in fast-flowing reaches.

4.2. Biological, hydromorphological and hydrochemical quality

To summarize the analyses, the datasets of the papers I, III and IV were combined. Table 1 and Fig. 2 comprise the mean values of the biological quality metrics and their EQR-s for the four habitats: above-dam (reservoir), below-dam, channelized and reference. In Table 2, the EQR values are compared pairwise by either ANOVA or the Kruskal-Wallis test.

Table 1. Metrics of biological quality for the hydromorphologically modified (above-dam, below-dam and channelized) and for the reference habitats. Bold letters indicate arithmetic means, the Ecological Quality Ratio (EQR) values are in brackets. The EQR quality levels are the following: high > 0.9, good 0.7–0.9, moderate 0.4–0.7, poor and bad < 0.4. T – total taxon richness, EPT – taxon richness of Ephemeroptera, Plecoptera and Trichoptera larvae, DSFI – Danish Stream Fauna Index, ASPT – Average Score Per Taxon, H' – Shannon diversity, MMQ – multimetric quality, MESH – score of hydromorphology.

Metric	Arithmetic mean			
	Above-dam (n=42)	Below-dam (n=42)	Channelized (n=91)	Reference (n=115)
T	32.1 (1.20)	35.0 (1.09)	30.5 (1.08)	35.0 (1.19)
EPT	11.5 (0.81)	16.3 (1.07)	12.6 (0.95)	16.2 (1.16)
DSFI	4.90 (0.70)	6.21 (0.89)	5.58 (0.80)	6.53 (0.93)
ASPT	5.49 (0.82)	5.96 (0.88)	5.51 (0.84)	6.17 (0.92)
H'	2.68 (0.91)	2.64 (0.99)	2.49 (0.94)	3.07 (1.08)
MMQ	16.6 (0.67)	20.4 (0.82)	19.4 (0.77)	23.2 (0.95)
MESH	1.77	2.50	2.47	2.59

Total taxon richness (T) was quite similar for all habitats. There were no significant differences in T between any modified habitats or reference areas. Nevertheless, the mean values of T for the reference and below-dam habitats were the highest (Tables 1–2). The mean EQR (observed/expected ratio) of T revealed also

high quality for all habitats. Thus, the current study does not support total taxon richness as an appropriate tool to assess hydromorphological changes. At the above-dam sites, the rheophilic species inhabiting this area earlier had probably been replaced by limnophilic species (as indicated by MESH, see below).

In the channelized reaches, natural species had also been replaced by other, more tolerant natural taxa by stream drift and other ways of the self-recovery of the habitat. Like in this study, hydromorphological alteration was not followed by a substantial decline in whole-community richness in several streams of the Netherlands, Germany and Poland (Feld, 2004). Channelization of boreal streams for timber transport did not alter hydromorphological conditions sufficiently to have a strong impact on the taxon richness of macroinvertebrates in Finland (Turunen et al., 2016). However, several authors reported a significant decrease in taxon richness in channelized streams (Negishi et al., 2002; Horsák et al., 2008; Pliūraite and Kesminas, 2010).

The number of sensitive taxa (**EPT**) and the index of organic pollution (**DSFI**) were significantly lower for the reservoirs and channelized reaches, compared to the reference areas. However, this relationship was not significant for the below-dam sites. The mean EQR of EPT indicated good quality for the reservoirs and high quality for the other habitats, being only slightly lowered for the channelized reaches. At the same time, the EQR of DSFI was moderate for the reservoirs, good for the below-dam and channelized habitats, and high only for the reference areas. Conditions for pollution-sensitive macroinvertebrates were not significantly deteriorated in the above-dam areas in case mud accumulation was low (paper **I**). At the same time, in the below-dam areas, disturbance to sensitive species might have been masked by fast flow. Even some pollution-tolerant species were sensitive to low water velocity in the reservoirs (paper **III**). A decreased EPT for the dammed areas was also observed in streams of Illinois, USA, due to degraded habitat and/or lowered water quality (Santucci et al., 2005). The structure of the macroinvertebrate community downstream a dam in Ontario (Canada) seemed to suffer for stress similar to mild organic pollution (Spence & Hynes, 1971). The channelized sites in the current study had significantly lower EPT and DSFI compared those in the reference areas (paper **IV**).

Mean taxon sensitivity (**ASPT**), Shannon diversity (**H'**) and multimetric quality (**MMQ**) were all significantly lower in the three modified habitats compared to the reference area. The EQR of ASPT revealed high quality for the reference area and good quality elsewhere (however, the lowest value was observed for the reservoirs). The EQR of Shannon diversity was high for all habitats. Davy-Bowker & Furse (2006) and Friberg et al. (2009) also found that ASPT was correlated significantly negatively to channel modification. Rabeni (2000) considered diversity indices as the most sensitive to habitat changes for stream macroinvertebrates. In this study, the mean EQR of Shannon diversity (**H'**) indicated high biological quality for all habitats, although the lowest value was observed for the above-dam habitat, and the highest quality, for the reference area.

The EQR of **MMQ** (biological quality on the basis of the five above indices) was the lowest for the reservoirs (moderate), followed by the channelized (good), below-dam (good) and reference habitats (high) (Table 1). The effect of insensitive total taxon richness was surpassed by the effects of the other four sensitive indices.

The MESH was significantly lower for the reservoirs compared to the reference areas, as well as to the other habitats. For the below-dam and channelized habitats with fast flow and hard bottom, it was not significantly different from that for the reference areas (Fig. 2, Table 2). Nevertheless, in the dataset used in paper **III**, a significant difference in MESH between the below-dam and the reference areas was observed. At the same time, all hydrochemical and hydrophysical parameters (water temperature, dissolved oxygen (O₂), oxygen saturation (O%), pH and conductivity) were highly similar among these three habitats. Thus, some advantage of macroinvertebrate quality indicators over commonly approved abiotic parameters for streams was ascertained.




Figure 2. Comparison of the EQR values of the quality metrics (mean \pm SD). T – total taxon richness, EPT – taxon richness of Ephemeroptera, Plecoptera and Trichoptera larvae, DSFI – Danish Stream Fauna Index, ASPT – Average Score Per Taxon, H' – Shannon diversity, MMQ – multimetric quality.

Table 2. Comparison of Ecological Quality Ratios (EQR) of the metrics between different hydromorphological habitats (by ANOVA and Kruskal-Wallis (*) tests). T – total taxon richness, EPT – taxon richness of Ephemeroptera, Plecoptera and Trichoptera larvae, DSFI – Danish Stream Fauna Index, ASPT – Average Score Per Taxon, H' – Shannon diversity, MMQ – multimetric quality, MESH – score of hydromorphology. n.s. – not significant. The critical Bonferroni p-value is 0.007.

Metric	p-value					
	Above-dam – Below-dam	Above-dam – Channelized	Below-dam – Channelized	Above-dam – Reference	Below-dam – Reference	Channelized – Reference
EQR-T	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.
EQR-EPT	0.001	n.s.	n.s.	<0.0001	n.s.	<0.0001
EQR-DSFI*	<0.0001	0.004	0.003	<0.0001	n.s.	<0.0001
EQR-ASPT	0.004	n.s.	n.s.	<0.0001	0.002	<0.0001
EQR-H'	n.s.	n.s.	n.s.	0.007	0.003	0.002
EQR-MMQ	<0.0001	<0.0001	n.s.	<0.0001	0.0001	<0.0001
MESH*	<0.0001	<0.0001	n.s.	<0.0001	n.s.	n.s.

Although other specific hydromorphological indication systems were not applicable to conditions in the the Baltic region, correlation between scores of the common elements for individual taxa in MESH and the corresponding British index LIFE (Extence et al., 1999) was fair ($p < 0.05$, $r = -0.5987$, $n = 252$). After transforming the verbal categories of rheophily into the corresponding six classes after Schmedtje and Colling (1996) (ignoring the class “indifferent”), a relatively similar result was obtained ($r = -0.5829$, $n = 271$) (paper II).

In relation to hydromorphological stress, the analyzed indices can be arranged into three groups: 1) insensitive (total taxon richness); 2) the lowest value for the reservoirs, the slightly higher value for the below-dam habitat than for the channelized area, the highest value for the reference area (EPT, DSFI); 3) the lowest value for the reservoirs, a similar value for the below-dam and channelized areas, the highest value for the reference areas (ASPT, H', MESH). Although EPT, ASPT and H' indicated good or even high quality for hydromorphologically modified habitats, their lowest mean values were always registered for the most degraded (above-dam) areas and the highest values were registered for the reference areas. The high MESH values for the below-dam and channelized habitats indicate that these areas were near-natural in terms of bottom hardness and flow velocity.

5. CONCLUSIONS

Macroinvertebrate communities responded significantly to both main hydromorphological disturbances – damming and channelization – in streams of Estonia. Most macroinvertebrate-based indices of biological quality, belonging to the corresponding national multimetric index, were significantly and negatively influenced both by damming and channelization. Unlike the other tools, a new index (MESH) was specially developed to assess hydromorphological changes. The MESH distinguished reliably the above-dam macroinvertebrate species from those of the other areas but failed to distinguish the species of the channelized habitats and the species of the reference areas. However, the latter difference can be well estimated by other available indices (EPT, DSFI, ASPT and H'). Nevertheless, inclusion of MESH in the national system as a relevant indicator of damming can be recommended, including development of corresponding reference values and quality levels that are in line with the European Water Framework Directive.

SUMMARY

Alteration of hydromorphology as a consequence of human activities (e.g. damming and channelization) poses a significant threat to habitat quality and in turn to riverine communities. Taxonomic composition of benthic macroinvertebrates has been widely used as an indicator of running water quality. Macroinvertebrates are also considered a necessary biological quality measure of freshwaters according to the Water Framework Directive. However, the corresponding metrics have been mostly focused on organic pollution or acidification, and only seldom on hydromorphological modifications. Estonia (together with Latvia and Lithuania) forms a separate Baltic province ecoregion in terms of the Water Framework Directive. Despite the occurrence of a large number of dammed and channelized waterbodies in this region, the effects of these disturbances on macroinvertebrates have almost not been studied. Nor is it not clear whether the metrics included in the Estonian multimetric index of biological quality (total taxon richness T, sensitive taxon richness EPT, organic pollution index DSFI, mean sensitivity of taxon ASPT, and Shannon diversity H') reflect hydromorphological disturbances appropriately. To clarify this issue, the relationships of these indices with damming and channelization were estimated for four habitats: above-dam, below-dam, channelized, reference. In addition, a new index (MESH), based on the affinities of 394 macroinvertebrate species to flow velocity and bottom type, was developed.

The taxa found at the above-dam sites were usually associated with standing waters. Similarly, significant indicators of channelized stream reaches were typical of slow-flowing waters. The taxa of the below-dam areas often included numerous filter-feeding species, while the taxa of the undisturbed sites included many rheophiles. Total taxon richness was similar for all habitats, as in the modified areas sensitive species had been replaced by tolerant species. However, all other indices (including multimetric estimation MMQ) were significantly negatively influenced by damming and/or channelization. The MMQ was the lowest for the reservoirs (with moderate biological quality), followed by the channelized (good quality), below-dam (good quality) and reference habitats (high quality). In conclusion, macroinvertebrate communities responded significantly to both main hydromorphological disturbances – damming and channelization – in streams of Estonia.

SUMMARY IN ESTONIAN

Inimtekkeliste hüdro-morfoloogiliste muutuste mõju vooluvete seisundile suurselgrootute järgi Eestis (Euroopa, Balti ökoregioon)

Inimesed on muutnud väga paljude vooluveekogude hüdro-morfoloogiat: voolukiirust, põhja iseloomu ja voolusängi kuju. Sellised muutused on nende veekogude elustikku sageli oluliselt mõjutanud. Kõige silmapaistvam mõju on paisutamine. Ülalpool paisu asuvas jõelõigis on voolukiirus pidurdatud, mis soodustab muda kogunemist; vahetult allpool paisu aga see-eest tugevasti suurendatud. Paisutamise mõju vee-elustikule sõltub paisu kõrgusest, vanusest ja ehituslikest eripäradest. Mõned loomad (näiteks siirdekalad) ei pruugi paisudest mööda pääseda neile elutähtsatesse piirkondadesse. Vähem teatakse sellest, kuidas elustikku mõjutavad muutunud voolukiirus ja/või põhja iseloom.

Õgvendamise käigus kaevatakse kogu vooluveekogu või mõni selle osa sirgeks. Tulemuseks on voolukiiruse ja põhja iseloomu ühtlustumine ning vooluhulga sesoonse stabiilsuse vähenemine. Enamasti eemaldatakse õgvendustööde käigus ka looduslik kaldataimestik (sh mets), mis varem jõesängi varjutas ja oli oluliseks orgaanilise aine allikaks vees.

Suurselgrootud on palja silmaga nähtavad, veekogude põhjal, taimedel, setete sees või veepinnal elavad loomad. Neid on veekogude seisundi indikaatoritena kogu maailmas palju kasutatud, sest nad reageerivad ennustatavalt mitmesugusele inimtegevusele, eriti orgaanilisele ja happelisele veereostusele. Ka Eestis on vastav, Euroopa Veepoliitika Raamdirektiivi nõuetest tulenev indikatsioonisüsteem välja töötatud. See on samuti suunatud pigem orgaanilise reostuse ja/või tuvastamata stressiallikate mõju hindamisele, mitte hüdro-morfoloogilistele muutustele. Samas, ehkki väga suur osa Eesti vooluvetest on paisutatud või õgvendatud, pole seisundi hindamise mõttes uuritud, kas ja kuidas suurselgrootud sellele reageerivad.

Praeguse töö eesmärgid olid järgmised:

- 1) hinnata paisutamise ja õgvendamise mõju vooluvete seisundile, kasutades senist suurselgrootutel tuginevat hindamissüsteemi,
- 2) töötada välja uus indeks just hüdro-morfoloogiliselt muudetud vooluvete seisundi hindamiseks,
- 3) võrrelda varasemate indeksite ja uue indeksi tulemuslikkust paisutatud ja õgvendatud vooluvete seisundi hindamiseks.

Paisutamise ja õgvenduse mõju uurimiseks võeti suurselgrootute proovid vooluvetest enamasti kevadel: pärast suurvee taandumist, kuid enne, kui suurem osa putukatest oleks jõudnud veest välja lennata. Materjal koguti nelinurkse standardkahvaga jalaproovide ja kahvatõmmete abil. Iga proov koosnes 1,25 m² suurusest pindalapõhisest osast ning kvalitatiivsest osast. Et keskenduda just hüdro-morfoloogilistele muutustele, välditi otsese hüdrokeemilise reostusega alasid.

Praegu Eestis kehtiv vastav seisundisüsteem koosneb viiest indeksist: T (üldine taksonirikkus), EPT (tundlike taksonite rikkus), DSFI (orgaanilise reostuse tase), ASPT (taksoni keskmine tundlikkus) ning Shannoni erisus. Iga indeksi seisundiväärtus annab kindla arvu punkte, mis summeeritult moodustavad koondhinnangu MMQ. Kohapealse materjali põhjal loodud uus indeks (MESH) tugineb Eesti liikide tundlikkusel voolukiirusesse ning põhja iseloomu. Seega peaks ta eeldatavasti sobima nii paisutamise kui õgvendamise mõju hindamiseks suurselgrootute taksonoomilisele koosseisule.

Eriti selgelt reageerisid suurselgrootud paisjärvede mudastumisele, kus reofiilsed liigid olid asendunud seisuveeliste liikidega. Ülalpool paise olid muuhulgas väga arvukad näiteks seisuveelise ühepäevikulise *Cloeon dipterum* vastsed ning vesikakand *Asellus aquaticus*. Allpool paise, kus voolukiirus suur, aga vees leidub palju paisjärvedes moodustunud hõljumit, olid sagedad voolulembesed liigid, kelle seas palju filtreerijaid (teiste seas näiteks kihulaste (*Simuliidae*) või ehmeistiivalise *Hydropsyche pellucidula* vastsed). Õgvendatud jõelõike eelistasid keskmise või madala reostustundlikkusega või mitte eriti voolulembesed liigid (näiteks lamekaan *Glossiphonia complanata*, habesääsklaste (*Ceratopogonidae*) vastsed, ehmeistiivalise *Limnephilus flavicornis* vastsed). Hüdromorfoloogiliselt rikkumata kohtades aga olid tavalised kõrge reostustundlikkuse ning voolulembesusega liigid (näiteks vesiking *Theodoxus fluviatilis*, kärestikulutikas *Aphelochelirus aestivalis*, paljude ühepäevikuliste, kevikuliste ja ehmeistiivaliste vastsed).

Paisutamine ja õgvendamine mõjutasid enamikku võrreldud seisundiindeksitest (välja arvatud üldine taksonirikkus T, kuid kaasa arvatud uus indeks MESH). Paisjärvedes oli seisund suurselgrootute järgi kas oluliselt madalam või vähemalt absoluutväärtuselt kehvem kui muudes uuritud elupaikades (allpool paise, õgvendatud, mõjutamata alad). Nii allpool paise kui õgvendatud lõikudes oli koondseisund viie indeksi põhjal paisjärvedega võrreldes oluliselt parem, kuid mõjutamata aladega võrreldes oluliselt kehvem. Töö aluseks olevate artiklite põhjal kombineeritud andmebaasi alusel oli suurselgrootute keskmine koondseisund senise hindamissüsteemi järgi järgmine: paisjärvedes kesine, õgvendatud lõikudes ja allpool paise hea, ning mõjutamata aladel väga hea. MESH võimaldas usaldusväärset hinnata paisjärvede mõju, mõnikord paisualuste lõikude mõju, kuid mitte õgvenduse mõju suurselgrootutele.

Paisutamise suhtes osutasid suurselgrootud isegi tundlikumaks kui mõned tavapärased hüdrokeemilised ja -füüsikalised parameetrid (temperatuur, hapnikusisaldus, pH, elektrijuhtivus). Viimaste kevadel mõõdetud väärtused ei erinenud paisjärvede, paisualuste ning mõjutamata alade vahel oluliselt, samas kui loomastiku järgi sai neid elupaiku eristada. Enamik suurselgrootute põhjal koostatud seisundiindeksitest (nii need, mida kasutati seni hüdrokeemilise seisundi või tuvastamata mõjude hindamiseks, kui otseselt hüdromorfoloogiliste mõjude hindamiseks loodud MESH) osutasid Eesti vooluvete paisutamise ja/või õgvenduse suhtes oluliselt tundlikeks. MESH on soovitatav kaasata kehtivasse hindamissüsteemi kui paisutamise indikaator, kehtestades talle ühtlasi seisundi-klasside piirid.

ACKNOWLEDGEMENTS

I am very grateful to my supervisors Dr. Taavi Virro and Dr. Henn Timm for their guidance, advice and continuous support during the studies. I also thank Dr. H. Timm for assistance in studying macroinvertebrate taxonomy and for constructive criticism in the writing process. Many thanks belong to my colleagues at the Estonian University of Life Sciences for their ongoing support and help in fieldwork during the studies. Rein Järvekülg provided the appropriate data on dams. Ester Jaigma kindly revised the English text of the manuscript.

The study was carried out at the Estonian University of Life Sciences, Centre for Limnology. It was financed by the Estonian research projects SF0170011s08, SF0170006s08 and IUT 21-2; by ETF Grants 6580 and 7643 of the Estonian Ministry of Education and Research; and by the national programme „Humanitarian and natural science collections“. The work was also supported by the European Social Fund's Doctoral Studies and by the Internationalization Programme DoRa carried out by Foundation Archimedes.

REFERENCES

- Ahearn, D. S., Sheibley, R. W. & Dahlgren, R. A. 2005. Effects of river regulation on water quality in the lower Mokelumne River, California. *River Research and Applications* 21: 651–670.
- Armanini, D. G., Horrigan, N., Monk, W. A., Peters, D. L. & Baird, D. J. 2011. Development of a benthic macroinvertebrate flow sensitivity index for Canadian rivers. *River Research and Applications* 27: 723–737.
- Armitage, P. D., Moss, D., Wright, J. F. & Furse, M. T. 1983. The performance of a new biological water quality score system based on a wide range of unpolluted running-water sites. *Water Research* 17: 333–347.
- Arndt, E. & Domdei, J. 2012. Influence of beaver ponds on the macroinvertebrate benthic community in lowland brooks. *Polish Journal of Ecology* 59: 799–811.
- Balestrini, R., Cazzola, M. & Buffagni, A. 2004. Characterising hydromorphological features of selected Italian rivers: a comparative application of environmental indices. *Hydrobiologia* 516: 365–379.
- Boles, G. L. 1981. Macroinvertebrate colonization of replacement substrate below a hypolimnial release reservoir. *Hydrobiologia* 78: 133–146.
- Bruno, M. C., Cashman, M. J., Maiolini, B., Biffi, S. & Zolezzi, G. 2016. Responses of benthic invertebrates to repeated hydropeaking in semi-natural flume simulations. *Ecohydrology* 9: 68–82.
- Bunn, S. E. & Arthington, A. H. 2002. Basic Principles and Ecological Consequences of Altered Flow Regimes for Aquatic Biodiversity. *Environmental Management* 30: 492–507.
- Cortes, R. M. V., Ferreira, M. T., Oliveira, S. V. & Godinho, F. 1998. Contrasting impact of small dams on the macroinvertebrates of two Iberian mountain rivers. *Hydrobiologia* 389: 51–61.
- Cranston, P. S., Fairweather P. & Clarke, G. 1996. Biological indicators of water quality. In: *Indicators of catchment health: a technical perspective* (Eds. Walker, J. & Reuter, D.J.) CSIRO Publishing, Melbourne, pp. 143–154.
- Dahl, J. & Johnson, R. K. 2004. A multimetric macroinvertebrate index for detecting organic pollution of streams in southern Sweden. *Archiv für Hydrobiologie* 160: 487–513.
- Davy-Bowker, J. & Furse, M. T. 2006. Hydromorphology – major results and conclusions from the STAR project. *Hydrobiologia* 188: 263–265.
- Dufrêne, M. & Legendre, P. 1997. Species Assemblages and Indicator Species: The Need for a Flexible Asymmetrical Approach. *Ecological Monographs* 67: 345–366.
- Estonian Environmental Register (<http://register.keskkonnainfo.ee/envreg/main#HTTPrLvA6YNYuMJpypcXHYYV0iCX2BGpk9C>), (in Estonian, access date: 25.12.2015)
- Etverk, I. 1974. Understanding and using the forest. In: *The forests of Estonia* (Eds. Valk, U. & Eilart, J.). Valgus, pp. 40–60 (in Estonian).
- European Committee for Standardization. 1994. Water quality – Methods for biological sampling – Guidance on handnet sampling of aquatic benthic macro-invertebrates. EN 27828. European Committee for Standardization, Brussels, Belgium.
- Extence, C. A., Balbi, D. M. & Chadd, R. P. 1999. River flow indexing using British benthic macroinvertebrates: a framework for setting hydroecological objectives. *Regulated Rivers: Research and Management* 15: 543–574.

- Feld, C. K. 2004. Identification and measure of hydromorphological degradation in Central European lowland streams. *Hydrobiologia* 516: 69–90.
- Fisher, S. G. & LaVoy, A. 1972. Differences in littoral fauna due to fluctuating water levels below a hydroelectric dam. *Journal of the Fisheries Research Board of Canada* 29: 1472–1476.
- Friberg, N., Sandin, L. & Pedersen, M. L. 2009. Assessing the effects of hydromorphological degradation on macroinvertebrate indicators in rivers: examples, constraints, and outlook. *Integrated Environmental Assessment and Management* 5: 86–96.
- Greenwood, M. J., Booker, D. J., Smith, B. J. & Winterbourn, M. J. 2016. A hydrologically sensitive invertebrate community index for New Zealand rivers. *Ecological Indicators* 61: 1000–1010.
- Horsák, M., Bojková, J., Zahrádková, S., Omesová, M. & Helešik, J. 2009. Impact of reservoirs and channelization on lowland river macroinvertebrates: A case study from Central Europe. *Limnologica* 39: 140–151.
- Hortle, K. G. & Lake, P.S. Macroinvertebrate Assemblages in Channelized and Unchannelized Sections of the Bunyip River, Victoria. *Australian Journal of Marine and Freshwater Research* 33: 1071–1082.
- Johnson, R. K. 1999. Benthic macroinvertebrates. Wiederholm, T. (ed). *Bedömningsgrunder för miljö kvalitet. Sjöar och vattendrag. Bakgrundsrapport 2. Biologiska parametrar.* Naturvårdsverket Förlag, pp. 85–166.
- Kennedy, T. L. & Turner, T. F. 2011. River channelization reduces nutrient flow and macroinvertebrate diversity at the aquatic terrestrial transition zone. *Ecosphere* 2: 1–13.
- Kong, D.-S. 2016. Development of Benthic Macroinvertebrates Streambed Index (BMSI) for Bioassessment of Stream Physical Habitat. *Journal of Korean Society on Water Environment* 32: 1–14.
- Lenat, D. R. 1988. Water quality assessment of streams using a qualitative collection method for benthic macroinvertebrates. *Journal of the North American Benthological Society* 7: 222–233.
- Lessard, J. L. & Hayes, D. B. 2003. Effects of elevated water temperature on fish and macroinvertebrate communities below small dams. *River Research and Applications* 19: 721–732.
- Liermann, C. R., Nilsson, C., Robertson, J. & Ng, R. Y. 2012. Implications of dam obstruction for global freshwater fish diversity. *BioScience* 62: 539–548.
- Lindström, K. & Koff, T. 2015. Antropotseen-inimeste ajastu. *Horisont* 5: 26–34. (in Estonian).
- Loopmann, A. 1979. The list of streams in the Estonian S.S.S.R. *Valgus*. (in Estonian).
- Lorenz, A., Hering, D., Feld, C. K. & Rolauuffs, P. 2004. A new method for assessing the impact of hydromorphological degradation on the macroinvertebrate fauna of five German stream types. *Hydrobiologia* 516: 107–127.
- Maxted, J. R., Barbour, M. T., Gerritsen, J., Poretti, V., Primrose, N., Silvia, A., Penrose, D. & Renfrow, R. 2000. Assessment framework for mid-Atlantic coastal plain streams using benthic macroinvertebrates. *Journal of the North American Benthological Society* 19: 128–144.
- Medin, M., Ericsson, U., Nilsson, C., Sundberg, I. & Nilsson, P. A. 2001. *Bedömningsgrunder för bottenfaunaundersökningar.* Medins Sjö- och Åbiologi AB, Mölnlycke.

- Munn, M. D. & Brusven, M. A. 1991. Benthic macroinvertebrate communities in nonregulated and regulated waters of the clearwater river, Idaho, U.S.A. *Regulated Rivers: Research and Management* 6: 1–11.
- Nakamura, F. & Yamada, H. 2005. The effects of pasture development on the ecological functions of riparian forests in Hokkaido in Northern Japan. *Journal of Ecological Engineering* 24: 539–55.
- Nakano, D., Nagayama, S., Kawaguchi, Y. & Nakamura, F. 2008. River restoration for macroinvertebrate communities in lowland rivers: insights from restorations of the Shibetsu River, north Japan. *Landscape and Ecological Engineering* 4: 63–68.
- Navarro-Llácer, C., Baeza, D. & de las Heras, J. 2010. Assessment of regulated rivers with indices based on macroinvertebrates, fish and riparian forest in the southeast of Spain. *Ecological Indicators* 10: 935–942.
- Negishi, J. N., Inoue, M. & Nunokawa, M. 2002. Effects of channelization on stream habitat in relation to a spate and flow refugia for macroinvertebrates in northern Japan. *Freshwater Biology* 47: 1515–1529.
- Ofenböck, T., Moog, O., Gerritsen, J. & Barbour, M. 2004. A stressor specific multi-metric approach for monitoring running waters in Austria using benthic macroinvertebrates. *Hydrobiologia* 516: 251–268.
- Petts, G. E. 1984. *Impounded Rivers. Perspectives for Ecological Management*. John Wiley & Sons Ltd., Chichester, UK.
- Pinto, P., Rosado, J., Morais, M. & Antunes, I. 2004. Assessment methodology for southern siliceous basins in Portugal. *Hydrobiologia* 516: 191–214.
- Pliūraite, V. & Kesminas, V. 2010. Response of benthic macroinvertebrates to agricultural pollution and channelization in the Mūša River, Lithuania. *Fresenius Environmental Bulletin* 19: 2929–2934.
- Pliūraite, V. & Kesminas, V. 2012. Ecological impact of Eurasian beaver (*Castor fiber*) activity on macroinvertebrate communities in Lithuanian trout streams. *Central European Journal of Biology* 7: 101–114.
- Poff, N. L., Allan, J. D., Bain, M. B., Karr, J. R., Prestegard, K. L., Richter, B. D., Sparks, R. E. & Stromberg, J. C. 1997. The Natural Flow Regime. *BioScience* 47: 769–784.
- Poff, N. L. & Zimmerman, J. K. H. 2009. Ecological responses to altered flow regimes: a literature review to inform the science and management of environmental flows. *Freshwater Biology* 55: 194–205.
- Porto, L. M., McLaughlin, R. L. & Noakes, D. L. G. 1999. Low-Head Barrier Dams Restrict the Movements of Fishes in Two Lake Ontario Streams. *North American Journal of Fisheries Management* 19: 1028–1036.
- Pozo, J., Orive, E., Fraile, H. & Basaguren, A. 1997. Effects of the Cernadilla-Valparaiso reservoir system on the river Tera. *Regulated Rivers: Research and Management* 13: 57–73.
- R Development Core Team, Vienna, Austria. 2016. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL: <http://www.R-project.org>.
- Rabeni, C. F. 2000. Evaluating physical habitat integrity in relation to the biological potential of streams. *Hydrobiologia* 422/423: 245–256.
- Raesaar, P. 2005. Resource and utilization of Estonian hydropower. *Oil Shale* 22: 233–241.

- Raukas, A. & Rõuk, A. M. 1995. Topography and its Origin. In: Estonian Nature (Ed. Raukas, A.). Tallinn, Valgus and Eesti Entsüklopeediakirjastus, pp. 120–175 (in Estonian).
- Rényi, A. 1970. Probability Theory. Elsevier, New York, Appendix, Sec. 4.
- Rosenberg, D. M., McCully, P. & Pringle, C. M. 2000. Global-scale environmental effects of hydrological alternations: introduction. *Bioscience* 50: 746–752.
- Rosenberg, D. M. & Resh, V. H. (eds.). 1993. Freshwater biomonitoring and benthic macroinvertebrates. Chapman & Hall, New York – London.
- Santucci, V. J. Jr., Gephard, S. R. & Pescitelli, S. M. 2005. Effects of Multiple Low-Head Dams on Fish, Macroinvertebrates, Habitat, and Water Quality in the Fox River, Illinois. *North American Journal of Fisheries Management* 25: 975–992.
- SAS 9.1.3 Help and Documentation. 2004. SAS Institute Inc., Cary, NC, USA.
- Schmedtje, U. & Colling, E. 1996. Ökologische Typisierung der aquatischen Makrofauna. 11 Informationsberichte des Bayerischen Landesamtes für Wasserwirtschaft, Heft 4/96.
- Shao, M. L., Xie, Z. C., Han, X. Q., Cao, M. & Cai, Q. H. 2008. Macroinvertebrate Community Structure in Three-Gorges Reservoir, China. *International Review of Hydrobiology* 93: 175–187.
- Silliman, C., Overton, S. & Baker, M. 2007. The effect of man-made and beaver dams on benthic invertebrate populations. <https://deepblue.lib.umich.edu/handle/2027.42/57586> (Access date: 25.12.2015)
- Skriver, J., Friberg, N. & Kirkegaard, J. 2000. Biological assessment of watercourse quality in Denmark: Introduction of the Danish Stream Fauna Index (DSFI) as the official biomonitoring method. *Verhandlungen des Internationalen Verein Limnologie* 27: 1822–1830.
- Smiley, P. C. Jr. & Dibble, E. D. 2008. Influence of spatial resolution on assessing channelization impacts on fish and macroinvertebrate communities in a warmwater stream in the southeastern United States. *Environmental Monitoring and Assessment* 138: 17–29.
- Spence, J. A. & Hynes, H. B. N. 1971. Differences in Benthos Upstream and Downstream of an Impoundment. *Journal of the Fisheries Research Board of Canada* 28: 35–43.
- StatSoft, Inc. 2011. STATISTICA (data analysis software system), version 10. Website: www.statsoft.com.
- Status classes and class boundaries for surface water bodies and the procedure of classification, 2009. Regulation of the Ministry of the Environment of Estonia. Website: <https://www.riigiteataja.ee/ert/act.jsp?id=13210253&replstring=33> (in Estonian). (Access date: 25.12.2015).
- Takao, A., Kawaguchi, Y., Minagawa, T., Kayaba, Y. & Morimoto, Y. 2008. The relationship between benthic macroinvertebrates and biotic and abiotic environmental characteristics downstream of the Yahagi Dam, Central Japan, and the state change caused by inflow from a tributary. *River Research and Applications* 24: 580–597.
- Timm H. & Vilbaste, S. 2010. Methods to estimate ecological quality of surface waters on the basis of biological elements. Communities of benthic diatoms in running waters. Communities of macroinvertebrates in running and standing waters (in Estonian). Estonian University of Life Sciences, Report to Estonian Ministry of Environment.

- Turunen, J., Muotka, T., Vuori, K.-M., Karjalainen, S. M., Raapysjärvi, J., Sutela, T. & Aroviita, J. 2016. Disentangling the responses of boreal stream assemblages to low stressor levels of diffuse pollution and altered channel morphology. *Science of the Total Environment* 544: 954–962.
- Tuulmets, H. & Aasalo, L. 1980. Streams and lakes. In: *Põllumajandusmaastik Eestis*. Tallinn, pp. 45–50 (in Estonian).
- Vaikre, M., Remm, L. & Rannap, R. 2015. Macroinvertebrates in woodland pools and ditches and their response to artificial drainage in Estonia. *Hydrobiologia* 762: 157–168.
- Van de Bund, W. (ed). 2009. Water Framework Directive intercalibration technical report. Part 1: Rivers. JRC Scientific and Technical Reports, EUR 23838 EN/1.
- Vlek, H. E., Verdonschot, P. F. M. & Nijboer, R. C. 2004. Towards a multimetric index for the assessment of Dutch streams using benthic macroinvertebrates. *Hydrobiologia* 516: 173–189.
- Water Framework Directive. 2002. The European Parliament and the Council of the European Union. Directive 2000/60/EC.

APPENDIX

Reference conditions and quality levels for macroinvertebrate indices in Estonian streams (according to Status classes..., 2009; adapted). R – reference, H – high, G – good, M – moderate, P – poor and B – bad.

Metric	Catchment area, flow velocity and bedrock	R	H	G	M	P or B
Total taxon richness	<100 km ² , fast-flowing	29	>26	23–26	17–22	<17
Total taxon richness	<100 km ² , slow-flowing	18	>16	14–16	11–13	<11
Total taxon richness	100–1,000 km ² , fast-flowing	35	>32	28–32	21–27	<21
Total taxon richness	100–1,000 km ² , slow-flowing	29	>26	23–26	17–22	<17
Total taxon richness	>1,000 km ²	33.5	>30	27–30	20–26	<20
EPT	<100 km ² , fast-flowing	13	>12	10–12	8–9	<8
EPT	<100 km ² , slow-flowing	9	>8	7–8	5–6	<5
EPT	>100 km ²	16,5	>15	13–15	10–12	<10
Shannon diversity	<100 km ² , limestone	2.4	>2.1	1.9–2.1	<1.9–1.4	<1.4
Shannon diversity	<100 km ² , sandstone and >100 km ²	3	>2.7	2.4–2.7	<2.4–1.8	<1.8
ASPT	<100 km ² , slow-flowing	6.1	>5.5	4.9–5.5	<4.9–3.7	<3.7
ASPT	<100 km ² , fast-flowing	6.6	>5.9	5.3–5.9	<5.3–4	<4
ASPT	>100 km ²	6.9	>6.2	5.5–6.2	<5.5–4.1	<4.1
DSFI	<10,000 km ²	7	6–7	5	4	<4

PUBLICATIONS

CURRICULUM VITAE

Name: Kairi Käiro
Date of birth: 30.12.1984
Citizenship: Estonian
Address: University of Tartu, Department of Zoology, Institute of Ecology and Earth Sciences, Vanemuise 46, Tartu, 51014, Estonia
E-mail, phone: kairi.kairo@ut.ee, (+372) 5058497

Education

2003 Puhja Gymnasium
2003–2006 University of Life Sciences, bachelor's degree in Applied Hydrobiology
2006–2008 University of Life Sciences, master degree in Applied Hydrobiology, *cum laude*
2008 –... University of Tartu, Department of Zoology, Institute of Ecology and Earth Sciences, doctoral studies
11.2011–03.2012 Aarhus University, Department of Bioscience, Silkeborg, Denmark, a semester abroad.

Language skills: Estonian, English, German

Professional career:

2012 – ... Estonian University of Life Sciences, Institute of Agricultural and Environmental Sciences, Centre for Limnology, specialist
2006 – 2012 Estonian University of Life Sciences, Institute of Agricultural and Environmental Sciences, Centre for Limnology, laboratory assistant

Research interests: Ecology of benthic macroinvertebrates

Publications:

Timm, H., Käiro, K., Mardi, K. & Timm, T. 2009. Haruldaste suurselgrootute leide Eesti sisevetest. Eesti Looduseuurijate Seltsi Aastaraamat 86: 212–215.
Pall, P., Vilbaste, S., Kõiv, T., Kõrs, A., Käiro, K., Laas, A., Nõges, P., Nõges, T., Piirsoo, K., Toomsalu, L. & Viik, M. 2011. Fluxes of carbon and nutrients through the inflows and outflow of Lake Võrtsjärv, Estonia. *Estonian Journal of Ecology* 60: 39–53, eco.2011.1.04.
Käiro, K., Möls, T., Timm, H., Virro, T. & Järvekülg, R. 2011. The effect of damming on biological quality according to macroinvertebrates in some Estonian streams, Central – Baltic Europe: a pilot study. *River Research and Applications* 27: 895–907, rra.1406.
Timm, H., Käiro, K., Möls, T. & Virro, T. 2011. An index to assess hydromorphological quality of Estonian surface waters based on macroinvertebrate taxonomic composition. *Limnologica* 41: 398–410, j.limno.2011.09.006.

- Käiro, K., Timm, H., Haldna, M. & Virro, T. 2012. Biological Quality on the Basis of Macroinvertebrates in Dammed Habitats of Some Estonian Streams, Central-Baltic Europe. *International Review of Hydrobiology* 97: 497–508, iroh.201111530.
- Piirsoo, K., Viik, M., Kõiv, T., Käiro, K., Laas, A., Nõges, T., Pall, P., Selberg, A., Toomsalu, L. & Vilbaste, S. 2012. Characteristics of dissolved organic matter in the inflows and in the outflow of Lake Võrtsjärv, Estonia. *Journal of Hydrology* 475: 306–313, 10.1016/j.jhydrol.2012.10.015.
- Kõrs, A., Vilbaste, S., Käiro, K., Pall, P., Piirsoo, K., Truu, J. & Viik, M. 2012. Temporal changes in the composition of macrophyte communities and environmental factors governing the distribution of aquatic plants in an unregulated lowland stream, the River Emajõgi (Estonia). *Boreal Environment Research* 17: 460–472.

Conference presentations:

- Käiro, K. (2006) Biological quality of the Selja and Pühajõgi Streams on the basis of taxonomical composition of macroinvertebrates in Estonia (poster presentation). Biodiversity and functioning of aquatic ecosystems in the Baltic Sea region. Lithuania, Klaipeda 2006, October 7–8.
- Käiro, K. (2007) Relationships between macroinvertebrates, benthic diatoms and water chemistry in Estonian running waters (poster presentation). Nordic Benthological Meeting VI. Kalmar, Sweden, 2007, June 11–13.
- Käiro, K., Timm, H. & Virro, T. (2009). Differences between stream reservoirs and lakes, according to littoral macroinvertebrates (poster presentation). A benthic perspective to restoration, mitigation and adaptation of freshwaters: Nordic Benthological Meeting VII. Tartu, Estonia, September 7–10 2009.
- Käiro, K., Timm, H. & Virro, T. (2011). Biological quality of dammed streams in Estonia according to macroinvertebrates (oral presentation). Ecosystem services in soil and water research: Focus on Soils and Water Symposium. 7–10 June 2011 Uppsala, Sweden.
- Timm, H., Käiro, K., Möls, T. & Virro, T. (2011). Relationships of macroinvertebrates with flow velocity and bottom type in freshwaters of Estonia (oral presentation). Food webs and climate change: Nordic Benthological Meeting VIII. Aalborg, Denmark 9–12 May, 2011.

Administrative responsibilities:

- September 2009 7th meeting of NORBS (Nordic Benthological Society) in Tartu, Estonia, co-ordinator

Awards and scholarships:

- 2011 ESF Dora T6 scholarship, Participation of doctoral students' in doctoral studies and research at foreign universities and research institutions
- 2011 Travelling grant from the Doctoral School of Earth Sciences and Ecology
- 2006 The third award in undergraduates sciences work competition in bio- and environmental science

ELULOOKIRJELDUS

Nimi: Kairi Käiro
Sünniaeg: 30.12.1984
Kodakondsus: Estonian
Aadress: Tartu Ülikool, Zooloogia osakond, Ökoloogia ja Maateaduste Instituut, Vanemuise 46, Tartu, 51014, Eesti
E-post, telefon: kairi.kairo@ut.ee, (+372) 5058497

Haridus:
2003 Puhja Gümnaasium
2003–2006 BSc hüdrobioloogia erialal, Eesti Maaülikool
2006–2008 MSc hüdrobioloogia erialal, Eesti Maaülikool, *cum laude*
2008–... doktorantuur, Tartu Ülikool, Zooloogia osakond, Ökoloogia ja Maateaduste Instituut
11.2011–03.2012 Aarhushi Ülikool, Silkeborg, Bioteaduste osakond; semester välismaal

Keelteoskus: eesti, inglise, saksa

Teenistuskäik:
2012–... Eesti Maaülikool, põllumajandus- ja keskkonnainstituut, limnoloogiakeskus, spetsialist
2006–2012 Eesti Maaülikool, põllumajandus- ja keskkonnainstituut, limnoloogiakeskus, laborant

Teadustöö põhisuunad: suurselgrootute ökoloogia

Publikatsioonid:

- Timm, H., Käiro, K., Mardi, K. & Timm, T. 2009. Haruldaste suurselgrootute leide Eesti sisevetest. Eesti Looduseuurijate Seltsi Aastaraamat 86: 212–215.
- Pall, P., Vilbaste, S., Kõiv, T., Kõrs, A., Käiro, K., Laas, A., Nõges, P., Nõges, T., Piirsoo, K., Toomsalu, L. & Viik, M. 2011. Fluxes of carbon and nutrients through the inflows and outflow of Lake Võrtsjärv, Estonia. *Estonian Journal of Ecology* 60: 39–53, eco.2011.1.04.
- Käiro, K., Möls, T., Timm, H., Virro, T. & Järvekülg, R. 2011. The effect of damming on biological quality according to macroinvertebrates in some Estonian streams, Central – Baltic Europe: a pilot study. *River Research and Applications* 27: 895–907, rra.1406.
- Timm, H., Käiro, K., Möls, T. & Virro, T. 2011. An index to assess hydro-morphological quality of Estonian surface waters based on macroinvertebrate taxonomic composition. *Limnologica* 41: 398–410, j.limno.2011.09.006.
- Käiro, K., Timm, H., Haldna, M. & Virro, T. 2012. Biological Quality on the Basis of Macroinvertebrates in Dammed Habitats of Some Estonian Streams, Central–Baltic Europe. *International Review of Hydrobiology* 97: 497–508, iroh.201111530.

- Piirsoo, K., Viik, M., Kõiv, T., Käiro, K., Laas, A., Nõges, T., Pall, P., Selberg, A., Toomsalu, L. & Vilbaste, S. 2012. Characteristics of dissolved organic matter in the inflows and in the outflow of Lake Võrtsjärv, Estonia. *Journal of Hydrology* 475: 306–313, 10.1016/j.jhydrol.2012.10.015.
- Kõrs, A., Vilbaste, S., Käiro, K., Pall, P., Piirsoo, K., Truu, J. & Viik, M. 2012. Temporal changes in the composition of macrophyte communities and environmental factors governing the distribution of aquatic plants in an unregulated lowland stream, the River Emajõgi (Estonia). *Boreal Environment Research* 17: 460–472.

Konverentsiettekanded:

- Käiro, K. (2006) Biological quality of the Selja and Pühajõgi Streams on the basis of taxonomical composition of macroinvertebrates in Estonia (poster ettekanne). Biodiversity and functioning of aquatic ecosystems in the Baltic Sea region. Klaipeda, Leedu, 7–8. oktoober 2006.
- Käiro, K. (2007) Relationships between macroinvertebrates, benthic diatoms and water chemistry in Estonian running waters (poster ettekanne). Nordic Benthological Society Meeting VI, Kalmar, Rootsi, 11–13. juuni 2007.
- Käiro, Timm, H. & Virro, T. (2009). Differences between stream reservoirs and lakes, according to littoral macroinvertebrates (poster ettekanne). A benthic perspective to restoration, mitigation and adaptation of freshwaters: Nordic Benthological Society Meeting VII Tartu, Eesti, 7–10. september 2009.
- Käiro, K., Timm, H. & Virro, T. (2011). Biological quality of dammed streams in Estonia according to macroinvertebrates. (suuline ettekanne). Ecosystem services in soil and water research: Focus on Soils and Water Symposium. Uppsala, Rootsi, 7–10. juuni 2011.
- Timm, H., Käiro, K., Möls, T., Virro, T. (2011). Relationships of macroinvertebrates with flow velocity and bottom type in freshwaters of Estonia. (suuline ettekanne) Food webs and climate change: Nordic Benthological Meeting VIII. Aalborg, Taani, 9–12. mai 2011.

Teadusorganisatsiooniline ja -administratiivne tegevus:

September 2009 NORBS (Nordic Benthological Society) 7. konverentsi kaaskorraldaja, 7–10. september, Tartu, Eesti.

Saadud uurimistoetused ja tunnustus:

- 2011 ESF Dora T6 stipendium, doktorantide semester välismaal
- 2011 Maateaduste ja ökoloogia doktorikooli välissõidutoetus
- 2006 Üliõpilaste teadustööde riikliku konkursi III preemia rakenduskõrghariduse ja bakalaureuseõppe üliõpilaste astmes.

DISSERTATIONES BIOLOGICAE UNIVERSITATIS TARTUENSIS

1. **Toivo Maimets.** Studies of human oncoprotein p53. Tartu, 1991, 96 p.
2. **Enn K. Seppet.** Thyroid state control over energy metabolism, ion transport and contractile functions in rat heart. Tartu, 1991, 135 p.
3. **Kristjan Zobel.** Epifüütsete makrosamblike väärtus õhu saastuse indikaatoritena Hamar-Dobani boreaalsetes mägimetsades. Tartu, 1992, 131 lk.
4. **Andres Mäe.** Conjugal mobilization of catabolic plasmids by transposable elements in helper plasmids. Tartu, 1992, 91 p.
5. **Maia Kivisaar.** Studies on phenol degradation genes of *Pseudomonas* sp. strain EST 1001. Tartu, 1992, 61 p.
6. **Allan Nurk.** Nucleotide sequences of phenol degradative genes from *Pseudomonas* sp. strain EST 1001 and their transcriptional activation in *Pseudomonas putida*. Tartu, 1992, 72 p.
7. **Ülo Tamm.** The genus *Populus* L. in Estonia: variation of the species biology and introduction. Tartu, 1993, 91 p.
8. **Jaanus Remme.** Studies on the peptidyltransferase centre of the *E.coli* ribosome. Tartu, 1993, 68 p.
9. **Ülo Langel.** Galanin and galanin antagonists. Tartu, 1993, 97 p.
10. **Arvo Käär.** The development of an automatic online dynamic fluorescence-based pH-dependent fiber optic penicillin flowthrough biosensor for the control of the benzylpenicillin hydrolysis. Tartu, 1993, 117 p.
11. **Lilian Järvekülg.** Antigenic analysis and development of sensitive immunoassay for potato viruses. Tartu, 1993, 147 p.
12. **Jaak Palumets.** Analysis of phytomass partition in Norway spruce. Tartu, 1993, 47 p.
13. **Arne Sellin.** Variation in hydraulic architecture of *Picea abies* (L.) Karst. trees grown under different environmental conditions. Tartu, 1994, 119 p.
13. **Mati Reeben.** Regulation of light neurofilament gene expression. Tartu, 1994, 108 p.
14. **Urmas Tartes.** Respiration rhythms in insects. Tartu, 1995, 109 p.
15. **Ülo Puurand.** The complete nucleotide sequence and infections *in vitro* transcripts from cloned cDNA of a potato A potyvirus. Tartu, 1995, 96 p.
16. **Peeter Hõrak.** Pathways of selection in avian reproduction: a functional framework and its application in the population study of the great tit (*Parus major*). Tartu, 1995, 118 p.
17. **Erkki Truve.** Studies on specific and broad spectrum virus resistance in transgenic plants. Tartu, 1996, 158 p.
18. **Illar Pata.** Cloning and characterization of human and mouse ribosomal protein S6-encoding genes. Tartu, 1996, 60 p.
19. **Ülo Niinemets.** Importance of structural features of leaves and canopy in determining species shade-tolerance in temperature deciduous woody taxa. Tartu, 1996, 150 p.

20. **Ants Kurg.** Bovine leukemia virus: molecular studies on the packaging region and DNA diagnostics in cattle. Tartu, 1996, 104 p.
21. **Ene Ustav.** E2 as the modulator of the BPV1 DNA replication. Tartu, 1996, 100 p.
22. **Aksel Soosaar.** Role of helix-loop-helix and nuclear hormone receptor transcription factors in neurogenesis. Tartu, 1996, 109 p.
23. **Maido Remm.** Human papillomavirus type 18: replication, transformation and gene expression. Tartu, 1997, 117 p.
24. **Tiiu Kull.** Population dynamics in *Cypripedium calceolus* L. Tartu, 1997, 124 p.
25. **Kalle Olli.** Evolutionary life-strategies of autotrophic planktonic microorganisms in the Baltic Sea. Tartu, 1997, 180 p.
26. **Meelis Pärtel.** Species diversity and community dynamics in calcareous grassland communities in Western Estonia. Tartu, 1997, 124 p.
27. **Malle Leht.** The Genus *Potentilla* L. in Estonia, Latvia and Lithuania: distribution, morphology and taxonomy. Tartu, 1997, 186 p.
28. **Tanel Tenson.** Ribosomes, peptides and antibiotic resistance. Tartu, 1997, 80 p.
29. **Arvo Tuvikene.** Assessment of inland water pollution using biomarker responses in fish *in vivo* and *in vitro*. Tartu, 1997, 160 p.
30. **Urmas Saarma.** Tuning ribosomal elongation cycle by mutagenesis of 23S rRNA. Tartu, 1997, 134 p.
31. **Henn Ojaveer.** Composition and dynamics of fish stocks in the gulf of Riga ecosystem. Tartu, 1997, 138 p.
32. **Lembi Lõugas.** Post-glacial development of vertebrate fauna in Estonian water bodies. Tartu, 1997, 138 p.
33. **Margus Pooga.** Cell penetrating peptide, transportan, and its predecessors, galanin-based chimeric peptides. Tartu, 1998, 110 p.
34. **Andres Saag.** Evolutionary relationships in some cetrarioid genera (Lichenized Ascomycota). Tartu, 1998, 196 p.
35. **Aivar Liiv.** Ribosomal large subunit assembly *in vivo*. Tartu, 1998, 158 p.
36. **Tatjana Oja.** Isoenzyme diversity and phylogenetic affinities among the eurasian annual bromes (*Bromus* L., Poaceae). Tartu, 1998, 92 p.
37. **Mari Moora.** The influence of arbuscular mycorrhizal (AM) symbiosis on the competition and coexistence of calcareous grassland plant species. Tartu, 1998, 78 p.
38. **Olavi Kurina.** Fungus gnats in Estonia (*Diptera: Bolitophilidae, Kero-platidae, Macroceridae, Ditomyiidae, Diadocidiidae, Mycetophilidae*). Tartu, 1998, 200 p.
39. **Andrus Tasa.** Biological leaching of shales: black shale and oil shale. Tartu, 1998, 98 p.
40. **Arnold Kristjuhan.** Studies on transcriptional activator properties of tumor suppressor protein p53. Tartu, 1998, 86 p.
41. **Sulev Ingerpuu.** Characterization of some human myeloid cell surface and nuclear differentiation antigens. Tartu, 1998, 163 p.

42. **Veljo Kisand**. Responses of planktonic bacteria to the abiotic and biotic factors in the shallow lake Võrtsjärv. Tartu, 1998, 118 p.
43. **Kadri Põldmaa**. Studies in the systematics of hypomyces and allied genera (Hypocreales, Ascomycota). Tartu, 1998, 178 p.
44. **Markus Vetemaa**. Reproduction parameters of fish as indicators in environmental monitoring. Tartu, 1998, 117 p.
45. **Heli Talvik**. Prepatent periods and species composition of different *Oesophagostomum* spp. populations in Estonia and Denmark. Tartu, 1998, 104 p.
46. **Katrin Heinsoo**. Cuticular and stomatal antechamber conductance to water vapour diffusion in *Picea abies* (L.) karst. Tartu, 1999, 133 p.
47. **Tarmo Annilo**. Studies on mammalian ribosomal protein S7. Tartu, 1998, 77 p.
48. **Indrek Ots**. Health state indices of reproducing great tits (*Parus major*): sources of variation and connections with life-history traits. Tartu, 1999, 117 p.
49. **Juan Jose Cantero**. Plant community diversity and habitat relationships in central Argentina grasslands. Tartu, 1999, 161 p.
50. **Rein Kalamees**. Seed bank, seed rain and community regeneration in Estonian calcareous grasslands. Tartu, 1999, 107 p.
51. **Sulev Kõks**. Cholecystokinin (CCK) — induced anxiety in rats: influence of environmental stimuli and involvement of endopioid mechanisms and serotonin. Tartu, 1999, 123 p.
52. **Ebe Sild**. Impact of increasing concentrations of O₃ and CO₂ on wheat, clover and pasture. Tartu, 1999, 123 p.
53. **Ljudmilla Timofejeva**. Electron microscopical analysis of the synaptosomal complex formation in cereals. Tartu, 1999, 99 p.
54. **Andres Valkna**. Interactions of galanin receptor with ligands and G-proteins: studies with synthetic peptides. Tartu, 1999, 103 p.
55. **Taavi Virro**. Life cycles of planktonic rotifers in lake Peipsi. Tartu, 1999, 101 p.
56. **Ana Rebane**. Mammalian ribosomal protein S3a genes and intron-encoded small nucleolar RNAs U73 and U82. Tartu, 1999, 85 p.
57. **Tiina Tamm**. Cocksfoot mottle virus: the genome organisation and translational strategies. Tartu, 2000, 101 p.
58. **Reet Kurg**. Structure-function relationship of the bovine papilloma virus E2 protein. Tartu, 2000, 89 p.
59. **Toomas Kivisild**. The origins of Southern and Western Eurasian populations: an mtDNA study. Tartu, 2000, 121 p.
60. **Niilo Kaldalu**. Studies of the TOL plasmid transcription factor XylS. Tartu 2000. 88 p.
61. **Dina Lepik**. Modulation of viral DNA replication by tumor suppressor protein p53. Tartu 2000. 106 p.

62. **Kai Vellak**. Influence of different factors on the diversity of the bryophyte vegetation in forest and wooded meadow communities. Tartu 2000. 122 p.
63. **Jonne Kotta**. Impact of eutrophication and biological invasions on the structure and functions of benthic macrofauna. Tartu 2000. 160 p.
64. **Georg Martin**. Phytobenthic communities of the Gulf of Riga and the inner sea the West-Estonian archipelago. Tartu, 2000. 139 p.
65. **Silvia Sepp**. Morphological and genetical variation of *Alchemilla L.* in Estonia. Tartu, 2000. 124 p.
66. **Jaan Liira**. On the determinants of structure and diversity in herbaceous plant communities. Tartu, 2000. 96 p.
67. **Priit Zingel**. The role of planktonic ciliates in lake ecosystems. Tartu 2001. 111 p.
68. **Tiit Teder**. Direct and indirect effects in Host-parasitoid interactions: ecological and evolutionary consequences. Tartu 2001. 122 p.
69. **Hannes Kollist**. Leaf apoplastic ascorbate as ozone scavenger and its transport across the plasma membrane. Tartu 2001. 80 p.
70. **Reet Marits**. Role of two-component regulator system PehR-PehS and extracellular protease PrtW in virulence of *Erwinia Carotovora* subsp. *Carotovora*. Tartu 2001. 112 p.
71. **Vallo Tilgar**. Effect of calcium supplementation on reproductive performance of the pied flycatcher *Ficedula hypoleuca* and the great tit *Parus major*, breeding in Northern temperate forests. Tartu, 2002. 126 p.
72. **Rita Hõrak**. Regulation of transposition of transposon Tn4652 in *Pseudomonas putida*. Tartu, 2002. 108 p.
73. **Liina Eek-Piirsoo**. The effect of fertilization, mowing and additional illumination on the structure of a species-rich grassland community. Tartu, 2002. 74 p.
74. **Krõõt Aasamaa**. Shoot hydraulic conductance and stomatal conductance of six temperate deciduous tree species. Tartu, 2002. 110 p.
75. **Nele Ingerpuu**. Bryophyte diversity and vascular plants. Tartu, 2002. 112 p.
76. **Neeme Tõnisson**. Mutation detection by primer extension on oligonucleotide microarrays. Tartu, 2002. 124 p.
77. **Margus Pensa**. Variation in needle retention of Scots pine in relation to leaf morphology, nitrogen conservation and tree age. Tartu, 2003. 110 p.
78. **Asko Lõhmus**. Habitat preferences and quality for birds of prey: from principles to applications. Tartu, 2003. 168 p.
79. **Viljar Jaks**. p53 — a switch in cellular circuit. Tartu, 2003. 160 p.
80. **Jaana Männik**. Characterization and genetic studies of four ATP-binding cassette (ABC) transporters. Tartu, 2003. 140 p.
81. **Marek Sammul**. Competition and coexistence of clonal plants in relation to productivity. Tartu, 2003. 159 p.
82. **Ivar Ilves**. Virus-cell interactions in the replication cycle of bovine papillomavirus type 1. Tartu, 2003. 89 p.

83. **Andres Männik.** Design and characterization of a novel vector system based on the stable replicator of bovine papillomavirus type 1. Tartu, 2003. 109 p.
84. **Ivika Ostonen.** Fine root structure, dynamics and proportion in net primary production of Norway spruce forest ecosystem in relation to site conditions. Tartu, 2003. 158 p.
85. **Godrun Veldre.** Somatic status of 12–15-year-old Tartu schoolchildren. Tartu, 2003. 199 p.
86. **Ülo Väli.** The greater spotted eagle *Aquila clanga* and the lesser spotted eagle *A. pomarina*: taxonomy, phylogeography and ecology. Tartu, 2004. 159 p.
87. **Aare Abroi.** The determinants for the native activities of the bovine papillomavirus type 1 E2 protein are separable. Tartu, 2004. 135 p.
88. **Tiina Kahre.** Cystic fibrosis in Estonia. Tartu, 2004. 116 p.
89. **Helen Orav-Kotta.** Habitat choice and feeding activity of benthic suspension feeders and mesograzers in the northern Baltic Sea. Tartu, 2004. 117 p.
90. **Maarja Õpik.** Diversity of arbuscular mycorrhizal fungi in the roots of perennial plants and their effect on plant performance. Tartu, 2004. 175 p.
91. **Kadri Tali.** Species structure of *Neotinea ustulata*. Tartu, 2004. 109 p.
92. **Kristiina Tambets.** Towards the understanding of post-glacial spread of human mitochondrial DNA haplogroups in Europe and beyond: a phylogeographic approach. Tartu, 2004. 163 p.
93. **Arvi Jõers.** Regulation of p53-dependent transcription. Tartu, 2004. 103 p.
94. **Lilian Kadaja.** Studies on modulation of the activity of tumor suppressor protein p53. Tartu, 2004. 103 p.
95. **Jaak Truu.** Oil shale industry wastewater: impact on river microbial community and possibilities for bioremediation. Tartu, 2004. 128 p.
96. **Maire Peters.** Natural horizontal transfer of the *pheBA* operon. Tartu, 2004. 105 p.
97. **Ülo Maiväli.** Studies on the structure-function relationship of the bacterial ribosome. Tartu, 2004. 130 p.
98. **Merit Otsus.** Plant community regeneration and species diversity in dry calcareous grasslands. Tartu, 2004. 103 p.
99. **Mikk Heidema.** Systematic studies on sawflies of the genera *Dolerus*, *Empria*, and *Caliroa* (Hymenoptera: Tenthredinidae). Tartu, 2004. 167 p.
100. **Ilmar Tõnno.** The impact of nitrogen and phosphorus concentration and N/P ratio on cyanobacterial dominance and N₂ fixation in some Estonian lakes. Tartu, 2004. 111 p.
101. **Lauri Saks.** Immune function, parasites, and carotenoid-based ornaments in greenfinches. Tartu, 2004. 144 p.
102. **Siiri Rootsi.** Human Y-chromosomal variation in European populations. Tartu, 2004. 142 p.

103. **Eve Vedler.** Structure of the 2,4-dichloro-phenoxyacetic acid-degradative plasmid pEST4011. Tartu, 2005. 106 p.
104. **Andres Tover.** Regulation of transcription of the phenol degradation *pheBA* operon in *Pseudomonas putida*. Tartu, 2005. 126 p.
105. **Helen Udras.** Hexose kinases and glucose transport in the yeast *Hansenula polymorpha*. Tartu, 2005. 100 p.
106. **Ave Suija.** Lichens and lichenicolous fungi in Estonia: diversity, distribution patterns, taxonomy. Tartu, 2005. 162 p.
107. **Piret Lõhmus.** Forest lichens and their substrata in Estonia. Tartu, 2005. 162 p.
108. **Inga Lips.** Abiotic factors controlling the cyanobacterial bloom occurrence in the Gulf of Finland. Tartu, 2005. 156 p.
109. **Kaasik, Krista.** Circadian clock genes in mammalian clockwork, metabolism and behaviour. Tartu, 2005. 121 p.
110. **Juhan Javoš.** The effects of experience on host acceptance in ovipositing moths. Tartu, 2005. 112 p.
111. **Tiina Sedman.** Characterization of the yeast *Saccharomyces cerevisiae* mitochondrial DNA helicase Hmi1. Tartu, 2005. 103 p.
112. **Ruth Aguraiuja.** Hawaiian endemic fern lineage *Diellia* (Aspleniaceae): distribution, population structure and ecology. Tartu, 2005. 112 p.
113. **Riho Teras.** Regulation of transcription from the fusion promoters generated by transposition of Tn4652 into the upstream region of *pheBA* operon in *Pseudomonas putida*. Tartu, 2005. 106 p.
114. **Mait Metspalu.** Through the course of prehistory in india: tracing the mtDNA trail. Tartu, 2005. 138 p.
115. **Elin Lõhmussaar.** The comparative patterns of linkage disequilibrium in European populations and its implication for genetic association studies. Tartu, 2006. 124 p.
116. **Priit Kupper.** Hydraulic and environmental limitations to leaf water relations in trees with respect to canopy position. Tartu, 2006. 126 p.
117. **Heili Ilves.** Stress-induced transposition of Tn4652 in *Pseudomonas Putida*. Tartu, 2006. 120 p.
118. **Silja Kuusk.** Biochemical properties of Hmi1p, a DNA helicase from *Saccharomyces cerevisiae* mitochondria. Tartu, 2006. 126 p.
119. **Kersti Püssa.** Forest edges on medium resolution landsat thematic mapper satellite images. Tartu, 2006. 90 p.
120. **Lea Tummeleht.** Physiological condition and immune function in great tits (*Parus major* L.): Sources of variation and trade-offs in relation to growth. Tartu, 2006. 94 p.
121. **Toomas Esperk.** Larval instar as a key element of insect growth schedules. Tartu, 2006. 186 p.
122. **Harri Valdmann.** Lynx (*Lynx lynx*) and wolf (*Canis lupus*) in the Baltic region: Diets, helminth parasites and genetic variation. Tartu, 2006. 102 p.

123. **Priit Jõers.** Studies of the mitochondrial helicase Hmi1p in *Candida albicans* and *Saccharomyces cerevisia*. Tartu, 2006. 113 p.
124. **Kersti Lilleväli.** Gata3 and Gata2 in inner ear development. Tartu, 2007. 123 p.
125. **Kai Rünk.** Comparative ecology of three fern species: *Dryopteris carthusiana* (Vill.) H.P. Fuchs, *D. expansa* (C. Presl) Fraser-Jenkins & Jermy and *D. dilatata* (Hoffm.) A. Gray (Dryopteridaceae). Tartu, 2007. 143 p.
126. **Aveliina Helm.** Formation and persistence of dry grassland diversity: role of human history and landscape structure. Tartu, 2007. 89 p.
127. **Leho Tedersoo.** Ectomycorrhizal fungi: diversity and community structure in Estonia, Seychelles and Australia. Tartu, 2007. 233 p.
128. **Marko Mägi.** The habitat-related variation of reproductive performance of great tits in a deciduous-coniferous forest mosaic: looking for causes and consequences. Tartu, 2007. 135 p.
129. **Valeria Lulla.** Replication strategies and applications of Semliki Forest virus. Tartu, 2007. 109 p.
130. **Ülle Reier.** Estonian threatened vascular plant species: causes of rarity and conservation. Tartu, 2007. 79 p.
131. **Inga Jüriado.** Diversity of lichen species in Estonia: influence of regional and local factors. Tartu, 2007. 171 p.
132. **Tatjana Krama.** Mobbing behaviour in birds: costs and reciprocity based cooperation. Tartu, 2007. 112 p.
133. **Signe Saumaa.** The role of DNA mismatch repair and oxidative DNA damage defense systems in avoidance of stationary phase mutations in *Pseudomonas putida*. Tartu, 2007. 172 p.
134. **Reedik Mägi.** The linkage disequilibrium and the selection of genetic markers for association studies in european populations. Tartu, 2007. 96 p.
135. **Priit Kilgas.** Blood parameters as indicators of physiological condition and skeletal development in great tits (*Parus major*): natural variation and application in the reproductive ecology of birds. Tartu, 2007. 129 p.
136. **Anu Albert.** The role of water salinity in structuring eastern Baltic coastal fish communities. Tartu, 2007. 95 p.
137. **Kärt Padari.** Protein transduction mechanisms of transportans. Tartu, 2008. 128 p.
138. **Siiri-Lii Sandre.** Selective forces on larval colouration in a moth. Tartu, 2008. 125 p.
139. **Ülle Jõgar.** Conservation and restoration of semi-natural floodplain meadows and their rare plant species. Tartu, 2008. 99 p.
140. **Lauri Laanisto.** Macroecological approach in vegetation science: generality of ecological relationships at the global scale. Tartu, 2008. 133 p.
141. **Reidar Andreson.** Methods and software for predicting PCR failure rate in large genomes. Tartu, 2008. 105 p.
142. **Birgot Paavel.** Bio-optical properties of turbid lakes. Tartu, 2008. 175 p.
143. **Kaire Torn.** Distribution and ecology of charophytes in the Baltic Sea. Tartu, 2008, 98 p.

144. **Vladimir Vimberg**. Peptide mediated macrolide resistance. Tartu, 2008, 190 p.
145. **Daima Örd**. Studies on the stress-inducible pseudokinase TRB3, a novel inhibitor of transcription factor ATF4. Tartu, 2008, 108 p.
146. **Lauri Saag**. Taxonomic and ecologic problems in the genus *Lepraria* (*Stereocaulaceae*, lichenised *Ascomycota*). Tartu, 2008, 175 p.
147. **Ulvi Karu**. Antioxidant protection, carotenoids and coccidians in greenfinches – assessment of the costs of immune activation and mechanisms of parasite resistance in a passerine with carotenoid-based ornaments. Tartu, 2008, 124 p.
148. **Jaanus Remm**. Tree-cavities in forests: density, characteristics and occupancy by animals. Tartu, 2008, 128 p.
149. **Epp Moks**. Tapeworm parasites *Echinococcus multilocularis* and *E. granulosus* in Estonia: phylogenetic relationships and occurrence in wild carnivores and ungulates. Tartu, 2008, 82 p.
150. **Eve Eensalu**. Acclimation of stomatal structure and function in tree canopy: effect of light and CO₂ concentration. Tartu, 2008, 108 p.
151. **Janne Pullat**. Design, functionlization and application of an *in situ* synthesized oligonucleotide microarray. Tartu, 2008, 108 p.
152. **Marta Putrinš**. Responses of *Pseudomonas putida* to phenol-induced metabolic and stress signals. Tartu, 2008, 142 p.
153. **Marina Semtšenko**. Plant root behaviour: responses to neighbours and physical obstructions. Tartu, 2008, 106 p.
154. **Marge Starast**. Influence of cultivation techniques on productivity and fruit quality of some *Vaccinium* and *Rubus* taxa. Tartu, 2008, 154 p.
155. **Age Tats**. Sequence motifs influencing the efficiency of translation. Tartu, 2009, 104 p.
156. **Radi Tegova**. The role of specialized DNA polymerases in mutagenesis in *Pseudomonas putida*. Tartu, 2009, 124 p.
157. **Tsipe Aavik**. Plant species richness, composition and functional trait pattern in agricultural landscapes – the role of land use intensity and landscape structure. Tartu, 2009, 112 p.
158. **Kaja Kiiver**. Semliki forest virus based vectors and cell lines for studying the replication and interactions of alphaviruses and hepaciviruses. Tartu, 2009, 104 p.
159. **Meelis Kadaja**. Papillomavirus Replication Machinery Induces Genomic Instability in its Host Cell. Tartu, 2009, 126 p.
160. **Pille Hallast**. Human and chimpanzee Luteinizing hormone/Chorionic Gonadotropin beta (*LHB/CGB*) gene clusters: diversity and divergence of young duplicated genes. Tartu, 2009, 168 p.
161. **Ain Vellak**. Spatial and temporal aspects of plant species conservation. Tartu, 2009, 86 p.
162. **Triinu Remmel**. Body size evolution in insects with different colouration strategies: the role of predation risk. Tartu, 2009, 168 p.

163. **Jaana Salujõe**. Zooplankton as the indicator of ecological quality and fish predation in lake ecosystems. Tartu, 2009, 129 p.
164. **Ele Vahtmäe**. Mapping benthic habitat with remote sensing in optically complex coastal environments. Tartu, 2009, 109 p.
165. **Liisa Metsamaa**. Model-based assessment to improve the use of remote sensing in recognition and quantitative mapping of cyanobacteria. Tartu, 2009, 114 p.
166. **Pille Säälük**. The role of endocytosis in the protein transduction by cell-penetrating peptides. Tartu, 2009, 155 p.
167. **Lauri Peil**. Ribosome assembly factors in *Escherichia coli*. Tartu, 2009, 147 p.
168. **Lea Hallik**. Generality and specificity in light harvesting, carbon gain capacity and shade tolerance among plant functional groups. Tartu, 2009, 99 p.
169. **Mariliis Tark**. Mutagenic potential of DNA damage repair and tolerance mechanisms under starvation stress. Tartu, 2009, 191 p.
170. **Riinu Rannap**. Impacts of habitat loss and restoration on amphibian populations. Tartu, 2009, 117 p.
171. **Maarja Adojaan**. Molecular variation of HIV-1 and the use of this knowledge in vaccine development. Tartu, 2009, 95 p.
172. **Signe Altmäe**. Genomics and transcriptomics of human induced ovarian folliculogenesis. Tartu, 2010, 179 p.
173. **Triin Suvi**. Mycorrhizal fungi of native and introduced trees in the Seychelles Islands. Tartu, 2010, 107 p.
174. **Velda Lauringson**. Role of suspension feeding in a brackish-water coastal sea. Tartu, 2010, 123 p.
175. **Eero Talts**. Photosynthetic cyclic electron transport – measurement and variably proton-coupled mechanism. Tartu, 2010, 121 p.
176. **Mari Nelis**. Genetic structure of the Estonian population and genetic distance from other populations of European descent. Tartu, 2010, 97 p.
177. **Kaarel Krjutškov**. Arrayed Primer Extension-2 as a multiplex PCR-based method for nucleic acid variation analysis: method and applications. Tartu, 2010, 129 p.
178. **Egle Köster**. Morphological and genetical variation within species complexes: *Anthyllis vulneraria* s. l. and *Alchemilla vulgaris* (coll.). Tartu, 2010, 101 p.
179. **Erki Õunap**. Systematic studies on the subfamily Sterrhinae (Lepidoptera: Geometridae). Tartu, 2010, 111 p.
180. **Merike Jõesaar**. Diversity of key catabolic genes at degradation of phenol and *p*-cresol in pseudomonads. Tartu, 2010, 125 p.
181. **Kristjan Herkül**. Effects of physical disturbance and habitat-modifying species on sediment properties and benthic communities in the northern Baltic Sea. Tartu, 2010, 123 p.
182. **Arto Pulk**. Studies on bacterial ribosomes by chemical modification approaches. Tartu, 2010, 161 p.

183. **Maria Põllupüü**. Ecological relations of cladocerans in a brackish-water ecosystem. Tartu, 2010, 126 p.
184. **Toomas Silla**. Study of the segregation mechanism of the Bovine Papillomavirus Type 1. Tartu, 2010, 188 p.
185. **Gyaneshwer Chaubey**. The demographic history of India: A perspective based on genetic evidence. Tartu, 2010, 184 p.
186. **Katrin Kepp**. Genes involved in cardiovascular traits: detection of genetic variation in Estonian and Czech populations. Tartu, 2010, 164 p.
187. **Virve Sõber**. The role of biotic interactions in plant reproductive performance. Tartu, 2010, 92 p.
188. **Kersti Kangro**. The response of phytoplankton community to the changes in nutrient loading. Tartu, 2010, 144 p.
189. **Joachim M. Gerhold**. Replication and Recombination of mitochondrial DNA in Yeast. Tartu, 2010, 120 p.
190. **Helen Tammert**. Ecological role of physiological and phylogenetic diversity in aquatic bacterial communities. Tartu, 2010, 140 p.
191. **Elle Rajandu**. Factors determining plant and lichen species diversity and composition in Estonian *Calamagrostis* and *Hepatica* site type forests. Tartu, 2010, 123 p.
192. **Paula Ann Kivistik**. ColR-ColS signalling system and transposition of Tn4652 in the adaptation of *Pseudomonas putida*. Tartu, 2010, 118 p.
193. **Siim Sõber**. Blood pressure genetics: from candidate genes to genome-wide association studies. Tartu, 2011, 120 p.
194. **Kalle Kipper**. Studies on the role of helix 69 of 23S rRNA in the factor-dependent stages of translation initiation, elongation, and termination. Tartu, 2011, 178 p.
195. **Triinu Siibak**. Effect of antibiotics on ribosome assembly is indirect. Tartu, 2011, 134 p.
196. **Tambet Tõnissoo**. Identification and molecular analysis of the role of guanine nucleotide exchange factor RIC-8 in mouse development and neural function. Tartu, 2011, 110 p.
197. **Helin Räägel**. Multiple faces of cell-penetrating peptides – their intracellular trafficking, stability and endosomal escape during protein transduction. Tartu, 2011, 161 p.
198. **Andres Jaanus**. Phytoplankton in Estonian coastal waters – variability, trends and response to environmental pressures. Tartu, 2011, 157 p.
199. **Tiit Nikopensius**. Genetic predisposition to nonsyndromic orofacial clefts. Tartu, 2011, 152 p.
200. **Signe Värvi**. Studies on the mechanisms of RNA polymerase II-dependent transcription elongation. Tartu, 2011, 108 p.
201. **Kristjan Väik**. Gene expression profiling and genome-wide association studies of non-small cell lung cancer. Tartu, 2011, 98 p.
202. **Arno Põllumäe**. Spatio-temporal patterns of native and invasive zooplankton species under changing climate and eutrophication conditions. Tartu, 2011, 153 p.

203. **Egle Tammeleht**. Brown bear (*Ursus arctos*) population structure, demographic processes and variations in diet in northern Eurasia. Tartu, 2011, 143 p.
205. **Teele Jairus**. Species composition and host preference among ectomycorrhizal fungi in Australian and African ecosystems. Tartu, 2011, 106 p.
206. **Kessy Abarenkov**. PlutoF – cloud database and computing services supporting biological research. Tartu, 2011, 125 p.
207. **Marina Grigorova**. Fine-scale genetic variation of follicle-stimulating hormone beta-subunit coding gene (*FSHB*) and its association with reproductive health. Tartu, 2011, 184 p.
208. **Anu Tiitsaar**. The effects of predation risk and habitat history on butterfly communities. Tartu, 2011, 97 p.
209. **Elin Sild**. Oxidative defences in immunoeological context: validation and application of assays for nitric oxide production and oxidative burst in a wild passerine. Tartu, 2011, 105 p.
210. **Irja Saar**. The taxonomy and phylogeny of the genera *Cystoderma* and *Cystodermella* (Agaricales, Fungi). Tartu, 2012, 167 p.
211. **Pauli Saag**. Natural variation in plumage bacterial assemblages in two wild breeding passerines. Tartu, 2012, 113 p.
212. **Aleksei Lulla**. Alphaviral nonstructural protease and its polyprotein substrate: arrangements for the perfect marriage. Tartu, 2012, 143 p.
213. **Mari Järve**. Different genetic perspectives on human history in Europe and the Caucasus: the stories told by uniparental and autosomal markers. Tartu, 2012, 119 p.
214. **Ott Scheler**. The application of tmRNA as a marker molecule in bacterial diagnostics using microarray and biosensor technology. Tartu, 2012, 93 p.
215. **Anna Balikova**. Studies on the functions of tumor-associated mucin-like leukosialin (CD43) in human cancer cells. Tartu, 2012, 129 p.
216. **Triinu Kõressaar**. Improvement of PCR primer design for detection of prokaryotic species. Tartu, 2012, 83 p.
217. **Tuul Sepp**. Hematological health state indices of greenfinches: sources of individual variation and responses to immune system manipulation. Tartu, 2012, 117 p.
218. **Rya Ero**. Modifier view of the bacterial ribosome. Tartu, 2012, 146 p.
219. **Mohammad Bahram**. Biogeography of ectomycorrhizal fungi across different spatial scales. Tartu, 2012, 165 p.
220. **Annely Lorents**. Overcoming the plasma membrane barrier: uptake of amphipathic cell-penetrating peptides induces influx of calcium ions and downstream responses. Tartu, 2012, 113 p.
221. **Katrin Männik**. Exploring the genomics of cognitive impairment: whole-genome SNP genotyping experience in Estonian patients and general population. Tartu, 2012, 171 p.
222. **Marko Prous**. Taxonomy and phylogeny of the sawfly genus *Empria* (Hymenoptera, Tenthredinidae). Tartu, 2012, 192 p.

223. **Triinu Visnapuu.** Levansucrases encoded in the genome of *Pseudomonas syringae* pv. tomato DC3000: heterologous expression, biochemical characterization, mutational analysis and spectrum of polymerization products. Tartu, 2012, 160 p.
224. **Nele Tamberg.** Studies on Semliki Forest virus replication and pathogenesis. Tartu, 2012, 109 p.
225. **Tõnu Esko.** Novel applications of SNP array data in the analysis of the genetic structure of Europeans and in genetic association studies. Tartu, 2012, 149 p.
226. **Timo Arula.** Ecology of early life-history stages of herring *Clupea harengus membras* in the northeastern Baltic Sea. Tartu, 2012, 143 p.
227. **Inga Hiiesalu.** Belowground plant diversity and coexistence patterns in grassland ecosystems. Tartu, 2012, 130 p.
228. **Kadri Koorem.** The influence of abiotic and biotic factors on small-scale plant community patterns and regeneration in boreonemoral forest. Tartu, 2012, 114 p.
229. **Liis Andresen.** Regulation of virulence in plant-pathogenic pectobacteria. Tartu, 2012, 122 p.
230. **Kaupo Kohv.** The direct and indirect effects of management on boreal forest structure and field layer vegetation. Tartu, 2012, 124 p.
231. **Mart Jüssi.** Living on an edge: landlocked seals in changing climate. Tartu, 2012, 114 p.
232. **Riina Klais.** Phytoplankton trends in the Baltic Sea. Tartu, 2012, 136 p.
233. **Rauno Veeroja.** Effects of winter weather, population density and timing of reproduction on life-history traits and population dynamics of moose (*Alces alces*) in Estonia. Tartu, 2012, 92 p.
234. **Marju Keis.** Brown bear (*Ursus arctos*) phylogeography in northern Eurasia. Tartu, 2013, 142 p.
235. **Sergei Põlme.** Biogeography and ecology of *alnus*- associated ectomycorrhizal fungi – from regional to global scale. Tartu, 2013, 90 p.
236. **Liis Uusküla.** Placental gene expression in normal and complicated pregnancy. Tartu, 2013, 173 p.
237. **Marko Lõoke.** Studies on DNA replication initiation in *Saccharomyces cerevisiae*. Tartu, 2013, 112 p.
238. **Anne Aan.** Light- and nitrogen-use and biomass allocation along productivity gradients in multilayer plant communities. Tartu, 2013, 127 p.
239. **Heidi Tamm.** Comprehending phylogenetic diversity – case studies in three groups of ascomycetes. Tartu, 2013, 136 p.
240. **Liina Kangur.** High-Pressure Spectroscopy Study of Chromophore-Binding Hydrogen Bonds in Light-Harvesting Complexes of Photosynthetic Bacteria. Tartu, 2013, 150 p.
241. **Margus Leppik.** Substrate specificity of the multisite specific pseudouridine synthase RluD. Tartu, 2013, 111 p.
242. **Lauris Kaplinski.** The application of oligonucleotide hybridization model for PCR and microarray optimization. Tartu, 2013, 103 p.

243. **Merli Pärnoja**. Patterns of macrophyte distribution and productivity in coastal ecosystems: effect of abiotic and biotic forcing. Tartu, 2013, 155 p.
244. **Tõnu Margus**. Distribution and phylogeny of the bacterial translational GTPases and the Mqsr/YgiT regulatory system. Tartu, 2013, 126 p.
245. **Pille Mänd**. Light use capacity and carbon and nitrogen budget of plants: remote assessment and physiological determinants. Tartu, 2013, 128 p.
246. **Mario Plaas**. Animal model of Wolfram Syndrome in mice: behavioural, biochemical and psychopharmacological characterization. Tartu, 2013, 144 p.
247. **Georgi Hudjašov**. Maps of mitochondrial DNA, Y-chromosome and tyrosinase variation in Eurasian and Oceanian populations. Tartu, 2013, 115 p.
248. **Mari Lepik**. Plasticity to light in herbaceous plants and its importance for community structure and diversity. Tartu, 2013, 102 p.
249. **Ede Leppik**. Diversity of lichens in semi-natural habitats of Estonia. Tartu, 2013, 151 p.
250. **Ülle Saks**. Arbuscular mycorrhizal fungal diversity patterns in boreo-nemoral forest ecosystems. Tartu, 2013, 151 p.
251. **Eneli Oitmaa**. Development of arrayed primer extension microarray assays for molecular diagnostic applications. Tartu, 2013, 147 p.
252. **Jekaterina Jutkina**. The horizontal gene pool for aromatics degradation: bacterial catabolic plasmids of the Baltic Sea aquatic system. Tartu, 2013, 121 p.
253. **Helen Vellau**. Reaction norms for size and age at maturity in insects: rules and exceptions. Tartu, 2014, 132 p.
254. **Randel Kreitsberg**. Using biomarkers in assessment of environmental contamination in fish – new perspectives. Tartu, 2014, 107 p.
255. **Krista Takkis**. Changes in plant species richness and population performance in response to habitat loss and fragmentation. Tartu, 2014, 141 p.
256. **Liina Nagirnaja**. Global and fine-scale genetic determinants of recurrent pregnancy loss. Tartu, 2014, 211 p.
257. **Triin Triisberg**. Factors influencing the re-vegetation of abandoned extracted peatlands in Estonia. Tartu, 2014, 133 p.
258. **Villu Soon**. A phylogenetic revision of the *Chrysis ignita* species group (Hymenoptera: Chrysididae) with emphasis on the northern European fauna. Tartu, 2014, 211 p.
259. **Andrei Nikonov**. RNA-Dependent RNA Polymerase Activity as a Basis for the Detection of Positive-Strand RNA Viruses by Vertebrate Host Cells. Tartu, 2014, 207 p.
260. **Eele Õunapuu-Pikas**. Spatio-temporal variability of leaf hydraulic conductance in woody plants: ecophysiological consequences. Tartu, 2014, 135 p.
261. **Marju Männiste**. Physiological ecology of greenfinches: information content of feathers in relation to immune function and behavior. Tartu, 2014, 121 p.

262. **Katre Kets**. Effects of elevated concentrations of CO₂ and O₃ on leaf photosynthetic parameters in *Populus tremuloides*: diurnal, seasonal and interannual patterns. Tartu, 2014, 115 p.
263. **Küllli Lokko**. Seasonal and spatial variability of zoopsammon communities in relation to environmental parameters. Tartu, 2014, 129 p.
264. **Olga Žilina**. Chromosomal microarray analysis as diagnostic tool: Estonian experience. Tartu, 2014, 152 p.
265. **Kertu Lõhmus**. Colonisation ecology of forest-dwelling vascular plants and the conservation value of rural manor parks. Tartu, 2014, 111 p.
266. **Anu Aun**. Mitochondria as integral modulators of cellular signaling. Tartu, 2014, 167 p.
267. **Chandana Basu Mallick**. Genetics of adaptive traits and gender-specific demographic processes in South Asian populations. Tartu, 2014, 160 p.
268. **Riin Tamme**. The relationship between small-scale environmental heterogeneity and plant species diversity. Tartu, 2014, 130 p.
269. **Liina Remm**. Impacts of forest drainage on biodiversity and habitat quality: implications for sustainable management and conservation. Tartu, 2015, 126 p.
270. **Tiina Talve**. Genetic diversity and taxonomy within the genus *Rhinanthus*. Tartu, 2015, 106 p.
271. **Mehis Rohtla**. Otolith sclerochronological studies on migrations, spawning habitat preferences and age of freshwater fishes inhabiting the Baltic Sea. Tartu, 2015, 137 p.
272. **Alexey Reshchikov**. The world fauna of the genus *Lathrolestes* (Hymenoptera, Ichneumonidae). Tartu, 2015, 247 p.
273. **Martin Pook**. Studies on artificial and extracellular matrix protein-rich surfaces as regulators of cell growth and differentiation. Tartu, 2015, 142 p.
274. **Mai Kukumägi**. Factors affecting soil respiration and its components in silver birch and Norway spruce stands. Tartu, 2015, 155 p.
275. **Helen Karu**. Development of ecosystems under human activity in the North-East Estonian industrial region: forests on post-mining sites and bogs. Tartu, 2015, 152 p.
276. **Hedi Peterson**. Exploiting high-throughput data for establishing relationships between genes. Tartu, 2015, 186 p.
277. **Priit Adler**. Analysis and visualisation of large scale microarray data, Tartu, 2015, 126 p.
278. **Aigar Niglas**. Effects of environmental factors on gas exchange in deciduous trees: focus on photosynthetic water-use efficiency. Tartu, 2015, 152 p.
279. **Silja Laht**. Classification and identification of conopeptides using profile hidden Markov models and position-specific scoring matrices. Tartu, 2015, 100 p.
280. **Martin Kesler**. Biological characteristics and restoration of Atlantic salmon *Salmo salar* populations in the Rivers of Northern Estonia. Tartu, 2015, 97 p.

281. **Pratyush Kumar Das**. Biochemical perspective on alphaviral nonstructural protein 2: a tale from multiple domains to enzymatic profiling. Tartu, 2015, 205 p
282. **Priit Palta**. Computational methods for DNA copy number detection. Tartu, 2015, 130 p.
283. **Julia Sidorenko**. Combating DNA damage and maintenance of genome integrity in pseudomonads. Tartu, 2015, 174 p.
284. **Anastasiia Kovtun-Kante**. Charophytes of Estonian inland and coastal waters: distribution and environmental preferences. Tartu, 2015, 97 p.
285. **Ly Lindman**. The ecology of protected butterfly species in Estonia. Tartu, 2015, 171 p.
286. **Jaanis Lodjak**. Association of Insulin-like Growth Factor I and Corticosterone with Nestling Growth and Fledging Success in Wild Passerines. Tartu, 2016, 113 p.
287. **Ann Kraut**. Conservation of Wood-Inhabiting Biodiversity – Semi-Natural Forests as an Opportunity. Tartu, 2016, 141 p.
288. **Tiit Örd**. Functions and regulation of the mammalian pseudokinase TRIB3. Tartu, 2016, 182. p.