

---

## Effects of dose Timing on Fluid Excretion During Sodium-Aided Hyperhydration Protocols

SHELBY GREENE, ELIZAVETA ROSLANOVA, and DAVID M. MORIS

Kinesiology Lab; Department of Kinesiology; University of Texas – Permian Basin;  
Odessa, TX

---

*Category: Undergraduate*

*Advisor / Mentor: Morris, David (morris\_da@utpb.edu)*

### ABSTRACT

Co-consumption of sodium and water has been shown to be superior in promoting hyperhydration compared to consumption of an equal amount of water alone. Most sodium-aided hyperhydration studies have provided subjects with a bolus of fluid followed by a urine collection period. However the effect of providing equal amounts of fluid in a single vs. multiple doses over time on fluid retention has not been systematically studied. **PURPOSE:** To compare the effects of different dosing strategies on urine excretion levels following the consumption of consistent amounts of sodium and water. **METHODS:** Urine excretion was measured during five separate 2-hr hyperhydration protocols in 13 well hydrated male subjects ( $23 \pm 3$  yr,  $176.1 \pm 10.1$  cm,  $82.2 \pm 19.4$  kg) who were free from known renal, digestive, and cardiovascular disease. Each protocol began with a complete bladder void and assessment of urine specific gravity (USG). Subjects then consumed  $20 \text{ mL H}_2\text{O} \cdot \text{kg bm}^{-1}$  and  $110 \text{ mg NaCl} \cdot \text{kg bm}^{-1}$  in five different dosing strategies: the entire dose was consumed at the beginning of the period (1X),  $\frac{1}{2}$  of the dose was consumed at the beginning and  $\frac{1}{2}$  consumed after 60 min (2X), and  $\frac{1}{3}$  of the dose was consumed at the beginning and  $\frac{1}{3}$  was consumed after 45 and 90 min (3X),  $\frac{1}{4}$  of the dose was consumed at the beginning and after 30, 60, and 90 min (4X), and  $\frac{1}{7}$  of the dose was consumed at the beginning and after 15, 30, 45, 60, 75, 90 min (7X). Protocols were administered in a randomized, crossover fashion. Total urine excretions (TUE) during the 2 hr collection periods were expressed as a percent of the  $\text{H}_2\text{O}$  consumed. USG and TUE were compared using repeated-measures ANOVA and Sidak *post hoc* analyses. **RESULTS:** USGs were  $1.006 \pm 0.004$  (1X),  $1.007 \pm 0.003$  (2X),  $1.009 \pm 0.005$  (3X),  $1.007 \pm 0.004$  (4X), and  $1.007 \pm 0.005$  (7X) ( $P = 0.37 - 1.00$ ) indicating that subjects were well and similarly hydrated for each trial. TUE expressed as a percentage of  $\text{H}_2\text{O}$  consumed were  $75 \pm 18\%$  (1X),  $69 \pm 11\%$  (2X),  $52\% \pm 15\%$  (3X),  $59 \pm 15\%$  (4X), and  $60 \pm 16\%$  (7X). Significant differences in TUE were seen between 1X and 3X ( $P = 0.03$ ) and 2X and 3X ( $P = 0.006$ ). No significant difference in TUE was detected between any of the other protocols ( $P = 0.16 - 1.00$ ). **CONCLUSION:** The data suggest that hyperhydration is better achieved when water and sodium are consumed in three equal doses over 90 min when compared to consuming an equal amount of a sodium and water dose in a single bolus or in two equal doses over a 60 min period. Consuming water in four or seven equal doses over 90 min did not result in better fluid retention than consuming an equal amount of water in a single bolus or in two equal doses over a 60 min period.