



Crystal structures of a copper(II) and the isotopic nickel(II) and palladium(II) complexes of the ligand (*E*)-1-[(2,4,6-tribromophenyl)diazenyl]naphthalen-2-ol

Souheyla Chetioui,^{a*} Djamil-Azzeddine Rouag,^a Jean-Pierre Djukic,^b Christian G. Bochet,^c Rachid Touzani,^{d,e*} Corinne Bailly,^f Aurélien Crochet^g and Katharina M. Fromm^c

Received 7 May 2016

Accepted 4 July 2016

Edited by H. Stoeckli-Evans, University of Neuchâtel, Switzerland


Keywords: crystal structures; copper(II); nickel(II); palladium(II); isotopic complexes; Cu···Br short contact; C—H···Br hydrogen bonds; C—H···π interactions.

CCDC references: 1490056; 1490055; 1490054

Supporting information: this article has supporting information at journals.iucr.org/e

^aUnité de Recherche de Chimie de l'Environnement et Moléculaire Structurale (CHEMS), Faculté des Sciences Exactes, Département de Chimie, Université des Frères Mentouri Constantine, Constantine 25000, Algeria, ^bLaboratoire de Chimie et Systémique Organométallique (LCSOM), Institut de Chimie, Université de Strasbourg, UMR 7177., 4 rue Blaise Pascal, F-67070 Strasbourg Cedex, France, ^cChemistry Department, University of Fribourg, Chemin du Musée 9, CH-1700 Fribourg, Switzerland, ^dLaboratoire de Chimie Appliquée et Environnement, LCAE-URAC18, COSTE, Faculté des Sciences, Université Mohamed Premier, BP524, 60000 Oujda, Morocco, ^eFaculté Pluridisciplinaire Nador BP 300, Selouane 62702, Nador, Morocco, ^fService de Radiocristallographie, Institut de Chimie, Université de Strasbourg, UMR 7177, 67008 Strasbourg Cedex, France, and ^gFribourg Center for Nanomaterials, FriMat, University of Fribourg, Chemin du Musée 9, CH-1700 Fribourg, Switzerland. *Correspondence e-mail: souheilachetioui@yahoo.fr, touzani@yahoo.fr


In the copper(II) complex, bis{(*E*)-1-[(2,4,6-tribromophenyl)diazenyl]naphthalen-2-olato}copper(II), [Cu(C₁₆H₈Br₃N₂O)₂], (I), the metal cation is coordinated by two N atoms and two O atoms from two bidentate (*E*)-1-[(2,4,6-tribromophenyl)diazenyl]naphthalen-2-olate ligands, forming a slightly distorted square-planar environment. In one of the ligands, the tribromobenzene ring is inclined to the naphthalene ring system by 37.4 (5)°, creating a weak intramolecular Cu···Br interaction [3.134 (2) Å], while in the other ligand, the tribromobenzene ring is inclined to the naphthalene ring system by 72.1 (6)°. In the isotopic nickel(II) and palladium(II) complexes, namely bis{(*E*)-1-[(2,4,6-tribromophenyl)diazenyl]naphthalen-2-olato}nickel(II), [Ni(C₁₆H₈Br₃N₂O)₂], (II), and bis{(*E*)-1-[(2,4,6-tribromophenyl)diazenyl]naphthalen-2-olato}palladium(II), [Pd(C₁₆H₈Br₃N₂O)₂], (III), respectively, the metal atoms are located on centres of inversion, hence the metal coordination spheres have perfect square-planar geometries. The tribromobenzene rings are inclined to the naphthalene ring systems by 80.79 (18)° in (II) and by 80.8 (3)° in (III). In the crystal of (I), molecules are linked by C—H···Br hydrogen bonds, forming chains along [010]. The chains are linked by C—H···π interactions, forming sheets parallel to (011). In the crystals of (II) and (III), molecules are linked by C—H···π interactions, forming slabs parallel to (10 $\bar{1}$). For the copper(II) complex (I), a region of disordered electron density was corrected for using the SQUEEZE routine in *PLATON* [Spek (2015)]. *Acta Cryst. C* **71**, 9–18]. The formula mass and unit-cell characteristics of the disordered solvent molecules were not taken into account during refinement.



1. Chemical context

Recently, 1-phenylazo-2-naphthol derivatives have attracted attention because the phenylazo-naphtholate group can provide *N,O*-bidentate chelation to stabilize transition or main group metal complexes. Azo-metal chelates have also attracted increasing attention due to their interesting electronic and geometrical features in connection with their applications in molecular memory storage, non-linear optical elements and printing systems. Another advantage of

complexes involving azo DNO's (dyes and pigments) and transition metal ions is the possibility to obtain new compounds with biological activity (Thomas *et al.*, 2004; Reed *et al.*, 2006). Transition metals have also been used in the treatment of several diseases, as metal complexes which are capable of cleaving DNA under physiological conditions are of interest in the development of metal-based anticancer agents. This is an impetus for chemists to develop innovative strategies for the preparation of more effective, target-specific and preferably non-covalently bound anticancer drugs (Chen *et al.*, 2010; Cvek *et al.*, 2008).



Being interested in the synthesis and preparation of metal complexes bearing such ligands, we have successfully synthesized and structurally characterized Cu^{II} complexes with *N,O*-bidentate phenylazo-naphtholate ligands (Chetioui *et al.*, 2015*a,b*). In this work we are involved in the colour-generation mechanism of azo pigments typically characterized by the chromophore of the azo group ($-\text{N}=\text{N}-$) (Chetioui *et al.*, 2013*c,d*) in order to synthesize new complexes with $\text{Cu}(\text{OAc})_2 \cdot \text{H}_2\text{O}$, $\text{Ni}(\text{OAc})_2 \cdot \text{H}_2\text{O}$, and $\text{Pd}(\text{OAc})_2 \cdot \text{H}_2\text{O}$. We report herein on the synthesis and crystal structures of the title complexes, (I)–(III), of the ligand (*E*)-1-[(2,4,6-tribromophenyl)diazenyl]naphthalen-2-ol, whose crystal structure has been described previously (Chetioui *et al.*, 2013*c*).

2. Structural commentary

In all three compounds the ligand (*E*)-1-[(2,4,6-tribromophenyl)diazenyl]naphthalen-2-ol (Chetioui *et al.*, 2013*c*) coordinates in a *N,O*-bidentate manner. The metal atoms are coordinated by two oxygen atoms in a *trans* position of the

Table 1
Selected geometric parameters (\AA , $^\circ$) for (I).

Cu1—Br4	3.134 (2)	Cu1—O1	1.892 (9)
Cu1—N1	1.947 (12)	Cu1—O2	1.888 (8)
Cu1—N3	1.970 (11)		
O2—Cu1—O1	169.4 (4)	O2—Cu1—N3	87.6 (4)
O2—Cu1—N1	91.3 (4)	O1—Cu1—N3	92.1 (4)
O1—Cu1—N1	90.9 (4)	N1—Cu1—N3	169.3 (5)

$\text{C}-\text{O}^-$ function and two nitrogen atoms in a *trans* position of the $\text{N}=\text{N}$ function. In compound (I), Fig. 1, the values of the angles involving the copper and the two oxygen and two nitrogen atoms (Table 1) indicate that the geometry of the coordination polyhedron is distorted square-planar. It has a τ_4 value of 0.15 [Yang *et al.*, 2007; extreme configurations: 0.00 for square-pyramidal (SQP) and 1.00 for tetrahedral (TET); 0.85 for trigonal-pyramidal (TRP)]. In one of the ligands, the tribromobenzene ring (C17–C22) is inclined to the naphthalene ring system (C23–C32) by $37.4 (5)^\circ$, creating a weak intramolecular $\text{Cu} \cdots \text{Br}$ interaction [$\text{Cu1} \cdots \text{Br4} = 3.134 (2) \text{\AA}$]. In the other ligand, the tribromobenzene ring (C1–C6) is almost normal to the naphthalene ring system (C7–C16), making a dihedral angle of $72.1 (6)^\circ$. A similar short intramolecular metal–halogen contact has been observed in the centrosymmetric complex bis(1-[(*E*)-(2-chlorophenyl)diazenyl]naphthalen-2-olato)copper(II), *viz.* $\text{Cu} \cdots \text{Cl} = 3.153 (1) \text{\AA}$ (Benaouida *et al.*, 2013), and the chlorobenzene ring is inclined to the naphthalene ring system by $32.72 (12)^\circ$.




Figure 1
The molecular structure of compound (I), with atom labelling and 50% probability displacement ellipsoids. The intramolecular $\text{Cu} \cdots \text{Br}$ contact is shown as a dashed line (details are given in Table 1).




Figure 2
The molecular structure of compound (II), with atom labelling and 50% probability displacement ellipsoids. The unlabelled atoms are related to the labelled atoms by the symmetry operation $(-x + 1, -y + 1, -z + 1)$.

Compounds (II) and (III), the nickel(II) (Fig. 2, Table 2) and palladium(II) (Fig. 3, Table 3) complexes, respectively, are isotopic. The metal atoms are each located on inversion centres, coordinating in a bidentate fashion to the N and O




Figure 3
The molecular structure of compound (III), with atom labelling and 50% probability displacement ellipsoids. The unlabelled atoms are related to the labelled atoms by the symmetry operation $(-x, -y, -z)$.

Table 2
Selected geometric parameters (\AA , $^\circ$) for (II).

Ni—N1	1.876 (3)	Ni—O1	1.821 (3)
O1—Ni—O1 ⁱ	180	O1 ⁱ —Ni—N1	87.41 (14)
O1—Ni—N1	92.59 (14)	N1—Ni—N1 ⁱ	180

Symmetry code: (i) $-x + 1, -y + 1, -z + 1$.

Table 3
Selected geometric parameters (\AA , $^\circ$) for (III).

Pd1—N1	2.004 (5)	Pd1—O1	1.972 (5)
O1—Pd1—O1 ⁱ	180	O1 ⁱ —Pd1—N1	88.7 (2)
O1—Pd1—N1	91.3 (2)	N1—Pd1—N1 ⁱ	180

Symmetry code: (i) $-x, -y, -z$.

atoms of the ligand, hence the metal coordination spheres have perfect square-planar geometry. The tribromobenzene rings (C1–C6) are almost normal to the naphthalene ring systems (C7–C16) with a dihedral angle of $80.79 (18)^\circ$ in (II) and $80.8 (3)^\circ$ in (III).




Figure 4
The crystal packing of compound (I), viewed along the a axis. The intermolecular interactions are shown as dashed lines (see Table 4 for details), and for clarity only the H atoms involved in these interactions have been included.

Table 4
Hydrogen-bond geometry (Å, °) for (I).

Cg1 is the centroid of the C27–C32 ring.

<i>D</i> –H... <i>A</i>	<i>D</i> –H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> –H... <i>A</i>
C5–H5...Br6 ⁱ	0.95	2.75	3.546 (15)	142
C3–H3...Cg1 ⁱⁱ	0.95	2.99	3.729 (15)	136

Symmetry codes: (i) $x, y, z - 1$; (ii) $-x + 1, y + \frac{1}{2}, -z$.

3. Supramolecular features

As shown in Fig. 4, in the crystal of compound (I), molecules are linked by C–H...Br hydrogen bonds, forming chains along [001]. The chains are linked by C–H... π interactions, forming sheets lying parallel to (011). Details of these interactions are given in Table 4.

The crystal packing in compound (II) [and isotopic compound (III)] is illustrated in Fig. 5. Molecules are linked by C–H... π interactions, forming slabs lying parallel to (10 $\bar{1}$). Details of the intermolecular interactions are given in Table 5 for (II) and Table 6 for (III).

Table 5
Hydrogen-bond geometry (Å, °) for (II).

Cg2 is the centroid of the C1–C6 ring.

<i>D</i> –H... <i>A</i>	<i>D</i> –H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> –H... <i>A</i>
C10–H10...Cg2 ⁱⁱ	0.95	2.71	3.391 (5)	130

Symmetry code: (ii) $x + \frac{1}{2}, -y + \frac{3}{2}, z + \frac{1}{2}$.

Table 6
Hydrogen-bond geometry (Å, °) for (III).

Cg2 is the centroid of the C1–C6 ring.

<i>D</i> –H... <i>A</i>	<i>D</i> –H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> –H... <i>A</i>
C10–H10...Cg2 ⁱⁱ	0.93	2.70	3.371 (8)	129

Symmetry code: (ii) $x + \frac{1}{2}, -y - \frac{1}{2}, z + \frac{1}{2}$.

4. Database survey

In the title ligand (*E*)-1-[(2,4,6-tribromophenyl)diazenyl]naphthalen-2-ol (CSD refcode AFOFIM; Chetioui *et al.*,





Figure 5
The crystal packing of compound (II), viewed along the normal to (10 $\bar{1}$). The intermolecular interactions are shown as dashed lines (see Table 5 for details), and for clarity only the H atoms involved in these interactions have been included.



LUQQIZ: $M = \text{Zn}, R^1 = R^2 = R^3 = \text{H}$

CBANAP: $M = \text{Cu}, R^1 = R^2 = R^3 = \text{H}$

NOTNUH: $M = \text{Ni}, R^1 = R^2 = R^3 = \text{H}$

AFATIM: $M = \text{Cu}, R^1 = \text{Cl}, R^2 = R^3 = \text{H}$

DURRIS: $M = \text{Pd}, R^1 = \text{CH}_3, R^2 = R^3 = \text{H}$

TOAZNI: $M = \text{Ni}, R^1 = \text{H}, R^2 = \text{CH}_3, R^3 = \text{H}$

NOTNOB: $M = \text{Cu}, R^1 = \text{CH}_3, R^2 = \text{H}, R^3 = \text{CH}_3$

Figure 6

The results of the database search (CSD; Groom *et al.*, 2016) for four-coordinate metal complexes of the ligand (*E*)-1-(phenyldiazenyl)naphthalen-2-ol and its derivatives.

2013c) the benzene ring is inclined to the naphthalene ring system by 33.80 (16)°. A search of the Cambridge Structural Database (Version 5.37, update February 2016; Groom *et al.*, 2016) for square-planar metal complexes of (*E*)-1-(phenyldiazenyl)naphthalen-2-ol and its derivatives gave seven hits (Fig. 6). They include a zinc(II) complex of the ligand (*E*)-1-(phenyldiazenyl)naphthalen-2-ol (LUQQIZ; Gallegos *et al.*, 2015), where the zinc atom has a distorted trigonal-pyramidal configuration with a τ_4 parameter of 0.77. In the two ligands, the phenyl rings are inclined to the naphthalene ring systems by 11.4 (2) and 9.2 (3)°. Among the other six complexes, in which the metal atoms are all located on inversion centres, there are three copper(II) complexes with the ligands (*E*)-1-(phenyldiazenyl)naphthalen-2-ol (refcode CBANAP; Jarvis, 1961), (*E*)-1-(2-chlorophenyl)diazenyl]naphthalen-2-ol (AFATIM; Benaouida *et al.*, 2013) and (*E*)-1-(2,4-dimethylphenyl)diazenyl]naphthalen-2-ol (NOTNOB; Ferreira *et al.*, 2015); two nickel complexes with the ligands (*E*)-1-(phenyldiazenyl)naphthalen-2-ol (NOTNUH; Ferreira *et al.*, 2015) and (*E*)-1-(3-methylphenyl)diazenyl]naphthalen-2-ol (TOAZNI; Alcock *et al.*, 1968); and one palladium complex

with the ligand [(*E*)-1-(2-methylphenyl)diazenyl]naphthalen-2-ol (DURRIS; Lin *et al.*, 2010). The orientation of the phenyl/benzene ring with respect to the naphthalene ring system varies quite considerably. In the palladium complex (DURRIS) and the copper complex (NOTNOB), where the benzene ring has a methyl group in the *ortho* position, the benzene ring is inclined to the naphthalene ring system by 74.41 (4) and 83.87 (6)°, respectively. In the other four complexes, the corresponding dihedral angles are 19.12 and 32.72 (12)° for the copper complexes CBANAP and AFATIM, respectively, and 24.06 (15) and *ca* 35.56 ° for the nickel complexes NOTNUH and TOAZNI, respectively.

5. Synthesis and crystallization

The title compounds were synthesized by the following procedure: (*E*)-1-[(2,4,6-tribromophenyl)diazenyl]naphthalen-2-ol (2.0 mmol) and $M(\text{OAc})_2 \cdot \text{H}_2\text{O}$ (1.0 mmol; where $M = \text{Cu}, \text{Ni}, \text{Pd}$) was stirred at 298 K in a mixture of THF/MeOH (10/10 ml) for 24 h. The solvents were removed under vacuum and the residue was washed twice with hexane to give dark solids. The resulting solids were crystallized from CH_2Cl_2 to yield red block-like crystals for (I), black prismatic crystals for (II) and dark-red plate-like crystals for (III).

6. Refinement details

Crystal data, data collection and structure refinement details are summarized in Table 7. For all three compounds the C-bound H atoms were included in calculated positions and refined as riding: C–H = 0.95 Å for (I) and (II) and 0.93 Å for (III), with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$. For the copper(II) complex (I), a region of disordered electron density was corrected for using the SQUEEZE routine in PLATON (Spek, 2015). The formula mass and unit-cell characteristics of the disordered solvent molecules were not taken into account during refinement. This complex crystallizes in the monoclinic space group $P2_1$, with the Flack parameter = -0.006 (14).

Acknowledgements

The authors are grateful to the Algerian Ministry of Higher Education and Scientific Research, the Algerian Directorate General for Scientific Research and Technological Development, and the University of Constantine for financial support.

References

- Alcock, N. W., Spencer, R. C., Prince, R. H. & Kennard, O. (1968). *J. Chem. Soc. A*, pp. 2382–2388.
- Altomare, A., Burla, M. C., Camalli, M., Cascarano, G. L., Giacovazzo, C., Guagliardi, A., Moliterni, A. G. G., Polidori, G. & Spagna, R. (1999). *J. Appl. Cryst.* **32**, 115–119.
- Benaouida, M. A., Benosmane, A., Bouguerria, H., Bouaoud, S. E. & Merazig, H. (2013). *Acta Cryst.* **E69**, m405.
- Chen, G. J., Qiao, X., Qiao, P. Q., Xu, G. J., Xu, J. Y., Tian, J. L., Gu, W., Liu, X. & Yan, S. P. (2010). *J. Inorg. Biochem.* **105**, 119–126.
- Chetoui, S., Boudraa, I., Bouacida, S., Bouchoul, A. & Bouaoud, S. E. (2013c). *Acta Cryst.* **E69**, o1250.

Table 7
Experimental details.

	(I)	(II)	(III)
Crystal data			
Chemical formula	[Cu(C ₁₆ H ₈ Br ₃ N ₂ O) ₂]	[Ni(C ₁₆ H ₈ Br ₃ N ₂ O) ₂]	[Pd(C ₁₆ H ₈ Br ₃ N ₂ O) ₂]
<i>M_r</i>	1031.49	1026.66	1074.35
Crystal system, space group	Monoclinic, <i>P</i> 2 ₁	Monoclinic, <i>P</i> 2 ₁ / <i>n</i>	Monoclinic, <i>P</i> 2 ₁ / <i>n</i>
Temperature (K)	173	173	200
<i>a</i> , <i>b</i> , <i>c</i> (Å)	11.9423 (7), 12.1314 (10), 12.8974 (10)	11.0909 (6), 12.4571 (6), 12.5382 (7)	11.1896 (8), 12.4540 (8), 12.5511 (9)
β (°)	107.032 (4)	107.820 (2)	107.749 (5)
<i>V</i> (Å ³)	1786.6 (2)	1649.17 (15)	1665.8 (2)
<i>Z</i>	2	2	2
Radiation type	Mo <i>K</i> α	Mo <i>K</i> α	Cu <i>K</i> α
μ (mm ⁻¹)	7.36	7.89	13.23
Crystal size (mm)	0.20 × 0.15 × 0.06	0.30 × 0.22 × 0.06	0.12 × 0.09 × 0.03
Data collection			
Diffractometer	Nonius KappaCCD	Nonius KappaCCD	STOE <i>IPDS</i> 2T
Absorption correction	Multi-scan (<i>MULABS</i> ; Spek, 2009)	Multi-scan (<i>MULABS</i> ; Spek, 2009)	Multi-scan (<i>MULABS</i> ; Spek, 2009)
<i>T</i> _{min} , <i>T</i> _{max}	0.311, 0.386	0.151, 0.317	0.360, 1.000
No. of measured, independent and observed [<i>I</i> > 2σ(<i>I</i>)] reflections	14985, 7819, 4785	11360, 3745, 2214	13003, 2895, 2371
<i>R</i> _{int}	0.077	0.094	0.142
(sin θ/λ) _{max} (Å ⁻¹)	0.650	0.649	0.600
Refinement			
<i>R</i> [<i>F</i> ² > 2σ(<i>F</i> ²)], <i>wR</i> (<i>F</i> ²), <i>S</i>	0.064, 0.140, 0.96	0.043, 0.096, 0.95	0.057, 0.170, 1.11
No. of reflections	7819	3745	2895
No. of parameters	388	205	206
No. of restraints	2	0	0
H-atom treatment	H-atom parameters constrained	H-atom parameters constrained	H-atom parameters constrained
Δρ _{max} , Δρ _{min} (e Å ⁻³)	0.59, -0.58	0.57, -0.66	0.88, -1.10
Absolute structure	Flack <i>x</i> determined using 1648 quotients [(<i>I</i> ⁺)-(<i>I</i>)]/[(<i>I</i> ⁺)+(<i>I</i>)] (Parsons <i>et al.</i> , 2013)	-	-
Absolute structure parameter	-0.006 (14)	-	-

Computer programs: *COLLECT* (Nonius, 1998), *X-AREA* and *X-RED32* (Stoe & Cie, 2002), *DENZO* (Otwinowski & Minor, 1997), *SIR97* (Altomare *et al.*, 1999), *SHELXS2014* (Sheldrick, 2008), *SHELXL2014* (Sheldrick, 2015), *Mercury* (Macrae *et al.*, 2008) and *PLATON* (Spek, 2009).

Chetioui, S., Boudraa, I., Bouacida, S., Bouchoul, A. & Bouaoud, S. E. (2013*d*). *Acta Cryst.* **E69**, o1322–o1323.
 Chetioui, S., Hamdouni, N., Bochet, C. G., Djukic, J.-P. & Bailly, C. (2015*a*). *Acta Cryst.* **E71**, m211–m212.
 Chetioui, S., Hamdouni, N., Rouag, D.-A., Bouaoud, S. E. & Merazig, H. (2015*b*). *Acta Cryst.* **E71**, m207–m208.
 Cvek, B., Milacic, V., Taraba, J. & Dou, Q. P. (2008). *J. Med. Chem.* **51**, 6256–6258.
 Ferreira, G. R., Marcial, B. L., Garcia, H. C., Faulstich, F. R. L., Dos Santos, H. F. & de Oliveira, L. F. C. (2015). *Supramol. Chem.* **27**, 13–20.
 Gallegos, C., Tabernero, V., Mosquera, M. E. G., Cuenca, T. & Cano, J. (2015). *Eur. J. Inorg. Chem.* pp. 5124–5132.
 Groom, C. R., Bruno, I. J., Lightfoot, M. P. & Ward, S. C. (2016). *Acta Cryst.* **B72**, 171–179.
 Jarvis, J. A. J. (1961). *Acta Cryst.* **14**, 961–964.
 Lin, M.-L., Tsai, C.-Y., Li, C.-Y., Huang, B.-H. & Ko, B.-T. (2010). *Acta Cryst.* **E66**, m1022.
 Macrae, C. F., Bruno, I. J., Chisholm, J. A., Edgington, P. R., McCabe, P., Pidcock, E., Rodriguez-Monge, L., Taylor, R., van de Streek, J. & Wood, P. A. (2008). *J. Appl. Cryst.* **41**, 466–470.

Nonius (1998). *COLLECT*. Nonius BV, Delft, The Netherlands.
 Otwinowski, Z. & Minor, W. (1997). *Methods in Enzymology*, Vol. 276, *Macromolecular Crystallography*, Part A, edited by C. W. Carter Jr & R. M. Sweet, pp. 307–326. New York: Academic Press.
 Parsons, S., Flack, H. D. & Wagner, T. (2013). *Acta Cryst.* **B69**, 249–259.
 Reed, J. E., Arnal, A. A., Neidle, S. & Vilar, R. (2006). *J. Am. Chem. Soc.* **128**, 5992–5993.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Sheldrick, G. M. (2015). *Acta Cryst.* **C71**, 3–8.
 Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.
 Spek, A. L. (2015). *Acta Cryst.* **C71**, 9–18.
 Stoe & Cie (2002). *X-AREA*, *X-RED32* and *X-SHAPE*. Stoe & Cie GmbH, Darmstadt, Germany.
 Thomas, A. M., Nethaji, M. & Chakravarty, A. R. J. (2004). *J. Inorg. Biochem.* **98**, 1087–1094.
 Yang, L., Powell, D. R. & Houser, R. P. (2007). *Dalton Trans.* pp. 955–964.

supporting information

Acta Cryst. (2016). E72, 1093-1098 [doi:10.1107/S205698901601080X]

Crystal structures of a copper(II) and the isotopic nickel(II) and palladium(II) complexes of the ligand (*E*)-1-[(2,4,6-tribromophenyl)diazenyl]naphthalen-2-ol

Souheyla Chetioui, Djamil-Azzeddine Rouag, Jean-Pierre Djukic, Christian G. Bochet, Rachid Touzani, Corinne Bailly, Aurélien Crochet and Katharina M. Fromm

Computing details

Data collection: *COLLECT* (Nonius, 1998) for (I), (II); *X-AREA* (Stoe & Cie, 2002) for (III). Cell refinement: *DENZO* (Otwinowski & Minor, 1997) for (I), (II); *X-AREA* (Stoe & Cie, 2002) for (III). Data reduction: *DENZO* (Otwinowski & Minor, 1997) for (I), (II); *X-RED32* (Stoe & Cie, 2002) for (III). Program(s) used to solve structure: *SHELXS2014* (Sheldrick, 2008) for (I), (II); *SIR97* (Altomare *et al.*, 1999) for (III). For all compounds, program(s) used to refine structure: *SHELXL2014* (Sheldrick, 2015); molecular graphics: *Mercury* (Macrae *et al.*, 2008); software used to prepare material for publication: *SHELXL2014* (Sheldrick, 2015) and *PLATON* (Spek, 2009).

(I) Bis{(E)-1-[(2,4,6-tribromophenyl)diazenyl]naphthalen-2-olato}copper(II)

Crystal data

[Cu(C₁₆H₈Br₃N₂O)₂]
M_r = 1031.49
 Monoclinic, *P2*₁
a = 11.9423 (7) Å
b = 12.1314 (10) Å
c = 12.8974 (10) Å
 β = 107.032 (4)°
V = 1786.6 (2) Å³
Z = 2

F(000) = 982
D_x = 1.917 Mg m⁻³
 Mo *K*α radiation, λ = 0.71073 Å
 Cell parameters from 19031 reflections
 θ = 1.0–27.5°
 μ = 7.36 mm⁻¹
T = 173 K
 Block, red
 0.20 × 0.15 × 0.06 mm

Data collection

Nonius KappaCCD
 diffractometer
 Radiation source: sealed tube
 Graphite monochromator
 phi and ω scans
 Absorption correction: multi-scan
 (MULABS; Spek, 2009)
T_{min} = 0.311, *T_{max}* = 0.386

14985 measured reflections
 7819 independent reflections
 4785 reflections with *I* > 2σ(*I*)
R_{int} = 0.077
 θ_{\max} = 27.5°, θ_{\min} = 2.4°
h = -15→15
k = -15→15
l = -16→16

Refinement

Refinement on *F*²
 Least-squares matrix: full
R[*F*² > 2σ(*F*²)] = 0.064
wR(*F*²) = 0.140
S = 0.96

7819 reflections
 388 parameters
 2 restraints
 Primary atom site location: structure-invariant
 direct methods

Secondary atom site location: difference Fourier map
 Hydrogen site location: inferred from neighbouring sites
 H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0639P)^2]$
 where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} < 0.001$
 $\Delta\rho_{\max} = 0.59 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.58 \text{ e } \text{\AA}^{-3}$
 Absolute structure: Flack x determined using 1648 quotients $[(I^+)-(I^-)]/[(I^+)+(I^-)]$ (Parsons *et al.*, 2013)
 Absolute structure parameter: -0.006 (14)

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.8599 (11)	1.0263 (11)	-0.0117 (12)	0.032 (3)
C2	0.7637 (10)	1.0951 (10)	-0.0438 (11)	0.029 (3)
C3	0.6867 (12)	1.0894 (11)	-0.1484 (12)	0.034 (3)
H3	0.6209	1.1370	-0.1709	0.041*
C4	0.7093 (12)	1.0134 (11)	-0.2171 (11)	0.036 (3)
C5	0.8009 (12)	0.9443 (13)	-0.1921 (12)	0.041 (4)
H5	0.8118	0.8927	-0.2438	0.049*
C6	0.8812 (11)	0.9500 (11)	-0.0865 (13)	0.038 (4)
C7	1.1218 (11)	1.0834 (10)	0.2004 (11)	0.037 (4)
C8	1.1126 (12)	1.0508 (12)	0.3013 (11)	0.038 (4)
C9	1.2090 (15)	1.0731 (12)	0.3969 (14)	0.050 (4)
H9	1.2028	1.0532	0.4662	0.060*
C10	1.3075 (13)	1.1215 (13)	0.3890 (14)	0.050 (3)
H10	1.3702	1.1346	0.4529	0.060*
C11	1.3195 (13)	1.1534 (13)	0.2865 (15)	0.050 (3)
C12	1.2277 (12)	1.1344 (11)	0.1930 (14)	0.042 (4)
C13	1.2435 (13)	1.1699 (12)	0.0916 (15)	0.051 (4)
H13	1.1810	1.1617	0.0267	0.061*
C14	1.3475 (13)	1.2156 (15)	0.0870 (17)	0.066 (5)
H14	1.3548	1.2383	0.0188	0.079*
C15	1.4404 (14)	1.2291 (16)	0.1775 (16)	0.060 (5)
H15	1.5122	1.2576	0.1713	0.072*
C16	1.4305 (13)	1.2026 (16)	0.274 (2)	0.075 (6)
H16	1.4947	1.2150	0.3369	0.090*
C17	0.9418 (11)	0.7896 (11)	0.3903 (12)	0.034 (3)
C18	1.0263 (12)	0.7317 (12)	0.3582 (12)	0.039 (4)
C19	1.1197 (13)	0.6785 (11)	0.4318 (13)	0.042 (4)
H19	1.1765	0.6396	0.4077	0.050*
C20	1.1273 (11)	0.6835 (12)	0.5348 (13)	0.039 (4)
C21	1.0499 (11)	0.7419 (11)	0.5769 (12)	0.038 (3)
H21	1.0600	0.7456	0.6526	0.046*
C22	0.9586 (11)	0.7938 (11)	0.5032 (12)	0.033 (3)

C23	0.6472 (5)	0.8679 (7)	0.2481 (6)	0.032 (3)
C24	0.6455 (5)	0.9058 (8)	0.1459 (7)	0.033 (3)
C25	0.5394 (7)	0.9308 (8)	0.0697 (6)	0.040 (4)
H25	0.5383	0.9567	-0.0001	0.047*
C26	0.4351 (5)	0.9179 (8)	0.0956 (6)	0.044 (4)
H26	0.3627	0.9350	0.0435	0.053*
C27	0.4369 (5)	0.8800 (8)	0.1978 (7)	0.035 (3)
C28	0.5429 (7)	0.8550 (7)	0.2740 (6)	0.035 (3)
C29	0.5396 (11)	0.8189 (12)	0.3776 (13)	0.042 (4)
H29	0.6110	0.7992	0.4299	0.050*
C30	0.4378 (14)	0.8113 (13)	0.4055 (14)	0.053 (4)
H30	0.4391	0.7902	0.4767	0.063*
C31	0.3284 (12)	0.8359 (12)	0.3248 (14)	0.048 (4)
H31	0.2568	0.8300	0.3424	0.057*
C32	0.3279 (12)	0.8675 (12)	0.2240 (12)	0.041 (4)
H32	0.2557	0.8815	0.1703	0.049*
N1	0.9383 (9)	1.0290 (9)	0.0989 (10)	0.035 (3)
N2	1.0368 (9)	1.0745 (9)	0.1038 (9)	0.032 (3)
N3	0.8497 (9)	0.8449 (9)	0.3100 (9)	0.031 (3)
N4	0.7460 (9)	0.8232 (8)	0.3214 (9)	0.031 (3)
O1	1.0249 (7)	0.9986 (8)	0.3181 (8)	0.040 (2)
O2	0.7408 (7)	0.9216 (8)	0.1119 (7)	0.038 (2)
Cu1	0.88988 (14)	0.94978 (13)	0.20990 (14)	0.0328 (4)
Br1	0.73408 (13)	1.19596 (12)	0.05505 (13)	0.0456 (4)
Br2	0.60688 (14)	1.00205 (14)	-0.36393 (13)	0.0535 (5)
Br3	1.01318 (13)	0.85864 (13)	-0.05111 (13)	0.0473 (4)
Br4	1.01312 (13)	0.72221 (13)	0.20691 (13)	0.0484 (4)
Br5	1.25913 (13)	0.61791 (13)	0.64036 (14)	0.0482 (4)
Br6	0.85805 (13)	0.88104 (13)	0.55982 (13)	0.0506 (4)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.024 (7)	0.031 (7)	0.042 (9)	-0.010 (6)	0.014 (6)	-0.003 (7)
C2	0.023 (6)	0.020 (7)	0.045 (9)	-0.008 (6)	0.012 (6)	-0.003 (6)
C3	0.037 (8)	0.033 (8)	0.032 (9)	0.002 (6)	0.010 (7)	0.005 (7)
C4	0.041 (8)	0.031 (8)	0.023 (8)	-0.005 (7)	-0.008 (6)	0.008 (7)
C5	0.043 (8)	0.046 (9)	0.035 (9)	-0.006 (7)	0.014 (7)	-0.007 (7)
C6	0.033 (7)	0.027 (7)	0.057 (11)	0.006 (6)	0.019 (7)	0.000 (7)
C7	0.033 (8)	0.027 (8)	0.050 (10)	0.009 (6)	0.009 (7)	-0.002 (7)
C8	0.032 (8)	0.038 (8)	0.038 (10)	0.008 (7)	0.000 (7)	-0.004 (7)
C9	0.073 (11)	0.040 (9)	0.042 (11)	-0.007 (8)	0.025 (9)	-0.002 (8)
C10	0.040 (6)	0.048 (7)	0.060 (8)	0.002 (5)	0.012 (6)	-0.010 (6)
C11	0.040 (6)	0.048 (7)	0.060 (8)	0.002 (5)	0.012 (6)	-0.010 (6)
C12	0.032 (8)	0.034 (9)	0.061 (12)	-0.005 (7)	0.014 (8)	-0.007 (8)
C13	0.042 (9)	0.037 (9)	0.071 (14)	0.007 (7)	0.014 (9)	-0.001 (8)
C14	0.044 (10)	0.072 (13)	0.088 (15)	-0.007 (9)	0.028 (10)	0.018 (12)
C15	0.039 (9)	0.080 (13)	0.064 (13)	-0.003 (9)	0.020 (9)	-0.009 (11)

C16	0.035 (9)	0.066 (12)	0.12 (2)	-0.018 (9)	0.020 (10)	-0.031 (13)
C17	0.032 (7)	0.025 (7)	0.046 (10)	0.002 (6)	0.014 (7)	0.013 (7)
C18	0.047 (8)	0.036 (8)	0.039 (9)	-0.004 (7)	0.022 (7)	0.005 (7)
C19	0.045 (9)	0.031 (8)	0.053 (11)	0.006 (7)	0.019 (8)	0.019 (8)
C20	0.027 (7)	0.042 (9)	0.049 (10)	0.000 (7)	0.012 (7)	0.024 (8)
C21	0.033 (7)	0.039 (8)	0.037 (9)	0.010 (7)	0.003 (7)	0.008 (7)
C22	0.039 (8)	0.025 (7)	0.043 (9)	0.003 (6)	0.024 (7)	-0.004 (7)
C23	0.028 (7)	0.028 (7)	0.045 (9)	0.006 (6)	0.018 (6)	-0.007 (7)
C24	0.025 (7)	0.034 (8)	0.034 (9)	-0.004 (6)	0.001 (6)	-0.007 (6)
C25	0.034 (8)	0.043 (9)	0.038 (9)	-0.013 (7)	0.006 (7)	-0.007 (7)
C26	0.029 (8)	0.036 (8)	0.059 (11)	-0.007 (7)	-0.001 (7)	0.004 (7)
C27	0.040 (8)	0.029 (7)	0.034 (8)	0.002 (7)	0.007 (6)	0.009 (7)
C28	0.044 (8)	0.024 (7)	0.034 (9)	-0.002 (6)	0.009 (7)	-0.008 (6)
C29	0.025 (7)	0.048 (9)	0.050 (10)	0.002 (7)	0.008 (6)	0.011 (7)
C30	0.072 (11)	0.045 (9)	0.050 (11)	0.004 (9)	0.031 (9)	0.006 (8)
C31	0.029 (8)	0.046 (10)	0.064 (12)	0.009 (7)	0.006 (7)	0.007 (8)
C32	0.043 (8)	0.033 (8)	0.037 (9)	-0.004 (7)	-0.003 (7)	-0.001 (7)
N1	0.032 (6)	0.038 (7)	0.032 (7)	0.000 (5)	0.005 (5)	0.005 (5)
N2	0.027 (6)	0.043 (7)	0.025 (7)	-0.008 (5)	0.006 (5)	-0.003 (5)
N3	0.029 (6)	0.034 (6)	0.030 (7)	0.001 (5)	0.005 (5)	0.006 (5)
N4	0.038 (7)	0.022 (6)	0.032 (7)	-0.002 (5)	0.011 (5)	0.000 (5)
O1	0.030 (5)	0.052 (6)	0.035 (6)	-0.007 (5)	0.004 (4)	0.003 (5)
O2	0.031 (5)	0.048 (6)	0.027 (6)	-0.009 (4)	-0.002 (4)	0.007 (4)
Cu1	0.0300 (8)	0.0336 (9)	0.0334 (10)	-0.0028 (7)	0.0071 (7)	0.0030 (8)
Br1	0.0499 (9)	0.0385 (8)	0.0471 (10)	0.0081 (7)	0.0122 (7)	-0.0023 (7)
Br2	0.0559 (9)	0.0580 (10)	0.0374 (10)	-0.0012 (9)	-0.0006 (7)	0.0009 (8)
Br3	0.0452 (8)	0.0488 (9)	0.0470 (10)	0.0147 (8)	0.0118 (7)	-0.0006 (8)
Br4	0.0449 (8)	0.0618 (11)	0.0390 (10)	0.0137 (8)	0.0131 (7)	-0.0007 (8)
Br5	0.0397 (8)	0.0465 (9)	0.0494 (11)	0.0064 (7)	-0.0013 (7)	0.0111 (8)
Br6	0.0586 (10)	0.0565 (10)	0.0397 (10)	0.0181 (8)	0.0188 (8)	0.0027 (8)

Geometric parameters (Å, °)

C1—C2	1.382 (17)	C18—Br4	1.914 (15)
C1—C6	1.413 (19)	C19—C20	1.31 (2)
C1—N1	1.460 (18)	C19—H19	0.9500
C2—C3	1.395 (19)	C20—C21	1.39 (2)
C2—Br1	1.874 (13)	C20—Br5	1.926 (13)
C3—C4	1.36 (2)	C21—C22	1.372 (18)
C3—H3	0.9500	C21—H21	0.9500
C4—C5	1.341 (19)	C22—Br6	1.900 (13)
C4—Br2	1.935 (13)	C23—N4	1.388 (12)
C5—C6	1.42 (2)	C23—C24	1.3900
C5—H5	0.9500	C23—C28	1.3900
C6—Br3	1.870 (13)	C24—O2	1.348 (10)
C7—N2	1.361 (16)	C24—C25	1.3900
C7—C8	1.394 (12)	C25—C26	1.3900
C7—C12	1.436 (19)	C25—H25	0.9500

C8—O1	1.296 (16)	C26—C27	1.3900
C8—C9	1.44 (2)	C26—H26	0.9500
C9—C10	1.34 (2)	C27—C28	1.3900
C9—H9	0.9500	C27—C32	1.445 (16)
C10—C11	1.42 (2)	C28—C29	1.418 (16)
C10—H10	0.9500	C29—C30	1.37 (2)
C11—C12	1.39 (2)	C29—H29	0.9500
C11—C16	1.50 (2)	C30—C31	1.44 (2)
C12—C13	1.44 (2)	C30—H30	0.9500
C13—C14	1.38 (2)	C31—C32	1.35 (2)
C13—H13	0.9500	C31—H31	0.9500
C14—C15	1.36 (2)	C32—H32	0.9500
C14—H14	0.9500	N1—N2	1.284 (14)
C15—C16	1.33 (3)	Cu1—Br4	3.134 (2)
C15—H15	0.9500	Cu1—N1	1.947 (12)
C16—H16	0.9500	N3—N4	1.315 (14)
C17—C18	1.389 (19)	Cu1—N3	1.970 (11)
C17—C22	1.411 (19)	Cu1—O1	1.892 (9)
C17—N3	1.437 (16)	Cu1—O2	1.888 (8)
C18—C19	1.39 (2)		
C2—C1—C6	119.5 (13)	C18—C19—H19	120.8
C2—C1—N1	121.2 (12)	C19—C20—C21	124.2 (13)
C6—C1—N1	119.3 (12)	C19—C20—Br5	120.0 (11)
C1—C2—C3	121.0 (13)	C21—C20—Br5	115.6 (11)
C1—C2—Br1	119.8 (11)	C22—C21—C20	116.6 (14)
C3—C2—Br1	119.2 (10)	C22—C21—H21	121.7
C4—C3—C2	117.6 (12)	C20—C21—H21	121.7
C4—C3—H3	121.2	C21—C22—C17	122.8 (12)
C2—C3—H3	121.2	C21—C22—Br6	117.0 (11)
C5—C4—C3	124.9 (13)	C17—C22—Br6	120.1 (10)
C5—C4—Br2	115.3 (11)	N4—C23—C24	123.3 (7)
C3—C4—Br2	119.8 (10)	N4—C23—C28	115.8 (7)
C4—C5—C6	118.3 (13)	C24—C23—C28	120.0
C4—C5—H5	120.8	O2—C24—C25	114.8 (7)
C6—C5—H5	120.8	O2—C24—C23	125.2 (7)
C1—C6—C5	118.7 (12)	C25—C24—C23	120.0
C1—C6—Br3	121.9 (11)	C26—C25—C24	120.0
C5—C6—Br3	119.3 (11)	C26—C25—H25	120.0
N2—C7—C8	126.2 (12)	C24—C25—H25	120.0
N2—C7—C12	114.1 (12)	C25—C26—C27	120.0
C8—C7—C12	119.7 (13)	C25—C26—H26	120.0
O1—C8—C7	125.7 (12)	C27—C26—H26	120.0
O1—C8—C9	115.4 (12)	C28—C27—C26	120.0
C7—C8—C9	118.8 (13)	C28—C27—C32	120.5 (8)
C10—C9—C8	120.9 (15)	C26—C27—C32	119.5 (8)
C10—C9—H9	119.6	C27—C28—C23	120.0
C8—C9—H9	119.6	C27—C28—C29	117.6 (7)

C9—C10—C11	121.1 (15)	C23—C28—C29	122.4 (7)
C9—C10—H10	119.5	C30—C29—C28	122.7 (13)
C11—C10—H10	119.5	C30—C29—H29	118.6
C12—C11—C10	119.5 (15)	C28—C29—H29	118.6
C12—C11—C16	118.0 (17)	C29—C30—C31	118.9 (15)
C10—C11—C16	122.4 (16)	C29—C30—H30	120.6
C11—C12—C7	120.0 (15)	C31—C30—H30	120.6
C11—C12—C13	117.2 (14)	C32—C31—C30	119.9 (14)
C7—C12—C13	122.8 (14)	C32—C31—H31	120.0
C14—C13—C12	121.3 (16)	C30—C31—H31	120.0
C14—C13—H13	119.3	C31—C32—C27	120.2 (12)
C12—C13—H13	119.3	C31—C32—H32	119.9
C15—C14—C13	121.9 (18)	C27—C32—H32	119.9
C15—C14—H14	119.0	N2—N1—C1	111.9 (11)
C13—C14—H14	119.0	N2—N1—Cu1	129.9 (9)
C16—C15—C14	120.1 (17)	C1—N1—Cu1	117.7 (8)
C16—C15—H15	120.0	N1—N2—C7	120.3 (12)
C14—C15—H15	120.0	N4—N3—C17	111.9 (10)
C15—C16—C11	121.3 (18)	N4—N3—Cu1	128.3 (8)
C15—C16—H16	119.4	C17—N3—Cu1	119.4 (8)
C11—C16—H16	119.4	N3—N4—C23	119.1 (10)
C18—C17—C22	115.5 (12)	C8—O1—Cu1	125.8 (9)
C18—C17—N3	119.4 (13)	C24—O2—Cu1	121.9 (7)
C22—C17—N3	125.1 (12)	O2—Cu1—O1	169.4 (4)
C17—C18—C19	122.5 (14)	O2—Cu1—N1	91.3 (4)
C17—C18—Br4	119.0 (11)	O1—Cu1—N1	90.9 (4)
C19—C18—Br4	118.5 (11)	O2—Cu1—N3	87.6 (4)
C20—C19—C18	118.4 (14)	O1—Cu1—N3	92.1 (4)
C20—C19—H19	120.8	N1—Cu1—N3	169.3 (5)
C6—C1—C2—C3	0.6 (19)	N3—C17—C22—C21	-178.2 (12)
N1—C1—C2—C3	-177.6 (12)	C18—C17—C22—Br6	174.4 (10)
C6—C1—C2—Br1	179.6 (10)	N3—C17—C22—Br6	-2.0 (18)
N1—C1—C2—Br1	1.3 (16)	N4—C23—C24—O2	-11.8 (10)
C1—C2—C3—C4	0.6 (19)	C28—C23—C24—O2	179.9 (10)
Br1—C2—C3—C4	-178.4 (10)	N4—C23—C24—C25	168.3 (9)
C2—C3—C4—C5	-1 (2)	C28—C23—C24—C25	0.0
C2—C3—C4—Br2	-179.4 (9)	O2—C24—C25—C26	-179.9 (9)
C3—C4—C5—C6	0 (2)	C23—C24—C25—C26	0.0
Br2—C4—C5—C6	178.5 (10)	C24—C25—C26—C27	0.0
C2—C1—C6—C5	-1.6 (19)	C25—C26—C27—C28	0.0
N1—C1—C6—C5	176.7 (12)	C25—C26—C27—C32	179.7 (10)
C2—C1—C6—Br3	177.2 (10)	C26—C27—C28—C23	0.0
N1—C1—C6—Br3	-4.5 (17)	C32—C27—C28—C23	-179.7 (10)
C4—C5—C6—C1	1 (2)	C26—C27—C28—C29	178.7 (10)
C4—C5—C6—Br3	-177.5 (11)	C32—C27—C28—C29	-1.0 (12)
N2—C7—C8—O1	6 (2)	N4—C23—C28—C27	-169.2 (9)
C12—C7—C8—O1	-175.9 (13)	C24—C23—C28—C27	0.0

N2—C7—C8—C9	-175.9 (13)	N4—C23—C28—C29	12.2 (11)
C12—C7—C8—C9	2.2 (19)	C24—C23—C28—C29	-178.6 (11)
O1—C8—C9—C10	176.4 (14)	C27—C28—C29—C30	-2.0 (18)
C7—C8—C9—C10	-2 (2)	C23—C28—C29—C30	176.6 (11)
C8—C9—C10—C11	1 (2)	C28—C29—C30—C31	3 (2)
C9—C10—C11—C12	0 (2)	C29—C30—C31—C32	-1 (2)
C9—C10—C11—C16	-177.0 (15)	C30—C31—C32—C27	-2 (2)
C10—C11—C12—C7	0 (2)	C28—C27—C32—C31	2.9 (17)
C16—C11—C12—C7	177.5 (14)	C26—C27—C32—C31	-176.7 (11)
C10—C11—C12—C13	179.0 (14)	C2—C1—N1—N2	-107.0 (14)
C16—C11—C12—C13	-4 (2)	C6—C1—N1—N2	74.8 (15)
N2—C7—C12—C11	176.9 (12)	C2—C1—N1—Cu1	80.7 (14)
C8—C7—C12—C11	-1 (2)	C6—C1—N1—Cu1	-97.6 (12)
N2—C7—C12—C13	-1.8 (19)	C1—N1—N2—C7	-179.3 (11)
C8—C7—C12—C13	179.9 (13)	Cu1—N1—N2—C7	-8.1 (18)
C11—C12—C13—C14	3 (2)	C8—C7—N2—N1	-3 (2)
C7—C12—C13—C14	-178.0 (15)	C12—C7—N2—N1	179.0 (12)
C12—C13—C14—C15	0 (3)	C18—C17—N3—N4	131.8 (12)
C13—C14—C15—C16	-3 (3)	C22—C17—N3—N4	-52.0 (17)
C14—C15—C16—C11	2 (3)	C18—C17—N3—Cu1	-53.8 (15)
C12—C11—C16—C15	1 (3)	C22—C17—N3—Cu1	122.4 (12)
C10—C11—C16—C15	178.3 (17)	C17—N3—N4—C23	-178.2 (10)
C22—C17—C18—C19	1.8 (19)	Cu1—N3—N4—C23	8.0 (16)
N3—C17—C18—C19	178.4 (12)	C24—C23—N4—N3	20.3 (13)
C22—C17—C18—Br4	-179.3 (9)	C28—C23—N4—N3	-170.9 (8)
N3—C17—C18—Br4	-2.7 (17)	C7—C8—O1—Cu1	2 (2)
C17—C18—C19—C20	0 (2)	C9—C8—O1—Cu1	-176.2 (9)
Br4—C18—C19—C20	-178.8 (11)	C25—C24—O2—Cu1	154.8 (5)
C18—C19—C20—C21	-2 (2)	C23—C24—O2—Cu1	-25.1 (11)
C18—C19—C20—Br5	-176.8 (10)	C24—O2—Cu1—O1	-52 (3)
C19—C20—C21—C22	2 (2)	C24—O2—Cu1—N1	-153.4 (9)
Br5—C20—C21—C22	176.9 (10)	C24—O2—Cu1—N3	37.3 (9)
C20—C21—C22—C17	0 (2)	C8—O1—Cu1—O2	-110 (2)
C20—C21—C22—Br6	-176.3 (10)	C8—O1—Cu1—N1	-8.2 (11)
C18—C17—C22—C21	-1.8 (19)	C8—O1—Cu1—N3	161.5 (11)

Hydrogen-bond geometry (\AA , $^\circ$)

Cg1 is the centroid of the C27–C32 ring.

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
C5—H5 \cdots Br6 ⁱ	0.95	2.75	3.546 (15)	142
C3—H3 \cdots Cg1 ⁱⁱ	0.95	2.99	3.729 (15)	136

Symmetry codes: (i) $x, y, z-1$; (ii) $-x+1, y+1/2, -z$.

(II) Bis{(E)-1-[(2,4,6-tribromophenyl)diazanyl]naphthalen-2-olato}nickel(II)

Crystal data

[Ni(C₁₆H₈Br₃N₂O)₂]
M_r = 1026.66
 Monoclinic, *P*2₁/*n*
a = 11.0909 (6) Å
b = 12.4571 (6) Å
c = 12.5382 (7) Å
 β = 107.820 (2)°
V = 1649.17 (15) Å³
Z = 2

F(000) = 980
D_x = 2.067 Mg m⁻³
 Mo *K*α radiation, λ = 0.71073 Å
 Cell parameters from 7253 reflections
 θ = 1.0–27.5°
 μ = 7.89 mm⁻¹
T = 173 K
 Prism, black
 0.30 × 0.22 × 0.06 mm

Data collection

Nonius KappaCCD
 diffractometer
 Radiation source: sealed tube
 Graphite monochromator
 phi and ω scans
 Absorption correction: multi-scan
 (MULABS; Spek, 2009)
T_{min} = 0.151, *T_{max}* = 0.317

11360 measured reflections
 3745 independent reflections
 2214 reflections with *I* > 2σ(*I*)
R_{int} = 0.094
 θ_{\max} = 27.5°, θ_{\min} = 2.9°
h = -13→14
k = -13→16
l = -16→16

Refinement

Refinement on *F*²
 Least-squares matrix: full
R [*F*² > 2σ(*F*²)] = 0.043
wR(*F*²) = 0.096
S = 0.95
 3745 reflections
 205 parameters
 0 restraints
 Primary atom site location: structure-invariant
 direct methods

Secondary atom site location: difference Fourier
 map
 Hydrogen site location: inferred from
 neighbouring sites
 H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.034P)^2]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} < 0.001$
 $\Delta\rho_{\max} = 0.57 \text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.66 \text{ e \AA}^{-3}$

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²)

	<i>x</i>	<i>y</i>	<i>z</i>	<i>U_{iso}</i> */ <i>U_{eq}</i>
C1	0.2389 (4)	0.4906 (3)	0.3565 (3)	0.0232 (10)
C2	0.2302 (4)	0.5055 (4)	0.2452 (4)	0.0287 (11)
C3	0.1396 (4)	0.4527 (4)	0.1602 (4)	0.0321 (12)
H3	0.1357	0.4619	0.0840	0.038*
C4	0.0551 (4)	0.3862 (3)	0.1896 (4)	0.0295 (12)
C5	0.0609 (4)	0.3679 (4)	0.3004 (4)	0.0318 (12)
H5	0.0028	0.3212	0.3192	0.038*
C6	0.1547 (4)	0.4204 (4)	0.3819 (3)	0.0277 (11)

C7	0.3408 (4)	0.6940 (3)	0.5623 (4)	0.0283 (11)
C8	0.4729 (4)	0.6805 (4)	0.6238 (4)	0.0293 (11)
C9	0.5307 (4)	0.7555 (4)	0.7110 (4)	0.0335 (12)
H9	0.6165	0.7462	0.7549	0.040*
C10	0.4637 (4)	0.8392 (4)	0.7310 (4)	0.0323 (12)
H10	0.5056	0.8894	0.7871	0.039*
C11	0.3320 (4)	0.8562 (4)	0.6717 (4)	0.0296 (11)
C12	0.2707 (4)	0.7808 (4)	0.5879 (3)	0.0272 (11)
C13	0.1395 (4)	0.7939 (4)	0.5345 (4)	0.0308 (12)
H13	0.0955	0.7435	0.4795	0.037*
C14	0.0748 (5)	0.8782 (4)	0.5608 (4)	0.0386 (13)
H14	-0.0135	0.8852	0.5240	0.046*
C15	0.1371 (5)	0.9547 (4)	0.6414 (4)	0.0416 (13)
H15	0.0916	1.0131	0.6591	0.050*
C16	0.2644 (5)	0.9436 (4)	0.6940 (4)	0.0342 (12)
H16	0.3075	0.9964	0.7466	0.041*
N1	0.3320 (3)	0.5483 (3)	0.4453 (3)	0.0251 (9)
N2	0.2773 (3)	0.6275 (3)	0.4773 (3)	0.0278 (9)
O1	0.5435 (3)	0.6061 (2)	0.6047 (2)	0.0328 (8)
Ni	0.5000	0.5000	0.5000	0.0271 (2)
Br1	0.16663 (5)	0.39355 (4)	0.53300 (4)	0.04164 (17)
Br2	0.34269 (5)	0.60009 (4)	0.20681 (4)	0.04697 (18)
Br3	-0.07417 (5)	0.31689 (4)	0.07466 (4)	0.05057 (19)

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.014 (2)	0.027 (2)	0.024 (2)	0.003 (2)	-0.0014 (19)	-0.004 (2)
C2	0.023 (2)	0.031 (3)	0.027 (2)	-0.007 (2)	0.000 (2)	-0.002 (2)
C3	0.029 (3)	0.037 (3)	0.025 (2)	0.004 (2)	0.001 (2)	0.000 (2)
C4	0.020 (2)	0.027 (3)	0.033 (3)	0.002 (2)	-0.005 (2)	-0.008 (2)
C5	0.028 (3)	0.030 (3)	0.034 (3)	-0.002 (2)	0.005 (2)	0.002 (2)
C6	0.025 (3)	0.031 (3)	0.022 (2)	0.006 (2)	0.000 (2)	0.001 (2)
C7	0.024 (3)	0.030 (3)	0.025 (2)	0.001 (2)	-0.002 (2)	-0.004 (2)
C8	0.025 (3)	0.034 (3)	0.025 (2)	-0.006 (2)	0.002 (2)	-0.004 (2)
C9	0.023 (3)	0.037 (3)	0.033 (3)	-0.003 (2)	-0.003 (2)	-0.009 (2)
C10	0.032 (3)	0.034 (3)	0.027 (2)	-0.009 (2)	0.002 (2)	-0.002 (2)
C11	0.033 (3)	0.028 (3)	0.026 (2)	-0.002 (2)	0.006 (2)	-0.001 (2)
C12	0.024 (3)	0.032 (3)	0.024 (2)	0.003 (2)	0.004 (2)	-0.002 (2)
C13	0.029 (3)	0.029 (3)	0.031 (2)	0.006 (2)	0.004 (2)	-0.003 (2)
C14	0.025 (3)	0.050 (3)	0.035 (3)	0.009 (3)	0.001 (2)	0.004 (3)
C15	0.043 (3)	0.040 (3)	0.042 (3)	0.017 (3)	0.011 (3)	0.001 (3)
C16	0.042 (3)	0.031 (3)	0.030 (3)	0.000 (2)	0.011 (2)	-0.003 (2)
N1	0.018 (2)	0.029 (2)	0.0214 (19)	-0.0025 (17)	-0.0034 (16)	-0.0038 (17)
N2	0.026 (2)	0.025 (2)	0.026 (2)	0.0012 (18)	-0.0019 (18)	-0.0021 (17)
O1	0.0247 (18)	0.0333 (19)	0.0316 (18)	0.0022 (15)	-0.0044 (15)	-0.0132 (14)
Ni	0.0197 (4)	0.0297 (5)	0.0255 (4)	0.0004 (4)	-0.0026 (4)	-0.0040 (4)
Br1	0.0424 (3)	0.0495 (4)	0.0289 (3)	-0.0039 (3)	0.0049 (2)	0.0067 (2)

Br2	0.0386 (3)	0.0609 (4)	0.0378 (3)	-0.0170 (3)	0.0063 (3)	0.0049 (3)
Br3	0.0419 (3)	0.0534 (4)	0.0427 (3)	-0.0125 (3)	-0.0073 (3)	-0.0172 (3)

Geometric parameters (Å, °)

C1—C2	1.382 (6)	C9—H9	0.9500
C1—C6	1.385 (6)	C10—C11	1.437 (6)
C1—N1	1.455 (5)	C10—H10	0.9500
C2—C3	1.387 (6)	C11—C16	1.399 (7)
C2—Br2	1.883 (5)	C11—C12	1.419 (6)
C3—C4	1.382 (7)	C12—C13	1.413 (6)
C3—H3	0.9500	C13—C14	1.367 (6)
C4—C5	1.389 (6)	C13—H13	0.9500
C4—Br3	1.901 (4)	C14—C15	1.407 (7)
C5—C6	1.381 (6)	C14—H14	0.9500
C5—H5	0.9500	C15—C16	1.370 (7)
C6—Br1	1.888 (4)	C15—H15	0.9500
C7—N2	1.363 (5)	C16—H16	0.9500
C7—C12	1.425 (6)	N1—N2	1.285 (5)
C7—C8	1.441 (6)	Ni—N1	1.876 (3)
C8—O1	1.281 (5)	Ni—O1	1.821 (3)
C8—C9	1.431 (6)	Ni—O1 ⁱ	1.821 (3)
C9—C10	1.348 (6)	Ni—N1 ⁱ	1.877 (3)
C2—C1—C6	118.3 (4)	C16—C11—C12	119.9 (4)
C2—C1—N1	121.3 (4)	C16—C11—C10	122.2 (4)
C6—C1—N1	120.4 (4)	C12—C11—C10	117.8 (4)
C1—C2—C3	121.5 (4)	C13—C12—C11	117.7 (4)
C1—C2—Br2	119.7 (3)	C13—C12—C7	122.4 (4)
C3—C2—Br2	118.8 (4)	C11—C12—C7	119.9 (4)
C4—C3—C2	118.0 (4)	C14—C13—C12	121.0 (4)
C4—C3—H3	121.0	C14—C13—H13	119.5
C2—C3—H3	121.0	C12—C13—H13	119.5
C3—C4—C5	122.5 (4)	C13—C14—C15	121.0 (4)
C3—C4—Br3	119.0 (4)	C13—C14—H14	119.5
C5—C4—Br3	118.5 (4)	C15—C14—H14	119.5
C6—C5—C4	117.1 (4)	C16—C15—C14	119.0 (5)
C6—C5—H5	121.4	C16—C15—H15	120.5
C4—C5—H5	121.4	C14—C15—H15	120.5
C5—C6—C1	122.5 (4)	C15—C16—C11	121.3 (4)
C5—C6—Br1	117.7 (4)	C15—C16—H16	119.4
C1—C6—Br1	119.8 (3)	C11—C16—H16	119.4
N2—C7—C12	116.7 (4)	N2—N1—C1	109.0 (3)
N2—C7—C8	123.0 (4)	N2—N1—Ni	130.0 (3)
C12—C7—C8	120.2 (4)	C1—N1—Ni	120.9 (3)
O1—C8—C9	117.3 (4)	N1—N2—C7	122.0 (4)
O1—C8—C7	124.2 (4)	C8—O1—Ni	128.1 (3)
C9—C8—C7	118.5 (4)	O1—Ni—O1 ⁱ	180

C10—C9—C8	120.3 (4)	O1—Ni—N1	92.59 (14)
C10—C9—H9	119.9	O1 ⁱ —Ni—N1	87.41 (14)
C8—C9—H9	119.9	O1—Ni—N1 ⁱ	87.41 (14)
C9—C10—C11	123.2 (4)	O1 ⁱ —Ni—N1 ⁱ	92.59 (14)
C9—C10—H10	118.4	N1—Ni—N1 ⁱ	180
C11—C10—H10	118.4		
C6—C1—C2—C3	0.3 (7)	C10—C11—C12—C7	-2.3 (7)
N1—C1—C2—C3	-178.4 (4)	N2—C7—C12—C13	4.6 (7)
C6—C1—C2—Br2	-180.0 (3)	C8—C7—C12—C13	-176.3 (4)
N1—C1—C2—Br2	1.3 (6)	N2—C7—C12—C11	-177.0 (4)
C1—C2—C3—C4	1.7 (7)	C8—C7—C12—C11	2.1 (7)
Br2—C2—C3—C4	-178.0 (3)	C11—C12—C13—C14	1.7 (7)
C2—C3—C4—C5	-2.3 (7)	C7—C12—C13—C14	-179.9 (5)
C2—C3—C4—Br3	177.6 (3)	C12—C13—C14—C15	0.3 (8)
C3—C4—C5—C6	0.8 (7)	C13—C14—C15—C16	-0.1 (8)
Br3—C4—C5—C6	-179.1 (3)	C14—C15—C16—C11	-2.2 (8)
C4—C5—C6—C1	1.4 (7)	C12—C11—C16—C15	4.2 (7)
C4—C5—C6—Br1	-178.4 (3)	C10—C11—C16—C15	-175.9 (4)
C2—C1—C6—C5	-1.9 (7)	C2—C1—N1—N2	100.7 (5)
N1—C1—C6—C5	176.8 (4)	C6—C1—N1—N2	-78.1 (5)
C2—C1—C6—Br1	177.8 (3)	C2—C1—N1—Ni	-82.2 (5)
N1—C1—C6—Br1	-3.4 (5)	C6—C1—N1—Ni	99.1 (4)
N2—C7—C8—O1	0.5 (7)	C1—N1—N2—C7	178.0 (4)
C12—C7—C8—O1	-178.6 (4)	Ni—N1—N2—C7	1.1 (6)
N2—C7—C8—C9	179.6 (4)	C12—C7—N2—N1	178.2 (4)
C12—C7—C8—C9	0.5 (7)	C8—C7—N2—N1	-0.9 (7)
O1—C8—C9—C10	176.2 (5)	C9—C8—O1—Ni	-179.4 (3)
C7—C8—C9—C10	-2.9 (7)	C7—C8—O1—Ni	-0.3 (7)
C8—C9—C10—C11	2.8 (7)	C8—O1—Ni—N1	0.4 (4)
C9—C10—C11—C16	179.9 (5)	C8—O1—Ni—N1 ⁱ	-179.6 (4)
C9—C10—C11—C12	-0.2 (7)	N2—N1—Ni—O1	-0.8 (4)
C16—C11—C12—C13	-3.9 (7)	C1—N1—Ni—O1	-177.3 (3)
C10—C11—C12—C13	176.2 (4)	N2—N1—Ni—O1 ⁱ	179.2 (4)
C16—C11—C12—C7	177.6 (4)	C1—N1—Ni—O1 ⁱ	2.7 (3)

Symmetry code: (i) $-x+1, -y+1, -z+1$.

Hydrogen-bond geometry (\AA , $^\circ$)

Cg2 is the centroid of the C1—C6 ring.

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
C10—H10 \cdots Cg2 ⁱⁱ	0.95	2.71	3.391 (5)	130

Symmetry code: (ii) $x+1/2, -y+3/2, z+1/2$.

(III) Bis{(E)-1-[(2,4,6-tribromophenyl)diazenyl]naphthalen-2-olato}palladium(II)

Crystal data

[Pd(C₁₆H₈Br₃N₂O)₂]
M_r = 1074.35
 Monoclinic, *P*2₁/*n*
a = 11.1896 (8) Å
b = 12.4540 (8) Å
c = 12.5511 (9) Å
 β = 107.749 (5)°
V = 1665.8 (2) Å³
Z = 2

F(000) = 1016
D_x = 2.142 Mg m⁻³
 Cu *K*α radiation, λ = 1.54186 Å
 Cell parameters from 3651 reflections
 θ = 2.1–22.3°
 μ = 13.23 mm⁻¹
T = 200 K
 Square plate, dark red
 0.12 × 0.09 × 0.03 mm

Data collection

STOE IPDS 2T
 diffractometer
 Radiation source: Genix-Cu,3D
 Graphite monochromator
 Detector resolution: 6.67 pixels mm⁻¹
 rotation method scans
 Absorption correction: multi-scan
 (MULABS; Spek, 2009)
T_{min} = 0.360, *T_{max}* = 1.000

13003 measured reflections
 2895 independent reflections
 2371 reflections with *I* > 2σ(*I*)
R_{int} = 0.142
 θ_{\max} = 67.7°, θ_{\min} = 5.5°
h = -13→12
k = -14→14
l = -14→15

Refinement

Refinement on *F*²
 Least-squares matrix: full
R[*F*² > 2σ(*F*²)] = 0.057
wR(*F*²) = 0.170
S = 1.11
 2895 reflections
 206 parameters
 0 restraints
 Primary atom site location: structure-invariant
 direct methods
 Secondary atom site location: difference Fourier
 map

Hydrogen site location: inferred from
 neighbouring sites
 H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.1019P)^2 + 0.8121P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} < 0.001$
 $\Delta\rho_{\max} = 0.88 \text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -1.10 \text{ e \AA}^{-3}$
 Extinction correction: SHELXL,
 $F_c^* = kF_c[1 + 0.001x F_c^2 \lambda^3 / \sin(2\theta)]^{-1/4}$
 Extinction coefficient: 0.0040 (4)

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²)

	<i>x</i>	<i>y</i>	<i>z</i>	<i>U_{iso}</i> */ <i>U_{eq}</i>
Br1	0.16311 (8)	0.39498 (7)	0.53262 (6)	0.0659 (3)
Br2	0.33399 (8)	0.60480 (8)	0.20631 (7)	0.0737 (4)
Br3	-0.07036 (8)	0.31295 (7)	0.07577 (7)	0.0754 (4)
Pd1	0.5000	0.5000	0.5000	0.0456 (3)
O1	0.5428 (4)	0.6130 (4)	0.6155 (4)	0.0573 (12)
N1	0.3223 (5)	0.5521 (4)	0.4432 (4)	0.0468 (12)

N2	0.2730 (5)	0.6310 (4)	0.4798 (4)	0.0486 (12)
C1	0.2323 (6)	0.4941 (5)	0.3582 (5)	0.0471 (15)
C2	0.2246 (6)	0.5074 (5)	0.2447 (5)	0.0482 (15)
C3	0.1354 (7)	0.4532 (6)	0.1598 (5)	0.0563 (17)
H3	0.1311	0.4629	0.0836	0.068*
C4	0.0544 (7)	0.3855 (5)	0.1889 (6)	0.0532 (16)
C5	0.0597 (6)	0.3675 (5)	0.2996 (6)	0.0495 (15)
H5	0.0032	0.3195	0.3184	0.059*
C6	0.1513 (6)	0.4227 (5)	0.3817 (5)	0.0474 (14)
C7	0.3371 (6)	0.6961 (5)	0.5647 (5)	0.0464 (14)
C8	0.4672 (6)	0.6842 (6)	0.6279 (5)	0.0523 (15)
C9	0.5192 (7)	0.7634 (6)	0.7146 (6)	0.0570 (17)
H9	0.6039	0.7562	0.7601	0.068*
C10	0.4511 (7)	0.8468 (6)	0.7323 (6)	0.0604 (18)
H10	0.4904	0.8983	0.7877	0.072*
C11	0.3214 (7)	0.8605 (6)	0.6705 (6)	0.0536 (16)
C12	0.2640 (7)	0.7831 (5)	0.5895 (5)	0.0505 (15)
C13	0.1345 (7)	0.7932 (6)	0.5352 (6)	0.0583 (17)
H13	0.0933	0.7427	0.4793	0.070*
C14	0.0678 (8)	0.8748 (7)	0.5619 (7)	0.070 (2)
H14	-0.0202	0.8783	0.5263	0.084*
C15	0.1257 (8)	0.9538 (7)	0.6406 (6)	0.071 (2)
H15	0.0780	1.0109	0.6573	0.085*
C16	0.2509 (8)	0.9469 (7)	0.6922 (7)	0.0652 (19)
H16	0.2917	1.0010	0.7438	0.078*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Br1	0.0649 (6)	0.0817 (6)	0.0436 (5)	-0.0081 (4)	0.0055 (4)	0.0092 (3)
Br2	0.0614 (6)	0.0997 (7)	0.0531 (5)	-0.0261 (4)	0.0070 (4)	0.0073 (4)
Br3	0.0648 (6)	0.0834 (6)	0.0603 (6)	-0.0172 (4)	-0.0073 (4)	-0.0226 (4)
Pd1	0.0353 (4)	0.0550 (4)	0.0369 (4)	-0.0014 (3)	-0.0032 (3)	-0.0038 (3)
O1	0.042 (3)	0.065 (3)	0.052 (3)	0.003 (2)	-0.005 (2)	-0.009 (2)
N1	0.042 (3)	0.053 (3)	0.037 (3)	-0.007 (2)	-0.001 (2)	-0.004 (2)
N2	0.040 (3)	0.054 (3)	0.042 (3)	-0.004 (2)	-0.003 (2)	0.000 (2)
C1	0.038 (3)	0.054 (3)	0.037 (3)	0.007 (3)	-0.007 (3)	0.001 (3)
C2	0.036 (3)	0.061 (4)	0.037 (3)	0.000 (3)	-0.005 (3)	0.002 (3)
C3	0.049 (4)	0.073 (4)	0.038 (3)	0.004 (3)	-0.001 (3)	-0.001 (3)
C4	0.045 (4)	0.056 (4)	0.048 (4)	-0.005 (3)	-0.002 (3)	-0.005 (3)
C5	0.043 (4)	0.047 (3)	0.054 (4)	-0.002 (3)	0.008 (3)	0.002 (3)
C6	0.042 (3)	0.049 (3)	0.040 (3)	-0.002 (3)	-0.003 (3)	0.000 (3)
C7	0.035 (3)	0.057 (3)	0.042 (3)	-0.001 (3)	0.004 (3)	-0.005 (3)
C8	0.045 (4)	0.063 (4)	0.040 (3)	-0.007 (3)	0.001 (3)	0.001 (3)
C9	0.046 (4)	0.070 (4)	0.047 (4)	-0.004 (3)	0.002 (3)	-0.012 (3)
C10	0.059 (4)	0.070 (4)	0.043 (4)	-0.014 (4)	0.002 (3)	-0.010 (3)
C11	0.053 (4)	0.060 (4)	0.046 (4)	0.001 (3)	0.012 (3)	-0.002 (3)
C12	0.048 (4)	0.057 (4)	0.043 (3)	-0.007 (3)	0.008 (3)	-0.001 (3)

C13	0.046 (4)	0.070 (4)	0.052 (4)	0.000 (3)	0.004 (3)	-0.010 (3)
C14	0.056 (5)	0.080 (5)	0.064 (5)	0.002 (4)	0.004 (4)	-0.003 (4)
C15	0.074 (6)	0.072 (5)	0.060 (5)	0.019 (4)	0.009 (4)	0.001 (4)
C16	0.069 (5)	0.065 (4)	0.057 (4)	-0.002 (4)	0.013 (4)	0.002 (4)

Geometric parameters (Å, °)

Br1—C6	1.889 (6)	C5—H5	0.9500
Br2—C2	1.887 (7)	C7—C8	1.437 (9)
Br3—C4	1.890 (7)	C7—C12	1.448 (9)
Pd1—O1	1.972 (5)	C8—C9	1.452 (10)
Pd1—O1 ⁱ	1.972 (5)	C9—C10	1.346 (10)
Pd1—N1 ⁱ	2.004 (5)	C9—H9	0.9500
Pd1—N1	2.004 (5)	C10—C11	1.432 (10)
O1—C8	1.268 (8)	C10—H10	0.9500
N1—N2	1.279 (8)	C11—C12	1.406 (10)
N1—C1	1.422 (8)	C11—C16	1.409 (11)
N2—C7	1.356 (8)	C12—C13	1.406 (10)
C1—C6	1.365 (9)	C13—C14	1.362 (11)
C1—C2	1.410 (9)	C13—H13	0.9500
C2—C3	1.393 (9)	C14—C15	1.405 (12)
C3—C4	1.367 (10)	C14—H14	0.9500
C3—H3	0.9500	C15—C16	1.355 (11)
C4—C5	1.390 (10)	C15—H15	0.9500
C5—C6	1.394 (9)	C16—H16	0.9500
O1—Pd1—O1 ⁱ	180.0	N2—C7—C12	114.7 (6)
O1—Pd1—N1 ⁱ	88.7 (2)	C8—C7—C12	120.1 (6)
O1 ⁱ —Pd1—N1 ⁱ	91.3 (2)	O1—C8—C7	127.3 (6)
O1—Pd1—N1	91.3 (2)	O1—C8—C9	115.9 (6)
O1 ⁱ —Pd1—N1	88.7 (2)	C7—C8—C9	116.8 (6)
N1 ⁱ —Pd1—N1	180.0	C10—C9—C8	122.0 (7)
C8—O1—Pd1	124.6 (4)	C10—C9—H9	119.0
N2—N1—C1	112.0 (5)	C8—C9—H9	119.0
N2—N1—Pd1	127.7 (4)	C9—C10—C11	122.3 (6)
C1—N1—Pd1	120.1 (4)	C9—C10—H10	118.9
N1—N2—C7	123.9 (5)	C11—C10—H10	118.9
C6—C1—C2	117.1 (6)	C12—C11—C16	120.3 (7)
C6—C1—N1	122.3 (6)	C12—C11—C10	118.3 (6)
C2—C1—N1	120.6 (6)	C16—C11—C10	121.4 (7)
C3—C2—C1	121.7 (6)	C13—C12—C11	117.7 (6)
C3—C2—Br2	119.0 (5)	C13—C12—C7	121.9 (6)
C1—C2—Br2	119.3 (5)	C11—C12—C7	120.4 (6)
C4—C3—C2	118.2 (6)	C14—C13—C12	120.6 (7)
C4—C3—H3	120.9	C14—C13—H13	119.7
C2—C3—H3	120.9	C12—C13—H13	119.7
C3—C4—C5	122.5 (6)	C13—C14—C15	121.7 (8)
C3—C4—Br3	119.5 (5)	C13—C14—H14	119.1

C5—C4—Br3	118.0 (5)	C15—C14—H14	119.1
C4—C5—C6	117.2 (6)	C16—C15—C14	118.7 (8)
C4—C5—H5	121.4	C16—C15—H15	120.7
C6—C5—H5	121.4	C14—C15—H15	120.7
C1—C6—C5	123.3 (6)	C15—C16—C11	120.9 (7)
C1—C6—Br1	119.2 (5)	C15—C16—H16	119.5
C5—C6—Br1	117.5 (5)	C11—C16—H16	119.5
N2—C7—C8	125.2 (6)		
C1—N1—N2—C7	176.7 (6)	Pd1—O1—C8—C9	-177.0 (5)
Pd1—N1—N2—C7	1.8 (9)	N2—C7—C8—O1	0.2 (11)
N2—N1—C1—C6	-77.8 (7)	C12—C7—C8—O1	-179.5 (6)
Pd1—N1—C1—C6	97.6 (6)	N2—C7—C8—C9	179.0 (6)
N2—N1—C1—C2	103.1 (7)	C12—C7—C8—C9	-0.7 (9)
Pd1—N1—C1—C2	-81.6 (6)	O1—C8—C9—C10	176.2 (7)
C6—C1—C2—C3	2.2 (9)	C7—C8—C9—C10	-2.7 (10)
N1—C1—C2—C3	-178.6 (6)	C8—C9—C10—C11	2.8 (11)
C6—C1—C2—Br2	-180.0 (5)	C9—C10—C11—C12	0.7 (11)
N1—C1—C2—Br2	-0.8 (8)	C9—C10—C11—C16	178.4 (7)
C1—C2—C3—C4	-0.1 (10)	C16—C11—C12—C13	-2.3 (10)
Br2—C2—C3—C4	-178.0 (5)	C10—C11—C12—C13	175.4 (7)
C2—C3—C4—C5	-1.4 (11)	C16—C11—C12—C7	178.2 (6)
C2—C3—C4—Br3	178.6 (5)	C10—C11—C12—C7	-4.1 (10)
C3—C4—C5—C6	0.8 (10)	N2—C7—C12—C13	4.8 (9)
Br3—C4—C5—C6	-179.2 (5)	C8—C7—C12—C13	-175.4 (6)
C2—C1—C6—C5	-2.9 (10)	N2—C7—C12—C11	-175.7 (6)
N1—C1—C6—C5	177.9 (6)	C8—C7—C12—C11	4.1 (9)
C2—C1—C6—Br1	176.4 (5)	C11—C12—C13—C14	-0.8 (11)
N1—C1—C6—Br1	-2.8 (8)	C7—C12—C13—C14	178.7 (7)
C4—C5—C6—C1	1.5 (10)	C12—C13—C14—C15	2.6 (13)
C4—C5—C6—Br1	-177.8 (5)	C13—C14—C15—C16	-1.2 (13)
N1—N2—C7—C8	-2.2 (10)	C14—C15—C16—C11	-2.0 (12)
N1—N2—C7—C12	177.6 (6)	C12—C11—C16—C15	3.8 (11)
Pd1—O1—C8—C7	1.8 (10)	C10—C11—C16—C15	-173.9 (7)

Symmetry code: (i) $-x+1, -y+1, -z+1$.

Hydrogen-bond geometry (\AA , $^\circ$)

Cg2 is the centroid of ring C1-C6.

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
C10—H10 \cdots Cg2 ⁱⁱ	0.95	2.70	3.371 (8)	128

Symmetry code: (ii) $x+1/2, -y+3/2, z+1/2$.