

SCIENTIFIC REPORTS

OPEN

Novel positive allosteric modulators of GABA_A receptors with anesthetic activity

Maria C. Maldifassi¹, Roland Baur¹, David Pierce², Anahita Nourmahad², Stuart A. Forman² & Erwin Sigel¹

Received: 05 March 2016

Accepted: 25 April 2016

Published: 20 May 2016

GABA_A receptors are the main inhibitory neurotransmitter receptors in the brain and are targets for numerous clinically important drugs such as benzodiazepines, anxiolytics and anesthetics. We previously identified novel ligands of the classical benzodiazepine binding pocket in $\alpha_1\beta_2\gamma_2$ GABA_A receptors using an experiment-guided virtual screening (EGVS) method. This screen also identified novel ligands for intramembrane low affinity diazepam site(s). In the current study we have further characterized compounds 31 and 132 identified with EGVS as well as 4-O-methylhonokiol. We investigated the site of action of these compounds in $\alpha_1\beta_2\gamma_2$ GABA_A receptors expressed in *Xenopus laevis* oocytes using voltage-clamp electrophysiology combined with a benzodiazepine site antagonist and transmembrane domain mutations. All three compounds act mainly through the two β +/ α - subunit transmembrane interfaces of the GABA_A receptors. We then used concatenated receptors to dissect the involvement of individual β +/ α - interfaces. We further demonstrated that these compounds have anesthetic activity in a small aquatic animal model, *Xenopus laevis* tadpoles. The newly identified compounds may serve as scaffolds for the development of novel anesthetics.

The search for novel anesthetics has been triggered by the rising age of patients and increasing use of anesthesia outside the operating room^{1,2}. A key site of action of the potent anesthetics propofol and etomidate is the major inhibitory receptor in the mammalian central nervous system, the γ -aminobutyric acid type A (GABA_A) receptor. These receptors are composed of five homologous subunits organized around a central Cl⁻ selective channel³. Each subunit contains a large N-terminal extracellular domain (ECD), a transmembrane domain (TMD) with four alpha-helices (TM1 to TM4), and a variable-length intracellular domain (ICD) between TM3 and TM4. 19 subunits of the GABA_A receptor have been cloned (for review see: 4–6), denoting that numerous types of receptor isoforms exist⁵. The most abundant GABA_A receptor in the brain comprises α_1 , β_2 and γ_2 subunits^{3–5,7}. The receptor possesses a 2 α :2 β :1 γ subunit stoichiometry^{8–11}, with a subunit arrangement of $\gamma\beta\alpha\beta\alpha$ anti-clockwise as seen from the synaptic cleft^{10–13}. The receptor composition and arrangement influence its pharmacological properties^{14,15}.

Benzodiazepines modulate $\alpha_1\beta_2\gamma_2$ GABA_A receptor function by binding to a high affinity site located at the α +/ γ - ECD interface, homologous to the agonist binding sites at β +/ α - ECD interfaces^{16,17}. In addition to the high affinity binding site for benzodiazepines (site 1), there are other low affinity sites. One of these, site 2, is located at the ECD α +/ β - interface^{18,19}. Others, together designated as site 3, are located in the TMD, based on abolition of benzodiazepine effects by combined isoleucine substitutions at the homologous residues α_1 S269, β_2 N265, and γ_2 S280²⁰.

GABA_A receptors are also targets for potent intravenous anesthetics, including barbiturates, propofol and etomidate^{21–25}. Interestingly, receptor sensitivity to intravenous anesthetics is affected by benzodiazepine site 3 mutations^{26–30}. Diverse anesthetics not only potentiate GABA-induced Cl⁻ currents, but additionally at high concentrations directly activate GABA_A receptors^{31,32}. Photo-affinity labeling has located allosteric sites for the intravenous anesthetics etomidate and propofol to the TM1 of α and TM2, TM3 of β subunits^{23,24,33}.

Previously, we reported a new method to identify ligands of the high affinity benzodiazepine pocket, experimental-guided virtual screening (EGVS), integrating experimental data with homology modeling of the

¹Institute of Biochemistry and Molecular Medicine, University of Bern, CH-3012 Bern, Switzerland. ²Department of Anesthesia, Critical Care and Pain Medicine, Massachusetts General Hospital, Boston, 02114 Massachusetts, USA. Correspondence and requests for materials should be addressed to E.S. (email: sigel@ibmm.unibe.ch)




Figure 1. Chemical structure of compounds 31, 132, 4-O-methylhonokiol, and the high-affinity benzodiazepine antagonist Ro 15-1788.

GABA_A receptor³⁴. EGVS identified some ligands that only recognized site 1, others that recognized both site 1 and site 3³⁵, and another set that only recognize site 3.

Here we describe the actions of two compounds identified by EGVS, 31 and 132³⁴ and 4-O-methylhonokiol³⁶. Using mutations and concatenated receptors we determined that the three compounds act mainly through the TMD $\beta+$ / $\alpha-$ interfaces (site 3), and particularly the $\gamma\beta+$ / $\alpha-\beta$ site. The anesthetic action of these drugs was explored *in vivo*, revealing potencies similar to propofol.

Results

In an attempt to find novel ligands for the high affinity site for benzodiazepines on GABA_A receptors we screened 198 compounds for displacement of the high affinity benzodiazepine site (called site 1 previously) antagonist [³H]-Ro 15-1788 at receptors expressed in HEK-cells. Many high affinity ligands were identified³⁴. One compound, SJM3 acted as antagonist with high affinity at site 1, but allosterically potentiated receptor activation through sites in the membrane (called sites 3 previously)³⁵. Compounds 31 and 132 either did not or weakly displaced [³H]-Ro 15-1788 from site 1 but potently enhanced GABA_A receptor activation. 4-O-methylhonokiol shares these characteristics, potentiating $\alpha_1\beta_2\gamma_2$ receptors with an EC₅₀ of $5.4 \pm 1.8 \mu\text{M}$, independent of the high affinity site for benzodiazepines³⁶. Here, we report mechanistic and animal studies of these three compounds.

Compounds 31 and 132 are allosteric modulators of $\alpha_1\beta_2\gamma_2$ GABA_A receptors. First, we investigated if compounds 31 and 132 were able to act as agonists. Figure 1 shows their chemical structures compounds. Both compounds at 3 and 30 μM elicited only very small currents by themselves in $\alpha_1\beta_2\gamma_2$ GABA_A receptors expressed in *Xenopus* oocytes. These compounds elicited at the concentration of 3 μM and 30 μM currents amounting to $0.1 \pm 0.06\%$ (mean \pm SD, $n = 3$) and $0.3 \pm 0.13\%$ (mean \pm SD, $n = 3$), respectively of the maximal current amplitude elicited by GABA in the same oocytes. Thus neither of the compounds tested acts as an appreciable agonist on $\alpha_1\beta_2\gamma_2$ receptors.

Both compounds strongly enhanced currents elicited by GABA. We established the concentration response curves of this positive allosteric modulation. After two applications of GABA at a concentration eliciting 0.5–1.5% of the maximal current amplitude, the same concentration of GABA was co-applied with increasing concentrations of the tested compounds. Figure 2a,b show current traces demonstrating positive modulation by different concentrations of compounds 31 and 132, respectively. Figure 2c summarizes the results of three of such experiments. Both compounds potentiate GABA elicited currents in oocytes expressing $\alpha_1\beta_2\gamma_2$ receptors. For compound 31, at high concentrations apparent desensitization was observed, that could be partly due to open channel blocker effect. For both compounds no saturation at the highest concentration was obtained, because of poor solubility we could not test higher concentrations.

Potentiation by compounds 31 and 132 is not affected by Ro 15-1788. From the binding data we did not expect that the two compounds are acting through the high affinity benzodiazepine binding site 1 in $\alpha_1\beta_2\gamma_2$ receptors. Nevertheless, we tested whether 1 μM of the site 1 antagonist Ro 15-1788 inhibits potentiation of GABA currents by compounds 31 or 132. Either compound (3 μM) strongly potentiated currents elicited by GABA. Potentiation by either drug was not inhibited by 1 μM Ro 15-1788 (Fig. 3). Relative current amplitudes in the presence vs. absence of the antagonist were $112 \pm 8\%$ (mean \pm SD, $n = 4$, $p > 0.05$, *t test*) for compound 31, and $115 \pm 14\%$ (mean \pm SD, $n = 4$, $p > 0.05$, *t test*) for compound 132. This confirms that potentiation by compounds 31 and 132 does not result from action at the benzodiazepine site 1.

Compounds 31 and 132 and 4-O-methylhonokiol act at the low affinity benzodiazepine site 3 in $\alpha_1\beta_2\gamma_2$ receptors. We next investigated whether benzodiazepine site 3 TMD mutations affect potentiation by compounds 31 and 132. Combining three homologous site 3 mutations in $\alpha_1\beta_2\gamma_2$ receptors, α_1 S269I, β_2 N265I and γ_2 S280I, eliminated the potentiation by high concentrations of diazepam²⁰. We investigated the effects of these mutations individually and combined, abbreviating them as α_1 M, β_2 M and γ_2 M. Recently, we described the




Figure 2. Compounds 31 and 132 stimulate GABA currents mediated by $\alpha_1\beta_2\gamma_2$ receptors in a concentration-dependent manner. $\alpha_1\beta_2\gamma_2$ receptors were expressed in *Xenopus* oocytes and electrophysiological experiments were performed. Original current traces of an experiment with compound 31 (a) and with compound 132 (b). Numbers indicate applied concentrations of the respective compounds. (c) Concentration dependence of the positive allosteric modulation by compound 31 (closed circles) and compound 132 (closed squares) in oocytes expressing $\alpha_1\beta_2\gamma_2$ receptors. Mean data \pm SD for both compounds is shown, $n = 3$.

potency of GABA to activate all of these receptor subtypes expressed in *Xenopus* oocytes³⁷. In order to exclude general gating effects caused by these mutations we showed that potentiation by low concentrations of diazepam and by THDOC are not affected³⁷. For illustration, the localization of the mutations is shown in the crystalized homomeric β_3 receptor³⁸ where some of the β_3 subunits were renamed α_1 , β_2 and γ_2 (Fig. 4a).

Wild-type $\alpha_1\beta_2\gamma_2$, $\alpha_1\text{M}\beta_2\gamma_2$, $\alpha_1\beta_2\text{M}\gamma_2$, $\alpha_1\beta_2\gamma_2\text{M}$ and $\alpha_1\text{M}\beta_2\text{M}\gamma_2\text{M}$ receptors were expressed in *Xenopus* oocytes. Using electrophysiological techniques we determined the effect of these point mutations on the potentiation by compounds 31, 132 and 4-O-methylhonokiol, normalizing to the potentiation in wild type $\alpha_1\beta_2\gamma_2$ receptors.

Figure 4b summarizes the results obtained. The single mutations in the α_1 and in the γ_2 subunits did not significantly alter the degree of potentiation by 3 μM of either compound 31 or 132. Relative to wild-type $\alpha_1\beta_2\gamma_2$ receptors, modulation in mutated $\alpha_1\text{M}\beta_2\gamma_2$ receptors by compound 31 amounted to $77 \pm 24\%$ (mean \pm SD, $n = 3$, $p > 0.05$, Tukey posthoc test), and by compound 132 to $118 \pm 44\%$ (mean \pm SD, $n = 3$, $p > 0.05$, Tukey posthoc test). Modulation in $\alpha_1\beta_2\gamma_2\text{M}$ receptors by compound 31 was $87 \pm 52\%$ (mean \pm SD, $n = 3$, $p > 0.05$, Tukey posthoc test), and by compound 132 was $91 \pm 21\%$ (mean \pm SD, $n = 3$, $p > 0.05$, Tukey posthoc test). In contrast, modulation by both compounds was strongly impaired in $\alpha_1\beta_2\text{M}\gamma_2$ and triply mutated $\alpha_1\text{M}\beta_2\text{M}\gamma_2\text{M}$ receptors. Potentiation by 3 μM compound 31 in $\alpha_1\beta_2\text{M}\gamma_2$ was $10.4 \pm 8.5\%$ (mean \pm SD, $n = 3$, $p = 0.007$, Tukey posthoc test), and in $\alpha_1\text{M}\beta_2\text{M}\gamma_2\text{M}$ receptors was $2.3 \pm 0.8\%$ (mean \pm SD, $n = 3$, $p = 0.0043$, Tukey posthoc test). Potentiation by compound 132 relative to that in $\alpha_1\beta_2\gamma_2$ was also dramatically reduced in $\alpha_1\beta_2\text{M}\gamma_2$ $7.3 \pm 6.8\%$ (mean \pm SD, $n = 3$, $p = 0.003$, Tukey posthoc test) and $\alpha_1\text{M}\beta_2\text{M}\gamma_2\text{M}$ $-0.4 \pm 5.1\%$ (mean \pm SD, $n = 3$, $p = 0.0021$, Tukey posthoc test), respectively.

Potentiation by 1 μM 4-O-methylhonokiol was also significantly reduced only in $\alpha_1\beta_2\text{M}\gamma_2$ and $\alpha_1\text{M}\beta_2\text{M}\gamma_2\text{M}$ receptors, similar to compounds 31 and 132. Relative to wild-type $\alpha_1\beta_2\gamma_2$ receptors, modulation in $\alpha_1\text{M}\beta_2\gamma_2$ mutated amounted to $63 \pm 6\%$ (mean \pm SD, $n = 3$, $p > 0.05$, Tukey posthoc test), and in $\alpha_1\beta_2\gamma_2\text{M}$ receptors to $79 \pm 16\%$ (mean \pm SD, $n = 3$, $p > 0.05$, Tukey posthoc test). Relative to $\alpha_1\beta_2\gamma_2$, residual potentiation in $\alpha_1\beta_2\text{M}\gamma_2$ receptors amounted to $15.6 \pm 1.5\%$ (mean \pm SD, $n = 3$, $p = 0.0022$, Tukey posthoc test), and to $2.2 \pm 0.2\%$ (mean \pm SD, $n = 3$, $p = 0.0018$, Tukey posthoc test) in $\alpha_1\text{M}\beta_2\text{M}\gamma_2\text{M}$.




Figure 3. Compounds 31 and 132 do not act at the classical high affinity site for benzodiazepines. GABA at a concentration eliciting 0.5% of the maximal current amplitude ($EC_{0.5}$, single bars) was applied until a stable response was obtained. Subsequently, the same concentration of GABA was co-applied with 3 μ M of compounds 31 (a) or 132 (b), which resulted both in a large increase of current amplitude. Co-application of Ro 15-1788 with compound and GABA did not reduce the degree of modulation in both cases. Experiments were repeated 4 times, with three different batches of oocytes, with a similar outcome.




Figure 4. (a) Model structure of the GABA_A receptor transmembrane domain. The major isoform of the GABA_A receptor is composed of two α_1 , two β_2 , and one γ_2 subunits. The model structure depicts the crystalized homomeric β_3 GABA_A receptor (PDB structure 4COF)³⁸. In this figure, some of the β_3 subunits were renamed α_1 (yellow), β_2 (blue) and γ_2 (red); structures are shown in ribbon representation. The mutated residues α_1 S269, β_2 N265, and γ_2 S280 are located at the interfaces between subunits. (b) Potentiation of the GABA response by compound 31 (3 μ M), compound 132 (3 μ M), and 4-O-methylhonokiol (1 μ M, abbreviated Mh) in wild-type $\alpha_1\beta_2\gamma_2$, single mutant (α_1M , β_2M , γ_2M), and triple mutant receptors expressed in *Xenopus* oocytes. The bars indicate mean \pm SD, n = 3.

The above data suggests that the modulatory site(s) for the three compounds we studied is located in one or both of the $\beta_+/ \alpha-$ TMD subunit interfaces on $\alpha_1\beta_2\gamma_2$ GABA_A receptors.




Figure 5. Individual roles of the two β +/ α - interfaces in channel modulation by compounds 31, 132 and 4-O-methylhonokiol. (a) Scheme showing the four concatenated wild-type and mutant receptors. 1 and 2 refer to the two different β +/ α - subunit interfaces, interface 1 and interface 2. The location of the β_2 N265I mutations is indicated in red color. Concatenated receptors were prepared containing no mutation (α_1 - β_2 - α_1 / γ_2 - β_2 , non M), a mutation at interface 1 (α_1 - β_2 - α_1 / γ_2 - β_2 M, interface 1 M), a mutation at interface 2 (α_1 - β_2 M- α_1 / γ_2 - β_2 , interface 2 M), or mutations in both sites (α_1 - β_2 M- α_1 / γ_2 - β_2 M, double M). Interface 2 harbors a binding site for GABA with higher apparent affinity for channel gating than the one positioned at the interface 1⁵⁴. (b) Potentiation of GABA for each concatenated receptor subtype. Bars indicate mean \pm SD, n = 3.

Role of the individual β +/ α - subunit interfaces in channel modulation by compounds 31, 132 and 4-O-methylhonokiol.

Each GABA_A receptor contains two β +/ α - subunit interfaces. Combined mutation at these interfaces greatly reduces the modulatory effects of compounds 31, 132 and 4-O-methylhonokiol. Using α_1 - β_2 - α_1 and γ_2 - β_2 subunit concatemers, we studied the effects of individual mutated interfaces. We designated receptors containing the mutant in the γ_2 - β_2 construct interface 1 M, and the mutation in the α_1 - β_2 - α_1 construct interface 2 M (Fig. 5a). The α_1 - β_2 M- α_1 and γ_2 - β_2 M constructs were built, and were co-expressed with non-mutated dual or triple subunit constructs forming α_1 - β_2 - α_1 / γ_2 - β_2 M and α_1 - β_2 M- α_1 / γ_2 - β_2 receptors. Both constructs were expressed together to form the double mutant receptor α_1 - β_2 M- α_1 / γ_2 - β_2 M.

Wild type concatenated receptors α_1 - β_2 - α_1 / γ_2 - β_2 were also expressed. This receptor has an EC₅₀ for GABA of approximately 120 μ M¹². Results are standardized to the potentiation observed in α_1 - β_2 - α_1 / γ_2 - β_2 concatenated receptors. As shown in Fig. 5b, in the double mutant concatemeric receptors (α_1 - β_2 M- α_1 / γ_2 - β_2 M) potentiation for all three compounds was abolished. Relative to α_1 - β_2 - α_1 / γ_2 - β_2 concatenated receptors, in the double mutant α_1 - β_2 M- α_1 / γ_2 - β_2 M receptor modulation by compound 31 was reduced to $1 \pm 5\%$ (mean \pm SD, n = 3, p < 0.0005, Tukey posthoc test), by compound 132 to $-2 \pm 0.4\%$ (mean \pm SD, n = 3, p = 0.0001, Tukey posthoc test), and by 4-O-methylhonokiol to $1 \pm 0.6\%$ (mean \pm SD, n = 3, p = 0.0001, Tukey posthoc test). In receptors containing only one mutation, either interface 1 or interface 2, modulation by compound 31 was reduced significantly compared to concatenated wild type receptors α_1 - β_2 - α_1 / γ_2 - β_2 . With interface 1 M, residual relative potentiation was $28 \pm 6\%$ (mean \pm SD, n = 3, p = 0.0005, Tukey posthoc test) and with interface 2 M it was $51 \pm 3\%$ (mean \pm SD, n = 3, p = 0.0059, Tukey posthoc test) of wild-type. Potentiation of interface 1 M and interface 2 M receptors differed significantly (p = 0.0035, t test). Therefore, both TMD β +/ α - sites seem to contribute differently to modulation by compound 31, interface 1 being more efficacious.

Likewise, modulation by compound 132 was sensitive to the mutations at both sites. In interface 1 M receptors relative potentiation was reduced to $8 \pm 4\%$ (mean \pm SD, n = 3, p = 0.0002, Tukey posthoc test), and in interface 2 M receptors to $39 \pm 19\%$ (mean \pm SD, n = 3, p = 0.0035, Tukey posthoc test). Again, interface 1 M produced a larger impact than 2 M, although the difference was at the statistical limit (p = 0.0510, t test). Modulation by 4-O-methylhonokiol in interface 1 M and 2 M receptors was reduced to $11 \pm 6\%$ (mean \pm SD, n = 3, p < 0.0001, Tukey posthoc test) and $28 \pm 8\%$ (mean \pm SD, n = 3, p = 0.0004, Tukey posthoc test), respectively. The two mutations produced significantly different effects (p = 0.043, t test), with the interface 1 M effect again larger.

Therefore, potentiation by all three compounds displayed similar sensitivity patterns with homologous mutations in distinct β +/ α - TMD sites of α_1 - β_2 - α_1 / γ_2 - β_2 receptors. Both sites are necessary for full modulation, while interface 1 produces a larger impact than interface 2.

Effect of the β_2 N265S mutation. The β_2 N265 residue is important for allosteric modulation of GABA_A receptors by many compounds acting through the TMD. This residue was initially described as a determinant for the modulatory action of loreclezole, where the β_2 N265S mutation created a receptor unresponsive to this compound³⁹. Mutations in this residue also abolish potentiation by the anesthetics etomidate and propofol^{28,40}.




Figure 6. Effect of the β_2N265S mutation on modulation by compounds 31 and 132. Wild-type $\alpha_1\beta_2\gamma_2$ and mutated $\alpha_1\beta_2N265S\gamma_2$ receptors were expressed in *Xenopus* oocytes and studied. Potentiation of GABA currents was determined using $3\ \mu\text{M}$ of compound 31 or 132. Bars indicate mean \pm SD, $n = 4-11$.




Figure 7. Subunit specificity of current potentiation in different GABA_A receptors. Different subunit combinations were expressed in *Xenopus* oocytes. Potentiation of GABA currents at a GABA concentration eliciting 0.5–1.5% of the maximal current amplitude was determined using $3\ \mu\text{M}$ of compound 31 (a) or 132 (b). Bars indicate mean \pm SD, $n = 4-11$.

As shown in Fig. 6, the β_2N265S mutation in $\alpha_1\beta_2\gamma_2$ receptors significantly reduced potentiation by $3\ \mu\text{M}$ compound 31, from $485 \pm 230\%$ in wild-type receptors (mean \pm SD, $n = 11$), to $125 \pm 25\%$ in the mutated receptor (mean \pm SD, $n = 4$, $p = 0.009$, *Tukey posthoc test*). In contrast, this mutation did not significantly reduce potentiation by $3\ \mu\text{M}$ compound 132: $415 \pm 126\%$ in wild-type receptors (mean \pm SD $n = 6$) versus $280 \pm 36\%$ in the mutated receptor (mean \pm SD, $n = 4$, $p > 0.05$, *Tukey posthoc test*). We have shown earlier that potentiation by 4-O-methylhonokiol in receptors carrying the β_2N265S mutation was greatly reduced to about 40%³⁶.

Subunit specificity of compounds 31 and 132. From the above experiments we inferred that the β -/ α -TMD subunit interfaces mediate potentiation of compounds 31 and 132. We also wanted to know if potentiation by these compounds depends on subunit isoforms. First we replaced the α_1 subunit by different α subunit isoforms: $\alpha_1\beta_2\gamma_2$, $\alpha_2\beta_2\gamma_2$, $\alpha_3\beta_2\gamma_2$, $\alpha_4\beta_2\gamma_2$, $\alpha_5\beta_2\gamma_2$ and $\alpha_6\beta_2\gamma_2$ (Fig. 7). Compound 31 displayed a similar degree of potentiation in $\alpha_1\beta_2\gamma_2$ receptors $485 \pm 230\%$ (mean \pm SD, $n = 11$), $\alpha_2\beta_2\gamma_2$ receptors, $407 \pm 172\%$ (mean \pm SD, $n = 4$, $p > 0.05$, *Tukey posthoc test*), $\alpha_3\beta_2\gamma_2$ receptors, $735 \pm 234\%$, (mean \pm SD, $n = 5$, $p > 0.05$, *Tukey posthoc test*),

a	<u>M1</u>	b	<u>M3</u>
α_1	KRKIGYFVIQTYLPCI	β_1	IPYVKAIDIYLMGCFV
α_2	KRKIGYFVIQTYLPCI	β_2	IPYVKAIDMYLMGCFV
α_3	KRKIGYFVIQTYLPCI	β_3	IPYVKAIDMYLMGCFV
α_4	RRKMGYFMIQTYIPCI		
α_5	KRKIGYFVIQTYLPCI		
α_6	QRKMGYFMIQIYTPCI		

Figure 8. Alignment of the rat amino acid residue sequences of different α (a) and β (b) subunit isoforms. (a) Sequences preceding M1 and the first part of M1 in α subunits are shown. (b) Sequences preceding M3 and the first part of M3 in β subunits are shown.

and $\alpha_4\beta_2\gamma_2$ receptors, $412 \pm 175\%$ (mean \pm SD, $n = 4$, $p > 0.05$, *Tukey posthoc test*). The $\alpha_5\beta_2\gamma_2$ receptor showed a significant decrease in potentiation compared to $\alpha_1\beta_2\gamma_2$ receptors to $253 \pm 92\%$ (mean \pm SD, $n = 8$; $p = 0.023$, *Tukey posthoc test*), and the $\alpha_6\beta_2\gamma_2$ receptor an increase in potentiation, amounting to $780 \pm 236\%$ (mean \pm SD, $n = 4$; $p = 0.048$, *Tukey posthoc test*). This discrepancy in modulation of $\alpha_5\beta_2\gamma_2$ and $\alpha_6\beta_2\gamma_2$ receptors may be explained by the fact that 3 of the first 7 residues of M1 located at the minus side of the α subunit are different (Fig. 8a). Compound 132 produced similar potentiation in receptors with all α subunits tested. Amounting to $415 \pm 126\%$ (mean \pm SD, $n = 6$) in $\alpha_1\beta_2\gamma_2$ receptors, $694 \pm 293\%$ (mean \pm SD, $n = 4$, $p > 0.05$, *Tukey posthoc test*) in $\alpha_2\beta_2\gamma_2$ receptors, $652 \pm 215\%$ (mean \pm SD, $n = 4$, $p > 0.05$, *Tukey posthoc test*) in $\alpha_3\beta_2\gamma_2$ receptors, $476 \pm 152\%$ (mean \pm SD, $n = 4$, $p > 0.05$, *Tukey posthoc test*) in $\alpha_4\beta_2\gamma_2$ receptors, $399 \pm 147\%$ (mean \pm SD, $n = 4$, $p > 0.05$, *Tukey posthoc test*) in $\alpha_5\beta_2\gamma_2$ receptors, and $542 \pm 175\%$ (mean \pm SD, $n = 4$, $p > 0.05$, *Tukey posthoc test*) in $\alpha_2\beta_2\gamma_2$ receptor type. These results indicate that although the type of α subunit has differential effects on potentiation between compounds 31 and 132, these compounds modulate all receptor subtypes studied.

Next, we examined the role of the β subunit, replacing the β_2 by β_1 or β_3 . For compounds 31 and 132, $\alpha_1\beta_3\gamma_2$ receptors showed a similar potentiation as $\alpha_1\beta_2\gamma_2$ receptors. Amounting to $381 \pm 115\%$ for compound 31 and $555 \pm 169\%$ for compound 132 (mean \pm SD, $n = 4$, $p > 0.05$, *Tukey posthoc test*). In the case of $\alpha_1\beta_1\gamma_2$ receptors, potentiation by both compounds was significantly reduced compared to that in $\alpha_1\beta_2\gamma_2$, $41 \pm 13\%$ for compound 31 and $37 \pm 7\%$ for compound 132 ($n = 4$; $p = 0.0023$, *Tukey posthoc test* for compound 31; $n = 4$; $p = 0.0004$, *Tukey posthoc test*, for compound 132). These results indicate that the type of β subunit is important for the potentiation by both compounds. It is interesting to note in this context that β_1 and β_2/β_3 differ not only in the residue 265, but also in the fourth residue of M3 predicted to be close to the latter residue (Fig. 8b).

When the γ_2 subunit was omitted, no statistical difference was observed for either compound tested. Compound 31 displayed a similar degree of potentiation between $\alpha_1\beta_2\gamma_2$ receptors and $\alpha_1\beta_2$ receptors, $339 \pm 73\%$ (mean \pm SD, $n = 4$, $p > 0.05$, *Tukey posthoc test*). Compound 132 also showed a similar potentiation between both receptors, potentiation in $\alpha_1\beta_2$ receptors amounting to $625 \pm 219\%$ (mean \pm SD, $n = 4$, $p > 0.05$, *Tukey posthoc test*). When the γ_2 subunit was replaced by a δ subunit, in the case of $\alpha_1\beta_2\delta$ receptors, potentiation was affected only in the case of compound 31, where a significant reduction was observed relative to $\alpha_1\beta_2\gamma_2$ receptors. Where potentiation amounted to $168 \pm 48\%$ (mean \pm SD, $n = 4$, $p = 0.0026$, *Tukey posthoc test*) for compound 31, and $640 \pm 275\%$ (mean \pm SD, $n = 4$, $p > 0.05$, *Tukey posthoc test*) for compound 132. In $\alpha_4\beta_3\delta$ receptors, potentiation by both compounds was statistically reduced relative to $\alpha_4\beta_3\gamma_2$ receptors. For compound 31 potentiation amounted to $345 \pm 77\%$ (mean \pm SD, $n = 4$) in $\alpha_4\beta_3\gamma_2$ receptors, and was reduced to $125 \pm 22\%$ (mean \pm SD, $n = 4$, $p = 0.0015$, *Tukey posthoc test*) in $\alpha_4\beta_3\delta$ receptors. Potentiation of compound 132 in $\alpha_4\beta_3\gamma_2$ receptors was $728 \pm 203\%$ (mean \pm SD, $n = 4$), and was reduced to $214 \pm 71\%$ (mean \pm SD, $n = 4$, $p = 0.0031$ *Tukey posthoc test*, for compound 132) in $\alpha_4\beta_3\delta$ receptors.

Previous work³⁶ showed that potentiation by 4-O-methylhonokiol was dependent on the α subunit in a similar fashion as compound 31, since modulation was reduced by receptors containing α_5 and α_6 subunits. The type of β subunit was also important, as the presence of the β_1 subunit strongly reduced potentiation by this compound. On the contrary, the presence of a γ or a δ subunit did not affect potentiation.

Anesthetic activity in tadpoles. The anesthetic activity for compounds 31, 132 and 4-O-methylhonokiol was determined as loss of righting reflex (LoRR) in *Xenopus* tadpoles. Fig. 9 shows the concentration dependence curve for each. Compound 31 yielded an EC_{50} of $2.7 \mu\text{M}$ (95% confidence interval = 2.0 to $3.7 \mu\text{M}$), while the EC_{50} compound 132 was $1.2 \mu\text{M}$ (95% confidence interval = 0.73 to $2.0 \mu\text{M}$), and 4-O-methylhonokiol $EC_{50} = 1.0 \mu\text{M}$ (95% confidence interval = 0.46 to $2.2 \mu\text{M}$). For comparison, the EC_{50} for propofol-induced LoRR in tadpoles is $1.3 \mu\text{M}$ ⁴¹. Anesthesia was fully reversible for compound 31; for animals tested with compound 132, recovery was minimal at concentrations above $3 \mu\text{M}$. For 4-O-methylhonokiol, animals tested at a concentration of $10 \mu\text{M}$ did not recover, whereas recovery was complete at $3 \mu\text{M}$ and lower concentrations.




Figure 9. Concentration-response curves for loss of righting reflexes (LORR) in tadpoles for compounds 31 (closed circle), 132 (closed square), and 4-O-methylhonokiol (closed triangle). The percent of animals anesthetized is plotted against aqueous anesthetic concentration, overlaid with logistic fits. Each point represents data from ten animals. Data were fitted to a Hill equation.

Discussion

Here we functionally characterized compounds 31, 132 and further investigated the properties of 4-O-methylhonokiol. All three compounds are potent allosteric potentiators of $\alpha_1\beta_2\gamma_2$ GABA_A receptors that do not act at site 1. We suspected that they instead act through the low affinity TMD site(s) for benzodiazepines (site 3). Indeed, potentiation by compounds 31, 132 and 4-O-methylhonokiol was abolished in the triple mutant receptor $\alpha_1S269I\beta_2N265I\gamma_2S280I$ as well as by the single mutation β_2N265I , but unaffected by the homologous mutations α_1S269I and γ_2S280I . Assuming that these TM2 mutations alter drug actions through local steric effects in adjacent TMD interfacial sites, our results indicate that of the five such sites, only the two $\beta+/α-$ interfaces 1 and 2 mediate the potentiating effects of these three compounds.

We further dissected the contribution of the individual $\beta+/α-$ subunit interfaces using concatenated subunit assemblies. The $\gamma\beta+/α-\beta$ interface (interface 1) and $\alpha\beta+/α-\gamma$ interface (2) participated differently in modulation by the three compounds studied. For all compounds the contribution of the interface 1 to drug modulation is apparently greater than that of the interface 2.

We and others have shown that the intravenous anesthetics etomidate, propofol and pentobarbital also act via TMD interfacial sites^{26–29,37}. In $\alpha_1\beta_2\gamma_2$ receptors, β_2N265I reduced potentiation by all compounds, α_1S269I reduced potentiation exclusively by pentobarbital, and the γ_2S280I mutation increased potentiation by etomidate, while reducing potentiation by propofol and pentobarbital³⁷. Different sets of residues located at subunit interfaces have been photo-labeled by etomidate, barbiturate, and propofol analogs, revealing that some anesthetics selectively bind within different TMD interfaces^{23,24,33,42,43}. Additionally, mutations of residues at the $\beta+/α-$ subunit interface affecting anesthetic action have been shown to affect modulation by valerianic acid. This suggests that the binding pocket for this compound is also at or near the anesthetics binding site⁴⁴. Other subunit interfaces were not investigated.

Work by our group using receptor concatenation determined that both $\beta+/α-$ subunit interfaces participated equally in modulation by propofol. In contrast, modulation by etomidate was found to be more affected by the $\gamma\beta+/α-\beta$ interface site (interface 1) than the $\alpha\beta+/α-\gamma$ site (interface 2)³⁷. Interestingly, studies using another mutation (β_2M286W) and different concatenated subunit assemblies suggest that etomidate interactions are equivalent in the two $\beta+/α-$ sites of $\alpha_1\beta_2\gamma_2$ receptors⁴⁵.

The homologs of β_2N265 in β_1 and β_3 are serine and asparagine, respectively, and this single residue dramatically influences sensitivity to loreclezole³⁹, etomidate and propofol^{26,28,29,39,40}. Potentiation by compound 31 was strongly affected by this mutation while that by compound 132 was affected less. On the other hand, potentiation by 132 was severely reduced in $\alpha_1\beta_1\gamma_2$ receptors, compared to $\alpha_1\beta_2\gamma_2$ receptors. Potentiation by 4-O-methylhonokiol is also reduced by the β_2N265S mutation or substitution of β_1 for β_2 ³⁶.

In subunit specificity studies, compounds 31 and 132 potentiated receptors containing the α_4 and α_6 subunits, contrasting with benzodiazepine site 1 agonists. Both compounds also potentiated receptors carrying the δ subunit, although to different degrees. Thus, these compounds not only acted at receptors shown to be located synaptically as $\alpha_1\beta_{2-3}\gamma_2$, $\alpha_2\beta_2\gamma_2$ and $\alpha_3\beta_2\gamma_2$ ⁴⁶, but also at $\alpha_5\beta_2\gamma_2$ receptors and receptors containing the δ subunit, which are all located extra-synaptically^{5,46,47}. Similarly, SJM-3 modulates both synaptic and extrasynaptic receptors³⁵.

The similarities between the three compounds we studied and the clinical anesthetics propofol, etomidate and pentobarbital suggested their possible use as sedative-hypnotics in animals. Indeed, all three compounds induced reversible loss of righting reflexes (LoRR) in *Xenopus laevis* tadpoles with EC₅₀s comparable to the anesthetics propofol⁴¹, and etomidate⁴⁸. However, LoRR was not reversible with high concentrations of compound 132 and 4-O-methylhonokiol.

In summary, the newly identified compounds 31 and 132 modulate both synaptic and extrasynaptic GABA_A receptors at molecular sites different from the classical benzodiazepine pocket. These compounds, together with 4-O-methylhonokiol, act through $\beta+/α-$ TMD interfaces, with strongest effects through the interface 1. These

compounds potently produced LoRR in aquatic animals and thus may be useful lead compounds in the search for novel anesthetic, sedative-hypnotic or anxiolytic drugs.

Methods

Construction of mutated receptor subunits. The point mutations α_1 S269I, β_2 N265I, β_2 N265S and γ_2 S280I were prepared using the QuikChange™ mutagenesis kit (Stratagene, Agilent Technologies, Basel, Switzerland).

Construction of concatenated subunits. Construction of tandem and triple subunit cDNAs. The tandem construct γ_2 - β_2 , and triple construct α_1 - β_2 - α_1 has been described previously¹². Site-directed mutagenesis of β_2 N265 to I was done in the tandem construct and the triple construct using the QuikChange™ mutagenesis kit (Stratagene, Agilent Technologies, Basel, Switzerland).

Expression of GABA_A receptors in *Xenopus oocytes*. Capped cRNAs were synthesized (Ambion, Austin, TX, USA) from the linearized plasmids with a cytomegalovirus promoter (pCMV vectors) containing the different subunits, respectively. A poly-A tail of about 400 residues was added to each transcript using yeast poly-A polymerase (United States Biologicals, Cleveland, OH, USA). The concentration of the cRNA was quantified on a formaldehyde gel using Radiant Red stain (Bio-Rad) for visualization of the RNA. Known concentrations of RNA ladder (Invitrogen) were loaded as standard on the same gel. cRNAs were precipitated in ethanol/isoamylalcohol 19:1, the dried pellet dissolved in water and stored at -80°C . cRNA mixtures were prepared from these stock solutions and stored at -80°C .

Animal experiments were carried out in strict accordance to the Swiss ethical guidelines, and have been approved by the local committee of the Canton Bern Kantonstierarzt, Kantonaler Veterinärmedizinischer Dienst Bern (BE85/15). Surgery was done under anesthesia, and all efforts were made to diminish animal suffering. *Xenopus laevis* oocytes were prepared, injected and defolliculated as described previously^{49,50}. Oocytes were injected with 50 nL of the cRNA solution containing wild type or mutated rat α_1 , β_2 and γ_2 subunits of the GABA_A receptors at a concentration of 10 nM:10 nM:50 nM⁵¹. For concatenated tandem and triple constructs, cRNA combinations ratios of 25:25 nM were used. Injected oocytes were incubated in modified Barth's solution at 18°C for at least 24 h before the measurements.

Functional characterization of GABA_A receptors. Currents were measured using a modified two-electrode voltage clamp amplifier Oocyte clamp OC-725 (Warner Instruments) in combination with a XY-recorder (90% response time 0.1 s) or digitized at 100 Hz using a PowerLab 2/20 (AD Instruments) using the computer programs Chart (ADInstruments GmbH, Spechbach, Germany). Tests with a model oocyte were performed to ensure linearity in the larger current range. The response was linear up to $15\ \mu\text{A}$. The holding potential was $-80\ \text{mV}$. The perfusion medium contained 90 mM NaCl, 1 mM KCl, 1 mM MgCl₂, 1 mM CaCl₂, and 5 mM Na-HEPES (pH 7.4). Concentration response curves for the compounds were fitted with the equation $I(c) = I_{\text{max}}/[1 + (\text{EC}_{50}/c)^n]$, where c is the concentration of the compound, EC_{50} the concentration eliciting half-maximal current amplitude, I_{max} is the maximal current amplitude, I the current amplitude, and n is the Hill coefficient. Maximal current amplitudes (I_{max}) were obtained from the fits of the concentration-response curves. For all receptors studied, modulation was measured at a GABA concentration eliciting 0.5–1.5% of the maximal GABA current amplitude. GABA was applied twice alone for 20–60 s, and then in combination with the different compounds for 45 s or 1 min. The duration of washout periods was 4 min in between agonist or agonist/drug applications to prevent receptor desensitization. At the beginning of the experiments, GABA applications were repeated when the elicited current amplitude altered by $>5\%$. Potentiation was calculated by the following equation: $(I_{\text{Modulator + GABA}}/I_{\text{GABA}} - 1) * 100\%$. The perfusion solution was applied through a glass capillary with an inner diameter of 1.35 mm, the mouth of which was placed about 0.4 mm from the surface of the oocyte. This allowed fast changes in agonist concentration around the oocyte. The rate of change was estimated 70% in less than $0.5\ \text{s}$ ¹⁴. The perfusion system was cleaned between drug applications by washing with DMSO to avoid contamination. All media contained a final concentration of 0.5% DMSO (v/v) to ensure drug solubility.

All data are from at least two different batches of oocytes. Data represent mean \pm SD. An unpaired t test was used to compare two means. One-way analysis of variance (ANOVA) was used for multiple comparisons followed by a Tukey post hoc test. * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$.

Loss of righting reflex assay in *Xenopus tadpole*. Animals were used and experiments were carried out with approval and according to the guidelines of the MGH Institutional Animal Care and Use Committee. General anesthetic potency was assessed in *Xenopus laevis* tadpoles as previously described^{41,52,53}. In brief, groups of 10 tadpoles were placed in aqueous solutions containing compound 31, 132, or 4-O-methylhonokiol, and tested every five minutes for loss of righting reflexes (LoRR). Each animal was assigned a score of either awake or LoRR, and the percent of animals anesthetized was plotted against the concentration of the compound tested. Concentration-response data was fitted by non-linear least squares to logistic functions of the form $Y = 1/(1 + 10^{(\text{LogEC}_{50} - \text{Log}[\text{Drug}]) * \text{HillSlope}})$ using Graphpad Prism 6. Results are reported as EC_{50} s and 95% confidence intervals.

References

- Forman, S. A. Molecular approaches to improving general anesthetics. *Anesthesiol. Clin.* **28**, 761–771 (2010).
- Chitilian, H. V., Eckenhoff, R. G. & Raines, D. E. Anesthetic drug development: novel drugs and new approaches. *Surg. Neurol. Int.* **4**, S2–S10 (2013).
- Macdonald, R. L. & Olsen, R. W. GABA_A receptor channels. *Annu. Rev. Neurosci.* **17**, 569–602 (1994).
- Barnard, E. A. *et al.* International union of pharmacology. XV. Subtypes of gamma-aminobutyric acid A receptors: classification on the basis of subunit structure and receptor function. *Pharmacol. Rev.* **50**, 291–313 (1998).

5. Olsen, R. W. & Sieghart, W. International union of pharmacology. LXX. Subtypes of γ -aminobutyric acid type A receptors: classification on the basis of subunit composition, pharmacology, and function. Update. *Pharmacol. Rev.* **60**, 243–260 (2008).
6. Sigel, E. & Steinmann, M. E. Structure, function, and modulation of GABA_A receptors. *J. Biol. Chem.* **287**, 40224–40231 (2012).
7. Rabow L. E., Russek, S. J. & Farb, D. H. From ion currents to genomic analysis: recent advances in GABA_A receptor research. *Synapse* **21**, 189–274 (1995).
8. Chang, Y., Wang, R., Barot, S. & Weiss, D. S. Stoichiometry of a recombinant GABA_A receptor. *J. Neurosci.* **16**, 5415–5424 (1996).
9. Farrar, S. J., Whiting, P. J., Bonnert, T. P. & McKernan, R. M. Stoichiometry of a ligand-gated ion channel determined by fluorescence energy transfer. *J. Biol. Chem.* **274**, 10100–10104 (1999).
10. Tretter, V., Ehya, N., Fuchs & Sieghart, K. W. Stoichiometry and assembly of a recombinant GABA_A receptor subtype. *J. Neurosci.* **17**, 2728–2737 (1997).
11. Baumann, S. W., Baur, R. & Sigel, E. Subunit arrangement of gamma aminobutyric acid type A receptors. *J. Biol. Chem.* **276**, 36275–36280 (2001).
12. Baumann, S. W., Baur, R. & Sigel, E. Forced subunit assembly in $\alpha_1\beta_2\gamma_2$ GABA_A receptors. Insight into the absolute arrangement. *J. Biol. Chem.* **277**, 46020–46025 (2002).
13. Baur, R., Minier, F. & Sigel, E. A GABA_A receptor of defined subunit composition and positioning: concatenation of five subunits. *FEBS Lett.* **580**, 1616–1620 (2006).
14. Sigel, E., Baur, R., Trube, G., Mohler, H. & Malherbe, P. The effect of subunit combination of rat brain GABA_A receptors on channel function. *Neuron.* **5**, 703–711 (1990).
15. Minier, F. & Sigel, E. Positioning of the α -subunit isoforms confers a functional signature to γ -aminobutyric acid type A receptors. *Proc. Natl. Acad. Sci. USA* **101**, 7769–7774 (2004).
16. Sigel, E. & Lüscher, B. P. A closer look at the high affinity benzodiazepine binding site on GABA_A receptors. *Curr. Top. Med. Chem.* **11**, 241–246 (2011).
17. Smith, G. B. & Olsen, R. W. Functional domains of GABA_A receptors. *Trends Pharmacol. Sci.* **16**, 162–168 (1995).
18. Baur, R. *et al.* Covalent modification of GABA_A receptor isoforms by a diazepam analogue provides evidence for a novel benzodiazepine binding site that prevents modulation by these drugs. *J. Neurochem.* **106**, 2353–2363 (2008).
19. Ramerstorfer, J. R. *et al.* The GABA_A receptor α +/ β - interface: a novel target for subtype selective drugs. *J. Neurosci.* **31**, 870–877 (2011).
20. Walters, R. J., Hadley, S. H., Morris, K. D. & Amin, J. Benzodiazepines act on GABA_A receptors via two distinct and separable mechanisms. *Nat. Neurosci.* **12**, 1274–1281 (2000).
21. Nishikawa, K., Jenkins, A., Paraskevakis, I. & Harrison, N. L. Volatile anesthetic actions on the GABA_A receptors: contrasting effects of α_1 (S270) and β_2 (N265) point mutations. *Neuropharmacol.* **42**, 337–345 (2002).
22. Chang, C. S., Olcese, R. & Olsen, R. W. A single M1 residue in the β_2 subunit alters channel gating of GABA_A receptor in anesthetic modulation and direct activation. *J. Biol. Chem.* **278**, 42821–42828 (2003).
23. Li, G. D. *et al.* Identification of a GABA_A receptor anesthetic binding site at subunit interfaces by photolabeling with an etomidate analog. *J. Neurosci.* **26**, 11599–11605 (2006).
24. Yip, G. M. *et al.* A propofol binding site on mammalian GABA_A receptors identified by photolabeling. *Nat. Chem. Biol.* **9**, 715–720 (2013).
25. Rudolph, U. & Antkowiak, B. Molecular and neuronal substrates for general anaesthetics. *Nat. Rev. Neurosci.* **5**, 709–720 (2004).
26. Belelli, D., Lambert, J. J., Peters, J. A., Wafford, K. & Whiting, P. J. The interaction of the general anesthetic etomidate with the gamma-aminobutyric acid type A receptor is influenced by a single amino acid. *Proc. Natl. Acad. Sci. USA* **94**, 11031–11036 (1997).
27. Krasowski, M. D. *et al.* Propofol and other intravenous anesthetics have sites of action on the gamma-aminobutyric acid type A receptor distinct from that for isoflurane. *Mol. Pharmacol.* **53**, 530–538 (1998).
28. Sieghart, R., Krahenbuhl, K., Lambert, S. & Rudolph, U. Mutational analysis of molecular requirements for the actions of general anaesthetics at the γ -aminobutyric acid type A receptor subtype, $\alpha_1\beta_2\gamma_2$. *BMC Pharmacol.* **3**, 13 (2003).
29. Stewart, D. S., Pierce, D. W., Hotta, M., Stern, A. T. & Forman, S. A. Mutations at β N265 in γ -aminobutyric acid type A receptors alter both binding affinity and efficacy of potent anesthetics. *PLoS One* **9**, e111470 (2014).
30. Jurd, R. *et al.* General anesthetic actions *in vivo* strongly attenuated by a point mutation in the GABA(A) receptor β_3 subunit. *FASEB J.* **2**, 250–252 (2002).
31. Hill-Venning, C., Belelli, D., Peters, J. A. & Lambert, J. J. Subunit-dependent interaction of the general anaesthetic etomidate with the gamma-aminobutyric acid type A receptor. *Br. J. Pharmacol.* **120**, 749–756 (1997).
32. Franks, N. P. General anaesthesia: from molecular targets to neuronal pathways of sleep and arousal. *Nat. Rev. Neurosci.* **5**, 370–386 (2008).
33. Jayakar, S. S. *et al.* Multiple propofol-binding sites in a γ -aminobutyric acid type A receptor (GABA_AR) identified using a photoreactive propofol analog. *J. Biol. Chem.* **289**, 27456–27468 (2014).
34. Middendorp, S. J., Puthenkalam, R., Baur, R., Ernst, M. & Sigel, E. Accelerated discovery of novel benzodiazepine ligands by experiment-guided virtual screening. *ACS Chem. Biol.* **9**, 1854–1859 (2014).
35. Middendorp, S. J., Maldifassi, M. C., Baur, R. & Sigel, E. Positive modulation of synaptic and extrasynaptic GABA_A receptors by an antagonist of the high affinity benzodiazepine binding site. *Neuropharmacol.* **95**, 459–467 (2015).
36. Baur, R., Schuehly, W. & Sigel, E. Moderate concentrations of 4-O-methylhonokiol potentiate GABA_A receptor currents stronger than honokiol. *Biochim. Biophys. Acta.* **1840**, 3017–3021 (2014).
37. Maldifassi, M. C., Baur, R. & Sigel, E. Functional sites involved in modulation of the GABA_A receptor channel by the intravenous anesthetics propofol, etomidate and pentobarbital. *Neuropharmacol.* **105**, 207–214 (2016).
38. Miller, P. S. & Aricescu, A. R. Crystal structure of a human GABA_A receptor. *Nature* **512**, 270–275 (2014).
39. Wingrove, P. B., Wafford, K. A., Bain, C. & Whiting, P. J. The modulatory action of loreclezole at the γ -aminobutyric acid type A receptors is determined by a single amino acid in the β_2 and β_3 subunit. *Proc. Natl. Acad. Sci. USA* **91**, 4569–4573 (1994).
40. Fernandez, S. P. *et al.* Flavan-3-ol esters: new agents for exploring modulatory sites on GABA_A receptors. *Br. J. Pharmacol.* **165**, 965–977 (2012).
41. Stein, M. *et al.* Azo-propofols: photochromic potentiators of GABA_A receptors. *Angew. Chem. Int. Ed. Engl.* **51**, 10500–10504 (2012).
42. Chiara, D. C. *et al.* Mapping general anesthetic binding site(s) in human $\alpha_1\beta_3\gamma$ -aminobutyric acid type A receptors with [³H]TDBzl-etomidate, a photoreactive etomidate analogue. *Biochemistry* **51**, 836–847 (2012).
43. Chiara, D. C. *et al.* Specificity of intersubunit general anesthetic binding sites in the transmembrane domain of the human $\alpha_1\beta_3\gamma_2$ GABA_A receptor. *J. Biol. Chem.* **288**, 19343–19357 (2013).
44. Luger, D. *et al.* Identification of the putative binding pocket of valerenic acid on GABA_A receptors using docking studies and site-directed mutagenesis. *Br. J. Pharmacol.* **22**, 5403–5413 (2015).
45. Guitchounts, G., Stewart, D. S. & Forman, S. A. The two etomidate sites in $\alpha_1\beta_3\gamma_2$ gamma-aminobutyric acid type A receptors contribute equally and noncooperatively to modulation of channel gating. *Anesthesiology.* **116**, 1235–1244 (2012).
46. Jacob, T. C., Moss, S. J. & Jurd, R. GABA_A receptor trafficking and its role in the dynamic modulation of neuronal inhibition. *Nat. Rev. Neurosci.* **9**, 331–343 (2008).
47. Farrant, M. & Nusser, Z. Variations on an inhibitory theme: Phasic and tonic activation of GABA_A receptors. *Nature Rev. Neurosci.* **6**, 215–229 (2005).

48. Husain, S. S. *et al.* 2-(3-Methyl-3H-diaziren-3-yl)ethyl 1-(1-phenylethyl)-1H-imidazole-5-carboxylate: a derivative of the stereoselective general anesthetic etomidate for photolabeling ligand-gated ion channels. *J. Med. Chem.* **46**, 1257–1265 (2003).
49. Sigel, E. Properties of single sodium channels translated by *Xenopus* oocytes after injection with messenger ribonucleic acid. *J. Physiol. Lond.* **386**, 73–90 (1987).
50. Sigel, E. & Minier, F. The *Xenopus* oocyte: system for the study of functional expression and modulation of proteins. *Mol. Nutr. Food Res.* **49**, 228–234 (2005).
51. Boileau, A. J., Baur, R., Sharkey, L. M., Sigel, E. & Czajkowski, C. The relative amount of cRNA coding for γ_2 subunits affects stimulation by benzodiazepines in GABA_A receptors expressed in *Xenopus* oocytes. *Neuropharmacol.* **43**, 695–700 (2002).
52. Desai, R. *et al.* Contrasting actions of a convulsant barbiturate and its anticonvulsant enantiomer on the $\alpha_1\beta_3\gamma_2$ GABA_A receptor account for their *in vivo* effects. *J. Physiol.* **593**, 4943–4961 (2015).
53. Ge, R. L. *et al.* Pharmacological studies of methoxycarbonyl etomidate's carboxylic acid metabolite. *Anesth. Analg.* **115**, 305–308 (2012).
54. Baumann, S. W., Baur, R. & Sigel, E. Individual properties of the two functional agonist sites in GABA_A receptors. *J. Neurosci.* **23**, 11158–11166 (2003).

Acknowledgements

This work was supported by the Swiss National Science Foundation grant 315230_156929/1. M.C.M. is a recipient of a fellowship (Beca Chile Postdoctorado from CONICYT, Ministerio de Educacion, Chile). Andrea Szabo (Department of Anesthesia Critical Care & Pain Medicine at Massachusetts General Hospital, Boston, MA, USA) helped to perform tadpole loss of righting reflexes studies. 4-O-methylhonokiol was a kind gift by Dr. Wolfgang Schuehly (Karl-Franzens-University, Graz, Austria).

Author Contributions

M.C.M. and R.B. performed electrophysiological experiments. D.P. and A.N. performed LoRR experiments. M.C.M., S.A.F. and E.S. designed the experiments, analysed the data, and wrote the manuscript.

Additional Information

Competing financial interests: The authors declare no competing financial interests.

How to cite this article: Maldifassi, M. C. *et al.* Novel positive allosteric modulators of GABA_A receptors with anesthetic activity. *Sci. Rep.* **6**, 25943; doi: 10.1038/srep25943 (2016).



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>