

Copyright © 2016 Sheppard et al.

This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International license.

1 [Capsular typing method for *Streptococcus agalactiae* using whole](#)

2 [genome sequence data](#)

3 Running title: GBS capsular typing using whole genome sequence data

4 Anna E Sheppard,^{1#} Alison Vaughan,¹ Nicola Jones,¹ Paul Turner,^{2,3} Claudia

5 Turner,^{2,3} Androulla Efstratiou,^{4,5} Darshana Patel,⁴ the Modernising Medical

6 Microbiology (MMM) Informatics Group,¹ A Sarah Walker,¹ James A Berkley,^{2,6}

7 Derrick W Crook,¹ Anna C Seale.^{2,6#}

8 1. Modernising Medical Microbiology Consortium, Nuffield Department of Clinical
9 Medicine, University of Oxford, UK.

10 2. Centre for Tropical Medicine, Nuffield Department of Clinical Medicine, University
11 of Oxford, UK.

12 3. Shoklo Malaria Research Unit, Thailand

13 4. Microbiology Reference Division, Public Health England, London, UK

14 5. Imperial College, London, UK

15 6. KEMRI-Wellcome Trust Research Programme, Kilifi, Kenya.

16

17 #Corresponding authors: Anna E Sheppard (anna.sheppard@ndm.ox.ac.uk) or Anna

18 C Seale (aseale@nhs.net).

19

20 Word count:

21 Abstract 47/50

22 Main text 933/1000

23 [Abstract](#)

24 Group B streptococcus (GBS) capsular serotype is a major determinant of virulence,
25 and affects potential vaccine coverage. Here we report a whole genome sequencing-
26 based method for GBS serotype assignment. This shows high agreement
27 ($\kappa=0.92$) with conventional methods, and increased serotype assignment
28 (100%) to all ten capsular types.

29

30 [Main text](#)

31 *Streptococcus agalactiae*, or Group B Streptococcus (GBS), is an important
32 pathogen in neonates (1-3), with early infection acquired from the maternal genito-
33 urinary tract (4). In addition, GBS is now recognised as an increasingly important
34 pathogen in high-income regions in immunosuppressed and elderly individuals (5, 6).

35 GBS expresses a capsular polysaccharide, which is involved in virulence and
36 immune evasion. Ten different variants, or serotypes, have been described (Ia, Ib, II,
37 III, IV, V, VI, VII, VIII and IX), which differ in their disease-causing ability. Conjugate
38 vaccines targeting the most common disease-causing serotypes are currently in
39 development (7). Establishing vaccine serotype coverage is important, as is
40 surveillance post-introduction to monitor for potential serotype replacement, as has
41 been seen following the introduction of other conjugate vaccines (8).

42 Current methods for GBS serotype allocation rely on latex agglutination assays or
43 PCR (9). Recent advances in whole genome sequencing (WGS) have enabled the
44 development of approaches that can be used in place of traditional microbiological
45 methods, such as strain typing and antibiotic susceptibility profiling (10-12). A major
46 advantage of this approach is that the cost of sequencing can be mitigated by the

47 ability to use the same data to generate multiple outputs. Given the decreasing cost
48 of WGS (13), and the ongoing increase in WGS data generation, we sought to
49 establish and validate a WGS-based method for GBS capsular typing.

50 We developed an algorithm for serotype assignment on the basis of sequence
51 similarity between a given *de novo* assembly and capsular gene sequences of the
52 ten GBS serotypes. For nine serotypes, published sequences were used as
53 references (Table 1), while for serotype IX, only a partial capsular locus sequence
54 has been published (14). A suitable reference for the full capsular locus region was
55 therefore determined by WGS of a serotype IX isolate obtained from the Statens
56 Serum Institute, Denmark.

57 To assign serotype for a given isolate, a BLAST database was generated from the
58 *de novo* assembly and queried with the variable region of the capsular locus
59 sequence for each serotype (*cpsG-cpsK* for serotypes Ia-VII and IX, *cpsR-cpsK* for
60 serotype VIII) using BLASTn with an evalue threshold of 1e-100 and otherwise
61 default parameters. A serotype was considered as correct if it showed $\geq 95\%$
62 sequence identity over $\geq 90\%$ of the sequence length. These thresholds were chosen
63 on the basis of being stringent enough to provide differentiation between the various
64 reference sequences, while maximising serotype allocation on an initial test set of
65 publicly available GBS WGS data, where serotype information was not available (so
66 we had no way of knowing whether the assigned serotypes were in fact correct).

67 This sequence-based method for serotype allocation was validated using WGS on a
68 set of 223 colonising or invasive human isolates from Canada, Latin America,
69 Singapore, UK, USA, and Thailand, for which serotype had previously been
70 determined using conventional latex agglutination assays, with PCR used to confirm

71 weak positives or negatives in a subset (15-17). For two rare serotypes (Serotype
72 VIII and IX), one isolate of each was obtained from the Statens Serum Institute,
73 Denmark. GBS isolates stored at -80°C were sub-cultured on Columbia blood agar
74 for 24-48 hours, followed by DNA extraction from a single colony using a commercial
75 kit (QuickGene, Fujifilm, Tokyo, Japan). High throughput sequencing was
76 undertaken at the Wellcome Trust Centre for Human Genetics (Oxford University,
77 UK) using the Illumina HiSeq2500 platform, generating 150 base paired-end reads.
78 *De novo* assembly was performed using Velvet and VelvetOptimiser (18, 19).
79 Agreement between serotype allocations was tested with the Kappa statistic.

80 High quality sequence data were obtained for all 223 GBS isolates (median read
81 number: 2,975,508, range: 1,798,744-13,073,718; median contig number: 46, range
82 16-106; median assembly length: 2.05 Mb, range: 1.94-2.22 Mb). Each isolate was
83 allocated to a single serotype using the WGS data (Table 2). Three isolates that did
84 not have a capsular type assigned by latex agglutination methods had serotypes Ib,
85 VI and VIII assigned. For all previously serotyped GBS isolates with a known capsule
86 type, the kappa statistic (0.92) indicated very high agreement between WGS-
87 predicted and conventional serotype. There were nine discordant isolates. In each
88 case there was strong support for the sequence-allocated serotype, with >98%
89 sequence identity over 100% of the reference length in all nine cases (Figure 1).
90 Across all isolates, differences in relatedness between the capsular locus sequences
91 of the different serotypes led to characteristic relationships between the allocated
92 (best match) serotype and the second-best match. For example, all isolates
93 assigned as serotype Ia had serotype III as the second-best match. In all cases, the
94 second-best match was substantially poorer than the best match, demonstrating that
95 there was no ambiguity in predicted serotype (Figure 1, Table 3).

96 The nine discordant and three non-typeable isolates were retested by latex
97 agglutination (Table 4) and resequenced using the Illumina MiSeq platform, with
98 sequence processing and WGS-based serotype prediction performed as above. In
99 all cases, resequencing was consistent with the initial WGS classification. For 6/9
100 discordant isolates, the new latex agglutination results matched the WGS-based
101 prediction, suggesting that the initial discordance may have resulted from incorrect
102 latex agglutination typing or sample mislabelling. The other three initially discordant
103 isolates, and the three non-typeable isolates, were all classified as non-typeable on
104 retesting.

105 This WGS-based method for GBS serotyping, validated using 223 isolates typed
106 using conventional methods, was therefore highly accurate. Although WGS may not
107 currently be cost-effective for directly replacing traditional serotyping, costs are likely
108 to further decrease. Furthermore, WGS may already be the cheapest option for
109 combined studies, with possibilities to utilise the resulting data for additional
110 analyses such as multi-locus sequence typing, relatedness to other sequenced
111 isolates, and detailed phylogenetic analysis.

112

113 [Funding](#)

114 This publication presents independent research commissioned by The Wellcome
115 Trust (www.wellcome.ac.uk 093804, 098532] ACS, JAB), the Health Innovation
116 Challenge Fund (grant HICF-T5-358 and WT098615/Z/12/Z), a parallel funding
117 partnership between the Department of Health and the Wellcome Trust (DWC, AES
118 and ASW); the UK Clinical Research Collaboration (a parallel funding partnership
119 between the Medical Research Council [G0800778], Biotechnology and Biological
120 Sciences Research Council and the Wellcome Trust [087646/Z/08/Z]); and the
121 National Institute for Health Research (NIHR) Oxford Biomedical Research Centre.
122 DWC is a NIHR (UK) Senior Investigator. The views expressed in this publication are
123 those of the author(s) and not necessarily those of the funders.

124

125 [Acknowledgements](#)

126 We would like to thank The Wellcome Trust Centre for Human Genetics, University
127 of Oxford where the whole genome sequencing was done, and we thank the library
128 and sequencing teams. We thank the Shoklo Malaria Research Unit, Thailand for
129 providing GBS isolates.

130

131 [Modernising Medical Microbiology \(MMM\) informatics group](#)

132 Jim Davies, Charles Crichton, Milind Acharya, Carlos del Ojo Elias

133

134 [References](#)

- 135 1. **Baker CJ, Barrett FF, Gordon RC, Yow MD.** 1973. Suppurative meningitis due to streptococci
136 of Lancefield group B: a study of 33 infants. *J Pediatr* **82**:724-729.
- 137 2. **Barton LL, Feigin RD, Lins R.** 1973. Group B beta hemolytic streptococcal meningitis in
138 infants. *J Pediatr* **82**:719-723.
- 139 3. **Communicable Disease Surveillance Centre London.** 1985. Neonatal meningitis: a review of
140 routine national data 1975-83. *Br Med J (Clin Res Ed)* **290**:778-779.
- 141 4. **Dillon HC, Jr., Gray E, Pass MA, Gray BM.** 1982. Anorectal and vaginal carriage of group B
142 streptococci during pregnancy. *J Infect Dis* **145**:794-799.
- 143 5. **Schuchat A.** 1998. Epidemiology of group B streptococcal disease in the United States:
144 shifting paradigms. *Clin Microbiol Rev* **11**:497-513.
- 145 6. **Phares CR, Lynfield R, Farley MM, Mohle-Boetani J, Harrison LH, Petit S, Craig AS, Schaffner**
146 **W, Zansky SM, Gershman K, Stefonek KR, Albanese BA, Zell ER, Schuchat A, Schrag SJ,**
147 **Active Bacterial Core surveillance/Emerging Infections Program N.** 2008. Epidemiology of
148 invasive group B streptococcal disease in the United States, 1999-2005. *JAMA* **299**:2056-
149 2065.
- 150 7. **Madhi SA, Dangor Z, Heath PT, Schrag S, Izu A, Sobanjo-Ter Meulen A, Dull PM.** 2013.
151 Considerations for a phase-III trial to evaluate a group B Streptococcus polysaccharide-
152 protein conjugate vaccine in pregnant women for the prevention of early- and late-onset
153 invasive disease in young-infants. *Vaccine* **31 Suppl 4**:D52-57.
- 154 8. **Mulholland K, Satzke C.** 2012. Serotype replacement after pneumococcal vaccination.
155 *Lancet* **379**:1387; author reply 1388-1389.
- 156 9. **Imperi M PM, Alfarone G, Baldassarri L, Orefici G, Creti R.** 2010. A multiplex PCR assay for
157 the direct identification of the capsular type (Ia to IX) of *Streptococcus agalactiae*. *J*
158 *Microbiol Methods* **80**:212-214.

- 159 10. **Gordon NC, Price JR, Cole K, Everitt R, Morgan M, Finney J, Kearns AM, Pichon B, Young B,**
160 **Wilson DJ, Llewelyn MJ, Paul J, Peto TE, Crook DW, Walker AS, Golubchik T.** 2014.
161 Prediction of *Staphylococcus aureus* antimicrobial resistance by whole-genome sequencing.
162 *J Clin Microbiol* **52**:1182-1191.
- 163 11. **Stoesser N, Batty EM, Eyre DW, Morgan M, Wyllie DH, Del Ojo Elias C, Johnson JR, Walker**
164 **AS, Peto TE, Crook DW.** 2013. Predicting antimicrobial susceptibilities for *Escherichia coli*
165 and *Klebsiella pneumoniae* isolates using whole genomic sequence data. *J Antimicrob*
166 *Chemother* **68**:2234-2244.
- 167 12. **Walker TM, Kohl TA, Omar SV, Hedge J, Del Ojo Elias C, Bradley P, Iqbal Z, Feuerriegel S,**
168 **Niehaus KE, Wilson DJ, Clifton DA, Kapatai G, Ip CL, Bowden R, Drobniewski FA, Allix-**
169 **Beguec C, Gaudin C, Parkhill J, Diel R, Supply P, Crook DW, Smith EG, Walker AS, Ismail N,**
170 **Niemann S, Peto TE, Modernizing Medical Microbiology Informatics G.** 2015. Whole-
171 genome sequencing for prediction of *Mycobacterium tuberculosis* drug susceptibility and
172 resistance: a retrospective cohort study. *Lancet Infect Dis* doi:10.1016/S1473-
173 3099(15)00062-6.
- 174 13. **Loman NJ, Constantinidou C, Chan JZ, Halachev M, Sergeant M, Penn CW, Robinson ER,**
175 **Pallen MJ.** 2012. High-throughput bacterial genome sequencing: an embarrassment of
176 choice, a world of opportunity. *Nat Rev Microbiol* **10**:599-606.
- 177 14. **Slotved HC, Kong F, Lambertsen L, Sauer S, Gilbert GL.** 2007. Serotype IX, a Proposed New
178 *Streptococcus agalactiae* Serotype. *J Clin Microbiol* **45**:2929-2936.
- 179 15. **Bisharat N, Crook DW, Leigh J, Harding RM, Ward PN, Coffey TJ, Maiden MC, Peto T, Jones**
180 **N.** 2004. Hyperinvasive neonatal group B streptococcus has arisen from a bovine ancestor. *J*
181 *Clin Microbiol* **42**:2161-2167.
- 182 16. **Jones N, Oliver K, Jones Y, Haines A, Crook D.** 2006. Carriage of group B streptococcus in
183 pregnant women from Oxford, UK. *J Clin Pathol* **59**:363-366.

- 184 17. **Davies HD, Jones N, Whittam TS, Elsayed S, Bisharat N, Baker CJ.** 2004. Multilocus sequence
185 typing of serotype III group B streptococcus and correlation with pathogenic potential. *J*
186 *Infect Dis* **189**:1097-1102.
- 187 18. **Gladman S, Seeman T.** 2012. VelvetOptimiser.
188 <http://bioinformatics.net.au/software.velvetoptimiser.shtml>
- 189 19. **Zerbino DR, Birney E.** 2008. Velvet: algorithms for de novo short read assembly using de
190 Bruijn graphs. *Genome Res* **18**:821-829.
- 191 20. **Yamamoto S, Miyake K, Koike Y, Watanabe M, Machida Y, Ohta M, Iijima S.** 1999.
192 Molecular characterization of type-specific capsular polysaccharide biosynthesis genes of
193 *Streptococcus agalactiae* type Ia. *J Bacteriol* **181**:5176-5184.
- 194 21. **Watanabe M, Miyake K, Yanae K, Kataoka Y, Koizumi S, Endo T, Ozaki A, Iijima S.** 2002.
195 Molecular characterization of a novel beta1,3-galactosyltransferase for capsular
196 polysaccharide synthesis by *Streptococcus agalactiae* type Ib. *J Biochem* **131**:183-191.
- 197 22. **Martins ER, Melo-Cristino J, Ramirez M.** 2007. Reevaluating the serotype II capsular locus of
198 *Streptococcus agalactiae*. *J Clin Microbiol* **45**:3384-3386.
- 199 23. **Chaffin DO, Beres SB, Yim HH, Rubens CE.** 2000. The serotype of type Ia and III group B
200 streptococci is determined by the polymerase gene within the polycistronic capsule operon.
201 *J Bacteriol* **182**:4466-4477.
- 202 24. **Cieslewicz MJ, Chaffin D, Glusman G, Kasper D, Madan A, Rodrigues S, Fahey J, Wessels**
203 **MR, Rubens CE.** 2005. Structural and genetic diversity of group B streptococcus capsular
204 polysaccharides. *Infect Immun* **73**:3096-3103.
- 205

206 **Table 1.** Reference sequences used for sequence-based serotype allocation

Serotype	Accession	Region	Reference
la	AB028896.2	6982-11695	Yamamoto et al.(20)
lb	AB050723.1	2264-6880	Watanabe et al.(21)
II	EF990365.1	1915-8221	Martins et al.(22)
III	AF163833.1	6592-11193	Chaffin et al.(23)
IV	AF355776.1	6417-11656	Cieslewicz et al.(24)
V	AF349539.1	6400-12547	Cieslewicz et al.(24)
VI	AF337958.1	6437-10913	Cieslewicz et al.(24)
VII	AY376403.1	3403-8666	Cieslewicz et al.(24)
VIII	AY375363.1	2971-7340	Cieslewicz et al.(24)
IX	NA	NA	This study

207

208 **Table 2.** Serotype allocation by WGS to serotype allocation by latex agglutination

		Serotype allocated by WGS										
		Ia	Ib	II	III	IV	V	VI	VII	VIII	IX	Total
Serotype by latex agglutination	Ia	34	0	0	1	0	0	0	0	0	0	35
	Ib	0	9	1	0	0	0	0	0	0	0	10
	II	0	0	25	0	0	0	0	0	0	0	25
	III	3	0	0	111	0	0	0	0	0	1	115
	IV	0	0	0	0	1	0	1	0	0	0	2
	V	0	0	0	0	0	16	0	0	0	0	16
	VI	0	0	0	0	0	1	8	0	0	0	9
	VII	0	0	0	0	0	0	0	5	0	0	5
	VIII	0	0	0	0	0	0	0	0	1*	0	1
	IX	0	1	0	0	0	0	0	0	0	1*	2
	Non- typeable	0	1	0	0	0	0	1	1	0	0	3
	Total	37	11	26	112	1	17	10	6	1	2	223

209 *Reference GBS isolates from Statens Serum Institute serotypes VIII and IX

210

211 **Table 3.** Relationship between allocated serotype and second-best match (see also Figure 1)

Allocated serotype	% match	Second-best serotype	% match
la	93.91-100	III	64.56
III	100	la	62.98
V	100	IX	36.26
IX	100	V	31.05
VI	100	III	26.68
IV	100	la	20.3
lb	99.61-100	VI	15.55
II	99.86-100	IV	9.45
VII	100	lb	6.95
VIII	100	none	0

212

213

214 **Table 4.** Retyping of discordant and non-typable isolates

Isolate	Reason for retyping	Latex agglutination		WGS	
		Initial	Repeat	Initial	Repeat
CB466	Discordant	III	Ia	Ia	Ia
IW8194	Discordant	III	IX	IX	IX
IW8466	Discordant	Ia	III	III	III
IW8471	Discordant	III	Ia	Ia	Ia
IW7157	Discordant	Ib	II	II	II
SMRU1	Discordant	VI	V	V	V
SMRU25	Discordant	IV	NT	VI	VI
SMRU4	Discordant	IX	NT	Ib	Ib
SMRU59	Discordant	III	NT	Ia	Ia
Z41	Non-typeable	NT	NT	Ib	Ib
UK22	Non-typeable	NT	NT	VII	VII
IW2723	Non-typeable	NT	NT	VI	VI
CB454	Control	III	III	III	III
IW4445	Control	Ia	Ia	Ia	Ia
IW4077	Control	II	II	II	II

215

216

217 **Figure 1 Discordant isolates show high support for sequence-based serotype allocation.**

218 For each isolate, the percentage of the capsular locus region present ($\geq 95\%$ sequence
219 identity) for the assigned serotype is shown on the X axis, and that for the serotype showing
220 the next best match on the Y axis. Isolates showing agreement between sequence-based
221 and conventional serotyping are shown in grey, those classified as non-typeable by
222 conventional methods in blue, and discordant isolates in red. Small circles represent single
223 isolates, the large circle represents 100 isolates. For each serotype, the second-best match is
224 identical in all cases, leading to the observed horizontal banding (details in Table 3).

