

How environment and self-motion combine in neural representations of space

Talfan Evans^{1,2,3,4,*}, Andrej Bicanski^{2,3,*}, Daniel Bush^{2,3}, Neil Burgess^{2,3}

*These authors contributed equally

¹UCL Centre for Mathematics and Physics in the Life Sciences and Experimental Biology, Gower Street, London, WC1E 6BT, UK

²UCL Institute of Cognitive Neuroscience, 17 Queen Square, London, WC1N 3AR, UK

³UCL Institute of Neurology, Queen Square, London, WC1N 3BG, UK

⁴UCL Department of Neuroscience, Physiology and Pharmacology, Gower Street, London, WC1E 6BT, UK

Correspondence: n.burgess@ucl.ac.uk

Keywords: Hippocampus, path integration, grid cell, head-direction cell, place cell, boundary vector cell.

Abstract: Estimates of location can be constructed solely from sensory information representing environmental cues. In unfamiliar or sensory-poor environments, these estimates can also be maintained and updated by integrating self-motion information. However, the accumulation of error means that updated representations of heading direction and location become progressively less reliable over time, and must be corrected by environmental sensory inputs when available. Here, we review anatomical, electrophysiological and behavioural evidence which indicates that angular and translational path integration contributes to the firing of head direction cells and grid cells respectively. We then review evidence concerning the mechanisms by which environmental sensory inputs contribute to the firing properties of these cells to reduce cumulative spatial errors, potentially mediated by boundary vector cells and place cells in the case of grid cell firing. Finally, we discuss how both types of information are combined together to support a coherent representation of the animal's location and orientation within its environment.

Introduction

Knowing one's location relative to the environment is crucial to the survival of most mobile creatures, and explicitly relies on environmental sensory input. Many organisms also use their own self-motion to maintain an estimate of location and orientation in the absence of sensory input, a process referred to as 'spatial updating' or 'path integration' (e.g., Barry and Burgess, 2014; Etienne and Jeffrey 2004). However, the cumulative nature of path integration dictates that the underlying representations will also integrate error, making the estimate of location and orientation less reliable over time. These location and orientation estimates therefore need to be corrected by environmental sensory information whenever possible, rectifying any misalignment between the internally derived representation and the current perceptual estimate.

The hippocampal formation contains several well described classes of neuron whose activity correlates with an animal's location or orientation. These include place cells (PCs) in the hippocampus proper, which fire in specific locations (O'Keefe and Dostrovsky, 1971); head direction cells (HDCs), found along Papez's circuit, which signal the orientation of an animal's head with respect to the environment (Ranck, 1984; Taube et al., 1990a,b); grid cells (GCs) in the medial entorhinal cortex (mEC), pre- and para- subiculum, which exhibit multiple firing fields arranged on a hexagonal grid (Hafting et al., 2005; Boccara et al., 2010); and border or boundary vector cells (BVCs) in the mEC and subiculum, which fire due to the presence of an extended boundary in a fixed allocentric direction (Barry et al., 2006; Savelli et al., 2008; Solstad et al., 2008; Lever et al., 2009).

In particular, HDCs and GCs are thought to provide the neural bases of angular and translational path-integration, respectively (McNaughton et al., 2006; Taube 2007). If this is the case, then these cells must receive both environmental and self-motion related sensory information, allowing them to update their activity according to path integration and eliminate accumulated error according to perceptual information. Here we review current knowledge about how environmental sensory inputs interact with spatial representations of location and orientation to reduce accumulated error.

Sensory reset of angular path integration

The firing rates of individual HDCs are strongly tuned to the direction that an animal's head is facing relative to the environment, and specifically to landmarks placed at or beyond the edge of the environment (Taube et al., 1990a). The resultant tuning curves are Gaussian or triangular in shape and centered on a distinct direction for each cell (the so-called preferred direction, cf. Fig. 1A,B). HDCs are found along Papez circuit, e.g. in the dorsal tegmental nucleus, lateral mammillary nucleus, anterior dorsal thalamus, and dorsal presubiculum (also called postsubiculum, for review see, Taube 2007), as well as in the mEC (Sargolini et al., 2006) and retrosplenial cortex (Chen et al., 1994). Lesion studies suggest that the generative circuitry is restricted to the dorsal tegmental nucleus and the mammillary bodies (Blair et al., 1998; Bassett et al., 2007), while little is known about the computations carried out by the downstream structures.

Several findings support the hypothesis that HDCs perform angular path integration. First, differences in preferred direction between cells are conserved after environmental manipulations (i.e. displacement of a salient landmark), suggesting that the entire ensemble of HDCs forms a coherent unit (Taube et al., 1990b). Second, aside from drift due to accumulating error or bias (i.e., within 2-3 minutes), HDCs maintain stable directional firing after the removal of prominent visual cues and in darkness (Taube et al., 1990b; Mizumori and Williams, 1993; Goodridge et al., 1998), demonstrating that the networks which support their firing are in principle capable of sustaining the head direction signal internally. Third, rotating an environment in darkness, below the vestibular threshold of an animal, will lead to errors in the homing vector equivalent to the amount of rotation (Mittelstaedt and Mittelstaedt, 1980; Valerio and Taube, 2012).

There is strong evidence that a continuous ring attractor circuit generates the head direction signal (see e.g., Zhang, 1996; Peyrache et al., 2015). Attractor networks can sustain persistent activity (the so-called activity packet or activity bump on topographically organized cells) by virtue of translation invariant recurrent connectivity; most often local excitation in combination with long-range inhibition (Fig. 1C). Unlike discrete attractor networks, which can occupy a finite set of stable states, the variable encoded by a continuous attractor circuit can in principle assume

any value within its range (e.g. 1 to 360 degrees for the head direction system), constraining patterns of activity to a continuous subspace corresponding to these values. The sensitivity of continuous attractor networks to noise (in external drive, network topology and neuronal parameters) explains the observation that HDCs preferred directions begin to drift in darkness (corresponding to drift of the activity packet in the attractor network), rendering the angular path integration system unreliable over time. Despite the fact that rodents have poor vision, drift of the head direction signal in darkness is a well-established experimental finding (Mizumori and Williams, 1993; Goodridge et al., 1998) underlining the strong influence of visual information. A small amount of drift can even be observed in light (Yoder et al., 2015).

Correcting for drift in the head direction circuit is closely tied to the notion of cue control, where a salient landmark in the environment gains control over the head direction ensemble (Taube et al., 1990a,b). Shifting the landmark in the absence of the animal leads to corresponding shifts in the head direction representation (measured in the laboratory reference frame) when the animal is reintroduced into the arena. Cue control can be established within the first few minutes of exploration in a novel environment (Goodridge et al., 1998), implying a fast acting plasticity mechanism. Similarly, gradually shifting the angular position of the reference landmark in the presence of the animal leads to a corresponding shift in head direction. Interestingly, however, the shift in HDC firing lags behind the movement of the cue, suggesting that plasticity may also act during the shift, leading to a stabilization of head direction activity offset towards the previously learned value (Knight et al., 2013).

Learning the correct mapping between visual representations and head direction is a dynamic process, which takes several minutes, and will most likely be refined with prolonged exposure to the same environment. Associating the head direction representation with environmental input will also rely on the spatiotemporal stability of external cues, with animals preferring to use stable landmarks as reference points (Knierim et al., 1995). While environmental boundaries can anchor head direction, ensembles of distal landmarks, which are more stable in the visual field, appear to exert the strongest influence (Jeffery et al., 1998; Zugaro et al., 2001; Clark et al., 2012).

In models of head direction, both cue control and sensory reset of the activity packet in the ring attractor circuit rely on a separate sensory representation, usually in the form of a ring of visual cells (or an equivalent implementation) that represent the angular bearing of a distal reference landmark (Skaggs et al., 1995). Topographic connections between this visual ring and corresponding cells in the head direction attractor ensure stability and cue control (cf. Fig. 1D). Suppose, for example, that facing magnetic North corresponds to a head direction of 90 degrees. When the animal is facing North, visual cells firing in response to a distal landmark located straight ahead will drive the 90 degree head direction cells in the ring attractor, shifting the activity bump to that position if it was misaligned. In the absence of this visual input (i.e. in darkness), drift of the activity bump is no longer overridden by the comparatively strong visual signal and will re-emerge due to the inherent noise-accumulating properties of attractor networks.

It is not entirely clear where such a visual representation may reside. Anatomical considerations suggest that retrosplenial cortex and postsubiculum are the main gateways for visual information onto the head direction circuit (Vann et al., 2009; Taube, 2007). Postsubiculum neurons receive direct input from visual area 17, which may represent salient visual landmarks. Retrosplenial cortex is reciprocally connected to two main downstream loci of HDCs (the anterior dorsal thalamic nucleus and postsubiculum, i.e. dorsal presubiculum; Vann and Aggleton, 2005), and appears to be crucial for spatial orientation in rodents. Lesions to the dysgranular area of retrosplenial cortex lead rats to switch to a motor turn strategy instead of relying on visual landmarks (Vann and Aggleton, 2005), suggesting landmark-based navigation is impaired after retrosplenial cortex lesions. More recently, Yoder et al. (2015) have shown that an intact postsubiculum is crucial for stable head direction tuning in the lateral mammillary nucleus. Most likely both retrosplenial cortex and postsubiculum are necessary processing stages for visual feedback to reach the generative head direction circuitry in upstream structures.

Sensory reset of translational path integration

Whereas the head direction system appears to be responsible for the angular component of path integration, the translational component is thought to be mediated

by grid cells (GCs), which exhibit a unique hexagonal pattern of firing fields that tile a given environment (Hafting et al., 2005). These grid patterns are characterized by their spatial scale (distance between grid fields) and orientation relative to the environment (Fig. 2A). Moving dorsoventrally along the mEC, GCs increase their scale in discrete steps (Barry et al., 2007; Stensola et al., 2012; Stensola et al., 2015). GCs with similar scales share the same orientation relative to their environment but have offset firing fields, providing uniform spatial coverage (Hafting et al., 2005; Fig. 2B).

Multiple mechanisms have been suggested for the means by which the GC firing patterns might track the movement of the animal. These include asymmetric interactions reflecting direction and speed of travel, analogous to those proposed for head-direction circuits (Zhang, 1996) and mediated by ‘directional’ or ‘conjunctive’ GCs (Sargolini et al., 2006; McNaughton et al., 2006; Fuhs and Touretzky, 2006). Alternatively, movement of the bump of activity may be driven by interference of velocity-controlled oscillators in the theta frequency band (Burgess et al., 2007) consistent with the theta-phase precession effect observed in GC firing (Hafting et al., 2008) and PC firing (O’Keefe and Recce, 1993), see (Bush and Burgess, 2014).

What evidence is there that GCs represent the neural substrate for path integration? Firstly, unlike place cells (PCs), which undergo re-mapping between different environments (Muller and Kubie, 1987), GCs preserve their firing patterns across all environments visited by the animal (Hafting et al., 2005). Importantly, the relative spatial offset of firing fields recorded from cells in the same module is maintained across different environments (Hafting et al., 2005; Yoon et al., 2013). These properties suggest that GCs whose firing fields share a similar scale and orientation form cohesive functional modules (Stensola et al., 2012). Moreover, in situations where the grids are unlikely to be directly driven by common environmental cues (Barry et al., 2012), this invariance in spatial offset between grid firing fields within the same module also points to a shared path integration mechanism (Yoon et al., 2013). Secondly, the spatial stability of grid firing fields decreases over time in darkness, in the supposed absence of ‘corrective’ sensory inputs (Hafting et al., 2005). Finally, lesion studies suggest that navigation tasks that rely specifically on path

integration mechanisms can be impaired after lesions of mEC (van Cauter et al., 2013).

The fact that the location of GC firing fields are stable between visits to an environment, and that grid orientation rotates (along with the tuning curves of HDCs) with a prominent visual cue (Fig. 2A; Hafting et al., 2005; Sargolini et al., 2006) indicates that GCs also receive environmental sensory inputs. More dramatically, reshaping a familiar rectangular environment along one axis can cause the parametric deformation of previously stable GC firing patterns along the same axis, resulting in both elliptical firing fields and grid patterns (Barry et al., 2007; Stensola et al., 2012, Fig. 2C). This suggests that, in familiar environments, sensory inputs can shift and deform the intrinsic metric generated by self-motion information. Interestingly, with repeated reshaping of the environment the expansion and contraction of GC firing patterns is diminished, suggesting that sensory inputs that are known to be unstable are eventually ignored mirroring similar results for orientation, described above (Knierim et al., 1995).

How might environmental sensory inputs be used to correct drift in GC firing? Hebbian associations from PCs to GCs have been proposed as a likely route, given the convenient overlap between the PC firing fields and single nodes of a grid (O'Keefe and Burgess, 2005), with PCs mediating the effect of boundary manipulations via their proposed BVC inputs (Hartley et al., 2000). However, it is equally possible that BVCs directly stabilize GC firing (Solstad et al., 2008). Several theoretical studies have demonstrated that input from either PCs or BVCs is sufficient to ameliorate the effects of noisy self-motion input to the GC network (e.g. Fuhs and Touretzky, 2006; Bush and Burgess, 2014; Hardcastle et al., 2015). The spatial stability and parametric deformation of grid firing fields described above, for example, could be mediated by direct excitatory input from BVCs or PCs with firing fields either along the borders or distributed evenly throughout the environment, 'correcting' the uniform hexagonal firing produced by self-motion inputs (Bush and Burgess, 2014).

Exploring the hypothesis that boundary input acts to stabilize path integration, Hardcastle et al. (2015) analyzed the stability of GC firing patterns recorded in vivo with respect to rats' encounters with boundaries. They showed that 'error' in the grid

fields (the distance between spikes and the center-of-mass of their nearest firing field, measured across the whole recording session) increases with both time and distance since the last boundary encounter. Interestingly, both the magnitude and direction of the error was also consistent across all recorded GCs, suggesting a coherent drift of the grid pattern relative to the environment, as opposed to an increase in the size of individual grid fields. Moreover, error correction was greater in the direction perpendicular to the last boundary encounter, as might be expected if the grid were responding to BVCs that provide information about distance perpendicular to that boundary. These findings suggest that positional uncertainty may manifest itself as environmental misalignment of an intrinsically stable GC firing pattern, rather than a breakdown of GC firing patterns altogether. Furthermore, the hypothesis that GC firing patterns are stabilized by environmental boundaries predicts that time-averaged grid fields should be smaller and more elliptical near to the boundary, and that this effect would be more pronounced in larger environments.

Notwithstanding the possibility of direct boundary-related input to GCs, there is also evidence that input from PCs contributes to stable GC firing. For example, inactivation of the hippocampus rapidly disrupts the spatial periodicity of GC firing patterns, decreases the mean firing rate of GCs and increases their directional tuning (Bonnievie et al., 2013). Moreover, stable PC, BVC and HDC firing patterns are known to precede stable GC firing patterns in the developmental timeline, suggesting that input from each of these cell types might be required for the generation of stable GC firing patterns (Langston et al., 2010; Wills et al., 2010). Interestingly, it has recently been shown that the stability of PC firing fields is inversely correlated with distance to the nearest environmental boundary in pre-weanling rats, but that this relationship disappears in adult animals (Muessig et al., 2015). This suggests that, prior to the development of GCs, inputs from BVCs active around the boundaries of an environment also serve to stabilize PC firing (O'Keefe and Burgess, 1996; Hartley et al., 2000). Later in development, the appearance of stable place fields in the center of an environment coincides with the establishment of stable GC firing (Muessig et al., 2015). However, whether path integration input from GCs is required to stabilize place fields, or environmental sensory input from PCs is required to eliminate grid field drift away from environmental boundaries is yet to be determined.

Adaptation over multiple timescales

The discussion so far has concerned interactions between place or boundary vector and grid cell populations according to pre-established neural connections. However, in new environments these connections must develop through synaptic plasticity over longer timescales, and may be associated with gradual changes observed in GC firing patterns with increasing experience of an environment. Notably, the spatial scale of grid firing patterns reduces and their spatial regularity increases with experience of a novel environment (Barry et al., 2012). At the same time, the orientation of the grid firing patterns relative to the boundaries of a square environment changes from an initially parallel alignment to develop a small ($\sim 7.5 - 8.8^\circ$) angular offset (Stensola et al., 2015; Krupic et al., 2015; Fig. 2D). In larger square environments, local 7.5° orientation to multiple boundaries is observed, and is associated with a departure from the globally regular hexagonal firing pattern, resulting in local ellipticity (the skew of an ellipse tracing the innermost six peaks of a grid, see Fig. 2E).

In smaller square environments, both the angular offset of the grid firing pattern to local boundaries and its ellipticity can be modelled as a uniaxial shearing transformation, suggesting that both are the result of a common underlying process (Stensola et al., 2015). In larger square environments, the effects can only be fully removed by a biaxial shearing centered about ‘anchoring’ points in opposite corners. Whether or not the observed angular offsets and distortions of the grid fields that develop with experience, as well as the initial decrease in grid scale, serve to distort the perception of space, or offer some functional advantage, are questions for future empirical studies. In either case, the influence of environmental boundaries on GC firing patterns seems likely to arise as an inextricable consequence of functional BVC inputs to GCs which are required to counteract error accumulation (Hardcastle et al., 2015; Stensola et al., 2015).

In addition to changes in GC firing patterns that may reflect their association to environmental sensory inputs, both PC and GC firing patterns are known to exhibit more gradual changes that reflect the slow disambiguation of perceptually similar environments. For example, on initial exposure to square and circular environments, place fields share preferred firing locations, measured relative to both a cue card and environmental boundaries, consistent with an important role of BVC inputs (Lever et

al., 2002). However, over a period of days, firing locations in each environment diverge, either as a result of the gradual shifting of firing fields, or through the respective disappearance and emergence of the original and new fields.

Similarly, when rats are allowed to commute freely between two perceptually identical environments, GC firing patterns that are initially identical in both boxes gradually shift to form a unified, globally consistent representation with experience (Carpenter et al., 2015). The alignment and spatial phase of the initially local GC firing patterns are consistent with conserved boundary-related inputs, and the subsequent shift suggests that, during navigation between the two environments, path integration input slowly forces the sensory associations to be relearned. Interestingly, global (i.e. path integration consistent) tuning of HDCs in connected environments emerges on similar timescales (Dudchenko and Zinkyuk, 2005). Thus, longer-term adaptation may be important both in disambiguating perceptually similar environments, and in embedding these localized representations within a consistent global framework.

Origins of the sensory inputs to grid cells and place cells

The firing patterns of place cells in CA1 are robust to the lesioning of many different input streams, including CA3 (Brun et al., 2002), mEC (Miller and Best, 1980; Brun et al., 2008), IEC (Lu et al., 2013), pre- and para- subiculum (Liu et al., 2004), and subcortical structures (Miller and Best, 1980). Thus, it seems likely that PC firing patterns are supported by a wide range of multi-modal inputs that converge on the hippocampus from a variety of different cortical structures. Accordingly, PC firing fields are modulated by contextual inputs such as wall color and odor (Bostock et al., 1991; Anderson and Jeffery, 2003; Lu et al., 2013), manifested in place cell “remapping”. It has been hypothesized that BVCs are a potential source for the environmental determination of the spatial firing locations of PCs, but it is important to note that BVC firing patterns are allocentric representations, e.g. responding to a boundary to the North independent of the orientation of the animal. By contrast, sensory representations are necessarily egocentric in nature. Hence, environmental sensory inputs to BVCs will at some stage require an egocentric-allocentric coordinate transformation.

One such mechanism for transformation between population codes in different frames of reference has been proposed, involving an expanded “gain field” representation formed by the product of both individual sets of basis functions (Salinas and Abbott, 1996; Pouget and Sejnowski, 1997; Burgess et al., 2001; Byrne et al., 2007). To account for the allocentric firing patterns of BVCs, it has been hypothesized that BVCs in the hippocampal formation have an egocentric analog in medial parietal areas (Burgess et al., 2001; Byrne et al., 2007). These putative cells would signal the presence of boundaries at given distances in an egocentric frame of reference via receptive fields in peripersonal space. This egocentric representation could be driven directly by sensory input and transformed to an allocentric representation via a ‘gain field’ circuit in posterior parietal or retrosplenial cortex. This egocentric-allocentric transformation depends on the head direction system to correctly map egocentric into allocentric direction (e.g. North corresponds to left when facing East).

Thus, boundary-related input would arrive at the hippocampal formation from the dorsal processing stream (Goodale and Milner, 1992; Ungerlieder and Mishkin, 1982) via the parietal, retrosplenial, parahippocampal and medial entorhinal cortices. By contrast, contextual and object-related inputs (Deshmukh and Knierim, 2011; Lu et al., 2013; Tsao et al. 2013) likely arrive from the ventral processing stream via perirhinal and lateral entorhinal cortices. Hence, these pathways carry complementary sensory signals (see color code in Fig. 3), which provide input to PCs and GCs independently, increasing the robustness of the global path-integration system.

Discussion

We have discussed how an effective spatial navigation system requires complementary information both from environmental sensory perception and self-motion inputs, and reviewed evidence that HDCs and GCs are the neural substrates of angular (i.e. directional) and translational (i.e. location) path integration, respectively. In this review, we have focused primarily on unidirectional processing pathways, i.e. the transmission of environmental information directly to HDCs, or via BVCs and PCs to GCs. However, it is important to note that GCs are also likely to project to PCs in the hippocampus (van Strien et al., 2009), and it has been suggested that GCs can partially support and update PC firing when sensory inputs are reduced (e.g. in darkness). Accordingly, evidence indicates that when visual and path integration

information are dissociated, the resulting PC firing is best described by a balanced combination of these inputs (Gothard et al., 1996; Chen et al., 2013).

More generally, it makes sense that our perception of spatial location and orientation combines both types of input. Behavioural evidence in humans further suggests that the two input streams are integrated into a single representation (Tcheang et al., 2012), and are possibly weighted to reflect the relative reliability of the available sources of information (Nardini et al., 2008). Thus, reciprocal connectivity between PCs and GCs is likely to underlie the concurrent development of stable PC and GC firing fields away from the boundaries of an environment (Muessig et al., 2015), in order to form a single multimodal representation of location. Similarly, HDC input is critical for the transformation of egocentric sensory input to allocentric BVC and PC firing patterns, all of which rotate coherently with HDC firing patterns when a single prominent cue is shifted in a circular arena (O'Keefe and Conway, 1978; Muller and Kubie, 1987; Knierim et al., 1995; Sargolini et al., 2006). Conversely, spatial information from PCs and GCs is likely involved in the development of coherent HD representations across connected environments (Dudchenko and Zinyuk, 2005). To summarise, these data suggest that the different spatial representations found in and around the hippocampal formation are mutually interdependent and support a coherent unitary representation of space based on both environmental sensory and path integration inputs.

Acknowledgements

The authors would like to thank James Bisby and Aidan Horner for helpful comments. This work was supported by the EPSRC, Wellcome Trust, Medical Research Council UK and the EU FP7 FET project SpaceCog.




Figure 1: Properties of Head Direction Cells. **(A)** Head direction (HD) cells code for the allocentric direction of an animal's head with reference to the normal vector of a salient landmark. **(B)** A prototypical head direction tuning curve (head direction Θ , firing rate r). **(C)** The topology of a canonical ring attractor circuit. Only connections from one cell (filled circle) are shown. Short range excitatory connections (ending in filled triangles) and long range inhibitory connections (ending in filled circles) assure a local self-sustaining 'packet of activity'. **(D)** The standard model of visual feedback in the head direction system. Each head direction is associated with a unique visual activity profile. Topographic connections (black arrows) excite the correct portion of the head direction ring to prevent drift.




Figure 2: The Grid Cell Path Integration System. **(A)** Rotation of a black cue card causes rotation of the grid pattern. Φ shows the grid orientation, λ the grid scale (adapted from Hafting et al., 2005). **(B)** Topologically neighboring grid cells (red, green, blue) have different spatial phases, but equal scale and orientation (adapted from Hafting et al., 2005). **(C)** Animals were placed in an environment with moveable walls (adapted from Barry et al., 2007). Grid cell firing patterns responded to uniaxial (top-right, bottom-left) and biaxial (bottom-right) contraction and returned to their original configuration (top-left). **(D)** Grids have three measures of orientation corresponding to their three axes. In a square environment, the angle from the wall is defined as the smallest of the three axis angles from their nearest wall. The smallest of these three angle clusters bi-modally about 7.5° (adapted from Stensola et al., 2015). **(E)** Left: An example binned spike map in a 2.2m² box. Middle: Ellipses fitted to the innermost six peaks of the local autocorrelogram (different to right). Right: Ellipticity

and gridness scores in different sections of a 2.2m² box, showing greater distortion near the edges (adapted from Stensola et al., 2015).




Figure 3: Main pathways for sensory signals necessary to reset path integration systems. Colored arrows: known/hypothesized pathways for sensory information. Purple: the head direction system receives projections from visual area 17 and from retrosplenial cortex (RSC). Head direction potentially gates an egocentric-alloentric transformation (red) between parietal and retrosplenial cortices, necessary to transform egocentric boundary distance (sensory input) to allocentric boundary distance (BVCs). Blue and green: complementary pathways conveying sensory information to grid cells, sources attributed to the dorsal and ventral visual streams (VS) respectively. Black arrows/arrow heads: pathways for idiothetic information, i.e. path integration signals. Abbreviations: place cells (PCs), boundary vector cells (BVCs), grid cells (GCs), head direction cells (HDCs), medial and lateral entorhinal cortex (m/IEC). Note, some known projections are omitted for clarity, e.g. from retrosplenial cortex to mEC.

References

- Anderson, M.I., and Jeffery, K.J. (2003). Heterogeneous modulation of place cell firing by changes in context. *The Journal of Neuroscience* *23*, 8827–8835.
- Barry, C., and Burgess, N. (2014). Neural Mechanisms of Self-Location. *Current Biology* *24*, R330–R339.
- Barry, C., Ginzberg, L.L., O’Keefe, J., and Burgess, N. (2012). Grid cell firing patterns signal environmental novelty by expansion. *Proceedings of the National Academy of Sciences* *109*, 17687–17692.
- Barry, C., Lever, C., Hayman, R., Hartley, T., Burton, S., O’Keefe, J., Jeffery, K., and Burgess, N. (2006). The boundary vector cell model of place cell firing and spatial memory. *Rev Neurosci* *17*, 71–97.
- Bassett, J.P., Tullman, M.L., and Taube, J.S. (2007). Lesions of the Tegmentomammillary Circuit in the Head Direction System Disrupt the Head Direction Signal in the Anterior Thalamus. *J. Neurosci.* *27*, 7564–7577.
- Blair, H.T., Cho, J., and Sharp, P.E. (1998). Role of the lateral mammillary nucleus in the rat head direction circuit: a combined single unit recording and lesion study. *Neuron* *21*, 1387–1397.
- Boccaro, C.N., Sargolini, F., Thoresen, V.H., Solstad, T., Witter, M.P., Moser, E.I., and Moser, M.-B. (2010). Grid cells in pre- and parasubiculum. *Nature Neuroscience* *13*, 987–994.
- Bonnevie, T., Dunn, B., Fyhn, M., Hafting, T., Derdikman, D., Kubie, J.L., Roudi, Y., Moser, E.I., and Moser, M.-B. (2013). Grid cells require excitatory drive from the hippocampus. *Nature Neuroscience* *16*, 309–317.
- Bostock, E., Muller, R.U., and Kubie, J.L. (1991). Experience-dependent modifications of hippocampal place cell firing. *Hippocampus* *1*, 193–205.
- Brun, V.H., Leutgeb, S., Wu, H.-Q., Schwarcz, R., Witter, M.P., Moser, E.I., and Moser, M.-B. (2008). Impaired Spatial Representation in CA1 after Lesion of Direct Input from Entorhinal Cortex. *Neuron* *57*, 290–302.

Brun, V.H., Otnæss, M.K., Molden, S., Steffenach, H.-A., Witter, M.P., Moser, M.-B., and Moser, E.I. (2002). Place Cells and Place Recognition Maintained by Direct Entorhinal-Hippocampal Circuitry. *Science* 296, 2243–2246.

Burgess, N., Barry, C., and O’Keefe, J. (2007). An oscillatory interference model of grid cell firing. *Hippocampus* 17, 801–812.

Burgess, N., Becker, S., King, J.A., and O’Keefe, J. (2001). Memory for events and their spatial context: models and experiments. *Philosophical Transactions of the Royal Society B: Biological Sciences* 356, 1493–1503.

Byrne, P., Becker, S., and Burgess, N. (2007). Remembering the past and imagining the future: A neural model of spatial memory and imagery. *Psychological Review* 114, 340–375.

Carpenter, F., Manson, D., Jeffery, K., Burgess, N., and Barry, C. (2015). Grid Cells Form a Global Representation of Connected Environments. *Current Biology* 25, 1-7

Chen, G., King, J.A., Burgess, N., and O’Keefe, J. (2013). How vision and movement combine in the hippocampal place code. *Proc Natl Acad Sci U S A* 110, 378–383.

Chen, L.L., Lin, L.-H., Green, E.J., Barnes, C.A., and McNaughton, B.L. (1994). Head-direction cells in the rat posterior cortex. *Experimental Brain Research* 101, 8–23.

Clark, B.J., Harris, M.J., and Taube, J.S. (2012). Control of anterodorsal thalamic head direction cells by environmental boundaries: Comparison with conflicting distal landmarks. *Hippocampus* 22, 172–187.

Dudchenko, P.A., and Zinyuk, L.E. (2005). The Formation of Cognitive Maps of Adjacent Environments: Evidence From the Head Direction Cell System. *Behavioral Neuroscience* 119, 1511–1523.

Etienne, A.S., and Jeffery, K.J. (2004). Path integration in mammals. *Hippocampus* 14, 180–192.

Fuhs, M.C., and Touretzky (2006). A Spin Glass Model of Path Integration in Rat Medial Entorhinal Cortex. *Journal of Neuroscience* 26, 4266–4276.

Goodale, M.A., and Milner, A.D. (1992). 1 2 Separate Visual Pathways for Perception and Action. *Essential Sources in the Scientific Study of Consciousness* 175.

Goodridge, J.P., Dudchenko, P.A., Worboys, K.A., Golob, E.J., and Taube, J.S. (1998). Cue control and head direction cells. *Behavioral Neuroscience* 112, 749.

Gothard, K.M., Skaggs, W.E., and McNaughton, B.L. (1996). Dynamics of mismatch correction in the hippocampal ensemble code for space: interaction between path integration and environmental cues. *The Journal of Neuroscience* 16, 8027–8040.

Hafting, T., Fyhn, M., Molden, S., Moser, M.-B., and Moser, E.I. (2005). Microstructure of a spatial map in the entorhinal cortex. *Nature* 436, 801–806.

Hardcastle, K., Ganguli, S., and Giocomo, L.M. (2015). Environmental Boundaries as an Error Correction Mechanism for Grid Cells. *Neuron* 86, 1-13

Hartley, T., Burgess, N., Lever, C., Cacucci, F., and O'keefe, J. (2000). Modeling place fields in terms of the cortical inputs to the hippocampus. *Hippocampus* 10, 369–379.

Jeffery, K.J. (1998). Learning of landmark stability and instability by hippocampal place cells. *Neuropharmacology* 37, 677–687.

Knierim, J.J., Kudrimoti, H.S., and McNaughton, B.L. (1995). Place cells, head direction cells, and the learning of landmark stability. *The Journal of Neuroscience* 15, 1648–1659.

Knight, R., Piette, C.E., Page, H., Walters, D., Marozzi, E., Nardini, M., Stringer, S., and Jeffery, K.J. (2013). Weighted cue integration in the rodent head direction system. *Philosophical Transactions of the Royal Society B: Biological Sciences* 369, 20120512–20120512.

Krupic, J., Bauza, M., Burton, S., Barry, C., and O'Keefe, J. (2015). Grid cell symmetry is shaped by environmental geometry. *Nature* 518, 232–235.

- Langston, R.F., Ainge, J.A., Couey, J.J., Canto, C.B., Bjerknes, T.L., Witter, M.P., Moser, E.I., and Moser, M.-B. (2010). Development of the Spatial Representation System in the Rat. *Science* 328, 1576–1580.
- Lever, C., Burton, S., Jeewajee, A., O’Keefe, J., and Burgess, N. (2009). Boundary Vector Cells in the Subiculum of the Hippocampal Formation. *Journal of Neuroscience* 29, 9771–9777.
- Lever, C., Wills, T., Cacucci, F., Burgess, N., and O’Keefe, J. (2002). Long-term plasticity in hippocampal place-cell representation of environmental geometry. *Nature* 416, 90–94.
- Liu, L., Wong, T.P., Pozza, M.F., Lingenhoehl, K., Wang, Y., Sheng, M., Auberson, Y.P., and Wang, Y.T. (2004). Role of NMDA receptor subtypes in governing the direction of hippocampal synaptic plasticity. *Science* 304, 1021–1024.
- Lu, L., Leutgeb, J.K., Tsao, A., Henriksen, E.J., Leutgeb, S., Barnes, C.A., Witter, M.P., Moser, M.-B., and Moser, E.I. (2013). Impaired hippocampal rate coding after lesions of the lateral entorhinal cortex. *Nature Neuroscience* 16, 1085–1093.
- McNaughton, B.L., Battaglia, F.P., Jensen, O., Moser, E.I., and Moser, M.-B. (2006). Path integration and the neural basis of the “cognitive map.” *Nature Reviews Neuroscience* 7, 663–678.
- Mishkin, M., and Ungerlieder, L.G. (1982). Contribution of striate inputs to the visuospatial functions of parieto-preoccipital cortex in monkeys. *Behavioural Brain Research* 6, 57–77.
- Mittelstaedt, M.-L., and Mittelstaedt, H. (1980). Homing by path integration in a mammal. *Naturwissenschaften* 67, 566–567.
- Mizumori, S.J., and Williams, J.D. (1993). Directionally selective mnemonic properties of neurons in the lateral dorsal nucleus of the thalamus of rats. *J. Neurosci.* 13, 4015–4028.
- Muessig, L., Hauser, J., Wills, T.J., and Cacucci, F. (2015). A Developmental Switch in Place Cell Accuracy Coincides with Grid Cell Maturation. *Neuron* 86, 1167–1173.

- Muller, R.U., and Kubie, J.L. (1987). The effects of changes in the environment on the spatial firing of hippocampal complex-spike cells. *J Neurosci* 7, 1951–1968.
- Muller, V.M., and Best, P.J. (1980). Spatial correlates of hippocampal unit activity are altered by lesions of the fornix and entorhinal cortex. *Brain Research* 311–323.
- Nardini, M., Jones, P., Bedford, R., and Braddick, O. (2008). Development of Cue Integration in Human Navigation. *Current Biology* 18, 689–693.
- O’Keefe, J., and Burgess, N. (1996). Geometric determinants of the place fields of hippocampal neurons. *Nature* 381, 425–428.
- O’Keefe, J., and Burgess, N. (2005). Dual phase and rate coding in hippocampal place cells: Theoretical significance and relationship to entorhinal grid cells. *Hippocampus* 15, 853–866.
- O’Keefe, J., and Conway, D.H. (1978). Hippocampal place units in the freely moving rat: why they fire where they fire. *Experimental Brain Research* 31, 573–590.
- O’Keefe, J., and Dostrovsky, J. (1971). The hippocampus as a spatial map. Preliminary evidence from unit activity in the freely-moving rat. *Brain Research* 34, 171–175.
- O’Keefe, J., and Recce, M.L. (1993). Phase relationship between hippocampal place units and the EEG theta rhythm. *Hippocampus* 3, 317–330.
- Peyrache, A., Lacroix, M.M., Petersen, P.C., and Buzsáki, G. (2015). Internally organized mechanisms of the head direction sense. *Nature Neuroscience* 18, 569–575.
- Pouget, A., and Sejnowski, T.J. (1997) A new view of hemineglect based on the response properties of parietal neurones. *Philosophical Transactions of the Royal Society B* 352: 1449–1459
- Ranck Jr, J. B. (1984). Head direction cells in the deep cell layer of dorsal presubiculum in freely moving rats. *Soc Neurosci Abstr.* Vol. 10. No. 176.12.
- Salinas, E., and Abbott, L.F. (1996). A model of multiplicative neural responses in parietal cortex. *Proceedings of the National Academy of Sciences* 93, 11956–11961.

- Sargolini, F. (2006). Conjunctive Representation of Position, Direction, and Velocity in Entorhinal Cortex. *Science* 312, 758–762.
- Savelli, F., Yoganarasimha, D., and Knierim, J.J. (2008). Influence of boundary removal on the spatial representations of the medial entorhinal cortex. *Hippocampus* 18, 1270–1282.
- Skaggs, W.E., Knierim, J.J., Kudrimoti, H.S., and McNaughton, B.L. (1995) A Mode of the Neural Basis of the Rat's Sense of Direction. *Advances in Neural Information Processing Systems 7*: 173-180
- Solstad, T., Boccara, C.N., Kropff, E., Moser, M.-B., and Moser, E.I. (2008). Representation of Geometric Borders in the Entorhinal Cortex. *Science* 322, 1865–1868.
- Stensola, H., Stensola, T., Solstad, T., Frøland, K., Moser, M.-B., and Moser, E.I. (2012). The entorhinal grid map is discretized. *Nature* 492, 72–78.
- Stensola, T., Stensola, H., Moser, M.-B., and Moser, E.I. (2015). Shearing-induced asymmetry in entorhinal grid cells. *Nature* 518, 207–212.
- Taube, J.S. (2007). The Head Direction Signal: Origins and Sensory-Motor Integration. *Annual Review of Neuroscience* 30, 181–207.
- Taube, J.S., Muller, R.U., and Ranck, J.B. (1990a). Head-direction cells recorded from the postsubiculum in freely moving rats. II. Effects of environmental manipulations. *The Journal of Neuroscience* 10, 436–447.
- Taube, J.S., Muller, R.U., and Ranck, J.B. (1990b). Head-direction cells recorded from the postsubiculum in freely moving rats. I. Description and quantitative analysis. *The Journal of Neuroscience* 10, 420–435.
- Tcheang, L., Bulthoff, H.H., and Burgess, N. (2011). Visual influence on path integration in darkness indicates a multimodal representation of large-scale space. *Proceedings of the National Academy of Sciences* 108, 1152–1157.
- Tsao, A., Moser, M.-B., and Moser, E.I. (2013). Traces of Experience in the Lateral Entorhinal Cortex. *Current Biology* 23, 399–405.

- Valerio, S., and Taube, J.S. (2012). Path integration: how the head direction signal maintains and corrects spatial orientation. *Nat Neurosci* *15*, 1445–1453.
- Van Cauter, T., Camon, J., Alvernhe, A., Elduayen, C., Sargolini, F., and Save, E. (2013). Distinct Roles of Medial and Lateral Entorhinal Cortex in Spatial Cognition. *Cerebral Cortex* *23*, 451–459.
- van Strien, N.M., Cappaert, N.L.M., and Witter, M.P. (2009). The anatomy of memory: an interactive overview of the parahippocampal–hippocampal network. *Nature Reviews Neuroscience* *10*, 272–282.
- Vann, S.D., Aggleton, J.P., and Maguire, E.A. (2009). What does the retrosplenial cortex do? *Nature Reviews Neuroscience* *10*, 792–802.
- Vann, S.D., and Aggleton, J.P. (2005). Selective Dysgranular Retrosplenial Cortex Lesions in Rats Disrupt Allocentric Performance of the Radial-Arm Maze Task. *Behavioral Neuroscience* *119*, 1682–1686.
- Wills, T.J., Cacucci, F., Burgess, N., and O’Keefe, J. (2010). Development of the Hippocampal Cognitive Map in Preweanling Rats. *Science* *328*, 1573–1576.
- Yoder, R.M., Peck, J.R., and Taube, J.S. (2015). Visual Landmark Information Gains Control of the Head Direction Signal at the Lateral Mammillary Nuclei. *Journal of Neuroscience* *35*, 1354–1367.
- Yoon, K., Buice, M.A., Barry, C., Hayman, R., Burgess, N., and Fiete, I.R. (2013). Specific evidence of low-dimensional continuous attractor dynamics in grid cells. *Nature Neuroscience* *16*, 1077–1084.
- Zhang, K. (1996). Representation of spatial orientation by the intrinsic dynamics of the head-direction cell ensemble: a theory. *The Journal of Neuroscience* *16*, 2112–2126.
- Zugaro, M.B., Berthoz, A., and Wiener, S.I. (2001). Background, But Not Foreground, Spatial Cues Are Taken as References for Head Direction Responses by Rat Anterodorsal Thalamus Neurons. *J. Neurosci.* *21*, RC154–RC154.