

On DLA's η

Hitoshi Kawakatsu^a, Jean-Paul Montagner^b and Teh-Ru Alex Song^c

^a*Earthquake Research Institute, The University of Tokyo,
1-1-1 Yayoi, Bunkyo-ku, Tokyo 113-0032 JAPAN (hitosi@eri.u-tokyo.ac.jp)*

^b*Institut de Physique du Globe de Paris, Universite Paris-Diderot,
1 rue Jussieu 75238-Paris cedex 05 FRANCE (jpm@ipgp.fr)*

^c*Department of Earth Sciences, University College London,
Gower Street, WC1E 6BT, London, United Kingdom (tehrusong@gmail.com)*

January 29, 2015

1 SUMMARY

2 In his pioneering paper on seismic anisotropy in a layered earth, Anderson (1961) introduced a
3 parameter often referred in global seismology as η without providing any reasoning. This note
4 hopes to clarify the significance of η in the context of the dependence of bodywave velocities in
5 a transversely isotropic system on the angle of incidence, and also its relation with the other
6 well-known anisotropic parameters introduced by Thomsen (1986).

7 **Key words:** Seismic anisotropy, transverse isotropy, radial anisotropy.

8 Introduction

9 To describe a radially anisotropy (transversely isotropy with a vertical symmetry axis, VTI)
10 system, we employ the Love's original notation (Love, 1927), where stress and strain tensors
11 are related by

$$\begin{bmatrix} \tau_{11} \\ \tau_{22} \\ \tau_{33} \\ \tau_{23} \\ \tau_{31} \\ \tau_{12} \end{bmatrix} = \begin{bmatrix} A & H & F & & & \\ & H & A & F & & \\ & F & F & C & & \\ & & & & L & \\ & & & & & L \\ & & & & & & N \end{bmatrix} \begin{bmatrix} e_{11} \\ e_{22} \\ e_{33} \\ 2e_{23} \\ 2e_{31} \\ 2e_{12} \end{bmatrix} \quad (1)$$

12 where $H = A - 2N$. There are five independent parameters, A , C , F , L , N , to describe this
13 system, while there are two, λ , μ , for the isotropic case, for which $A = C = \lambda + 2\mu$, $F = \lambda$,
14 $L = N = \mu$. For convenience, Anderson (1961) introduced the following "anisotropy factors":

$$\varphi = C/A = \alpha_V^2/\alpha_H^2 \quad (2)$$

$$\xi = (A - H)/2L = N/L = \beta_H^2/\beta_V^2 \quad (3)$$

$$\eta = (A - 2L)/F, \quad (4)$$

15 which are all equal to 1 for isotropic case ($\alpha_V = \sqrt{C/\rho}$, $\alpha_H = \sqrt{A/\rho}$, $\beta_H = \sqrt{N/\rho}$, $\beta_V =$
16 $\sqrt{L/\rho}$, where ρ gives the density).

17 While both φ and ξ have simple meanings (degree of anisotropy in P- and S-wave, re-
 18 spectively), the physical meaning of η is not so trivial. Takeuchi and Saito (1972), in their
 19 monograph on seismic surface waves, reversed the order of the denominator and numerator in
 20 the definition of η as,

$$\eta = F/(A - 2L), \quad (5)$$

21 without commenting on the physical meaning either. As the expression of Takeuchi and Saito
 22 (1972) is now commonly used in the global seismological community, we will use this notation
 23 and denote it as $\eta_{DLA} = F/(A - 2L)$ in the following. In his text book, Anderson (1989) called
 24 this η_{DLA} “the fifth parameter required to fully describe transverse isotropy”. In Dziewonski
 25 and Anderson (1981), by showing examples, the effect of η_{DLA} on the incident angle dependence
 26 of the phase velocity of P and S waves is discussed, and we generally think that η_{DLA} controls,
 27 to some extent, the incidence angle dependence of those bodywaves, as well as those of Rayleigh
 28 waves.

29 The purpose of this short note is provide simple theoretical background to how η_{DLA} affects
 30 the bodywave propagation.

31 Incidence angle dependence of bodywaves

32 By solving an eigenvalue problem of an appropriate Christoffel matrix, the incident angle, θ
 33 dependence of bodywave phase velocities can be obtained as

$$\rho v_P^2(\theta) = \frac{(L + C) + (A - C) \sin^2 \theta + \sqrt{S}}{2} \quad (6)$$

$$\rho v_{SV}^2(\theta) = \frac{(L + C) + (A - C) \sin^2 \theta - \sqrt{S}}{2} \quad (7)$$

$$\rho v_{SH}^2(\theta) = L + (N - L) \sin^2 \theta, \quad (8)$$

34 where v_P , v_{SV} , and v_{SH} denote phase velocities of pseudo- P, SV and SH waves respectively,
 35 and

$$S = \{(A - L) \sin^2 \theta - (C - L) \cos^2 \theta\}^2 + (F + L)^2 \sin^2 2\theta \quad (9)$$

$$= \{(A - L) \sin^2 \theta + (C - L) \cos^2 \theta\}^2 + \{(F + L)^2 - (C - L)(A - L)\} \sin^2 2\theta \quad (10)$$

$$= \{(C - L) + (A - C) \sin^2 \theta\}^2 + \{(F + L)^2 - (C - L)(A - L)\} \sin^2 2\theta \quad (11)$$

$$= (C - L)^2 + (A - C)(A + C - 2L) \sin^2 \theta + \{(F + L)^2 - (\frac{A + C}{2} - L)^2\} \sin^2 2\theta. \quad (12)$$

36 When the condition

$$(F + L)^2 = (C - L)(A - L) \quad (13)$$

is satisfied, equation (11) will be $S = \{(C - L) + (A - C) \sin^2 \theta\}^2$, and

$$\rho v_P^2(\theta) = C + (A - C) \sin^2 \theta \quad (14)$$

$$\rho v_{SV}^2(\theta) = L \quad (15)$$

$$\rho v_{SH}^2(\theta) = L + (N - L) \sin^2 \theta. \quad (16)$$

37 The condition (13) is called by Thomsen (1986) the elliptic condition, since, in the absence
 38 of the $\sin^2 2\theta$ term, the forms of the wave velocity surfaces as a function of incidence angle θ
 39 are elliptical with only a $\sin^2 \theta$ dependence. When condition (13) is not satisfied the presence
 40 of the $\sin^2 2\theta$ term means that the wavesurfaces can be either convex or concave. (The convex-
 41 ity/concavity of the P velocity is in the opposite sense to that of the SV velocity. This is an
 42 explicit consequence of the presence of the \sqrt{S} term in (6) and (7) with opposite signs.)

43 Thus if we were to introduce an additional parameter to characterize the incidence angle
 44 dependence of bodywaves, one reasonable choice may be

$$\eta_\kappa = \frac{F + L}{(A - L)^{1/2}(C - L)^{1/2}}, \quad (17)$$

45 and $\eta_\kappa = 1$ for the isotropic case.

46 Further considering

$$(A - L)(C - L) = \left(\frac{A + C}{2} - L\right)^2 - \left(\frac{A - C}{2}\right)^2,$$

47

$$\eta_{\kappa'} = \frac{F + L}{\frac{A + C}{2} - L} \quad (18)$$

48 may be another possibility that might make sense by looking at equation (12).

49 One of the good points of η_{DLA} is that it is simple and depends on just A and not C .
 50 Assuming P-wave anisotropy is small, if we substitute $\frac{A + C}{2}$ in (18) by A , we get

$$\eta_{\kappa''} = \frac{F + L}{A - L} \quad (19)$$

51 It is instructive how these parameters (η 's) behave when both P- and S-wave anisotropy is
 52 absent (i.e., $A = C$ and $L = N$). When these conditions are satisfied,

$$\begin{aligned} \rho v_P^2(\theta) &= \frac{(L + A) + \sqrt{S}}{2} \\ \rho v_{SV}^2(\theta) &= \frac{(L + A) - \sqrt{S}}{2} \\ \rho v_{SH}^2(\theta) &= L, \end{aligned}$$

53 and

$$S = \{(A - L)\}^2 + \{(F + L)^2 - (A - L)^2\} \sin^2 2\theta,$$

54 and $\sin^2 \theta$ dependence disappears. In this case, η_κ , $\eta_{\kappa'}$, and $\eta_{\kappa''}$ reduce to the same form. Also,
 55 it is easy to see that $\eta_{DLA} = 1$ gives the elliptic condition, and so in this sense, $\eta_{DLA} - 1$
 56 becomes a measure of a departure from the elliptic condition to dictate the convex/concave
 57 pattern.

58 For more general case, $\chi = \eta_{DLA} - 1$ is small for weak anisotropy,

$$\chi = \eta_{DLA} - 1 = \frac{F - A + 2L}{A - 2L}. \quad (20)$$

59 Similarly

$$\chi'' = \eta_{\kappa''} - 1 = \frac{F - A + 2L}{A - L} = \chi \times \frac{A - 2L}{A - L}, \quad (21)$$

60 and as long as $A - L > A - 2L > 0$ is satisfied, χ'' has the same sign as χ , and $\chi > \chi''$, indicating
 61 χ'' is also small. So in this respect, if anisotropy is weak (especially in P), η_{DLA} might be a
 62 good proxy for η_κ whose departure from unity provides a measures of the deviation from elliptic
 63 anisotropy and dictates the convex/concave pattern of the incidence angle dependence of v_P
 64 and v_{SV} .

65 Thomsen's parameters

66 Thomsen (1986) introduced three parameters for VTI system, now referred to as Thomsen's
67 parameters, and they are defined as

$$\varepsilon = \frac{A - C}{2C} = \frac{1}{2}(\varphi^{-1} - 1) \quad (22)$$

$$\gamma = \frac{N - L}{2L} = \frac{1}{2}(\xi - 1) \quad (23)$$

$$\delta = \frac{(F + L)^2 - (C - L)^2}{2C(C - L)}, \quad (24)$$

68 which are all small for weak anisotropy. While ε and γ are directly related to φ and ξ respec-
69 tively as shown above and thus to P- and S-wave anisotropy, δ was introduced such that it
70 dominates v_P in the case of near vertical incidence as in reflection profiling.

71 Considering that $\delta = \varepsilon$ is their condition for elliptical anisotropy, examination of $\varepsilon - \delta$ leads
72 to

$$\varepsilon - \delta = \frac{A - C}{2C} - \frac{(F + L)^2 - (C - L)^2}{2C(C - L)} \quad (25)$$

$$= \frac{(A - L)(C - L) - (F + L)^2}{2C(C - L)} \quad (26)$$

$$= (1 - \eta_\kappa^2) \frac{A - L}{2C}, \quad (27)$$

73 and we now see the connection between Thomsen's δ and η_κ introduced here. If η_{DLA} were a
74 proxy of η_κ for weak anisotropy, we might be able to say that a connection between η_{DLA} and
75 Thomsen's δ is established.

76 For weak anisotropy, the incidence angle dependence of bodywaves are, according to Thom-
77 sen (1986),

$$v_P(\theta) = \alpha_H (1 + \delta \sin^2 \theta \cos^2 \theta + \varepsilon \sin^4 \theta) \quad (28)$$

$$v_{SV}(\theta) = \beta_V \left[1 + \frac{\alpha_H^2}{\beta_V^2} (\varepsilon - \delta) \sin^2 \theta \cos^2 \theta \right] \quad (29)$$

$$v_{SH}(\theta) = \beta_V (1 + \gamma \sin^2 \theta), \quad (30)$$

78 and when the elliptic condition is satisfied

$$v_P(\theta) = \alpha_H (1 + \varepsilon \sin^2 \theta)$$

$$v_{SV}(\theta) = \beta_V$$

$$v_{SH}(\theta) = \beta_V (1 + \gamma \sin^2 \theta),$$

79 which show simple incidence angle dependences.

80 (28)(29)(30) may be expressed in terms of 2θ and 4θ to make the incidence angle dependence
81 more explicit:

$$v_P(\theta) = \alpha_H \left[1 + \frac{\varepsilon}{2}(1 - \cos 2\theta) - \frac{\omega}{2}(1 - \cos 4\theta) \right] \quad (31)$$

$$v_{SV}(\theta) = \beta_V \left[1 + \frac{\alpha_H^2 \omega}{\beta_V^2} \frac{\omega}{2}(1 - \cos 4\theta) \right] \quad (32)$$

$$v_{SH}(\theta) = \beta_V \left[1 + \frac{\gamma}{2}(1 - \cos 2\theta) \right], \quad (33)$$

82 where $\omega = (\varepsilon - \delta)/4$ is introduced. These equations show that $(\varepsilon - \delta)$ dictates the con-
83 vex/concave nature (i.e, $\cos 4\theta$ dependence) of v_P and v_{SV} .

84 η_{DLA} and η_{κ} for weakly anisotropic models

85 To finish up this short note, we compare distributions of η -related parameters for some of
 86 weakly anisotropic cases.

87 Millefeuille (isotropic layers) case

88 In the first example, we present a series of VTI models constructed by the Backus averaging
 89 (Backus, 1962) of a stack of two kinds of homogeneous isotropic layers: soft layers embedded in
 90 a background solid matrix (e.g., Kawakatsu et al., 2009). We parameterize (i) the proportional
 91 reduction of rigidity of soft layers to the background by a ($0 \leq a \leq 1$), (ii) the proportional
 92 reduction of the bulk modulus by $a/2$, and (iii) the volume fraction of soft layers by f ($0 \leq$
 93 $f \leq 1$). Both a and f are varied in intervals of 0.05. Figure 1(a) compares η_{κ} with η_{DLA}
 94 (blue circles) or $\eta_{\kappa'}$ (magenta crosses). While η_{κ} and $\eta_{\kappa'}$ give almost the same values, η_{DLA}




Figure 1: Comparison of η -related parameters for various weakly anisotropic models. (a) Millefeuille case, (b) general TI case, and (c) rotated A-type olivine case. Green asterisks correspond to η_{κ} vs. η_{DLA} for (a) $a = 0.9$, $f = 0.01$, (b) peak-to-peak anisotropy for both P and S waves is 1.5% with $\eta_{DLA} = 0.9$, 1.0, and (c) the A-type olivine fabric case whose fast axis lies in the horizontal plane, all for which examples of incident angle dependency of bodywaves are shown in Figure 2.

95 gives slightly smaller values. As $\eta_\kappa \leq 1$ is guaranteed (Berryman, 1979), all values appear
96 generally less than 1. Although η_{DLA} in this case slightly deviates from η_κ , nearly one-to-one
97 correspondence may be observed to make η_{DLA} a reasonable proxy to η_κ .

98 General case

99 For a more general case, we construct a series of VTI models which have a maximum of $\pm 5\%$
100 anisotropy in both $\alpha_{V,H}$ and $\beta_{V,H}$, and $0.5 < \eta_{DLA} < 1.5$ (Figure 1(b)). While η_κ and $\eta_{\kappa'}$ give
101 almost the same values, η_{DLA} deviates significantly from the corresponding η_κ .

102 A-type olivine case

103 As a third example, we construct a series of VTI models by azimuthal averaging (Montagner
104 and Nataf, 1986; Montagner and Anderson, 1989) of an arbitrarily rotated A-type olivine fabric
105 (Jung et al., 2006) (Figure 1(c)) (rotation is done with a 30-degree interval for each Euler angle).
106 In a similar way to the preceding cases, η_κ and $\eta_{\kappa'}$ have almost the same values, but η_{DLA}
107 deviates from corresponding η_κ .

108 Examples of the incidence angle dependence of representative VTI models (denoted by
109 green asterisks in Figure 1) are shown in Figure 2. Note that the convex pattern of v_{SV}
110 velocity occurs when $\eta_\kappa < 1$.

111 Discussion

112 The incidence angle dependence of bodywave phase velocities in a radially anisotropic system
113 has not been discussed much in the geophysical literature as it is a difficult effect to observe.
114 In the laboratory, on the other hand, the simple $\sin \theta$ and $\sin 2\theta$ dependence (e.g., (6) and (11))
115 has been used to obtain the fifth elastic constant from measurement along the angle 45 degrees
116 from the symmetric axes (e.g., Christensen and Crosson, 1968; Anderson, 1966). Song and
117 Kawakatsu (2012, 2013) recently suggested that such incident angle dependency in the Earth
118 may be constrained at subduction zones where the dip of the lithosphere/asthenosphere changes
119 along with the subduction, affecting the effective incidence angle of teleseismic bodywaves to
120 the system. If such analyses can be made generally, the new parameter η_κ (or $\eta_{\kappa'}$) might be
121 a useful tool in global seismology to characterize VTI (radially anisotropic) systems. How
122 η -related parameters might be constrained from Rayleigh wave dispersion needs also to be
123 understood (e.g., Anderson, 1966).


124 Acknowledgments

125 This work was conducted while the first author was a visiting professor at IPGP in 2014. He
126 thanks the members of IPGP, especially those of the seismology section for hospitality. We
127 thank Brian Kennett and Thorsten Becker for their excellent and prompt reviews, and the
128 editor, Gillian Foulger for her judgement. It is unfortunate that Professor Anderson's decease
129 robbed us of the opportunity to discuss with him directly how he came up with his η .


References

- 130
131 Anderson, D.L., 1961. Elastic wave propagation in layered anisotropic media. *J. Geophys. Res.*
132 9, 2953–2963.
- 133 Anderson, D.L., 1966. Recent evidence concerning the structure and composition of the Earth's
134 mantle, In *Physics and Chemistry of the Earth*. Pergamon, Oxford. volume 6. pp. 1–131.
- 135 Anderson, D.L., 1989. *Theory of the Earth*. Blackwell, Boston.
- 136 Backus, G.E., 1962. Long-wave elastic anisotropy produced by horizontal layering. *J. Geophys.*
137 *Res.* 67, 4427–4440.
- 138 Berryman, J.G., 1979. Long-wave elastic anisotropy in transversely isotropic media. *Geophysics*
139 44, 896–917.
- 140 Christensen, N., Crosson, R.S., 1968. Seismic anisotropy in the upper mantle. *Tectonophysics*
141 6, 93–107.
- 142 Dziewonski, A.M., Anderson, D.L., 1981. Preliminary reference Earth model. *Phys. Earth*
143 *Planet. Inter.* 25, 297–356.
- 144 Jung, H., Katayama, I., Jian, Z., Hiraga, T., Karato, S., 2006. Effect of water and stress on
145 the lattice-preferred orientation of olivine. *Tectonophysics* 421, 1–22.
- 146 Kawakatsu, H., Kumar, P., Takei, Y., Shinohara, M., Kanazawa, T., Araki, E., Suyehiro, K.,
147 2009. Seismic Evidence for Sharp Lithosphere-Asthenosphere Boundaries of Oceanic Plates.
148 *Science* 324, 499–502.
- 149 Love, A.E.H., 1927. *A treatise on the mathematical theory of elasticity*. Dover, New York.
- 150 Montagner, J.P., Anderson, D. L., 1989. Constraints on elastic combinations inferred from
151 petrological models. *Phys. Earth Planet. Inter.* 54, 82–105.
- 152 Montagner, J.P., Nataf, H. C., 1986. On the inversion of the azimuthal anisotropy of surface
153 waves. *J. Geophys. Res.* 91, 511–520.
- 154 Song, T.R.A., Kawakatsu, H., 2012. Subduction of oceanic asthenosphere: evidence from
155 sub-slab seismic anisotropy. *Geophys. Res. Lett.* 39, L17301. doi:10.1029/2012GL052639.
- 156 Song, T.R.A., Kawakatsu, H., 2013. Subduction of oceanic asthenosphere: A Critical Appraisal
157 in Central Alaska. *Earth Planetary Science Lett.* 367, 82–94.
- 158 Takeuchi, H., Saito, M., 1972. Seismic surface waves. In *Seismology: Surface Waves and Earth*
159 *Oscillations, Methods in Computational Physics*. Academic Press, New York. volume 11. pp.
160 217–295.
- 161 Thomsen, L., 1986. Weak elastic anisotropy. *Geophysics* 51, 1954–1966.

MF, $a = 0.9$, $f = 0.01$, $R_p = 0.54\%$, $R_s = 3.9\%$, $\eta_{DLA} = 0.889$, $\eta_\kappa = 0.941$



GE, $R_p = 3\%$, $R_s = 3\%$, $\eta_{DLA} = 0.9$, $\eta_\kappa = 1$, $\eta_\kappa = 0.983$, $\eta_\kappa = 1.04$



Averaged A-type, $R_p = 4\%$, $R_s = 1.7\%$, $\eta_{DLA} = 0.9$, $\eta_\kappa = 0.997$




Figure 2: Examples of the incidence angle dependency of bodywaves for the VTI models represented by asterisks in Figure 1. Blue (and magenta in middle) solid lines, red (and cyan) dash-dot lines, and green dashed lines are respectively for v_P , v_{SV} , and v_{SH} . Phase velocities are scaled by those of corresponding reference isotropic models. (Top), (middle), and (bottom) correspond to the models in (a), (b) and (c) in Figure 1. In the middle panel, v_P and v_{SV} shown by magenta and cyan lines are for $\eta_{DLA} = 1$, $\eta_\kappa = 1.04$ case, and v_{SH} behaves the same for two cases.