



ELSEVIER

Contents lists available at ScienceDirect

Free Radical Biology and Medicine

journal homepage: www.elsevier.com/locate/freeradbiomedPPAR γ as a therapeutic target to rescue mitochondrial function in neurological diseaseJuan Carlos Corona^{a,b}, Michael R. Duchen^{a,*}^a Department of Cell and Developmental Biology, University College London, London WC1E 6BT, United Kingdom^b Laboratory of Neurosciences, Hospital Infantil de México Federico Gómez, Mexico City, Mexico

ARTICLE INFO

Article history:

Received 7 March 2016

Received in revised form

20 June 2016

Accepted 21 June 2016

Available online 25 June 2016

Keywords:

PPAR γ agonists

Mitochondrial function

Neurodegenerative disorders

Neuroprotection

ABSTRACT

There is increasing evidence for the involvement of mitochondrial dysfunction and oxidative stress in the pathogenesis of many of the major neurodegenerative and neuroinflammatory diseases, suggesting that mitochondrial and antioxidant pathways may represent potential novel therapeutic targets. Recent years have seen a rapidly growing interest in the use of therapeutic strategies that can limit the defects in, or even to restore, mitochondrial function while reducing free radical generation. The peroxisome proliferation-activated receptor gamma (PPAR γ), a ligand-activated transcription factor, has a wide spectrum of biological functions, regulating mitochondrial function, mitochondrial turnover, energy metabolism, antioxidant defence and redox balance, immune responses and fatty acid oxidation. In this review, we explore the evidence for potential beneficial effects of PPAR γ agonists in a number of neurological disorders, including Parkinson's disease, Alzheimer's disease, Amyotrophic lateral sclerosis and Huntington's disease, ischaemia, autoimmune encephalomyelitis and neuropathic pain. We discuss the mechanisms underlying those beneficial effects in particular in relation to mitochondrial function, antioxidant defence, cell death and inflammation, and suggest that the PPAR γ agonists show significant promise as therapeutic agents in otherwise intractable neurological disease.

© 2016 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

1. Introduction

The peroxisome proliferator-activated receptor (PPAR) was originally cloned from liver peroxisomes, which are organelles that participate in the metabolism of fatty acids [77]. PPARs are ligand-

inducible transcription factors that belong to the hormone nuclear receptor superfamily. PPAR activation is involved in cellular differentiation and proliferation, angiogenesis, vascular protection and regulation of blood pressure. PPAR activation is also anti-apoptotic, anti-oxidant, anti-inflammatory, and limits cancer metastasis [13,19,24,41,44,129,160]. In addition to these effects, PPARs also play key roles in the regulation of metabolism, regulating insulin sensitivity, mitochondrial biogenesis, and glucose and lipid homeostasis [15,23,168]. As lipid sensors, PPARs modulate metabolism in response to dietary lipid intake and direct lipid storage and metabolism [106].

Three members of the PPAR family have been cloned, encoded by three distinct genes located on human chromosomes 22, 6 and 3: PPAR α (NR1C1), PPAR β/δ (NR1C2), and PPAR γ (NR1C3) respectively [48]. While the three PPAR isoforms share a high degree of structural similarity, they have distinct physiological roles, ligand specificity and tissue distribution. Mutations of each of the PPAR isoforms has been associated with hereditary disease. Some mutations of the PPARs lead to a loss of function, resulting in familial partial lipodystrophy, acanthosis nigricans or insulin resistance [10,26,173]. Dysfunction of each of the three PPAR isoforms has been specifically implicated in dysfunction in the central nervous system (CNS), suggesting key roles for these pathways in maintaining the integrity of the nervous system.

Abbreviations: PPAR γ , peroxisome proliferation-activated receptor gamma; 15D-PGJ₂, 15-deoxy- $\Delta^{12,14}$ -Prostaglandin J₂; TZDs, thiazolidinediones; CNS, central nervous system; WAT, white adipose tissue; BAT, brown adipose tissue; SRC, steroid receptor coactivator; CBP, CREB-binding protein; AF-1, transcriptional activation domain; PGC1 α , peroxisome proliferator-activated receptor gamma coactivator 1 α ; TFAM, mitochondrial transcription factor A; mtDNA, mitochondrial DNA; UCP-1, uncoupling protein 1; UCP-2, uncoupling protein 2; NRF1, nuclear respiratory factor 1; NRF2, nuclear respiratory factor 2; ROS, reactive oxygen species; SOD1, superoxide dismutase 1; TNF α , tumour necrosis factor α ; Bcl-2, B-cell lymphoma2; Bax, Bcl-2-associated X protein; MPP⁺, 1-methyl-4-phenylpyridinium ion; iNOS, inducible nitric oxide synthase; COX-2, cyclooxygenase-2; LPS, lipopolysaccharide; $\Delta\Psi_m$, mitochondrial membrane potential; ALS, amyotrophic lateral sclerosis; HD, huntington's disease; AD, alzheimer's disease; PD, parkinson's disease; MPTP, 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine; Nrf2, nuclear factor erythroid-derived 2-like 2; NF- κ B, nuclear factor- κ B; HO-1, haem oxygenase-1; TDP-43, TAR DNA-binding protein 43; BDNF, brain derived neurotrophic factor; A β , protein amyloid- β ; GSK-3 β , glycogen synthase kinase-3 β ; Cdk5, cyclin-dependent kinase 5; 6-OHDA, 6-hydroxydopamine

* Corresponding author.

E-mail address: m.duchen@ucl.ac.uk (M.R. Duchen).<http://dx.doi.org/10.1016/j.freeradbiomed.2016.06.023>0891-5849/© 2016 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

Table 1
Natural and synthetic ligands, physiological role, and tissue expression of PPAR γ .

Natural ligands	Refs.	Synthetic ligands	Refs.	Tissue expression	Refs.	Physiological role	Refs.
Unsaturated Fatty Acids	[44]	Pioglitazone	[88]	High-Adipose tissue (white and brown)	[1,155]	Glucose homeostasis	[42,133]
Prostaglandin A1	[8]	Rosiglitazone	[96]	Low-Intestines, Liver, Retina	[16]	Lipid storage	[155]
15-deoxy- $\Delta^{12,14}$ -PGJ ₂	[180]	Troglitazone	[96]	Low-Heart	[16]	Differentiation and maturation of adipocytes	[155]
Nitroalkenes	[140]	Ciglitazone	[122]	Low-Muscle	[162]	Fatty acid oxidation	[14,15]
Oxidized phospholipids	[39]	Glitazars	[2]	Low-Brain (oligodendrocytes, neurons, astrocytes, microglia)	[61,109]		
		PPAR γ modulators	[177]				
		Non-steroidal anti-inflammatory drugs	[95]				




Fig. 1. Chemical structures of TZD, 15-deoxy- $\Delta^{12,14}$ -PGJ₂, Rosiglitazone, Pioglitazone, Troglitazone and the antagonist GW9662.

PPAR α and PPAR δ are regulators of fatty acid oxidation in a wide variety of tissues [20,124]. PPAR α was cloned as the molecular target of fibrates, cholesterol-lowering compounds that increase hepatic fatty acid oxidation [77]. The expression pattern of PPAR α is restricted to tissues with a high capacity for fatty acid oxidation, such as heart, liver, brown adipose tissue (BAT), and oxidative muscle [16]. The function of PPAR α seems to be more restricted to fatty acid uptake, beta-oxidation, and ketogenesis. In contrast, PPAR δ is ubiquitously expressed, with higher levels in the digestive tract, heart, and BAT [52]. PPAR δ plays a role in controlling oxidative metabolism and fuel preference, mainly in relation to fatty acid oxidation, mitochondrial OXPHOS, and glucose utilization [92,98].

PPAR γ is probably the most intensively studied of the three isoforms [160]. Knockout of PPAR γ is embryonically lethal [7,143]. PPAR γ activation is associated with increased carbohydrate and lipid metabolism, reduction of plasma glucose concentration and stimulation of adipocyte differentiation [53]. PPAR γ agonists are currently in clinical use for the management of type 2 diabetes [11,168]. The physiological roles of PPAR γ are highlighted in Table 1.

All PPARs can be activated by polyunsaturated fatty acids, but with different affinities [11,44,87]. Specific PPAR γ ligands are divided into two major groups: [26,34,163]: (i) Endogenous or natural agonists and (ii) synthetic agonists. The endogenous agonists are divided into four subgroups: (i.1) The eicosanoids prostaglandin-A1 [8] and the cyclopentenone prostaglandin 15-deoxy- $\Delta^{12,14}$ -Prostaglandin J₂ (15D-PGJ₂), which is the most naturally occurring and is commonly used as a potent ligand for PPAR γ [180]; (i.2) unsaturated fatty acids such as Linoleic acid and Docosahexaenoic acid; (i.3) Nitroalkenes such as Nitrated oleic acid and Linoleic acid [140]; and (i.4). oxidized phospholipids such as hexadecyl azelaoyl phosphatidylcholine and Lysophosphatidic acid [39]. Non-steroidal anti-inflammatory drugs, as flufenamic acid, ibuprofen, fenopfen, and indomethacin A, are also PPAR γ ligands [11,12,95,178].

(ii) The synthetic agonists are divided into five subgroups: (ii.1) The thiazolidinedione (TZD) family, which include Rosiglitazone, Pioglitazone and Troglitazone [88,96]; (ii.2) the non-TZD agonists, such as Ciglitazone [122], Netoglitazone and Rivoglitazone; (ii.3) the dual α/γ agonists [2] also called glitazars (Include: Aleglitazar, Muraglitazar, Tesaglitazar and Saroglitazar); (ii.4) pan $\alpha/\delta/\gamma$ agonists like bezafibrate; (ii.5) The selective PPAR γ modulators (SPPAR γ M) such as metaglidasen/halofenate, PA-082, INT₁₃₁ besylate and the angiotensin receptor antagonists [177]. The natural and synthetic ligands of PPAR γ are listed in Table 1.

The TZDs were the first ligand family discovered to bind and activate PPAR γ with therapeutic benefit for diabetic patients [96]. However, the TZDs do have some adverse effects in man, including increased risk of heart failure, weight gain, increased cardiac load, peripheral oedema and haemodilution [136,170]. While no endogenous antagonists for PPAR γ have been reported, some synthetic antagonists are available, including GW9662, Bisphenol A Diglyceryl ether, LG100641, SR-202, and PDO68235. The structure of some PPAR γ agonist and antagonist are shown in Fig. 1.

2. Distribution of PPAR γ

PPAR γ is distributed in a number of tissues and cell types [16,43,133], demonstrated by in situ hybridization and by immunohistochemistry both in adult rat and in rat embryonic development [16,17]. In man, PPAR γ mRNA expression was abundant in adipose tissue, while lower level expression was seen in skeletal muscle [162]. PPAR γ protein and mRNA were expressed at low levels in the adult rodent CNS, including neurons, astrocytes, microglia and oligodendrocytes [12,16,38,61,109]. High levels of expression of PPAR γ have been found in discrete areas of the CNS, including the stellate cells of the cerebellar cortex, the olfactory tubercle, the piriform cortex, in rhomboid, centromedial, and parafascicular thalamic nuclei and in the reticular formation [61,109]. PPAR γ is highly expressed in dopaminergic cells in the basal ganglia [38,109]. Other studies have demonstrated the presence of PPAR γ protein by Western blot and immunohistochemistry in the midbrain [22,120].

In cortical neurons, PPAR γ immunoreactivity appears mainly as nuclear labelling, although cytoplasmic staining is sometimes detectable [109]. PPAR γ is present at lower levels in liver, skeletal muscle and heart, but it is abundantly expressed in white adipose tissue. PPAR γ is expressed in brown adipose tissue of rodents and in the CNS during foetal development [1,17,107]. The tissue expression of PPAR γ is shown in Table 1.

3. PPAR γ , adipose tissue, obesity and type 2 diabetes

The highest levels of PPAR γ are expressed in adipose tissue

[25,154]. The expression and activation of PPAR γ is sufficient to induce adipogenesis [155]. Adipose tissue can be divided into two major depots, white adipose tissue (WAT) and brown adipose tissue (BAT). In both WAT and BAT, PPAR γ is a key regulator of differentiation. PPAR γ plays an important role in regulating lipid metabolism in mature adipocytes. WAT stores excess energy as triglycerides, which can be mobilized by lipolysis to generate free fatty acids for use by other tissues, while BAT is the site of non-shivering thermogenesis, through activation of the brown adipocyte-specific protein, uncoupling protein 1 (UCP1). TZDs appear to co-ordinately activate gene expression leading to an increase in net lipid partitioning into adipocytes [157]. Target genes directly regulated by PPAR γ , include lipoprotein lipase [139], fatty-acid transport protein [58], and oxidized LDL receptor 1 [31], favouring adipocyte uptake of phosphoenolpyruvate carboxykinase [156,158], glycerol kinase [68], the glycerol transporter aquaporin 7 [73] and the circulating fatty acids.

Adipose tissue has generally been considered an organ used for storage and release of energy. Recent years have seen an explosive increase in the incidence of obesity worldwide, driving a resurgence of interest into the biology of adipose tissue, and of the adipocyte as regulators of body energy homeostasis. PPAR γ is a master regulator in the formation of fat cells and their normal function in the adult [134,135]. PPAR γ is induced during adipocyte differentiation, and forced expression of PPAR γ in non-adipogenic cells effectively drives their differentiation into mature adipocytes [155,157]. Adipose tissue accounts for about 10% of insulin-stimulated glucose disposal and has a key role in directing whole-body glucose homeostasis. The discovery that the insulin-sensitizing TZDs are potent agonists for PPAR γ [55,96], suggested that apart from being a fatty acid sensor, PPAR γ might serve an important role as a regulator of glucose metabolism. TZDs act by inhibiting hepatic glucose output and increasing insulin action to stimulate muscle glucose disposal in insulin-resistant animal models and patients [157].

Adipose tissue is dysfunctional in obesity, leading to progressive insulin resistance and risk for the development of type 2 diabetes. Phosphorylation of PPAR γ at serine 273 by cyclin-dependent kinase 5 (Cdk5) stimulates diabetogenic gene expression in adipose tissues [30]. Inhibition of this modification is a key therapeutic mechanism of the TZDs. Insulin signalling activates the Akt/PI3K and Grb2/Ras/MEK/ERK kinase cascades. Recently it was demonstrated that ERK directly phosphorylates serine 273 of PPAR γ and Cdk5 suppresses ERKs through direct action on a novel site in MEK, the ERK kinase. The pharmacological inhibition of MEK and ERK markedly improves insulin resistance in both obese and wild type ob/ob mice, and reversed the effects of the Cdk5 ablation, indicating that the ERK/Cdk5 axis controls the diabetogenic actions of PPAR γ [5].

3.1. PPAR γ coactivators

The binding of PPAR γ to specific DNA sequences requires heterodimerization with a second member of the nuclear receptor family, the retinoic X receptor. Binding of agonist ligands to PPAR γ triggers a conformational change that attracts transcriptional coactivators, which exist in multiprotein complexes and include members of the steroid receptor coactivator (SRC) family [103,104]. SRC1 was originally identified as a coactivator of the steroid receptor superfamily [118]. SRC1 also interacts with PPAR γ via LXXLL motifs in a ligand dependent manner [46]. The effects of SRC1 and SRC2 on metabolism are predominantly mediated by PPAR γ coactivator 1 α (PGC1 α). PGC1 α is upregulated in SRC2-depleted BAT. SRC1 and SRC2 control the energy balance between BAT and WAT. In BAT, SRC1 promotes energy expenditure via fatty acid oxidation [181]. SRC2 represses this process by activation of

PPAR γ in WAT, producing an increase in triglyceride accumulation in WAT and a decrease in free fatty acids.

PGC1 α is the master regulator of mitochondrial biogenesis and plays a central role in coordinating and driving energy metabolism, fatty-acid oxidation, gluconeogenesis, glucose transport, glycogenolysis, peroxisomal remodelling, and oxidative phosphorylation [125,126]. PGC1 α also integrates and coordinates the activity of multiple transcription factors, such as nuclear respiratory factor 1 and 2 (NRF1–2), PPAR α and mitochondrial transcription factor A (TFAM), which are all involved in mitochondrial biogenesis [47,127].

PGC1 α was originally identified in a two-hybrid screen using murine BAT cDNA and PPAR γ as bait [127]. The interaction with PPAR γ is ligand-independent and mediated through the N-terminal domain and an LXXLL motif [97,127]. PGC1 α also targets other transcription factors involved in metabolism, including FOXO1, SREBP, Foxa2 and Sox9 [85,126,171]. PGC1 α works in concert with the oestrogen-related receptors (ERR α , ERR β , ERR γ) which are members of the nuclear hormone receptor family which promote mitochondrial biogenesis in response to hormonal signals [47]. As well as promoting mitochondrial biogenesis, PGC1 α also regulates the expression of several enzymes involved in antioxidant defence, detoxifying reactive oxygen species (ROS). These include superoxide dismutases 1 and 2 (SOD1–2), catalase and glutathione peroxidase-1 [146].

Other nuclear receptor coactivators are the histone acetyltransferases p300 and CREB-binding protein (CBP) [6,117]. Both proteins interact with the ligand binding domain of PPAR γ in a ligand- and LXXLL-dependent manner, while at the same time they interact with the transcriptional activation domain (AF-1) of PPAR γ [64]. p300 and CBP enhance the transcriptional activity of both the AF-1 and AF-2 domains separately. CBP and p300 are also involved in adipogenesis.

PRDM16 is a BAT-specific transcription factor also involved in mitochondrial biogenesis [142]. It is a zinc finger protein which was first identified at a chromosomal breakpoint in myeloid leukaemia [116]. Overexpression of PRDM16 in cultured white pre-adipocytes resulted in activation of the brown adipose differentiation programme, the upregulation of mitochondrial genes, increase of mitochondrial biogenesis and uncoupled respiration [142]. Therefore, PRDM16 binds to PPAR γ in a ligand-independent manner, works as a DNA binding transcription factor and enhances the transcriptional activity of PPAR γ . PPAR γ activation is required for the adipogenic function of PRDM16.

4. Actions of PPAR γ in rescuing mitochondrial function

Growing evidence suggests the involvement of mitochondrial dysfunction in the pathogenesis of many diseases, and especially in a number of major neurodegenerative and neuroinflammatory disorders. This dysfunction is associated with defects in mitophagy, in bioenergetics, in mitochondrial fusion/fission and mitochondrial trafficking, frequently also in combination with oxidative stress. Therefore, pharmacological pathways that promise to enhance mitochondrial function, reduce generation of reactive oxygen species (ROS) or enhance antioxidant defence represent important new therapeutic prospects for these diseases.

4.1. Beneficial actions of PPAR γ on mitochondrial function

TZDs have been shown in many model systems to upregulate mitochondrial oxidative phosphorylation, mitochondrial biogenesis and antioxidant defence, serving together to protect cells from various forms of injury. Thus, TZDs were shown to protect lymphocytes from apoptosis following growth factor withdrawal,

preventing the loss of mitochondrial membrane potential ($\Delta\Psi_m$). The survival-enhancing effects were dependent on both the presence and activation of PPAR γ [165]. During the development of obesity, in white adipocytes from ob/ob mice, which were treated with rosiglitazone showed mitochondrial remodelling, enhanced oxygen consumption and increased energy expenditure in white fat, suggesting that enhanced lipid utilization in this tissue may affect whole-body energy homeostasis and insulin sensitivity [169]. In human adipocytes, both TZDs rosiglitazone and pioglitazone significantly increased mitochondrial DNA (mtDNA) copy number and expression of factors involved in mitochondrial biogenesis, including PGC1 α and TFAM while also increasing the expression of genes in the fatty acid oxidation pathway, including carnitine palmitoyltransferase-1, malonyl-CoA decarboxylase, increased the expression of UCP-1 and medium-chain acyl-CoA dehydrogenase [14,15]. A similar response was described in differentiated 3T3-L1 and C3H/10T1/2 adipocytes treated with rosiglitazone, which showed increased mitochondrial biogenesis, oxygen consumption and increased mitochondrial citrate synthase activity [132]. Additionally, pioglitazone stabilized MitoNEET, a protein that regulates oxidative capacity in 3T3-L1 cell line, rat brain mitochondria and COS-7 cells [33,119,167].

Protection by TZDs has particularly been explored in relation to the CNS, using a variety of model systems. Treatment with TZDs protected cortical astrocytes in culture against cell death induced by glucose deprivation, increasing glucose utilization, lactate production and mitochondrial membrane potential ($\Delta\Psi_m$), demonstrating modification of astrocyte metabolism and mitochondrial function following PPAR γ activation [42]. Pioglitazone and rosiglitazone prevented the death of differentiated SH-SY5Y cells (a neuroblastoma derived cell line), following glucose deprivation by promoting mitochondrial biogenesis, indicated by the up-regulation of PGC1 α , NRF1, TFAM, cytochrome c oxidase subunit CO-I and CO-IV, and an increased mtDNA copy number, while there was

no change in $\Delta\Psi_m$ [108].

In the human neuron-like NT2 cell line, pioglitazone treatment also increased mtDNA content and the expression of nuclear-encoded electron transport chain subunit proteins. Oxygen consumption was increased as well as complex I and complex IV V (max) activities. Pioglitazone treatment was also associated with increased cytoplasmic, but reduced generation of mitochondrial, peroxide [65]. The natural ligand 15D-PGJ₂ and rosiglitazone, both PPAR γ ligands, increased glucose metabolism in rat cortical slices and astrocytes, restored brain ATP levels and prevented the impairment of glutamate uptake induced by repeated stress in rat brain [60]. In a more recent study, the PPAR γ agonist ciglitazone protected the mitochondrial population of hippocampal neurons to oxidative stress (exposure to hydrogen peroxide). Protection was attributed to the modulation of mitochondrial fusion-fission events, suggesting that a PPAR γ based therapy could have beneficial effects mediated by simultaneous actions on several complementary cellular pathways [182]. An schematic overview of PPAR γ mechanisms is shown in Fig. 2.

4.2. Anti-oxidant and anti-apoptotic properties of PPAR γ

Besides the protective effects of PPAR γ on mitochondrial function, the protective effects of PPAR γ activation have also been attributed to their anti-oxidant and anti-apoptotic properties. Thus, in hepatocytes, rosiglitazone enhanced the antioxidant defence of the cells, increasing the rate of elimination of ROS while increasing the expression of nuclear factor erythroid-derived 2-like 2 (Nrf2) and the antioxidant enzyme haem oxygenase-1 (HO-1) through the PPAR γ -pathway [164].

Rosiglitazone was protective against neuronal damage by acetaldehyde, which causes oxidative neuronal death, inducing the generation of intracellular ROS and ultimately causing apoptotic cell death. The protection by rosiglitazone was attributed to the




Fig. 2. Activation of PPAR γ increases mitochondrial biogenesis, oxygen consumption, $\Delta\Psi_m$, antioxidant defences, regulates autophagy, and increases transcription factors, like PGC1 α , NRF1–2, TFAM. Also PPAR γ agonists regulate apoptosis and reduce inflammation. Thus, PPAR γ agonists like TZDs regulate the expression of several target genes involved in neuronal survival and protected against neuronal death in several neurodegenerative disorders, mainly improving mitochondrial function and increasing redox capacity.

induction of the expression of anti-oxidant enzymes including SOD and catalase and by regulating the expression of B-cell lymphoma 2 (Bcl-2) and Bcl-2-associated X protein (Bax) [83]. In another study, treatment with pioglitazone was neuroprotective in rats treated with lipopolysaccharide (LPS), decreasing degeneration of dopaminergic pathways, decreasing inflammation and restoring mitochondrial function while also attenuating oxidative stress [74]. Rosiglitazone or PPAR γ overexpression protected the neuronal N2A cell line against oxygen-glucose deprivation by maintaining $\Delta\Psi_m$ and preventing apoptosis [172].

Rosiglitazone treatment also protected the neuroblastoma derived cell line, SH-SY5Y cells, against cytotoxicity induced by chlorpyrifos (an acetylcholinesterase inhibitor, widely used as pesticide). The protection was apparently mediated again by attenuation of oxidative stress and by anti-apoptotic effects, as well as the inhibition of the inflammatory cascade via inactivation of signalling by p38, nuclear factor- κ B (NF- κ B) and c-Jun N-terminal kinase [94].

Finally, in a model of bronchial asthma, oral treatment with rosiglitazone significantly increased anti-oxidant activity in lung tissue, associated with improved lung function, improved histopathological features and a significant decrease in the serum levels of IL-5 and IgE [51].

5. Mechanisms of protection mediated by PPAR γ

Here, we will discuss the neuroprotective effects of PPAR γ in relation to ischaemia, spinal cord injury, chronic neuropathic pain and experimental autoimmune encephalomyelitis (EAE), a model of Multiple sclerosis. We will also explore the literature relating to several neurodegenerative disorders including Amyotrophic lateral sclerosis (ALS), Huntington's disease (HD), Alzheimer's disease (AD) and Parkinson's disease (PD), beyond their mechanisms on the mitochondrial function or on oxidative stress and apoptosis.

5.1. PPAR γ actions on CNS ischaemic injury

PPAR γ agonists may help to promote recovery after stroke. Endogenous neuronal PPAR γ may be important in limiting damage after cerebral ischaemia, as PPAR γ deficiency (PPAR γ knock-out mice) was associated with increased brain damage and oxidative stress, a reduced expression of glutathione S-transferase, UCP-1, catalase and SOD1 after cerebral ischaemia [179]. In contrast, activation of PPAR γ by the TZDs troglitazone or pioglitazone reduced inflammation and infarct volume and improved neurological function following middle cerebral artery occlusion in rats [149]. Treatment with rosiglitazone also reduced the infarct volume in a rat model of transient focal ischaemia [161]. Treatment with rosiglitazone significantly decreased DNA fragmentation, attenuated infarct volume, neurological deficits and neutrophilia after embolic focal cerebral ischaemia in rats [3]. Recently, 15D-PGJ2 a natural PPAR γ ligand decreased neuronal autophagy by decreasing the expression of proteins LC3-II, Beclin 1, cathepsin-B, and LAMP1 after cerebral ischaemia reperfusion injury [174]. A single injection of rosiglitazone immediately following global cerebral ischaemia significantly increased neuronal survival and functional recovery while reducing the cerebral infarct volume and brain oedema [144]. All these data suggest that intrinsic PPAR γ may be important in limiting stroke mediated injury and that further activation by TZDs may improve outcome.

5.2. PPAR γ actions on pain and spinal cord injury

Neuropathic pain can be a major problem after spinal cord injury. Several studies suggest that TZDs can limit chronic severe

pain and aid recovery from spinal injury. One study demonstrated that PPAR γ is essential for coupling ibuprofen to RhoA inhibition and the subsequent promotion of neurite growth in primary neurons. Also in the same study, a similar mechanism accounted for the activation of PPAR γ by ibuprofen in neuron-like PC12 and B104 cells, suggesting a molecular mechanism that could represent a novel therapeutic target in spinal cord injury [45]. Ajulemic acid, which directly interacts with PPAR γ and has a potent analgesic and anti-inflammatory activity, may also be a potential candidate for drug development in the treatment of pain [4].

In animal models, rosiglitazone and pioglitazone, improved motor function, prevented myelin loss and reduced long term pain after spinal injury in adult rats [121]. In a mouse model of neuropathic pain, the intrathecal administration of rosiglitazone or the natural ligand 15D-PGJ2 reduced nerve injury-induced allodynia and hyperalgesia [32]. Additionally, pioglitazone reduced tactile allodynia and thermal hyperalgesia in mice subjected to peripheral nerve injury through attenuation of proinflammatory cytokine secretion [100]. Intracerebroventricular administration of rosiglitazone or 15D-PGJ2 reduced peripheral inflammation in the brain, reduced inflammatory pain, induced the expression of Fos in the dorsal horn, and inhibited local oedema [110].

Treatment of animals subjected to spinal cord injury with rosiglitazone significantly increased the proliferation of neuronal precursor cells, accelerated locomotor recovery and reduced NF- κ B expression [105]. Local administration of rosiglitazone significantly reduced hypersensitivity to heat and mechanical stimuli, and paw swelling and reduced the development of post-incisional pain, possibly by regulating macrophage polarity at the inflamed site [71]. Attenuation of the development of inflammatory pain by rosiglitazone was attributed to the induction of haem oxygenase 1 (HO-1) in macrophages, and gene induction of endogenous opioid proenkephalin [70]. In rats with neuropathic pain, oral or intraperitoneal administration of pioglitazone limited multiple behavioural signs of somatosensory hypersensitivity and significantly reduced touch stimulus-evoked phospho-extracellular signal-related kinase (p-ERK) [111].

5.3. PPAR γ actions on EAE

Experimental allergic encephalomyelitis is often used as a model of demyelinating disease. The pathophysiology of the disease model is complex, but a significant body of data suggests key roles of mitochondrial dysfunction and oxidative and nitrosative stress [56,57,166]. In two models of EAE (immunized with myelin oligodendrocyte glycoprotein peptide and with myelin basic protein), treatment with pioglitazone delayed neurodegeneration by decreasing lymphocyte infiltration, reduced demyelination, chemokine and cytokine expression and increased inhibitor of kappa B expression in the brain [54].

6. Mechanisms of protection by PPAR γ in neurodegenerative diseases

6.1. Impact of PPAR γ in Amyotrophic lateral sclerosis

Amyotrophic lateral sclerosis (ALS, also known as motoneuron disease, MND), is a neurodegenerative disease that causes paralysis and shortened lifespan due to the selective death of motoneurons. Extensive data point to a major role for calcium mediated mitochondrial toxicity and oxidative stress in ALS including a major role for excitotoxicity, which is a prime example of calcium dependent mitochondrial toxicity [36,37,78]. The effects of the PPAR γ agonist troglitazone have been explored in rat motoneurons, which showed improved survival. However,

troglitazone did not promote the survival of sensory, sympathetic, septal or hippocampal neurons [115]. These results indicate specific neurotrophic activity of troglitazone for motoneurons through the activation of phosphatidylinositol 3-kinase (PI3-kinase) and supported the potential value of troglitazone for the treatment of motoneuron diseases such as ALS.

Mutations of SOD1 underlie MND in a rare subgroup of patients with familial disease. The specific mechanism of MND/ALS caused by mutations of SOD1 remain unclear – they are not simply due to loss of antioxidant function but seem to involve a toxic gain of function [137,176]. Pioglitazone protected transgenic mice carrying the ALS related SOD1-G93A mutation from motoneuron neurodegeneration, improving muscle strength and body weight and delaying disease onset [141]. In SOD1-G93A transgenic mice, treatment with pioglitazone extended survival, reduced gliosis and also reduced iNOS, NF- κ B, and 3-nitrotyrosine immunoreactivity in the spinal cord, suggesting a significant role against nitrosative stress [86]. In another study of the same mouse model, pioglitazone also protected motoneurons against p38-mediated neuronal death and NF- κ B-mediated glial inflammation [145]. The accumulation of critical concentrations of lipid peroxidation adducts during the progression of ALS leads to the activation of PPAR γ in motoneurons of SOD1-G93A mice. This in turn triggers self-protective mechanisms that involve the up-regulation of lipid detoxification enzymes, including lipoprotein lipase and glutathione S-transferase α -2 [9].

Recently, in a Drosophila model of ALS based on TAR DNA-binding protein 43 (TDP-43) that recapitulates several aspects of ALS pathophysiology, pioglitazone rescued TDP-43-dependent locomotor dysfunction in motoneurons and glia but not in muscles and also was neuroprotective when FUS, but not SOD1, is expressed in motoneurons [81].

Finally, a randomized, double blind, placebo-controlled trial of pioglitazone in combination with riluzole in ALS patients, disappointingly found that treatment with pioglitazone had no positive benefit, with no increase in the survival of ALS patients treated with riluzole [50].

6.2. Impact of PPAR γ in Huntington's disease

Huntington's disease (HD) is an autosomal dominant progressive neurodegenerative disorder associated with progressive extrapyramidal motor dysfunction. It is a genetic disease, caused by mutations that cause CAG trinucleotide expansion in the exon-1 region of the huntingtin gene. Extensive data links huntingtin mutations and CAG repeats to mitochondrial damage and oxidative stress [66,69,138].

The protective effects of PPAR γ agonists have been studied in several different models of HD. In striatal cells in culture expressing mutant huntingtin STHdh(Q111), rosiglitazone increased mitochondrial biogenesis and protected against calcium dependent (thapsigargin-induced) mitochondrial dysfunction [128]. Similarly, in cultured rat primary cortical neurons expressing mutant huntingtin STHdh(Q111), PPAR γ activation by rosiglitazone significantly attenuated thapsigargin-induced cell death, concomitant with reduced caspase activation, a delay in the loss of $\Delta\Psi_m$, and a reduction of ROS generation [80]. In addition, a reduction in mitochondrial mass associated with the expression of mutant huntingtin in N2A cells, was rescued by rosiglitazone [29]. Rosiglitazone also significantly increased survival in N2A cells expressing mutant huntingtin, while also upregulating the expression of PPAR γ , PGC1 α , NRF-1, TFAM and improving mitochondrial function in mutant cells. Rosiglitazone treatment also normalized endoplasmic reticulum stress sensors Bip, CHOP and ASK1 and significantly reduced mutant huntingtin aggregates that included ubiquitin and heat shock factor 1 and increased the levels of the

functional ubiquitin-proteasome system and heat shock protein 27/70 [28].

PPAR γ activation by pioglitazone protected striatal cells from mitochondrial dysfunction and oxidative stress in a 3-nitropropionic acid (3-NP)-induced experimental model of HD (3-NP, irreversibly inhibits succinate dehydrogenase and produces in animals behavioural, biochemical and morphologic changes considered to be similar to those occurring in HD). The beneficial effects on mitochondrial dysfunction were attributed to interference with the NF- κ B signalling pathway, which has been implicated in the pathogenesis of HD [112].

In a mouse model R6/2 of HD, treatment with rosiglitazone rescued progressive weight loss, motor deterioration, formation of mutant huntingtin aggregates, and reduced the expression of two neuroprotective proteins, the brain derived neurotrophic factor (BDNF) and Bcl-2 [27]. Bezafibrate, a pan-PPAR agonist, increased mitochondrial biogenesis, increased the numbers of muscle mitochondria, improved survival, improved the phenotype and reduced brain, muscle and BAT pathology in transgenic mouse R6/2 [82]. Treatment with rosiglitazone also reduced huntingtin aggregates and normalized the expression of PGC1 α , NRF1-2, and TFAM in the cortex of a transgenic mouse R6/2 [29]. Moreover, chronic administration of rosiglitazone to N171-82Q HD mice significantly improved motor function, attenuated hyperglycaemia, rescued BDNF deficiency in the cerebral cortex, prevented loss of orexin-A-immunopositive neurons in the hypothalamus, prevented PGC1 α reduction and increased Sirt6 protein levels [79].

6.3. Potential benefits of PPAR γ activators in Alzheimer's disease

Alzheimer's disease (AD) is a neurodegenerative disorder characterized by a progressive cognitive decline, associated with intracerebral aggregates of amyloid- β ($A\beta$) protein and the accumulation of the hyperphosphorylated tau protein. Substantial data point to oxidative stress and mitochondrial dysfunction as major contributors to the pathophysiology, although there is significant controversy about the relative roles of potential underlying mechanisms [21,49].

Acute treatment of APPV7171 transgenic mice model of AD with pioglitazone or ibuprofen significantly decreased the number of activated microglia and astrocytes in the hippocampus and cortex. Both drugs reduced the expression of the proinflammatory enzymes COX-2 and iNOS and reduced $A\beta$ deposits in the hippocampus and cortex [72]. In APPV7171 transgenic mice, pioglitazone treatment significantly attenuated astroglial activation, reduced signs of oxidative stress and normalized the cerebral blood flow and glucose uptake responses to increased neuronal activity, but failed to improve spatial memory [114]. In AD Apolipoprotein E (ApoE) knockout mice, rosiglitazone increased mitochondrial biogenesis as demonstrated by the induction of both oestrogen-stimulated related receptor alpha (ESRRA) mRNA and mtDNA copy number and improved glucose utilization [147]. Treatment with rosiglitazone improved cognition in the Tg2576 mice, another transgenic mouse model of AD which presents accumulation of $A\beta$, neuronal loss, and cognitive decline [131]. In the Tg2576 mice, rosiglitazone enhanced cognitive function and normalized dentate granule cell presynaptic function, through regulation of presynaptic vesicular proteins critical for glutamatergic neurotransmitter release, synaptic transmission, and short-term plasticity [113].

In another mouse model of AD, the APPswe/PS1 Δ e9 mice, treatment with pioglitazone reduced brain levels of soluble and insoluble $A\beta$ levels which correlated with the loss of both diffuse and dense-core plaques within the cortex [101]. In another study using the same APPswe/PS1 Δ e9 mice, the novel selective PPAR γ modulator, DSP-8658 and pioglitazone, improved spatial memory

performance and increased microglial A β phagocytosis which subsequently resulted in a reduction of cortical and hippocampal A β levels. The phagocytosis of A β was mediated through the up-regulation of scavenger receptor CD36 [175]. Activation of Wnt signalling by lithium and rosiglitazone reduced spatial memory impairment and neurodegeneration in brains of APPswe/PS1 Δ e9 mice, and both drugs also prevented changes in presynaptic and postsynaptic marker proteins [153].

In Cdk5 conditional knock-out mice, astrogliosis, microgliosis, neuronal loss and behavioural deficit were all improved by rosiglitazone. Cdk5 plays an important role in neurogenesis and there is some evidence that it participates in the pathogenesis of AD [159].

PPAR γ activation by troglitazone and rosiglitazone protected rat hippocampal neurons in culture against toxicity through exposure to synthetic A β _{1–40} peptide (5 μ M), resulting in the modulation of Wnt signalling components, including the increase of the cytoplasmic and nuclear β -catenin levels and the inhibition of glycogen synthase kinase-3 β (GSK-3 β), suggesting that GSK-3 β is responsible for the tau hyperphosphorylation in AD [75]. Furthermore, in cultured hippocampal neurons, rosiglitazone was protective against mitochondrial damage, oxidative stress and apoptosis induced by 5 μ M of synthetic A β _{1–40} peptide, through a mechanism involving elevated expression of Bcl-2. Also, PPAR γ overexpression prevented H₂O₂-induced ROS, stabilized mitochondria, and decreased A β -induced toxicity [59].

In a clinical trial, patients with mild-to-moderate AD were randomized to placebo or rosiglitazone treatment and the exploratory analyses suggested an improvement in cognitive and functional measures in ApoE ϵ 4 non-carriers, whereas no improvement and some decline was noted in the ApoE ϵ 4 allele carriers [130]. In a further randomized, double-blind, placebo-controlled phase III study in mild-to-moderate AD, rosiglitazone monotherapy failed to show any benefit in cognition or in global function in ApoE ϵ 4 negative subjects [67].

In addition, in a randomized pilot clinical trial of the safety of pioglitazone in non-diabetic patients with AD, pioglitazone was generally well tolerated with an increase in peripheral oedema (oedema is a commonly recognized effect of pioglitazone in clinical use and this potential adverse treatment effect was anticipated) and there were no other serious or unanticipated adverse events or clinical laboratory changes attributable to pioglitazone over a long-term exposure [63]. Some clinical trials of different PPAR γ agonist for the treatment of neurodegenerative diseases are shown in Table 2.

6.4. Impact of PPAR γ in Parkinson's disease

Parkinson's disease (PD) is characterized by the progressive

Table 2
Some clinical trials of TZDs for the treatment of different neurodegenerative diseases.

Agent	Disease	Outcome	Refs.
Rosiglitazone + Riluzole	ALS	Had no positive benefit	[50]
Rosiglitazone	AD	No improvement in ApoE ϵ 4 allele carriers	[130]
Rosiglitazone	AD	No benefit in cognition or in global function in ApoE ϵ 4 negative subjects	[67]
Pioglitazone	AD	Well tolerated but an increase in peripheral oedema	[63]
Glitazone	PD	A reduction in the incidence of PD in patients with diabetes	[18]
Pioglitazone	PD	Is unlikely to modify progression in early PD	[76]

neurodegeneration of dopaminergic neurons of the substantia nigra, causing a characteristic tremor, bradykinesia and a slower deterioration of cognitive function. There are extensive data pointing to impaired mitochondrial function and oxidative stress as key mechanisms in the pathophysiology of the disease, most notably pointing to complex I damage, impaired autophagy and calcium dependent mitochondrial free radical generation for reviews see: [49,99,148]. The neuroprotection of PPAR γ activation has been studied in a number of models of PD.

In human neuroblastoma cells in culture, rosiglitazone was protective against mitochondrial dysfunction produced by the complex I inhibitor 1-methyl-4-phenylpyridinium ion (MPP⁺), increasing $\Delta\Psi_m$, while also increasing the expression of SOD and catalase, and the expression of Bcl-2 and Bax, thus promoting both antioxidant defence and limiting apoptosis [84]. 15D-PGJ2 and pioglitazone preserved mitochondrial function of oligodendrocyte progenitors against the inhibitory activities of rotenone and cytokine tumour necrosis factor α (TNF α), by maintaining $\Delta\Psi_m$, reducing mitochondrial ROS production and by increasing the expression of PGC1 α , uncoupling protein 2 (UCP-2) and cytochrome oxidase subunit COX1. TNF α decreased Ca²⁺ oscillations, as well as the frequency of the oscillatory events and the TNF α effects were prevented by pioglitazone, also pioglitazone protected the oligodendrocyte progenitor differentiation to immature oligodendrocyte [40]. In differentiated neuroblastoma SH-SY5Y cells associated with chronic partial inhibition of complex I with rotenone and with loss of PINK1 function, as cellular models related to sporadic and familial PD respectively, rosiglitazone increased mitochondrial biogenesis, increased oxygen consumption, increased mitochondrial mass, $\Delta\Psi_m$, mtDNA copy number, decreased autophagy and suppressed free radical generation [35]. A novel thiobarbituric-like compound MDG548, which acts as a functional PPAR γ agonist, was tested in vitro and in vivo models of PD, MDG548 protected against H₂O₂ and MPP⁺ neurotoxicity in PC12 cells and in the mice treated with MPTP, the MDG548 reduced reactive microglia and iNOS induction in the substantia nigra [91].

1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine (MPTP) administration is used in animals to model PD, as it causes selective death of dopaminergic neurons. The oxidative stress produced by this toxin was reduced by pioglitazone treatment as evidenced by a reduction in the levels of malondialdehyde and increased glutathione levels [89]. In response to MPTP plus probenecid, treatment with rosiglitazone was effective in protection against partial degeneration of the substantia nigra and the decline of striatal dopamine. Also, the production of the pro-inflammatory cytokine TNF α by microglia was inhibited by rosiglitazone [22]. In another study, rhesus monkeys were subjected to an intracarotid injection of MPTP and pioglitazone was also shown to be neuroprotective when administered early. Behavioural recovery was associated with preservation of dopaminergic markers and reduced infiltration by CD68-positive macrophages in the nigrostriatal area and by a significant improvement in a clinical rating score [152].

A novel non-TZD partial PPAR γ agonist LSN862, was shown to be neuroprotective in MPTP-induced neurodegeneration, which was associated with the modulation of PPAR γ and PGC1 α expression, the downregulation of neuroinflammation and decreased oxidative stress [151]. The expression of PPAR γ in the dopaminergic neurons of the substantia nigra pars compacta and putamen of hemiparkinsonian nonhuman primates was prominent and the dopaminergic neurons were more likely to survive against MPTP-induced damage after activation by pioglitazone [150]. In the MPTP-probenecid mouse model, the PPAR γ agonists may act through an immunomodulatory action, since the anti-inflammatory microglia were stimulated and inflammatory microglia were inhibited in the substantia nigra compacta by

rosiglitazone [123]. Pioglitazone protected against deterioration of motor function and neuronal loss in the acute MPTP mouse model, but failed to protect in a rat model using 6-hydroxydopamine (6-OHDA) as a dopaminergic selective toxin, thought to reflect the severity of damage caused by 6-OHDA [90]. The activation of PPAR γ receptors by rosiglitazone in the 6-OHDA-lesioned rat, significantly increased GFAP expression in the striatum and attenuated COX-2 and TNF α expression [93].

In 2013, pioglitazone entered in a multicenter placebo-controlled phase II study, in early PD (NCT01280123) and depending on the study results, a subsequent phase III trial could follow. Recently, in a retrospective cohort study, the prescription of the PPAR γ agonist glitazone in patients with diabetes was associated with a reduction in the incidence of PD, suggesting that PPAR γ pathways may be a successful drug target in PD [18].

6.5. Impact of PPAR γ in Friedreich's ataxia

The effect of PPAR γ activation has also been studied in Friedreich's ataxia, an autosomal recessive disorder caused by mutations of the protein frataxin. The disease phenotype includes a progressive cardiomyopathy which leads eventually to heart failure. It was demonstrated that PPAR γ agonist increased both messenger RNA and protein levels of frataxin [102]. Pioglitazone, which seems to have fewer side effects than rosiglitazone is on a phase III clinical trial for its neuroprotective properties. The project, titled "Effect of pioglitazone administered to patients with Friedreich's ataxia: proof of concept". However, the antecedents and possible consequences should be taken into consideration when PPAR γ agonist drugs are considered as possible therapeutic agents for FRDA patients [62].

7. Conclusion

The modulation of molecular pathways that include mitochondrial dysfunction, oxidative stress, process of autophagy, inflammation and apoptosis, which are implicated in the pathophysiology of many major diseases including neurodegenerative disorders seem to present very valuable therapeutic targets. In a similar vein, PPAR γ agonists have been showed to have a wide range of activities that positively influence the pathology of diseases in several experimental models, and have the capacity to be neuroprotective by regulating the expression of genes involved in neuronal survival processes and therefore promote the health of neurons. PPAR γ agonists could be valuable potential therapeutic targets for several diseases including neurodegenerative disorders, since the results from patients and from in vivo and in vitro models of neurodegenerative disorders underline the beneficial effects of PPAR γ agonists for future clinical trials. However, further studies are needed if we are to understand the potential of PPAR γ agonists for patient benefit or the mechanisms by which PPAR γ exerts its protective actions.

References

- [1] B.D. Abbott, Review of the expression of peroxisome proliferator-activated receptors alpha (PPAR alpha), beta (PPAR beta), and gamma (PPAR gamma) in rodent and human development, *Reprod. Toxicol.* 27 (2009) 246–257.
- [2] R. Agrawal, The first approved agent in the Glitazar's Class: Saroglitazar, *Curr. Drug Targets* 15 (2014) 151–155.
- [3] M. Allahtavakoli, R. Moloudi, M.K. Arababadi, et al., Delayed post ischemic treatment with Rosiglitazone attenuates infarct volume, neurological deficits and neutrophilia after embolic stroke in rat, *Brain Res.* 1271 (2009) 121–127.
- [4] A.L. Ambrosio, S.M. Dias, I. Polikarpov, et al., Ajulemic acid, a synthetic nonpsychoactive cannabinoid acid, bound to the ligand binding domain of the human peroxisome proliferator-activated receptor gamma, *J. Biol. Chem.* 282 (2007) 18625–18633.
- [5] A.S. Banks, F.E. McAllister, J.P. Camporez, et al., An ERK/Cdk5 axis controls the diabetogenic actions of PPARgamma, *Nature* 517 (2015) 391–395.
- [6] A.J. Bannister, T. Kouzarides, The CBP co-activator is a histone acetyltransferase, *Nature* 384 (1996) 641–643.
- [7] Y. Barak, M.C. Nelson, E.S. Ong, et al., PPAR gamma is required for placental, cardiac, and adipose tissue development, *Mol. Cell* 4 (1999) 585–595.
- [8] L.C. Bell-Parikh, T. Ide, J.A. Lawson, et al., Biosynthesis of 15-deoxy-delta12,14-PGJ2 and the ligation of PPARgamma, *J. Clin. Investig.* 112 (2003) 945–955.
- [9] V. Benedusi, F. Martorana, L. Brambilla, et al., The peroxisome proliferator-activated receptor gamma (PPARgamma) controls natural protective mechanisms against lipid peroxidation in amyotrophic lateral sclerosis, *J. Biol. Chem.* 287 (2012) 35899–35911.
- [10] J. Berger, D.E. Moller, The mechanisms of action of PPARs, *Annu. Rev. Med.* 53 (2002) 409–435.
- [11] A. Bernardo, L. Minghetti, PPAR-gamma agonists as regulators of microglial activation and brain inflammation, *Curr. Pharm. Des.* 12 (2006) 93–109.
- [12] A. Bernardo, L. Minghetti, Regulation of glial cell functions by PPAR-gamma natural and synthetic agonists, *PPAR Res.* 2008 (2008) 864140.
- [13] F. Biscetti, G. Straface, D. Pitocco, et al., Peroxisome proliferator-activated receptors and angiogenesis, *Nutr. Metab. Cardiovasc. Dis.* 19 (2009) 751–759.
- [14] I. Bogacka, B. Ukropcova, M. McNeil, et al., Structural and functional consequences of mitochondrial biogenesis in human adipocytes in vitro, *J. Clin. Endocrinol. Metab.* 90 (2005) 6650–6656.
- [15] I. Bogacka, H. Xie, G.A. Bray, et al., Pioglitazone induces mitochondrial biogenesis in human subcutaneous adipose tissue in vivo, *Diabetes* 54 (2005) 1392–1399.
- [16] O. Braissant, F. Fougelle, C. Scotto, et al., Differential expression of peroxisome proliferator-activated receptors (PPARs): tissue distribution of PPAR-alpha, -beta, and -gamma in the adult rat, *Endocrinology* 137 (1996) 354–366.
- [17] O. Braissant, W. Wahli, Differential expression of peroxisome proliferator-activated receptor-alpha, -beta, and -gamma during rat embryonic development, *Endocrinology* 139 (1998) 2748–2754.
- [18] R. Brauer, K. Bhaskaran, N. Chaturvedi, et al., Glitazone treatment and incidence of Parkinson's Disease among people with diabetes: a retrospective cohort study, *PLoS Med.* 12 (2015) e1001854.
- [19] J.J. Bright, S. Kanakasabai, W. Chearwae, et al., PPAR regulation of inflammatory signaling in CNS diseases, *PPAR Res.* 2008 (2008) 658520.
- [20] E.M. Burkart, N. Sambandam, X. Han, et al., Nuclear receptors PPARbeta/delta and PPARalpha direct distinct metabolic regulatory programs in the mouse heart, *J. Clin. Investig.* 117 (2007) 3930–3939.
- [21] L. Canevari, A.Y. Abramov, M.R. Duchon, Toxicity of amyloid beta peptide: tales of calcium, mitochondria, and oxidative stress, *Neurochem. Res.* 29 (2004) 637–650.
- [22] A.R. Carta, L. Frau, A. Pisanu, et al., Rosiglitazone decreases peroxisome proliferator receptor-gamma levels in microglia and inhibits TNF-alpha production: new evidences on neuroprotection in a progressive Parkinson's disease model, *Neuroscience* 194 (2011) 250–261.
- [23] V. Chandra, P. Huang, Y. Hamuro, et al., Structure of the intact PPAR-gamma-RXR- nuclear receptor complex on DNA, *Nature* 456 (2008) 350–356.
- [24] A. Chawla, Y. Barak, L. Nagy, et al., PPAR-gamma dependent and independent effects on macrophage-gene expression in lipid metabolism and inflammation, *Nat. Med.* 7 (2001) 48–52.
- [25] A. Chawla, E.J. Schwarz, D.D. Dimaculangan, et al., Peroxisome proliferator-activated receptor (PPAR) gamma: adipose-predominant expression and induction early in adipocyte differentiation, *Endocrinology* 135 (1994) 798–800.
- [26] Y.C. Chen, J.S. Wu, H.D. Tsai, et al., Peroxisome proliferator-activated receptor gamma (PPAR-gamma) and neurodegenerative disorders, *Mol. Neurobiol.* 46 (2012) 114–124.
- [27] M.C. Chiang, C.M. Chen, M.R. Lee, et al., Modulation of energy deficiency in Huntington's disease via activation of the peroxisome proliferator-activated receptor gamma, *Hum. Mol. Genet.* 19 (2010) 4043–4058.
- [28] M.C. Chiang, Y.C. Cheng, C.J. Nicol, et al., Rosiglitazone activation of PPAR-gamma-dependent signaling is neuroprotective in mutant huntingtin expressing cells, *Exp. Cell Res.* 338 (2015) 183–193.
- [29] M.C. Chiang, Y. Chern, R.N. Huang, PPARgamma rescue of the mitochondrial dysfunction in Huntington's disease, *Neurobiol. Dis.* 45 (2012) 322–328.
- [30] J.H. Choi, A.S. Banks, J.L. Estall, et al., Anti-diabetic drugs inhibit obesity-linked phosphorylation of PPARgamma by Cdk5, *Nature* 466 (2010) 451–456.
- [31] P.C. Chui, H.P. Guan, M. Lehrke, et al., PPARgamma regulates adipocyte cholesterol metabolism via oxidized LDL receptor 1, *J. Clin. Investig.* 115 (2005) 2244–2256.
- [32] S.B. Churi, O.S. Abdel-Aleem, K.K. Tumber, et al., Intrathecal rosiglitazone acts at peroxisome proliferator-activated receptor-gamma to rapidly inhibit neuropathic pain in rats, *J. Pain* 9 (2008) 639–649.
- [33] J.R. Colca, W.G. McDonald, D.J. Waldon, et al., Identification of a novel mitochondrial protein ("mitoNEET") cross-linked specifically by a thiazolidinedione photoprobe, *Am. J. Physiol. Endocrinol. Metab.* 286 (2004) E252–260.
- [34] M. Collino, N.S. Patel, C. Thiemermann, PPARs as new therapeutic targets for the treatment of cerebral ischemia/reperfusion injury, *Ther. Adv. Cardiovasc. Dis.* 2 (2008) 179–197.
- [35] J.C. Corona, S.C. de Souza, M.R. Duchon, PPARgamma activation rescues mitochondrial function from inhibition of complex I and loss of PINK1, *Exp.*

- Neurol. 253 (2014) 16–27.
- [36] J.C. Corona, R. Tapia, Ca²⁺-permeable AMPA receptors and intracellular Ca²⁺ determine motoneuron vulnerability in rat spinal cord in vivo, *Neuropharmacology* 52 (2007) 1219–1228.
- [37] J.C. Corona, L.B. Tovar-y-Romo, R. Tapia, Glutamate excitotoxicity and therapeutic targets for amyotrophic lateral sclerosis, *Expert Opin. Ther. Targets* 11 (2007) 1415–1428.
- [38] T.E. Cullingford, K. Bhakoo, S. Peuchen, et al., Distribution of mRNAs encoding the peroxisome proliferator-activated receptor alpha, beta, and gamma and the retinoid X receptor alpha, beta, and gamma in rat central nervous system, *J. Neurochem.* 70 (1998) 1366–1375.
- [39] S.S. Davies, A.V. Pontsler, G.K. Marathe, et al., Oxidized alkyl phospholipids are specific, high affinity peroxisome proliferator-activated receptor gamma ligands and agonists, *J. Biol. Chem.* 276 (2001) 16015–16023.
- [40] C. De Nuccio, A. Bernardo, C. Cruciani, et al., Peroxisome proliferator activated receptor-gamma agonists protect oligodendrocyte progenitors against tumor necrosis factor-alpha-induced damage: effects on mitochondrial functions and differentiation, *Exp. Neurol.* 271 (2015) 506–514.
- [41] P. Delerive, J.C. Fruchart, B. Staels, Peroxisome proliferator-activated receptors in inflammation control, *J. Endocrinol.* 169 (2001) 453–459.
- [42] C. Dello Russo, V. Gavriluk, G. Weinberg, et al., Peroxisome proliferator-activated receptor gamma thiazolidinedione agonists increase glucose metabolism in astrocytes, *J. Biol. Chem.* 278 (2003) 5828–5836.
- [43] B. Desvergne, L. Michalik, W. Wahli, Be fit or be sick: peroxisome proliferator-activated receptors are down the road, *Mol. Endocrinol.* 18 (2004) 1321–1332.
- [44] B. Desvergne, W. Wahli, Peroxisome proliferator-activated receptors: nuclear control of metabolism, *Endocr. Rev.* 20 (1999) 649–688.
- [45] J. Dill, A.R. Patel, X.L. Yang, et al., A molecular mechanism for ibuprofen-mediated RhoA inhibition in neurons, *J. Neurosci.* 30 (2010) 963–972.
- [46] J. DiRenzo, M. Soderstrom, R. Kurokawa, et al., Peroxisome proliferator-activated receptors and retinoic acid receptors differentially control the interactions of retinoid X receptor heterodimers with ligands, coactivators, and corepressors, *Mol. Cell. Biol.* 17 (1997) 2166–2176.
- [47] J.E. Dominy, P. Puigserver, Mitochondrial biogenesis through activation of nuclear signaling proteins, *Cold Spring Harb. Perspect. Biol.* 5 (2013).
- [48] C. Dreyer, G. Krey, H. Keller, et al., Control of the peroxisomal beta-oxidation pathway by a novel family of nuclear hormone receptors, *Cell* 68 (1992) 879–887.
- [49] M.R. Duchon, Mitochondria, calcium-dependent neuronal death and neurodegenerative disease, *Pflug. Arch.* 464 (2012) 111–121.
- [50] L. Dupuis, R. Dengler, M.T. Heneka, et al., A randomized, double blind, placebo-controlled trial of pioglitazone in combination with riluzole in amyotrophic lateral sclerosis, *PLoS One* 7 (2012) e37885.
- [51] M.M. El-Naa, M.F. El-Refaei, W.A. Nasif, et al., In-vivo antioxidant and anti-inflammatory activity of rosiglitazone, a peroxisome proliferator-activated receptor-gamma (PPAR-gamma) agonists in animal model of bronchial asthma, *J. Pharm. Pharmacol.* 67 (2015) 1421–1430.
- [52] P. Escher, O. Braissant, S. Basu-Modak, et al., Rat PPARs: quantitative analysis in adult rat tissues and regulation in fasting and refeeding, *Endocrinology* 142 (2001) 4195–4202.
- [53] R.M. Evans, G.D. Barish, Y.X. Wang, PPARs and the complex journey to obesity, *Nat. Med.* 10 (2004) 355–361.
- [54] D.L. Feinstein, E. Galea, V. Gavriluk, et al., Peroxisome proliferator-activated receptor-gamma agonists prevent experimental autoimmune encephalomyelitis, *Ann. Neurol.* 51 (2002) 694–702.
- [55] B.M. Forman, P. Tontonoz, J. Chen, et al., 15-Deoxy-delta 12, 14-prostaglandin J2 is a ligand for the adipocyte determination factor PPAR gamma, *Cell* 83 (1995) 803–812.
- [56] M. Forte, B.G. Gold, G. Marracci, et al., Cyclophilin D inactivation protects axons in experimental autoimmune encephalomyelitis, an animal model of multiple sclerosis, *Proc. Natl. Acad. Sci. USA* 104 (2007) 7558–7563.
- [57] R.J. Franklin, C. French-Constant, J.M. Edgar, et al., Neuroprotection and repair in multiple sclerosis, *Nat. Rev. Neurol.* 8 (2012) 624–634.
- [58] B.I. Frohnert, T.Y. Hui, D.A. Bernlohr, Identification of a functional peroxisome proliferator-responsive element in the murine fatty acid transport protein gene, *J. Biol. Chem.* 274 (1999) 3970–3977.
- [59] K. Fuenzalida, R. Quintanilla, P. Ramos, et al., Peroxisome proliferator-activated receptor gamma up-regulates the Bcl-2 anti-apoptotic protein in neurons and induces mitochondrial stabilization and protection against oxidative stress and apoptosis, *J. Biol. Chem.* 282 (2007) 37006–37015.
- [60] B. Garcia-Bueno, J.R. Caso, B.G. Perez-Nievas, et al., Effects of peroxisome proliferator-activated receptor gamma agonists on brain glucose and glutamate transporters after stress in rats, *Neuropsychopharmacology* 32 (2007) 1251–1260.
- [61] B. Garcia-Bueno, J.L. Madrigal, I. Lizasoain, et al., Peroxisome proliferator-activated receptor gamma activation decreases neuroinflammation in brain after stress in rats, *Biol. Psychiatry* 57 (2005) 885–894.
- [62] J.L. Garcia-Gimenez, F. Sanchis-Gomar, F.V. Pallardo, Could thiazolidinediones increase the risk of heart failure in Friedreich's ataxia patients? *Mov. Disord.* 26 (2011) 769–771.
- [63] D.S. Geldmacher, T. Fritsch, M.J. McClendon, et al., A randomized pilot clinical trial of the safety of pioglitazone in treatment of patients with Alzheimer disease, *Arch. Neurol.* 68 (2011) 45–50.
- [64] L. Gelman, G. Zhou, L. Fajas, et al., p300 interacts with the N- and C-terminal part of PPARgamma2 in a ligand-independent and -dependent manner, respectively, *J. Biol. Chem.* 274 (1999) 7681–7688.
- [65] S. Ghosh, N. Patel, D. Rahn, et al., The thiazolidinedione pioglitazone alters mitochondrial function in human neuron-like cells, *Mol. Pharmacol.* 71 (2007) 1695–1702.
- [66] J. Gil-Mohapel, P.S. Brocardo, B.R. Christie, The role of oxidative stress in Huntington's disease: are antioxidants good therapeutic candidates? *Curr. Drug Targets* 15 (2014) 454–468.
- [67] M. Gold, C. Alderton, M. Zvartau-Hind, et al., Rosiglitazone monotherapy in mild-to-moderate Alzheimer's disease: results from a randomized, double-blind, placebo-controlled phase III study, *Dement Geriatr. Cogn. Disord.* 30 (2010) 131–146.
- [68] H.P. Guan, Y. Li, M.V. Jensen, et al., A futile metabolic cycle activated in adipocytes by anti-diabetic agents, *Nat. Med.* 8 (2002) 1122–1128.
- [69] P. Guedes-Dias, B.R. Pinho, T.R. Soares, et al., Mitochondrial dynamics and quality control in Huntington's disease, *Neurobiol. Dis.* 90 (2016) 51–57.
- [70] M. Hasegawa-Moriyama, T. Kurimoto, M. Nakama, et al., Peroxisome proliferator-activated receptor-gamma agonist rosiglitazone attenuates inflammatory pain through the induction of heme oxygenase-1 in macrophages, *Pain* 154 (2013) 1402–1412.
- [71] M. Hasegawa-Moriyama, T. Ohnno, K. Godai, et al., Peroxisome proliferator-activated receptor-gamma agonist rosiglitazone attenuates postincisional pain by regulating macrophage polarization, *Biochem. Biophys. Res. Commun.* 426 (2012) 76–82.
- [72] M.T. Heneka, M. Sastre, L. Dumitrescu-Ozimek, et al., Acute treatment with the PPARgamma agonist pioglitazone and ibuprofen reduces glial inflammation and Abeta1-42 levels in APPV7171 transgenic mice, *Brain* 128 (2005) 1442–1453.
- [73] T. Hibuse, N. Maeda, T. Funahashi, et al., Aquaporin 7 deficiency is associated with development of obesity through activation of adipose glycerol kinase, *Proc. Natl. Acad. Sci. USA* 102 (2005) 10993–10998.
- [74] R.L. Hunter, N. Dragicevic, K. Seifert, et al., Inflammation induces mitochondrial dysfunction and dopaminergic neurodegeneration in the nigrostriatal system, *J. Neurochem.* 100 (2007) 1375–1386.
- [75] N.C. Inestrosa, J.A. Godoy, R.A. Quintanilla, et al., Peroxisome proliferator-activated receptor gamma is expressed in hippocampal neurons and its activation prevents beta-amyloid neurodegeneration: role of Wnt signaling, *Exp. Cell Res.* 304 (2005) 91–104.
- [76] Investigators. NETIPDN-PF-Z, Pioglitazone in early Parkinson's disease: a phase 2, multicentre, double-blind, randomised trial, *Lancet Neurol.* 14 (2015) 8.
- [77] I. Issemann, S. Green, Activation of a member of the steroid hormone receptor superfamily by peroxisome proliferators, *Nature* 347 (1990) 645–650.
- [78] M.K. Jaiswal, Selective vulnerability of motoneuron and perturbed mitochondrial calcium homeostasis in amyotrophic lateral sclerosis: implications for motoneurons specific calcium dysregulation, *Mol. Cell Ther.* 2 (2014) 26.
- [79] J. Jin, J. Albertz, Z. Guo, et al., Neuroprotective effects of PPAR-gamma agonist rosiglitazone in N171-82Q mouse model of Huntington's disease, *J. Neurochem.* 125 (2013) 410–419.
- [80] Y.N. Jin, W.Y. Hwang, C. Jo, et al., Metabolic state determines sensitivity to cellular stress in Huntington disease: normalization by activation of PPAR-gamma, *PLoS One* 7 (2012) e30406.
- [81] A. Joardar, J. Menzl, T.C. Podolsky, et al., PPAR gamma activation is neuroprotective in a Drosophila model of ALS based on TDP-43, *Hum. Mol. Genet.* 24 (2015) 1741–1754.
- [82] A. Johri, N.Y. Calingasan, T.M. Hennessey, et al., Pharmacologic activation of mitochondrial biogenesis exerts widespread beneficial effects in a transgenic mouse model of Huntington's disease, *Hum. Mol. Genet.* 21 (2012) 1124–1137.
- [83] T.W. Jung, J.Y. Lee, W.S. Shim, et al., Rosiglitazone protects human neuroblastoma SH-SY5Y cells against acetaldehyde-induced cytotoxicity, *Biochem. Biophys. Res. Commun.* 340 (2006) 221–227.
- [84] T.W. Jung, J.Y. Lee, W.S. Shim, et al., Rosiglitazone protects human neuroblastoma SH-SY5Y cells against MPP+ induced cytotoxicity via inhibition of mitochondrial dysfunction and ROS production, *J. Neurol. Sci.* 253 (2007) 53–60.
- [85] Y. Kawakami, M. Tsuda, S. Takahashi, et al., Transcriptional coactivator PGC-1alpha regulates chondrogenesis via association with Sox9, *Proc. Natl. Acad. Sci. USA* 102 (2005) 2414–2419.
- [86] M. Kiaei, K. Kipiani, J. Chen, et al., Peroxisome proliferator-activated receptor-gamma agonist extends survival in transgenic mouse model of amyotrophic lateral sclerosis, *Exp. Neurol.* 191 (2005) 331–336.
- [87] G. Krey, O. Braissant, F. L'Horsset, et al., Fatty acids, eicosanoids, and hypolipidemic agents identified as ligands of peroxisome proliferator-activated receptors by coactivator-dependent receptor ligand assay, *Mol. Endocrinol.* 11 (1997) 779–791.
- [88] A. Krishnaswami, S. Ravi-Kumar, J.M. Lewis, Thiazolidinediones: a 2010 perspective, *Perm. J.* 14 (2010) 64–72.
- [89] P. Kumar, R.K. Kaundal, S. More, et al., Beneficial effects of pioglitazone on cognitive impairment in MPTP model of Parkinson's disease, *Behav. Brain Res.* 197 (2009) 398–403.
- [90] C. Laloux, M. Petraut, C. Lecoite, et al., Differential susceptibility to the PPAR-gamma agonist pioglitazone in 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine and 6-hydroxydopamine rodent models of Parkinson's disease, *Pharmacol. Res.* 65 (2012) 514–522.
- [91] D. Lecca, D.K. Nevin, G. Mulas, et al., Neuroprotective and anti-inflammatory

- properties of a novel non-thiazolidinedione PPAR γ agonist in vitro and in MPTP-treated mice, *Neuroscience* 302 (2015) 23–35.
- [92] C.H. Lee, P. Olson, A. Hevener, et al., PPAR δ regulates glucose metabolism and insulin sensitivity, *Proc. Natl. Acad. Sci. USA* 103 (2006) 3444–3449.
- [93] E.Y. Lee, J.E. Lee, J.H. Park, et al., Rosiglitazone, a PPAR- γ agonist, protects against striatal dopaminergic neurodegeneration induced by 6-OHDA lesions in the substantia nigra of rats, *Toxicol. Lett.* 213 (2012) 332–344.
- [94] J.E. Lee, J.H. Park, S.J. Jang, et al., Rosiglitazone inhibits chlorpyrifos-induced apoptosis via modulation of the oxidative stress and inflammatory response in SH-SY5Y cells, *Toxicol. Appl. Pharmacol.* 278 (2014) 159–171.
- [95] J.M. Lehmann, J.M. Lenhard, B.B. Oliver, et al., Peroxisome proliferator-activated receptors alpha and gamma are activated by indomethacin and other non-steroidal anti-inflammatory drugs, *J. Biol. Chem.* 272 (1997) 3406–3410.
- [96] J.M. Lehmann, L.B. Moore, T.A. Smith-Oliver, et al., An antidiabetic thiazolidinedione is a high affinity ligand for peroxisome proliferator-activated receptor gamma (PPAR gamma), *J. Biol. Chem.* 270 (1995) 12953–12956.
- [97] J. Lin, R. Yang, P.T. Tarr, et al., Hyperlipidemic effects of dietary saturated fats mediated through PGC-1 β coactivation of SREBP, *Cell* 120 (2005) 261–273.
- [98] S. Liu, B. Hatano, M. Zhao, et al., Role of peroxisome proliferator-activated receptor $\{\delta\}/\{\beta\}$ in hepatic metabolic regulation, *J. Biol. Chem.* 286 (2011) 1237–1247.
- [99] M.A. Lynch-Day, K. Mao, K. Wang, et al., The role of autophagy in Parkinson's disease, *Cold Spring Harb. Perspect. Med.* 2 (2012) a009357.
- [100] T. Maeda, N. Kiguchi, Y. Kobayashi, et al., Pioglitazone attenuates tactile allodynia and thermal hyperalgesia in mice subjected to peripheral nerve injury, *J. Pharmacol. Sci.* 108 (2008) 341–347.
- [101] S. Mandrekar-Colucci, J.C. Karlo, G.E. Landreth, Mechanisms underlying the rapid peroxisome proliferator-activated receptor- γ -mediated amyloid clearance and reversal of cognitive deficits in a murine model of Alzheimer's disease, *J. Neurosci.* 32 (2012) 10117–10128.
- [102] D. Marmolino, F. Acquaviva, M. Pinelli, et al., PPAR- γ agonist Azelaoyl PAF increases frataxin protein and mRNA expression: new implications for the Friedreich's ataxia therapy, *Cerebellum* 8 (2009) 98–103.
- [103] N.J. McKenna, B.W. O'Malley, Combinatorial control of gene expression by nuclear receptors and coregulators, *Cell* 108 (2002) 465–474.
- [104] N.J. McKenna, B.W. O'Malley, Minireview: nuclear receptor coactivators—an update, *Endocrinology* 143 (2002) 2461–2465.
- [105] Q.Q. Meng, X.J. Liang, P. Wang, et al., Rosiglitazone enhances the proliferation of neural progenitor cells and inhibits inflammation response after spinal cord injury, *Neurosci. Lett.* 503 (2011) 191–195.
- [106] L. Michalik, J. Auwerx, J.P. Berger, et al., International Union of Pharmacology. LXI. Peroxisome proliferator-activated receptors, *Pharmacol. Rev.* 58 (2006) 726–741.
- [107] L. Michalik, B. Desvergne, C. Dreyer, et al., PPAR expression and function during vertebrate development, *Int. J. Dev. Biol.* 46 (2002) 105–114.
- [108] G. Miglio, A.C. Rosa, L. Rattazzi, et al., PPAR γ stimulation promotes mitochondrial biogenesis and prevents glucose deprivation-induced neuronal cell loss, *Neurochem. Int.* 55 (2009) 496–504.
- [109] S. Moreno, S. Farioli-Vecchioli, M.P. Ceru, Immunolocalization of peroxisome proliferator-activated receptors and retinoid X receptors in the adult rat CNS, *Neuroscience* 123 (2004) 131–145.
- [110] J. Morgenweck, O.S. Abdel-Aleem, K.C. McNamara, et al., Activation of peroxisome proliferator-activated receptor gamma in brain inhibits inflammatory pain, dorsal horn expression of Fos, and local edema, *Neuropharmacology* 58 (2010) 337–345.
- [111] J. Morgenweck, R.B. Griggs, R.R. Donahue, et al., PPAR γ activation blocks development and reduces established neuropathic pain in rats, *Neuropharmacology* 70 (2013) 236–246.
- [112] M. Napolitano, L. Costa, R. Palermo, et al., Protective effect of pioglitazone, a PPAR γ ligand, in a 3 nitropropionic acid model of Huntington's disease, *Brain Res. Bull.* 85 (2011) 231–237.
- [113] M.N. Nenov, F. Laezza, S.J. Haidacher, et al., Cognitive enhancing treatment with a PPAR γ agonist normalizes dentate granule cell presynaptic function in Tg2576 APP mice, *J. Neurosci.* 34 (2014) 1028–1036.
- [114] N. Nicolakakis, T. Aboulkassim, B. Ongali, et al., Complete rescue of cerebrovascular function in aged Alzheimer's disease transgenic mice by antioxidants and pioglitazone, a peroxisome proliferator-activated receptor gamma agonist, *J. Neurosci.* 28 (2008) 9287–9296.
- [115] C. Nishijima, K. Kimoto, Y. Arakawa, Survival activity of troglitazone in rat motoneurons, *J. Neurochem.* 76 (2001) 383–390.
- [116] I. Nishikata, H. Sasaki, M. Iga, et al., A novel EVI1 gene family, MEL1, lacking a PR domain (MEL1S) is expressed mainly in t(1;3)(p36;q21)-positive AML and blocks G-CSF-induced myeloid differentiation, *Blood* 102 (2003) 3323–3332.
- [117] V.V. Ogryzko, R.L. Schiltz, V. Russanova, et al., The transcriptional coactivators p300 and CBP are histone acetyltransferases, *Cell* 87 (1996) 953–959.
- [118] S.A. Onate, S.Y. Tsai, M.J. Tsai, et al., Sequence and characterization of a coactivator for the steroid hormone receptor superfamily, *Science* 270 (1995) 1354–1357.
- [119] M.L. Paddock, S.E. Wiley, H.L. Axelrod, et al., MitoNEET is a uniquely folded 2Fe 2S outer mitochondrial membrane protein stabilized by pioglitazone, *Proc. Natl. Acad. Sci. USA* 104 (2007) 14342–14347.
- [120] K.S. Park, R.D. Lee, S.K. Kang, et al., Neuronal differentiation of embryonic midbrain cells by upregulation of peroxisome proliferator-activated receptor-gamma via the JNK-dependent pathway, *Exp. Cell Res.* 297 (2004) 424–433.
- [121] S.W. Park, J.H. Yi, G. Miranpuri, et al., Thiazolidinedione class of peroxisome proliferator-activated receptor gamma agonists prevents neuronal damage, motor dysfunction, myelin loss, neuropathic pain, and inflammation after spinal cord injury in adult rats, *J. Pharmacol. Exp. Ther.* 320 (2007) 1002–1012.
- [122] H.A. Pershadshingh, J. Szollosi, S. Benson, et al., Effects of ciglitazone on blood pressure and intracellular calcium metabolism, *Hypertension* 21 (1993) 1020–1023.
- [123] A. Pisanu, D. Lecca, G. Mulas, et al., Dynamic changes in pro- and anti-inflammatory cytokines in microglia after PPAR- γ agonist neuroprotective treatment in the MPTP mouse model of progressive Parkinson's disease, *Neurobiol. Dis.* 71 (2014) 280–291.
- [124] L. Poulsen, M. Siersbaek, S. Mandrup, PPARs: fatty acid sensors controlling metabolism, *Semin. Cell Dev. Biol.* 23 (2012) 631–639.
- [125] P. Puigserver, Tissue-specific regulation of metabolic pathways through the transcriptional coactivator PGC1- α , *Int. J. Obes.* 29 (Suppl. 1) (2005) S5–S9.
- [126] P. Puigserver, B.M. Spiegelman, Peroxisome proliferator-activated receptor-gamma coactivator 1 alpha (PGC-1 alpha): transcriptional coactivator and metabolic regulator, *Endocr. Rev.* 24 (2003) 78–90.
- [127] P. Puigserver, Z. Wu, C.W. Park, et al., A cold-inducible coactivator of nuclear receptors linked to adaptive thermogenesis, *Cell* 92 (1998) 829–839.
- [128] R.A. Quintanilla, Y.N. Jin, K. Fuenzalida, et al., Rosiglitazone treatment prevents mitochondrial dysfunction in mutant huntingtin-expressing cells: possible role of peroxisome proliferator-activated receptor-gamma (PPAR-gamma) in the pathogenesis of Huntington disease, *J. Biol. Chem.* 283 (2008) 25628–25637.
- [129] M. Ricote, A.C. Li, T.M. Willson, et al., The peroxisome proliferator-activated receptor-gamma is a negative regulator of macrophage activation, *Nature* 391 (1998) 79–82.
- [130] M.E. Risner, A.M. Saunders, J.F. Altman, et al., Efficacy of rosiglitazone in a genetically defined population with mild-to-moderate Alzheimer's disease, *Pharmacogenom. J.* 6 (2006) 246–254.
- [131] J. Rodriguez-Rivera, L. Denner, K.T. Dineley, Rosiglitazone reversal of Tg2576 cognitive deficits is independent of peripheral gluco-regulatory status, *Behav. Brain Res.* 216 (2011) 255–261.
- [132] J.X. Rong, J.L. Klein, Y. Qiu, et al., Rosiglitazone induces mitochondrial biogenesis in differentiated murine 3T3-L1 and C3H/10T1/2 adipocytes, *PPAR Res.* 2011 (2011) 179454.
- [133] E.D. Rosen, P. Sarraf, A.E. Troy, et al., PPAR gamma is required for the differentiation of adipose tissue in vivo and in vitro, *Mol. Cell* 4 (1999) 611–617.
- [134] E.D. Rosen, B.M. Spiegelman, Molecular regulation of adipogenesis, *Annu. Rev. Cell Dev. Biol.* 16 (2000) 145–171.
- [135] E.D. Rosen, C.J. Walkey, P. Puigserver, et al., Transcriptional regulation of adipogenesis, *Genes Dev.* 14 (2000) 1293–1307.
- [136] A. Rubenstrunk, R. Hanf, D.W. Hum, et al., Safety issues and prospects for future generations of PPAR modulators, *Biochim. Biophys. Acta* 1771 (2007) 1065–1081.
- [137] D. Sau, S. De Biasi, L. Vitellaro-Zuccarello, et al., Mutation of SOD1 in ALS: a gain of a loss of function, *Hum. Mol. Genet.* 16 (2007) 1604–1618.
- [138] A. Sawa, G.W. Wiegand, J. Cooper, et al., Increased apoptosis of Huntington disease lymphoblasts associated with repeat length-dependent mitochondrial depolarization, *Nat. Med.* 5 (1999) 1194–1198.
- [139] K. Schoonjans, J. Peinado-Onsurbe, A.M. Lefebvre, et al., PPAR α and PPAR γ activators direct a distinct tissue-specific transcriptional response via a PPRE in the lipoprotein lipase gene, *EMBO J.* 15 (1996) 5336–5348.
- [140] F.J. Schopfer, Y. Lin, P.R. Baker, et al., Nitrolinoleic acid: an endogenous peroxisome proliferator-activated receptor gamma ligand, *Proc. Natl. Acad. Sci. USA* 102 (2005) 2340–2345.
- [141] B. Schutz, J. Reimann, L. Dumitrescu-Ozimek, et al., The oral antidiabetic pioglitazone protects from neurodegeneration and amyotrophic lateral sclerosis-like symptoms in superoxide dismutase-G93A transgenic mice, *J. Neurosci.* 25 (2005) 7805–7812.
- [142] P. Seale, S. Kajimura, W. Yang, et al., Transcriptional control of brown fat determination by PRDM16, *Cell Metab.* 6 (2007) 38–54.
- [143] R.K. Semple, V.K. Chatterjee, S. O'Rahilly, PPAR gamma and human metabolic disease, *J. Clin. Investig.* 116 (2006) 581–589.
- [144] Z.Q. Shao, Z.J. Liu, Neuroinflammation and neuronal autophagic death were suppressed via Rosiglitazone treatment: new evidence on neuroprotection in a rat model of global cerebral ischemia, *J. Neurol. Sci.* 349 (2015) 65–71.
- [145] N. Shibata, M. Kawaguchi-Niida, T. Yamamoto, et al., Effects of the PPAR-gamma activator pioglitazone on p38 MAP kinase and I κ B α in the spinal cord of a transgenic mouse model of amyotrophic lateral sclerosis, *Neuropharmacology* 28 (2008) 387–398.
- [146] J. St-Pierre, S. Drori, M. Uldry, et al., Suppression of reactive oxygen species and neurodegeneration by the PGC-1 transcriptional coactivators, *Cell* 127 (2006) 397–408.
- [147] J.C. Strum, R. Shehee, D. Virley, et al., Rosiglitazone induces mitochondrial biogenesis in mouse brain, *J. Alzheimers Dis.* 11 (2007) 45–51.
- [148] D. Sulzer, D.J. Surmeier, Neuronal vulnerability, pathogenesis, and Parkinson's disease, *Mov. Disord.* 28 (2013) 41–50.
- [149] S. Sundararajan, J.L. Gamboa, N.A. Victor, et al., Peroxisome proliferator-activated receptor-gamma ligands reduce inflammation and infarction size in transient focal ischemia, *Neuroscience* 130 (2005) 685–696.
- [150] C. Swanson, M. Emborg, Expression of peroxisome proliferator-activated receptor-gamma in the substantia nigra of hemiparkinsonian nonhuman

- primates, *Neurol. Res.* 36 (2014) 634–646.
- [151] C.R. Swanson, E. Du, D.A. Johnson, et al., Neuroprotective Properties of a Novel Non-Thiazolidinedione Partial PPAR- γ Agonist against MPTP, *PPAR Res.* 2013 (2013) 582809.
- [152] C.R. Swanson, V. Joers, V. Bondarenko, et al., The PPAR- γ agonist pioglitazone modulates inflammation and induces neuroprotection in parkinsonian monkeys, *J. Neuroinflamm.* 8 (2011) 91.
- [153] E.M. Toledo, N.C. Inestrosa, Activation of Wnt signaling by lithium and rosiglitazone reduced spatial memory impairment and neurodegeneration in brains of an APP^{swe}/PSEN1 Δ E9 mouse model of Alzheimer's disease, *Mol. Psychiatry* 15 (272–285) (2010) 228.
- [154] P. Tontonoz, E. Hu, R.A. Graves, et al., mPPAR γ 2: tissue-specific regulator of an adipocyte enhancer, *Genes Dev.* 8 (1994) 1224–1234.
- [155] P. Tontonoz, E. Hu, B.M. Spiegelman, Stimulation of adipogenesis in fibroblasts by PPAR γ 2, a lipid-activated transcription factor, *Cell* 79 (1994) 1147–1156.
- [156] P. Tontonoz, E. Hu, B.M. Spiegelman, Regulation of adipocyte gene expression and differentiation by peroxisome proliferator activated receptor γ , *Curr. Opin. Genet. Dev.* 5 (1995) 571–576.
- [157] P. Tontonoz, B.M. Spiegelman, Fat and beyond: the diverse biology of PPAR- γ , *Annu. Rev. Biochem.* 77 (2008) 289–312.
- [158] J. Tordjman, W. Khazen, B. Antoine, et al., Regulation of glyceroneogenesis and phosphoenolpyruvate carboxylase by fatty acids, retinoic acids and thiazolidinediones: potential relevance to type 2 diabetes, *Biochimie* 85 (2003) 1213–1218.
- [159] E. Utreras, R. Hamada, M. Prochazkova, et al., Suppression of neuroinflammation in forebrain-specific Cdk5 conditional knockout mice by PPAR- γ agonist improves neuronal loss and early lethality, *J. Neuroinflamm.* 11 (2014) 28.
- [160] T. Varga, Z. Czimmerer, L. Nagy, PPARs are a unique set of fatty acid regulated transcription factors controlling both lipid metabolism and inflammation, *Biochim. Biophys. Acta* 1812 (2011) 1007–1022.
- [161] N.A. Victor, E.W. Wanderi, J. Gamboa, et al., Altered PPAR γ expression and activation after transient focal ischemia in rats, *Eur. J. Neurosci.* 24 (2006) 1653–1663.
- [162] A.J. Vidal-Puig, R.V. Considine, M. Jimenez-Linan, et al., Peroxisome proliferator-activated receptor gene expression in human tissues. Effects of obesity, weight loss, and regulation by insulin and glucocorticoids, *J. Clin. Investig.* 99 (1997) 2416–2422.
- [163] L. Villacorta, F.J. Schopfer, J. Zhang, et al., PPAR γ and its ligands: therapeutic implications in cardiovascular disease, *Clin. Sci.* 116 (2009) 205–218.
- [164] X. Wang, Z. Wang, J.Z. Liu, et al., Double antioxidant activities of rosiglitazone against high glucose-induced oxidative stress in hepatocyte, *Toxicol. In Vitro* 25 (2011) 839–847.
- [165] Y.L. Wang, K.A. Frauwirth, S.M. Rangwala, et al., Thiazolidinedione activation of peroxisome proliferator-activated receptor γ can enhance mitochondrial potential and promote cell survival, *J. Biol. Chem.* 277 (2002) 31781–31788.
- [166] J. Warne, G. Pryce, J.M. Hill, et al., Selective inhibition of the mitochondrial permeability transition pore protects against neurodegeneration in experimental multiple sclerosis, *J. Biol. Chem.* 291 (2016) 4356–4373.
- [167] S.E. Wiley, A.N. Murphy, S.A. Ross, et al., MitoNEET is an iron-containing outer mitochondrial membrane protein that regulates oxidative capacity, *Proc. Natl. Acad. Sci. USA* 104 (2007) 5318–5323.
- [168] T.M. Willson, M.H. Lambert, S.A. Kliewer, Peroxisome proliferator-activated receptor γ and metabolic disease, *Annu. Rev. Biochem.* 70 (2001) 341–367.
- [169] L. Wilson-Fritch, S. Nicoloso, M. Chouinard, et al., Mitochondrial remodeling in adipose tissue associated with obesity and treatment with rosiglitazone, *J. Clin. Investig.* 114 (2004) 1281–1289.
- [170] A.G. Winterstein, Rosiglitazone and the risk of adverse cardiovascular outcomes, *Clin. Pharmacol. Ther.* 89 (2011) 776–778.
- [171] C. Wolfrum, M. Stoffel, Coactivation of Foxa2 through Pgc-1 β promotes liver fatty acid oxidation and triglyceride/VLDL secretion, *Cell Metab.* 3 (2006) 99–110.
- [172] J.S. Wu, T.N. Lin, K.K. Wu, Rosiglitazone and PPAR- γ overexpression protect mitochondrial membrane potential and prevent apoptosis by upregulating anti-apoptotic Bcl-2 family proteins, *J. Cell Physiol.* 220 (2009) 58–71.
- [173] M.P. Wymann, R. Schreiner, Lipid signalling in disease, *Nat. Rev. Mol. Cell Biol.* 9 (2008) 162–176.
- [174] F. Xu, J. Li, W. Ni, et al., Peroxisome proliferator-activated receptor- γ agonist 15d-prostaglandin J2 mediates neuronal autophagy after cerebral ischemia-reperfusion injury, *PLoS One* 8 (2013) e55080.
- [175] M. Yamanaka, T. Ishikawa, A. Griep, et al., PPAR γ /RXR α -induced and CD36-mediated microglial amyloid- β phagocytosis results in cognitive improvement in amyloid precursor protein/presenilin 1 mice, *J. Neurosci.* 32 (2012) 17321–17331.
- [176] M.B. Yim, J.H. Kang, H.S. Yim, et al., A gain-of-function of an amyotrophic lateral sclerosis-associated Cu,Zn-superoxide dismutase mutant: An enhancement of free radical formation due to a decrease in Km for hydrogen peroxide, *Proc. Natl. Acad. Sci. USA* 93 (1996) 5709–5714.
- [177] F. Zhang, B.E. Lavan, F.M. Gregoire, Selective modulators of PPAR- γ activity: molecular aspects related to obesity and side-effects, *PPAR Res.* 2007 (2007) 32696.
- [178] X. Zhao, J.E. Quigley, J. Yuan, et al., PPAR- α activator fenofibrate increases renal CYP-derived eicosanoid synthesis and improves endothelial dilator function in obese Zucker rats, *Am. J. Physiol. Heart Circ. Physiol.* 290 (2006) H2187–2195.
- [179] X. Zhao, R. Strong, J. Zhang, et al., Neuronal PPAR γ deficiency increases susceptibility to brain damage after cerebral ischemia, *J. Neurosci.* 29 (2009) 6186–6195.
- [180] X. Zhao, Y. Zhang, R. Strong, et al., 15d-Prostaglandin J2 activates peroxisome proliferator-activated receptor- γ , promotes expression of catalase, and reduces inflammation, behavioral dysfunction, and neuronal loss after intracerebral hemorrhage in rats, *J. Cereb. Blood Flow Metab.* 26 (2006) 811–820.
- [181] Y. Zhu, C. Qi, C. Calandra, et al., Cloning and identification of mouse steroid receptor coactivator-1 (mSRC-1), as a coactivator of peroxisome proliferator-activated receptor γ , *Gene Expr.* 6 (1996) 185–195.
- [182] J.M. Zolezzi, C. Silva-Alvarez, D. Ordenes, et al., Peroxisome proliferator-activated receptor (PPAR) γ and PPAR α agonists modulate mitochondrial fusion-fission dynamics: relevance to reactive oxygen species (ROS)-related neurodegenerative disorders? *PLoS One* 8 (2013) e64019.