

HLA Immunogenotype Determines Persistent Human Papillomavirus Virus Infection in HIV-Infected Patients Receiving Antiretroviral Treatment

Rhonda Meys,¹ Karin J. Purdie,⁵ Maurits N. C. de Koning,⁸ Koen D. Quint,^{8,9} Ann-Margaret Little,^{6,7} Finnuuala Baker,⁶ Nick Francis,⁴ David Asboe,³ David Hawkins,³ Steven G. E. Marsh,^{6,7} Catherine A. Harwood,⁵ Frances M. Gotch,² and Christopher B. Bunker¹

¹Department of Dermatology, ²Department of Immunology, ³Department of HIV Medicine and Sexual Health, Chelsea and Westminster Hospital, ⁴Department of Histopathology, Charing Cross Hospital, Imperial College London, ⁵Centre for Cutaneous Research, Barts and the London School of Medicine and Dentistry, Queen Mary University of London, ⁶Anthony Nolan Research Institute, Royal Free Hospital, and ⁷Cancer Institute, University College London, United Kingdom; ⁸DDL Diagnostic Laboratory, Rijswijk, and ⁹Department of Dermatology, Leiden University Medical Centre, The Netherlands

Keywords. human papillomavirus; HPV; human immunodeficiency virus; HIV; human leukocyte antigen; HLA; immunogenetics; warts; cutaneous warts; genital warts.

Abstract

A proportion of human immunodeficiency virus (HIV)–infected patients develop persistent, stigmatizing human papillomavirus (HPV)–related cutaneous and genital warts and anogenital (pre)cancer. This is the first study to investigate immunogenetic variations that might account for HPV susceptibility and the largest to date to categorize the HPV types associated with cutaneous warts in HIV-positive patients. The HLA class I and II allele distribution was analyzed in 49 antiretroviral (ART)–treated HIV-positive patients with persistent warts, 42 noninfected controls, and 46 HIV-positive controls. The allele HLA-B*44 was more frequently identified in HIV-positive patients with warts ($P = .004$); a susceptible haplotype (HLA-B*44, HLA-C*05; $P = .001$) and protective genes (HLA-DQB1*06; $P = .03$) may also contribute. Cutaneous wart biopsy specimens from HIV-positive patients harbored common wart types HPV27/57, the unusual wart type HPV7, and an excess of Betapapillomavirus types ($P = .002$), compared with wart specimens from noninfected controls. These findings suggest that HLA testing might assist in stratifying those patients in whom vaccination should be recommended.

Introduction

Human immunodeficiency virus (HIV)–infected individuals are commonly afflicted with troublesome cutaneous and genital warts and genital precancer and cancer despite receipt of fully suppressive antiretroviral therapy (ART). The virus that causes warts, human papillomavirus (HPV), is classified on the basis of its DNA sequence into 5 genera, containing >120 different genotypes: Alphapapillomavirus (predominantly comprising genital HPV types but also including cutaneous wart types), Mupapillomavirus, Nupapillomavirus, Gammapapillomavirus (containing cutaneous wart types), and Betapapillomavirus (largely nonpathogenic cutaneous types) [1, 2]. HPV is ubiquitous; HPV types belonging to the Betapapillomavirus genus are detected with a high prevalence (>95%) in the general population, and it is estimated that around 75% of all individuals have been exposed to genital HPV types, mostly subclinically and without morbidity [3, 4]. HPV-related disease in HIV-infected patients is common and frequently results in a disabling and stigmatizing dermatosis, consuming significant clinical resources. HIV infection probably increases the prevalence of warts and other HPV-related diseases, and these diseases do not appear to abate with fully suppressive ART [5–8]. Persistent HPV genital disease in HIV-infected patients can lead to the development of precancerous lesions with the risk of progression to frank cancer. However, many individuals remain disease-free

despite presumed HPV exposure, whereas others are particularly troubled by HPV-associated diseases. This transition is likely to be augmented by long-standing HIV infection [9, 10]. The pattern and severity of skin disease in HIV-infected patients may be determined by factors that transcend their level of immunodysfunction, such as immunogenotype, of which CD4⁺ T-cell count is just one marker. We investigated the immunogenotype (specifically, the HLA type) of HIV-infected patients with persistent viral warts and identified the HPV types in HIV-positive individuals with cutaneous and genital warts.

Methods

Subjects

Ethics approval was sought and received from the Riverside NHS Research Ethics Committee (London, United Kingdom) before study commencement. Informed written consent was taken from all patients. Participants were white Caucasoid (ie, they reported parental and grandparental ancestry from Northern Europe, defined as the United Kingdom, Ireland, Netherlands, Belgium, France, Germany, Switzerland, and the Scandinavian countries). In brief, 3 patient groups were recruited: (1) HIV-positive cases, defined as HIV-positive individuals with HPV disease and a history of persistent (duration, ≥ 6 months) cutaneous/genital warts (with or without a history of HPV related anogenital precancer); (2) HIV-positive controls, defined as HIV-positive patients without a history of persistent HPV disease; and (3) HIV-negative cases, defined as individuals without HIV infection and with HPV disease (as defined above).

All HIV-positive individuals were male, were infected with HIV-1 and had been taking fully suppressive ART for ≥ 1 year (HIV load < 50 copies/mL for ≥ 6 months). HIV-positive controls had an absent or transitory past history (< 3 months) of warts and no evidence of other HPV-related diseases, based on their medical history, case note review, and dermatological examination findings. Females with cutaneous warts (with or without genital warts) were included in the HIV-negative case group, but females with a history of genital warts or cervical abnormalities alone were not. HIV-negative cases were individuals with persistent warts in good physical health with no clinical indicators and no other risk factors for HIV infection (as assessed by private interview and thorough dermatological examination) and/or a negative result of an HIV antibody test. To be included as HIV-negative cases, men who have sex with men (MSM) either had a recent negative result of an HIV test or consented to HIV antibody testing (all test results were negative for HIV antibody). From a cohort of approximately 5000 HIV-infected patients attending the HIV clinic at Chelsea and Westminster Hospital (London) and from other clinics in the same hospital, 137 Caucasoid individuals were recruited (49 HIV-positive cases, 42 HIV-negative cases, and 46 HIV-positive controls). These individuals were HLA typed. A subset of HLA-typed

patients with warts at the time of review provided fresh wart tissue for HPV typing, resulting in 106 wart specimens from 30 HIV-positive cases and 36 HIV-negative cases. Demographic information was collected for all patients and included ethnicity, age, sex, total wart duration for cases (volunteered by the patient and corroborated by patient case records), year of HIV infection diagnosis and ART commencement, nadir CD4⁺ T-cell count (corroborated by the year of ART initiation and defined as the lowest historical recorded CD4⁺ T-cell count).

HLA Typing

DNA was extracted from fresh frozen whole-blood specimens, using standard Qiagen protocols (Qiagen, Crawley, United Kingdom) [11] and underwent HLA typing by a validated laboratory (Anthony Nolan Histocompatibility Laboratories, London), using a commercial Luminex-based platform (One Lambda LABtype SSO typing, VH Bio, Gateshead, United Kingdom). HLA allele frequencies for Caucasoid individuals derived from UK based populations were also available [12–14]. HLA-A, HLA-B, and HLA-DRB1 allele frequencies were taken from a population of 5024 Caucasoid blood donors who attended a UK blood transfusion centre; HLA-C frequencies were derived from a subgroup of 2420 individuals from the same population. HLA-DQB1 frequencies were from 177 individuals included in a London anthropology study comprising subjects from England, Scotland, Wales, and Northern Ireland; HLA-B*44 allele subtypes were obtained from 298 Caucasoid blood donors from Liverpool, United Kingdom.

HPV Typing

Study subjects with warts were offered surgical treatment (curettage/shave biopsy or excision). Wart biopsy specimens were divided: half were sent in 10% formalin for confirmatory histopathologic analysis, and the remaining fresh tissue specimens were snap frozen and used for DNA extraction; one of the 106 samples was derived from recently processed formalin-fixed paraffin-embedded tissue. Clinically diagnosed (by a dermatologist) viral warts that were located on sites not easily amenable to surgical biopsy under local anesthetic (eg, plantar, periungual, and finger warts) were pared deeply with a scalpel, and the parings were used for HPV typing. DNA was extracted from tissue by using standard Qiagen protocols as described above [11], with the addition of the following stringent anti-contamination procedures: extended tissue lysis, a longer elution time, and a smaller elution volume. DNA from cutaneous wart associated HPV types was detected using 2 techniques: (1) nested PCR/sequencing designed to detect HPV types 3, 10, 28, 29, 77, 2, 27, 57, 1, 41, and 63, as well as other potentially novel types [15]; and (2) a newly developed Luminex-based platform (HSL-PCR/MPG), using broad-spectrum primers and probes to detect both the types specified above and the additional HPV types 7, 4, 40, 43, 48, 50, 60, 65, 88, 91, 94, and 95 [16]. Thus, 23 cutaneous wart associated HPV types were detectable. Genital and

Betapapillomavirus types were distinguished using 2 separate commercial reverse hybridization line probe assays. The genital assay [17, 18] identified 25 HPV types: high-risk HPV types 16, 18, 31, 33, 35, 39, 45, 51, 52, 56, 58, 59, 66, 68, and 70 and low risk HPV types 6, 11, 34, 40, 42–44, 53, 54, and 74 (HPV SPF₁₀-DEIA-LiPA₂₅ system, version 1 [based on licensed Innogenetics technology]; Labo Biomedical Products, Rijswijk, the Netherlands). The Betapapillomavirus assay [19] contained probes that detected 25 HPV types: HPV 5, 8, 9, 12, 14, 15, 17, 19, 20, 21, 22, 23, 24, 25, 36, 37, 38, 47, 49, 75, 76, 80, 92, 93, and 96 (Diassay B.V., Rijswijk, The Netherlands).

Statistical Analysis

Descriptive and analytical statistics for HPV data were computed with the use of the following standard software programs and with the advice of the Statistical Advisory Service and St Stephens Centre (Imperial College, London): Microsoft Excel; Statistical Package for the Social Sciences (SPSS), version 20; and Epi Info (Centers for Disease Control and Prevention; available at: <http://www.cdc.gov/epiinfo/>; accessed 20 January 2009). Nonparametric data were analyzed with the Mann–Whitney U test. Categorical data were analyzed with Yates' corrected χ^2 or Fisher exact tests (2-tailed). Advice regarding HLA allele counting and immunophenotypic analysis was provided by the Anthony Nolan Histocompatibility Laboratories. A substantial Bonferroni correction for multiple testing was applied to each HLA analysis, to account for all potential allele comparisons, and was accomplished by multiplying the calculated P value by 135 (denoted “P_{corrected}”).

Results

Patient Attributes

Results of HLA typing of DNA from blood specimens were available for 49 HIV-positive cases and 42 HIV-negative cases with persistent warts (Table 1). Of these, most (86% [42 of 49] and 69% [29 of 42], respectively) had a history of cutaneous warts; 39% (19 of 49) and 45% (19 of 42), respectively, had a history of genital warts, and 24% (12 of 49) and 14% (6 of 42), respectively, had a history of both. Three individuals among HIV-positive cases and 3 among HIV-negative cases also had a history of HPV-related penile or anal cancer in situ. The median duration of self-reported warts in this study was between 3 and 5 years for both HIV-positive cases and HIV-negative cases; this long duration is expected because individuals with persistent warts for six months were recruited. HIV-positive cases and HIV-positive controls included significantly more males (100%; P < .0001) and MSM (98%; P < .0001) than HIV-negative cases, a function of the demographic characteristics of HIV clinic attendees (largely male); among HIV-

negative cases, 29% (12 of 42) were female and 15% (6 of 42) were MSM. Females with cutaneous warts were also included as HIV-negative cases because cutaneous warts in males and females are probably comparable. There were no other significant differences between study subjects.

Molecular HLA data were reported as a standard string consisting of 2 allele fields (as defined by the assay manufacturer). The first field was used to determine the allele distribution, by phenotypic allele counting. Therefore, a homozygous individual counted only once toward the allele total (Figures 1 and 2 and Supplementary Table 1).

The main HLA allele of interest that emerged from this study was HLA-B*44 allele: 47% of HIV-positive cases (23 of 49) with HPV disease had this allele, compared with 17% of HIV-positive controls (8 of 46; $P = .004$; $P_{\text{corrected}} = .54$) and with 32% of the blood donor population (1608 of 5024; $P = .04$; $P_{\text{corrected}} = 1.0$; Figure 1). The HLA-B*44 allele count for HIV-negative cases was between that for HIV-positive cases and HIV-positive controls, at 33% (14 of 42). Further interpretation of the allele subgroups revealed that HLA-B*44:02 was the predominant HLA-B*44 subtype; it was present in 38% of HIV-positive cases (18 of 47), 7% of HIV-positive controls (3 of 46; $P = .0006$; $P_{\text{corrected}} = .04$), and 19% of individuals (57 of 298) from a comparative United Kingdom Caucasoid blood donor population ($P = .006$; $P_{\text{corrected}} = .43$). Another allele, HLA-C*05 was also seen more frequently in HIV-positive cases (33% [16 of 49]), compared with HIV-positive controls (9% [4 of 46]; $P = .009$; $P_{\text{corrected}} = .59$). HLA-B*44 and HLA-C*05 are in linkage disequilibrium and constitute a common Caucasoid haplotype. Thirty-seven percent of HIV-positive cases (18 of 49) had the haplotype HLA-B*44, -C*05, compared with only 7% of HIV-positive controls (3 of 46; $P = .001$; $P_{\text{corrected}} = .07$). The allele HLA-DQB1*06 may also be of possible of interest because it was detected more frequently in HIV-positive controls (which makes it potentially protective against HPV disease), compared with United Kingdom Caucasoid blood donors (61% [28 of 46] vs 42% [74 of 177]; $P = .03$; $P_{\text{corrected}} = .8$). There was no significant difference between the allele frequency of HLA-DQB1*06 among HIV-positive cases (43% [21 of 49]) and that among HIV-positive controls (61% [28 of 46]; $P = .12$).

There was no significant difference in the number of subjects from each study group who were homozygous for least 1 allele at HLA-A, -B, -C, -DRB1, and -DRB1 loci: 41% of HIV-positive cases (20 of 49) were homozygotes, compared with 43% of HIV-positive controls (20 of 46) and 57% of HIV-negative cases (24 of 42; $P = .24$).

HPV

The characteristics of individuals in the HIV-negative and HIV-positive case groups who contributed HPV typing data from wart tissue specimens were almost identical to those of individuals included in the HLA analysis. HPV typing was performed for 30 HIV-positive cases (using tissue specimens from 38 cutaneous warts and 10 genital warts) and 36 HIV-negative cases (using tissue specimens from 46 cutaneous warts and 12 genital warts). All 106 sampled warts were positive for at least 1 HPV type (Figures 3 and 4 and Supplementary Table 2). However, in specimens from 4% of cutaneous warts (3 of 84) and 5% of genital warts (1 of 22), no likely pathogenic HPV type was identified, although Betapapillomavirus types were detected at low levels.

Mixed HPV infections were common. For the cutaneous wart associated HPV types and genital HPV types combined, two or more HPV types from either group were detected in 29% (31 of 106) of warts. For the ubiquitous Betapapillomavirus HPV types, the detection of multiple types was higher in HIV-positive cases than in HIV-negative cases (60% [29 of 48] vs 38% [22 of 58]; $P = .03$). The median number of Betapapillomavirus types was also greater in HIV-positive cases than in HIV-negative cases (3 [interquartile range {IQR}, 0.3–8] vs 1 [IQR, 0–2]; $P = .002$). HPV 27 and 57 were the most commonly identified HPV types in all cutaneous warts (from both HIV-positive and HIV-negative cases; Figure 3). HPV 2 was more frequently detected in specimens of cutaneous warts from HIV-negative cases than those from HIV-positive cases (30% [14 of 46] vs 3% [1 of 38]; $P = .002$). HPV 7 was identified in specimens of 8% of cutaneous warts (4 of 38) from HIV-positive cases and in 0% (0 of 46) from HIV-negative individuals ($P = .04$). The HPV 7-positive warts were located on the face and head. After correction for multiple warts from a single individual, this difference for HPV 7 was not statistically significant, with detection in 3 of 27 HIV-positive cases and 0 of 26 HIV-negative cases ($P = .2$). HPV 11 and HPV 6 were identified in genital wart specimens from both HIV-positive and HIV-negative cases, and there was no difference in the HPV type prevalence; the number of genital warts examined was small.

Discussion

To our knowledge, this work is the first to examine the immunogenotypic profile of HIV-infected individuals with warts. The HLA class I allele groups HLA-B*44 (as the allele subtype HLA-B*44:02) and HLA-C*05 were more frequent in HIV-positive cases with persistent warts than in HIV-positive controls. The class II allele HLA-DQB1*06 was found more often in HIV-positive controls than in HIV-positive cases, implying it might have a moderate, possibly protective effect. However, after a conservative Bonferroni correction (ie, multiplication of the P value by 135), none of these associations were statistically significant, except for that of HLA-B*44:02 ($P_{\text{corrected}} = .04$).

The differences in the allele distributions suggest an HLA class I predisposition to HPV disease in HIV-infected individuals. The potential mechanisms behind HLA associations with HPV disease in individuals with treated HIV infection might operate at several levels; 2 viral pathogens are involved.

The best evidence to date for HPV diseases (ie, data on cervical cancer and recurrent respiratory papillomatosis) suggests that predominantly HLA class II alleles appear to govern the host response to HPV. HLA-DRB1*13:01/02 and -DQB1*06:03 have been associated with protection against HPV diseases, whereas HLA-DQB1*06:02 has been identified with discordant associations (protective, neutral, or deleterious) [20–22]. Other alleles, such as HLA-DRB1*03:01 and HLA-DQB1*03:01/02/03, and the haplotype HLA-DRB1*15:01, -DQB1*06:02 have been found to confer susceptibility [21, 23, 24]. There are few reports describing class I associations in HPV. However, HLA-B*44 was previously correlated, in a small number of subjects, with increased progression of HPV 16–related cervical dysplasia [21, 25]. More data are provided by 2 studies that examined HLA class II (but not HLA class I) in immunocompetent individuals with cutaneous warts. None, to our knowledge, have studied HIV-infected patients. HLA-DRB1*03 and HLA-DRB1*09 were found more frequently in Mexican individuals with cutaneous warts than in the general population, and HLA-DR6 (HLA-DRB1*13/14)

was found to be protective [26]. A German study compared 71 otherwise healthy individuals with persistent cutaneous warts lasting >18 months that were positive for HPV 2, 27, and 57 to 92 controls [27]. A statistically significant increase was seen in HLA-DQB1*03:01 and HLA-DQA1*03 in cases, compared with controls, consistent with previous studies in cervical cancer; nonsignificant trends were also identified for alleles, including HLA-DQB1*06:03 (protective). The data from the investigation herein in HIV-infected individuals suggest that HLA-DQB1*06 (all alleles combined) might be protective against HPV. Others have suggested this same allele may be protective in other HPV-related diseases, such as cervical cancer/precancer and recurrent respiratory papillomatosis [21]. One HPV-protective HLA class II allele may be insufficient to prevent HPV disease in HIV-infected individuals, and other protective alleles and/or fewer predisposing alleles may be necessary, such as the haplotype HLA-B*44, -C*05 in the absence of -DQB1 that was seen in some of our HIV-positive cases.

HIV disease progression, on the other hand, is thought to be HLA class I related. Alleles protective for HIV may be able to generate an effective CD8⁺ T-cell response to highly conserved HIV p24 peptides, while deleterious alleles may direct an ineffective immune response [28, 29]. HLA-B*27 and -B*57 have historically been most consistently associated with delayed HIV progression, and HLA-B*35 has been associated with HIV disease that progresses more rapidly.

Other work suggested that the HLA-Bw4 group of alleles (HLA-B*44 is one) may influence the function of natural killer receptors (KIRs) in HIV infection [30, 31]. However, alleles such as HLA-B*27:02, HLA-B*57 (rather than HLA-B*44) are preferential ligands for the KIRs KIR3DL1/KIRDS1, and therefore the effect of weaker ligands such as HLA-B*44 may be clinically insignificant. It is possible that the allele of interest, HLA-B*44:02, is not related to HPV disease and may be a marker of greater HIV disease progression that is not reflected in variables such as HIV infection duration or nadir CD4⁺ T-cell count. Additionally, class II HLA molecules might affect thymic T-cell education. The thymus may be of particular importance in HIV infection during ART receipt because of the recognized emergence of naive CD4⁺ T-cells during immune recovery; this ability declines with aging, as does HIV-related prognosis [32].

This study is also the first to comprehensively categorise the HPV type profile of warts from HIV-infected patients. It supports the limited evidence to date (from case reports) that HPV 7 may be an important pathogen in HIV-infected patients [33–35] and that HPV types commonly associated with warts (such as HPV 27/57 in cutaneous warts) were frequently detected in wart specimens from HIV-infected patients. The only other study that typed HPV from a substantial collection of warts from HIV-infected patients found HPV 7 in specimens from 4% of wart specimens (1 of 25) in HIV-infected individuals but not in 14 wart specimens from non-HIV-infected subjects [36]. In the general population, HPV 7 has been reported infrequently (in 0.5%–2% of individuals) and in unusual sites [16, 37–43]. Immunodysfunction may be important; HPV 7 was found in 17% of 23 wart specimens from 12 organ transplant recipients but was absent in several other studies that typed HPV in wart specimens from immunosuppressed patients [15, 44–47]. However, it is possible that rarer HPV types (such as HPV 7) may be overrepresented in our HIV-infected cohort owing to propagation of the infection between individuals in close contact within the cohort (ie, the cohort is epidemiologically closed). MSM may potentially carry more HPV types, particularly genital types, through greater exposure to HPV. HPV 1 has been reported to be a common HPV type in cutaneous warts [38] but was absent from both HIV-positive cases and HIV-negative cases in this study. A predilection of HPV 1 for children, plantar warts, and warts of short duration may have excluded HPV 1 from our study [37, 40, 43]. The excess of HPV 2 in warts from HIV-negative cases might be a compensatory phenomenon related to the smaller number of HIV-negative patients that harbor HPV 57.

What was also apparent from these results is the greater median number of Betapapillomavirus types in warts from HIV-infected patients and the higher proportion of warts with mixed Betapapillomavirus types. This excess of Betapapilloma-virus types in HIV-infected patients might be related to immunodysfunction (similar to the situation in organ transplant recipients) or due

to the number of close contacts or lifetime sex partners (as appears to be the case for genital HPVs).

Epidermodysplasia verruciformis–like eruptions have been reported in HIV-infected individuals [48], although none of the individuals included in this study had this clinical phenotype.

In summary, HIV-infected individuals with warts were more likely to be HLA-B*44 positive and have the haplotype HLA-B*44, -C*05 than either HIV-infected controls or the general population. The potentially protective allele HLA-DQB1*06 was increased in HIV-positive controls to lesser degree. However, the power of the study is too small to draw definite conclusions, and further work is necessary to validate our findings. We have also characterized the HPV profile in this same group of Caucasoid patients and identified some differences from a control population; in particular, our data suggest that HPV 7 may be an important pathogen in HIV-infected patients.

The emergent alleles of interest fit within the context of previous work and potentially may involve the interaction of HLA and KIR ligands. Immunogenotypic influences are unlikely to compose the entire picture, and other factors probably are important in HPV disease during treated HIV infection, such as as-yet unelucidated or unquantified immunodysfunction or variations in immune adaptivity. These might include thymic generation of naive CD4⁺ T cells, immunosenescence or host phenotypic factors such as CD4⁺ T-helper cell type 1/2 polarization, HIV-associated CD4⁺ T-cell predominant immune decimation and dysregulation, regulatory T-cell function, natural killer cells, and/or CD8⁺ T-cell immune responses.

Regarding the clinical usefulness of our HLA findings, they are consistent with the presence of immunosusceptible and immunoprotective genotype(s). This adds indirectly to the argument and rationale for prophylactic HPV vaccination in boys, as well as for postexposure HPV vaccination in the management of morbidity and diminution of mortality due to HPV-related disease in men (and women) in all settings. Although prophylactic vaccines may not be therapeutic, vaccination can impart cross-reactive immunity to other HPV types [49]. This is most beneficial among younger individuals (age, <26 years); MSM; persons at higher risk of HPV-related morbidity, such as HIV-infected patients [50], and perhaps those who immunogenetically predisposed to HPV infection. Our clinical practice is increasingly to recommend HPV vaccination in these settings.

Acknowledgments. We thank Prof Rino Cerio, for providing additional histopathological input; and Dr Bernard North and Dr Joseph Eliahoo (Statistical Advisory Service, Imperial College, London) and Ms Sundhya Mandalia (St Stephens' Centre, Imperial College, London), for providing statistical advice about categorical parametric and nonparametric analyses.

Financial support. This work was supported by the Joint Research Committee of the Chelsea and Westminster Hospital, the Chelsea and Westminster Health Charity, and the Skin Treatment and Research Trust.

Potential conflicts of interest. All authors: No reported conflicts. All authors have submitted the ICMJE Form for Disclosure of Potential Conflicts of Interest. Conflicts that the editors consider relevant to the content of the manuscript have been disclosed.

References

1. Stanley MA. Epithelial cell responses to infection with human papillomavirus. *Clin Microbiol Rev* 2012; 25:215–22.
2. Bernard HU, Burk RD, Chen Z, van Doorslaer K, zur Hausen H, de Villiers EM. Classification of papillomaviruses (PVs) based on 189 PV types and proposal of taxonomic amendments. *Virology* 2010; 401:70–9.
3. de Koning MN, Struijk L, Bavinck JN, et al. Betapapillomaviruses frequently persist in the skin of healthy individuals. *J Gen Virol* 2007; 88:1489–95.
4. Koutsky L. Epidemiology of genital human papillomavirus infection. *Am J Med* 1997; 102:3–8.
5. Price P, Mathiot N, Krueger R, Stone S, Keane NM, French MA. Immune dysfunction and immune restoration disease in HIV patients given highly active antiretroviral therapy. *J Clin Virol* 2001; 22:279–87.
6. Rodrigues LK, Baker T, Maurer T. Cutaneous warts in HIV-positive patients undergoing highly active antiretroviral therapy. *Arch Dermatol* 2001; 137: 1103–4.
7. Maurer T, Rodrigues LK, Ameli N, et al. The effect of highly active antiretroviral therapy on dermatologic disease in a longitudinal study of HIV type 1-infected women. *Clin Infect Dis* 2004; 38:579–84.
8. Dolev JC, Maurer T, Springer G, et al. Incidence and risk factors for verrucae in women. *AIDS* 2008; 22:1213–9.
9. Hagensee ME, Cameron JE, Leigh JE, Clark RA. Human papillomavirus infection and disease in HIV-infected individuals. *Am J Med Sci* 2004; 328:57–63.
10. Bernard JE, Butler MO, Sandweiss L, Weidner N. Anal intraepithelial neoplasia: correlation of grade with p16INK4a immunohistochemistry and HPV in situ hybridization. *Appl Immunohistochem Mol Morphol* 2008; 16:215–20.
11. Qiagen. DNeasy Blood & Tissue Handbook 07/2006. Crawley, UK: Qiagen, 2006.
12. Allele Frequencies in Worldwide Populations. The Allele Frequency Net Database. <http://www.allele-frequencies.net>. Accessed 1 April 2012.
13. Gonzalez-Galarza FF, Christmas S, Middleton D, Jones AR. Allele frequency net: a database and online repository for immune gene frequencies in worldwide populations. *Nucleic Acids Res* 2011; 39:D913–9.
14. Doherty DG, Vaughan RW, Donaldson PT, Mowat AP. HLA DQA, DQB, and DRB genotyping by

- oligonucleotide analysis: distribution of alleles and haplotypes in British caucasoids. *Hum Immunol* 1992; 34:53–63.
15. Harwood CA, Spink PJ, Suretheran T, et al. Degenerate and nested PCR: a highly sensitive and specific method for detection of human papillomavirus infection in cutaneous warts. *J Clin Microbiol* 1999; 37:3545–55.
 16. de Koning MN, ter Schegget J, Eekhof JA, et al. Evaluation of a novel broad-spectrum PCR-multiplex genotyping assay for identification of cutaneous wart-associated human papillomavirus types. *J Clin Microbiol* 2010; 48:1706–11.
 17. Kleter B, van Doorn LJ, ter Schegget J, et al. Novel short-fragment PCR assay for highly sensitive broad-spectrum detection of anogenital human papillomaviruses. *Am J Pathol* 1998; 153:1731–9.
 18. Kleter B, van Doorn LJ, Schrauwen L, et al. Development and clinical evaluation of a highly sensitive PCR-reverse hybridization line probe assay for detection and identification of anogenital human papillomavirus. *J Clin Microbiol* 1999; 37:2508–17.
 19. de Koning M, Quint W, Struijk L, et al. Evaluation of a novel highly sensitive, broad-spectrum PCR-reverse hybridization assay for detection and identification of beta-papillomavirus DNA. *J Clin Microbiol* 2006; 44:1792–800.
 20. Beskow AH, Josefsson AM, Gyllensten UB. HLA class II alleles associated with infection by HPV16 in cervical cancer in situ. *Int J Cancer* 2001; 93: 817–22.
 21. Hildesheim A, Wang SS. Host and viral genetics and risk of cervical cancer: a review. *Virus Res* 2002; 89:229–40.
 22. Madkan VK, Cook-Norris RH, Steadman MC, Arora A, Mendoza N, Tyring SK. The oncogenic potential of human papillomaviruses: a review on the role of host genetics and environmental cofactors. *Br J Dermatol* 2007; 157:228–41.
 23. Gelder CM, Williams OM, Hart KW, et al. HLA class II polymorphisms and susceptibility to recurrent respiratory papillomatosis. *J Virol* 2003; 77:1927–39.
 24. Cuzick J, Terry G, Ho L, et al. Association between high-risk HPV types, HLA DRB1* and DQB1* alleles and cervical cancer in British women. *Br J Cancer* 2000; 82:1348–52.
 25. Bontkes HJ, van Duin M, de Gruijl TD, et al. HPV 16 infection and progression of cervical intra-epithelial neoplasia: analysis of HLA polymorphism and HPV 16 E6 sequence variants. *Int J Cancer* 1998; 78:166–71.
 26. Garcia-Corona C, Vega-Memije E, Barquera R, Granados J. HLA-DR alleles associated with skin warts induced by human papillomavirus infection. *Int J Dermatol* 2010; 49:1376–9.
 27. Spelten B, Grussendorf-Conen EI, Rubben A. Human leukocyte antigen class II alleles and natural history of HPV 2/27/57-induced common warts. *Arch Dermatol Res* 2004; 296:105–11.
 28. Blackwell JM, Jamieson SE, Burgner D. HLA and infectious diseases. *Clin Microbiol Rev* 2009; 22:370–85.
 29. Pelak K, Need AC, Fellay J, et al. Copy number variation of KIR genes influences HIV-1 control. *PLoS Biol* 2011; 9:e1001208.
 30. Martin MP, Qi Y, Gao X, et al. Innate partnership of HLA-B and KIR3DL1 subtypes against HIV-1. *Nat Genet* 2007; 39:733–40.
 31. Martin MP, Gao X, Lee JH, et al. Epistatic interaction between KIR3DS1 and HLAB delays the progression to AIDS. *Nat Genet* 2002; 31:429–34.
 32. Janeway CA, Travers P, Walport M, Shlomchik MJ. *Immunobiology*. 6th ed. New York: Garland Science, 2005.
 33. Greenspan D, de Villiers EM, Greenspan JS, de Souza YG, zur Hausen H. Unusual HPV types in oral warts in association with HIV infection. *J Oral Pathol* 1988; 17:482–8.

34. deVilliers EM. Prevalence of HPV 7 papillomas in the oral mucosa and facial skin of patients with human immunodeficiency virus. *Arch Dermatol* 1989; 125:1590.
35. Volter C, He Y, Delius H, et al. Novel HPV types present in oral papillomatous lesions from patients with HIV infection. *Int J Cancer* 1996; 66:453–6.
36. Porro AM, Alchorne MM, Mota GR, Michalany N, Pignatari AC, Souza IE. Detection and typing of human papillomavirus in cutaneous warts of patients infected with human immunodeficiency virus type 1. *Br J Dermatol* 2003; 149:1192–9.
37. Rubben A, Kalka K, Spelten B, Grussendorf-Conen EI. Clinical features and age distribution of patients with HPV 2/27/57-induced common warts. *Arch Dermatol Res* 1997; 289:337–40.
38. Ritzkowsky A, Weissenborn S, Krieg T, Pfister H, Wieland U. Extensive human papillomavirus type 7-associated orofacial warts in an immunocompetent patient. *Acta Derm Venereol* 2001; 81:130–3.
39. Iftner A, Klug SJ, Garbe C, et al. The prevalence of human papillomavirus genotypes in nonmelanoma skin cancers of nonimmunosuppressed individuals identifies high-risk genital types as possible risk factors. *Cancer Res* 2003; 63:7515–9.
40. Hagiwara K, Uezato H, Arakaki H, et al. A genotype distribution of human papillomaviruses detected by polymerase chain reaction and direct sequencing analysis in a large sample of common warts in Japan. *J Med Virol* 2005; 77:107–12.
41. Matsukura T, Mitsuishi T, Sugase M, Kawashima M. Human papillomavirus type 7-associated condyloma. *Dermatology* 2010; 221:5–8.
42. Sun C, Chen K, Gu H, Chang B, Jiang M. Association of human papillomavirus 7 with warts in toe webs. *Br J Dermatol* 2010; 162:579–86.
43. Bruggink SC, de Koning MN, Gussekloo J, et al. Cutaneous wart-associated HPV types: prevalence and relation with patient characteristics. *J Clin Virol* 2012; 55:250–5.
44. Shamanin V, Glover M, Rausch C, et al. Specific types of human papillomavirus found in benign proliferations and carcinomas of the skin in immunosuppressed patients. *Cancer Res* 1994; 54:4610–3.
45. Stark LA, Arends MJ, McLaren KM, et al. Prevalence of human papillomavirus DNA in cutaneous neoplasms from renal allograft recipients supports a possible viral role in tumour promotion. *Br J Cancer* 1994; 69:222–9.
46. Kohler A, Meyer T, Stockfleth E, Nindl I. High viral load of human wart-associated papillomaviruses (PV) but not beta-PV in cutaneous warts independent of immunosuppression. *Br J Dermatol* 2009; 161:528–35.
47. Martelli-Marzagao F, Yamashiro AS, Ogawa MM, Santos GF Jr, Tomimori J, Porro AM. Clinical and histopathological characterization and typing of the human papillomavirus in common warts of kidney transplant recipients. *An Bras Dermatol* 2010; 85:743–6.
48. Jacobelli S, Laude H, Carlotti A, et al. Epidermodysplasia verruciformis in human immunodeficiency virus-infected patients: a marker of human papillomavirus-related disorders not affected by antiretroviral therapy. *Arch Dermatol* 2011; 147:590–6.
49. Herrero R. Human Papillomavirus (HPV) Vaccines: Limited Cross-Protection against Additional HPV Types. *J Infect Dis* 2009; 199:919–22.
50. Markowitz LE, Dunne EF, Saraiya M, et al. Human papillomavirus vaccination: recommendations of the Advisory Committee on Immunization Practices (ACIP). *MMWR Recomm Rep* 2014; 63(RR-05):1–30.

Table 1: Study Participant Demographics

	Group			HIV+ cases compared with HIV+ controls	HIV+ cases compared with NI cases
	HIV+ cases	HIV+ controls	Non-Infected cases	p=significance level	p=significance level
Number of individuals (n)	49	46	42	—	—
Age years median (IQR)	45 (41-54)	49.5 (44-56)	41 (32-52)	p=0.05	p=0.06
Sex (% male)	100%	100%	71%	p=1*	p<0.0001*
HIV Infection years median (IQR)	14 (8-19)	11 (7-20)	—	p=0.62	—
ART years median (IQR)	9.0 (5-13)	8.5 (5-12)	—	p=0.96	—
Risk factors for HIV (% MSM)	98%	98%	15%	p=1*	p<0.0001*
Current CD4 T-cell cells/μl median (IQR)	483 (343-693)	544 (431-683)	—	p=0.41	—
CD4 T-cell nadir cells/μl median (IQR)	148 (41-205)	137 (48-205)	—	p=0.86	—
Viral load % undetectable (<50 copies/ml)	100%	100%	—	p=1*	—
Wart disease years median (IQR)	4.0 (1.8-9)	—	3.5 (1.4-5)	—	p=0.42

Figure 1: HLA-B allele count distribution as percentage of individuals in each group




Figure 2: HLA-A, HLA-C, HLA-DRB1 and HLA-DQB1 allele count distributions as percentage of individuals in each group







Figure 3: Cutaneous warts - HPV type profile




Figure 4: Genital warts - HPV type profile

