

Table1: Pre, per and post-operative details of the study eyes (N=161). Percents given are out of the total for which records were available.

	present n (%)	missing record n
PRE-OPERATIVE		
secondary glaucoma (uveitis, ocular trauma or vascular accident)	13 (8%)	4
advanced glaucoma (= cup/disc $\geq 0,8$)	119 (79%)	10
very high preoperative IOP (≥ 40 mmHg)	38 (24%)	3
preoperative WHO-VA category		8
<i>normal vision</i>	43 (28%)	
<i>visual impairment</i>	52 (34%)	
<i>blind</i>	58 (38%)	
PER-OPERATIVE		
scleral flap		33
<i>fornix based</i>	120 (94%)	
<i>limbus based</i>	8 (6%)	
simultaneous cataract extraction	9 (7%)	33
use of releasable sutures	94 (60%)	3
operated by one surgeon (SDS)	99 (67%)	14
use of antimetabolite		16
<i>none</i>	17 (12%)	
<i>5-Fluorouracil</i>	33 (23%)	
<i>mitomycin C</i>	95 (66%)	
POST-OPERATIVE		
early complications (< 1 week postop)	27 (17%)	6
late complications (≥ 1 week postop)	21 (14%)	11
re-intervention needed	21 (13%)	4

Figure 1: Complications and re-interventions in the 161 study eyes.




Table 2: Ophthalmic findings at the time of the study visit (N=161). Percents given are out of the total for which records were available.

	present	missing records
	n (%)	n
postoperative IOP \leq 15mmHg	113 (71%)	1
postoperative IOP < 21mmHg	153 (96%)	1
IOP reduction of \geq 30%	133 (85%)	4
Bleb description		
<i>white</i>	61 (38%)	1
<i>elevated</i>	121 (76%)	2
lens status		
<i>phakic</i>	134 (85%)	3
<i>pseudophakic</i>	19 (12%)	
<i>aphakic</i>	5 (3%)	
chronic uveitis	1 (1%)	1
compliant to follow-up visits (\geq 5 visits after trabeculectomy)	69 (43%)	0
long total follow-up duration (\geq 2 years)	93 (58%)	0
postoperative WHO-visual acuity category		
<i>normal vision</i>	60 (37%)	0
<i>visual impairment</i>	37 (23%)	
<i>blind</i>	64 (40%)	

Table 3: Univariate and multivariate analysis of potential predictors for IOP failure (>20mmHg and/or <30% reduction) in first operated eyes (n= 118 eyes), expressed in absolute and relative number of eyes, odds ratio (OR), 95% confidence intervals (CI) and P-value.

Predictors	UNIVARIATE ANALYSIS			MULTIVARIATE ANALYSIS		
	OR	CI	P-value	OR	CI	P-value
compliant to follow-up visits (≥ 5 visits)	1.08	0.407-2.46	0.88			
male gender	1.087	0.407- 2.941	0.86	1.229	0.413-3.663	0.711
young age at operation (< 51 years)	0.741	0.224- 2.439	0.62	0.977	0.274-3.485	0.971
long total follow-up duration (≥ 2 years)	3.571	1.087- 12.5	0.03	2.90	0.864-9.730	0.085
use of releasable sutures	0.503	0.183- 1.370	0.17			
use of any antimetabolite	0.346	0.862- 1.333	0.11	0.353	0.089-1.402	0.139
use of mitomycine C compared to the rest	0.386	0.135- 1.111	0.07			
3 min duration of mitomycine C application	0.781	0.143- 4.348	0.78			
early complications (< 1 week postop)	0.758	0.198- 2.941	0.68			
late complications (≥ 1 week postop)	1.667	0.469- 5.882	0.42			
reintervention needed	1.087	0.277- 4.348	0.91			
advanced glaucoma (cup/disc ≥ 0,8)	1.587	0.498- 5.263	0.45			
very high preoperative IOP (≥ 40mmHg)	0.495	0.132- 1.852	0.29			
fornix based trabeculectomy technique	1.149	0.127- 10	0.90			
combined with cataract operation	0.735	0.083- 6.667	0.78			
secondary glaucoma	0.633	0.073- 5.556	0.67			
operated by one surgeon (SDS)	0.565	0.193- 1.639	0.29			

Figure 2: Cumulative probability to maintain good outcome (intraocular pressure IOP << 21mmHg and a 30% IOP reduction after trabeculectomy) for the first eyes only.

