

1 **Reported appetite, taste and smell changes following Roux-en-Y gastric bypass**  
2 **and sleeve gastrectomy: effect of gender, type 2 diabetes and relationship to**  
3 **post-operative weight loss**

4  
5 Janine M Makaronidis<sup>1, 2, 3\*</sup>, Sabrina Neilson<sup>1\*</sup>, Wui-Hang Cheung<sup>1, 2</sup>, Urszula Tymoszuk<sup>1</sup>,  
6 Andrea Pucci<sup>1, 2</sup>, Nicholas Finer<sup>1, 2</sup>, Jacqueline Doyle<sup>2</sup>, Majid Hashemi<sup>1, 2</sup>, Mohamed  
7 Elkalaawy<sup>1, 2, 3</sup>, Marco Adamo<sup>2</sup>, Andrew Jenkinson<sup>2</sup>, Rachel L Batterham<sup>1, 2, 4†</sup>

8  
9 \* joint first authors

10

11 **Author Affiliations**

12

13 <sup>1</sup>Centre for Obesity Research, Rayne Institute, Department of Medicine, University College  
14 London, London WC1E 6JF, United Kingdom.

15 <sup>2</sup>University College London Hospital Bariatric Centre for Weight Management and Metabolic  
16 Surgery, University College London Hospital, Ground Floor West Wing, 250 Euston Road,  
17 London NW1 2PG, United Kingdom.

18 <sup>3</sup>Clinical and Experimental Surgery Department, Medical Research Institute, University of  
19 Alexandria, Hadara, Alexandria 21561, Egypt.

20 <sup>4</sup>National Institute of Health Research, University College London Hospital Biomedical  
21 Research Centre, London W1T 7DN, United Kingdom.

22

23 †Correspondence to:

24

25 Professor Rachel L Batterham

26 Centre for Obesity Research

27 Department of Medicine

28 University College London

29 Rayne Building

30 5 University Street

31 London WC1E 6JF

32 Telephone: +44207 6790991

33 [r.batterham@ucl.ac.uk](mailto:r.batterham@ucl.ac.uk)

34

35

36 **Abstract**

37

38 Reduced energy intake drives weight loss following Roux-en-Y gastric bypass (RYGB) and  
39 sleeve gastrectomy (SG) procedures. Post-operative changes in subjective appetite, taste, and  
40 smell and food preferences are reported and suggested to contribute to reduced energy  
41 intake. We aimed to investigate the prevalence of these changes following RYGB and SG and  
42 to evaluate their relationship with weight loss.

43

44 98 patients post-RYGB and 155 post-SG from a single bariatric centre were recruited to a  
45 cross-sectional study. Participants completed a questionnaire, previously utilised in post-  
46 operative bariatric patients, to assess the prevalence of post-operative food aversions and  
47 subjective changes in appetite, taste and smell. Anthropometric data were collected and  
48 percentage weight loss (%WL) was calculated. The relationship between food aversions,  
49 changes in appetite, taste and smell and %WL was assessed. The influence of time post-  
50 surgery, gender and type 2 diabetes (T2D) were evaluated.

51

52 Following RYGB and SG the majority of patients reported food aversions (RYGB=62%,  
53 SG=59%), appetite changes (RYGB=91%, SG=91%) and taste changes (RYGB=64%, SG=59%).  
54 Smell changes were more common post-RYGB than post-SG (RYGB=41%, SG=28%,  $p=0.039$ ).  
55 No temporal effect was observed post-RYGB. In contrast, the prevalence of appetite changes  
56 decreased significantly with time following SG.

57

58 Post-operative appetite changes associated with and predicted higher %WL post-SG but not  
59 post-RYGB. Taste changes associated with and predicted higher %WL following RYGB but not  
60 post-SG. There was no gender effect post-RYGB. Post-SG taste changes were less common in  
61 males (female=65%, males=40%,  $p=0.008$ ). T2D status in females did not influence post-  
62 operative subjective changes. However, in males with T2D, taste changes were less common  
63 post-SG than post-RYGB together with lower %WL (RYGB= $27.5\pm 2.7$ , SG= $14.6\pm 2.1$ ,  $p=0.003$ ).  
64 Further research is warranted to define the biology underlying these differences and to  
65 individualise treatments.

66

67 **Keywords:** Obesity, Roux-en-Y gastric bypass, sleeve gastrectomy, appetite, taste, smell, food  
68 aversions, weight loss.

69

## 70 Introduction

71

72 Bariatric surgery is the most effective treatment for patients with severe obesity, leading to  
73 sustained weight reduction, improved obesity-associated co-morbidities and decreased  
74 mortality (Sjostrom, 2013). The most commonly performed bariatric procedures globally are  
75 the Roux-en-Y gastric bypass (RYGB) and the sleeve gastrectomy (SG), accounting for 45% and  
76 37% respectively of operations undertaken in 2013 (Angrisani et al., 2015). Observational  
77 studies and a limited number of randomised controlled studies suggest that RYGB and SG  
78 produce comparable health improvements in the short-term (Schauer et al., 2014; Sczepaniak,  
79 Owens, Shukla, Perlegos, & Garner, 2015). Other procedures, such as the adjustable gastric  
80 banding (AGB) are now less commonly performed (Angrisani et al., 2015).

81

82 Eating behaviour is a key determinant of the pathogenesis of obesity and weight loss achieved  
83 following bariatric surgery (Manning, Pucci, & Batterham, 2015; Scott & Batterham, 2011). An  
84 energy intake that consistently exceeds energy expenditure leads to weight gain and  
85 eventually obesity (Berthoud, 2011). Obese individuals subjectively rate energy-dense foods  
86 as more pleasant compared to lean individuals (Rissanen et al., 2002). In addition, weight gain  
87 and obesity have been linked to a reduction in taste sensitivity and smell perception (Miras &  
88 le Roux, 2010; Patel, DelGaudio, & Wise, 2015; A. C. Shin, Townsend, Patterson, & Berthoud,  
89 2011). Furthermore, neuroimaging studies have revealed that obese subjects exhibit altered  
90 neural responses within reward regions in response to food cues (Atalayer et al., 2014;  
91 Rissanen et al., 2002).

92

93 The gastrointestinal (GI) tract is established as a key regulator of energy and glucose  
94 homeostasis and it is now clear that changes in gut-derived signals as a consequence of  
95 altered GI anatomy following bariatric surgery play a key role in driving reduced energy intake  
96 and weight loss (Dirksen, Damgaard, et al., 2013; Manning et al., 2015; Scott & Batterham,  
97 2011). Following RYGB and SG patients report reduced hunger in the fasted state, increased  
98 post-meal satiety, changes in subjective taste and altered food preferences (Manning et al.,  
99 2015; Scott & Batterham, 2011). For example, a recent prospective study of 30 patients  
100 undergoing SG reported a 68% decrease in energy intake 6 months post-surgery sustained at  
101 24 months post-surgery. 75% of patients in this study reported reduced preference toward  
102 sweet and fatty foods (Coluzzi et al., 2016).

103

104 In order to assess changes in subjective taste following bariatric surgery Tichansky *et al.*  
105 developed a questionnaire comprised of 23 questions. They reported that subjective taste  
106 changes were more common post-RYGB than following AGB surgery (Tichansky, Boughter, &  
107 Madan, 2006). Subsequently, Graham and colleagues used Tichansky's questionnaire in a  
108 cross-sectional study to evaluate taste changes following RYGB in patients who were a median  
109 19 months post-RYGB (Graham, Murty, & Bowrey, 2014). They added 10 additional questions  
110 assessing subjective changes in appetite, smell and food aversions. They found that 93% of  
111 patients reported a change in appetite, 73% a change in taste, 42% a change in smell and 73%  
112 developed food aversions. Additionally, they reported that patients who developed food  
113 aversions achieved higher absolute post-operative weight loss and greater reduction in body  
114 mass index (BMI) (Graham et al., 2014). Zerrweck *et al.* using the questionnaire from Graham  
115 *et al.*, reported that appetite, taste, smell and food aversions were equally common following  
116 RYGB and SG at 10 months post-surgery (Zerrweck et al., 2015). However, it remains unclear

117 whether these subjective changes in appetite, taste and smell are a consequence of weight  
118 loss *per se* or if they are mediated by bariatric procedure-dependent physiological changes.  
119 The subjective hedonic value of sweet foods has been shown to reduce following RYGB  
120 (Ochner et al., 2011; Ochner et al., 2012; Scholtz et al., 2014). This effect was not observed  
121 in BMI-matched subjects following AGB, suggesting that post-RYGB hedonic responses to  
122 food change independent of weight loss (Scholtz et al., 2014).

123  
124 The impact of bariatric surgery on objectively assessed olfaction and taste sensitivity is  
125 controversial, in part due to methodological issues. There are reports of improved post-  
126 operative taste sensitivity for sweet, salty, sour and bitter (Altun et al., 2016; Holinski,  
127 Menenakos, Haber, Olze, & Ordemann, 2015), no taste sensitivity changes (Pepino et al.,  
128 2014) and improved olfactory sensitivity (Holinski et al., 2015). There is also a suggestion of a  
129 difference between RYGB and SG with improved olfactory sensitivity post-SG but not post-  
130 RYGB (Jurowich et al., 2014).

131  
132 Taste and smell perception are complex processes, integrating a range of sensory, cognitive  
133 and hormonal signals (Cummings, 2015; Miras & le Roux, 2010). Gender, obesity, presence of  
134 T2D and nutritional status (vitamin B12 and zinc levels) have all been reported to impact upon  
135 gustatory and olfactory function (Bustos-Saldana et al., 2009; Deglaire et al., 2015; Fabian,  
136 Beck, Fejerdy, Hermann, & Fabian, 2015; Hwang, Kang, Seo, Han, & Joo, 2016). The tendency  
137 to like fatty and salty tasting foods has been shown to have a linear relationship with  
138 increasing BMI in both males and females (A. C. Shin et al., 2011). However, a liking for sweet  
139 foods is more commonly reported by obese females compared to obese males (Deglaire et  
140 al., 2015) and females outperform males in their ability to detect certain odours (Doty &  
141 Cameron, 2009). T2D *per se* has been linked to impaired taste sensitivity, particularly for  
142 sweet stimuli and to impaired olfaction (Bustos-Saldana et al., 2009). Following RYGB,  
143 patients with T2D loose significantly less weight compared to patients who do not have T2D  
144 (Courcoulas et al., 2015). However, there are no reports comparing the prevalence of changes  
145 in subjective appetite, taste or smell following bariatric surgery in people with T2D compared  
146 to people without T2D.

147  
148 Following RYGB and SG, circulating gut hormone levels are markedly altered and these  
149 changes are suggested to contribute to post-operative appetite changes (Yousseif et al., 2014).  
150 Patients with a poor response to surgery experience increased hunger and reduced satiety  
151 levels. In addition, an attenuated gut hormone response is seen in poor weight loss  
152 responders compared to good weight loss responders (Dirksen, Jorgensen, et al., 2013;  
153 Manning et al., 2015). Interestingly, gut hormones are present in saliva and their cognate  
154 receptors are found on taste buds and olfactory neurons (Acosta et al., 2011; Cummings,  
155 2015; Loch, Breer, & Strotmann, 2015; Y. K. Shin et al., 2008). Hence, it is plausible that gut  
156 hormones mediate gustatory and olfactory changes following bariatric surgery through  
157 weight-independent mechanisms. Of note, RYGB and SG are anatomically very different and  
158 differentially impact upon circulating gut hormone levels (le Roux et al., 2007; Yousseif et al.,  
159 2014). These differences may in turn result in post-procedural differences in appetite, taste  
160 and smell. Whilst the development of food aversions following SG and RYGB has been linked  
161 to increased weight loss, it remains to be established whether subjective change in appetite  
162 taste or smell associate with weight loss (Graham et al., 2014).

163

164 We hypothesized that post-operative subjective changes in appetite, taste and smell would  
165 differ between SG and RYGB patients and be influenced by gender and the presence of T2D.  
166 In addition that appetite, taste and smell changes would associate with post-operative weight  
167 loss. Thus, we aimed to investigate prevalence of appetite, taste, smell changes and food  
168 aversions following RYGB and SG and their relationship to post-operative percentage weight  
169 loss (%WL). We also aimed to evaluate the influence of gender, T2D and time post-surgery  
170 upon these changes.

171

172

## 173 **Methods**

174

175 Patients who attended the University College London Hospitals (UCLH) Bariatric Centre for  
176 Weight Management and Metabolic Surgery for follow up appointments after primary RYGB  
177 or primary SG were invited to participate. Participation was voluntary and informed consent  
178 was obtained in person by a healthcare professional. Ethical approval was obtained from the  
179 National Health Service Research Ethics Committee (ID#09/H0715/65) and the study was  
180 undertaken in accordance with the Helsinki Declaration. Inclusion criteria were adult patients  
181 (18 years or older), following either primary RYGB or SG and proficient in spoken and written  
182 English.

183

184 Prior to surgery all patients had been assessed by a multidisciplinary team and fulfilled the  
185 criteria outlined by the National Institute for Health and Care Excellence (NICE, 2014). In RYGB  
186 stomach size is reduced by the creation of a small gastric pouch (~ 20cm<sup>3</sup>), ingested nutrients  
187 pass rapidly from the gastric pouch directly into the mid-jejunum, bypassing the majority of  
188 the stomach, the duodenum and the proximal jejunum (Olbers, Lonroth, Fagevik-Olsen, &  
189 Lundell, 2003). In SG, 80-90% of the total stomach volume is removed by transecting along  
190 the greater curvature of the stomach (Abu-Jaish & Rosenthal, 2010), the remainder of the GI  
191 tract is left intact and nutrient follow the normal anatomical route.

192

193 Weight was measured using a Walkthrough Platform by a trained health professional.  
194 Participants completed a 33-question questionnaire, developed by Tichansky *et al.*, and  
195 modified by Graham *et al.* (Graham *et al.*, 2014; Tichansky *et al.*, 2006) (appendix). Permission  
196 to use the questionnaire was obtained (Graham *et al.*, 2014). Clinical data including height  
197 and weight on the day of surgery and presence or absence of T2D and exclusion criteria were  
198 obtained from the patients' clinical records. Vitamin B12 and zinc levels were measured by  
199 the UCLH Department of Clinical Biochemistry using a competitive immunoassay (Roche) and  
200 colorimetric assay (Randox) respectively, as part of routine post-surgical care.

201

202 After RYGB or SG surgery, patients are advised to adhere to a soft diet for the first two post-  
203 operative weeks, followed by a soft diet with gradual reintroduction of solid food. Patients  
204 start eating meals of normal textured food 7 weeks after surgery. Thus, in order to eliminate  
205 the effect of early post-operative dietary restriction and allow for their eating behaviour to  
206 be established, patients less than 90 days post-surgery were excluded. Patients with factors  
207 affecting gustatory (including low zinc and low vitamin B12 levels) or olfactory function or  
208 who suffered a severe or debilitating illness, active malignancy and pregnant women were  
209 also excluded.

210

211 Percentage weight loss (%WL) was calculated by the weight difference between the day of  
 212 surgery and the day of questionnaire completion and expressed as percentage of the weight  
 213 on the day of surgery. Data were analysed using GraphPad Prism version 6 and STATA  
 214 statistical software version 13. Mean and standard error of mean (SEM) were calculated.  
 215 Continuous data was assessed for normality using D'Angostino and Pearson omnibus  
 216 normality test. Parametric (t-test or one-way analysis of variance (ANOVA) and non-  
 217 parametric tests (Mann-Whitney tests)) were used as appropriate. Chi-square tests were used  
 218 for categorical data. Furthermore, linear regression analyses were performed. Significance  
 219 was assumed below the 0.05 level.

220  
 221

## 222 Results

223  
 224

### 224 Patient demographics

225 253 patients were included in the final analysis, 98 following RYGB and 155 post-SG. 37  
 226 patients were excluded from the study due pregnancy (n=8), B12 and/or zinc deficiency (n=4),  
 227 symptomatic hypoglycaemia (n=3), conversion of SG to RYGB (n=8), active malignancy (n=4),  
 228 previous cranial radiotherapy (n=1), anosmia (n=2), intolerance to solid foods (n=2), severe  
 229 illness or reduced mobility (n=4) and inability to read English (n=1). Out of the included  
 230 patients, 199 (79%) were female and 54 (20%) male. The patient characteristics are presented  
 231 in Table 1. The RYGB and SG groups had a similar age and BMI but %WL and time post-surgery  
 232 were significantly greater in the RYGB group.

233  
 234  
 235

**Table 1: Patient characteristics**

	Age (years)	BMI (kg/m <sup>2</sup> )	%WL	Time post-surgery (days)
<b>RYGB</b> n=98 (M=19, F=79)	46.5 ± 1.1	44.7 ± 0.7	25.6 ± 0.9	769 ± 53
<b>SG</b> n=155 (M=35, F=120)	44.3 ± 1.0	46.1 ± 0.6	21.2 ± 0.8	593 ± 43
<b>p value</b>	0.120	0.260	0.0001	0.001

236 Mean age, Body Mass Index (BMI), percentage weight loss (%WL) and time post-surgery in patients following  
 237 Roux-en-Y gastric bypass (RYGB) and sleeve gastrectomy (SG)

238  
 239

### 239 Subjective changes in appetite, taste and smell post-surgery

240 The majority of patients reported post-operative change in appetite (Figure 1A, Question 1).  
 241 Changes in taste and to a lesser extent smell were observed following both RYGB and SG, as  
 242 were the development of food aversions (Figure 1B, 1C and 1D). While there were no  
 243 significant differences in appetite, taste and food aversions between the two groups, smell  
 244 changes were significantly more common following RYGB compared to SG (RYGB=41%,  
 245 SG=28%, p=0.039).

246

247 In view of evidence regarding difference in taste perception between males and females, we  
 248 examined the influence of gender upon the frequency of reported appetite, taste and smell  
 249 changes and development of food aversions. No significant differences were found when  
 250 comparing responses of female and male patients following RYGB (Table 2). However, within  
 251 the SG group, taste and smell changes were significantly more common in female compared  
 252 to male patients (Table 2). Furthermore, men post-SG lost significantly less weight compared  
 253 to men following RYGB (%WL: RYGB=26.7 ± 2.13, SG=18.7 ± 1.7 p=0.004)

254  
 255 (Figure 1 here)

256  
 257 **Table 2: Gender differences in the prevalence of subjective appetite, taste, smell changes**  
 258 **and food aversions**  
 259

	RYGB Group (% of patients/number reporting post-surgery change)			SG Group (% of patients/number reporting post-surgery change)		
	Females	Males	p value	Females	Males	p value
<b>Appetite changes</b>	92.4% (73)	84.2% (16)	0.370	91.7% (110)	88.6% (31)	0.521
<b>Taste changes</b>	62.0% (49)	73.7% (14)	0.341	65% (78)	40% (14)	0.008
<b>Smell changes</b>	41.8% (33)	36.8% (7)	0.695	31.7% (38)	14.3% (5)	0.043
<b>Food aversions</b>	45.7% (49)	66.7% (12)	0.713	62.3% (76)	45.7% (16)	0.062

260 Roux-en-Y gastric bypass (RYGB) and sleeve gastrectomy (SG)

261  
 262 **Influence of time post-surgery upon subjective changes in appetite, taste, smell, food**  
 263 **aversions and %WL**

264 In light of the cross-sectional nature of our study, we investigated the effect of time since  
 265 surgery on patients reporting perceived changes in appetite, taste and smell (Table 3).  
 266 Following RYGB, no differences were found in the frequency of changes in appetite, taste,  
 267 smell and food aversions at different time points. In contrast, following SG, the prevalence of  
 268 subjective appetite changes decreased after 3 years (Table 3). The prevalence of taste and  
 269 smell changes also reduced with time post-SG but these failed to reach significance. Following  
 270 RYGB, maximum %WL was achieved at 1-2 years post-surgery and was similar at 2-3 years  
 271 and 3-5 years post-RYGB (Table 4). Maximum %WL following SG was also observed at 1-2  
 272 years post-surgery and was comparable to that seen in the RYGB group. %WL decreased with  
 273 time post-surgery in the SG group (Table 4). Comparison of 2-5 year %WL between the RYGB  
 274 and SG groups revealed greater %WL post-RYGB (%WL; RYGB=26.2±1.3, SG=20.8 ±1.9,  
 275 p=0.023).

276  
 277  
 278  
 279  
 280

281 **Table 3: Effect of time post-surgery on the prevalence of food aversions, subjective**  
 282 **changes in appetite, taste and smell changes**  
 283

<b>RYGB Group</b>				
<b>Time post-surgery</b>	<b>Change in appetite</b>	<b>Change in taste</b>	<b>Change in smell</b>	<b>Food aversions</b>
<b>&lt; 180 days (n=6)</b>	100.0%	50.0%	50.0%	66.7%
<b>180-364 days (n=24)</b>	83.3%	62.5%	45.8%	54.2%
<b>p value</b>	0.468	0.458	0.855	0.709
<b>1-2 years (n=23)</b>	95.7%	73.9%	43.5%	65.2%
<b>p value</b>	0.267	0.289	0.817	0.528
<b>2-3 years (n=16)</b>	100%	62.5%	31.3%	62.5%
<b>p value</b>	0.202	0.795	0.292	0.879
<b>3-5 years (n=29)</b>	86.2%	62.1%	37.9%	65.5%
<b>p value</b>	0.306	0.767	0.706	0.665
<b>SG Group</b>				
<b>Time post-surgery</b>	<b>Change in appetite</b>	<b>Change in taste</b>	<b>Change in smell</b>	<b>Aversions</b>
<b>&lt; 180 days (n=36)</b>	97.2%	75.0%	44.4%	58.5%
<b>180-364 days (n=41)</b>	100.0%	65.9%	24.4%	68.3%
<b>p value</b>	0.283	0.321	0.063	0.879
<b>1-2 years (n=27)</b>	88.9%	48.1%	14.8%	59.3%
<b>p value</b>	0.072	0.065	0.067	0.485
<b>2-3 years (n=19)</b>	84.2%	57.9%	21.1%	52.6%
<b>p value</b>	0.722	0.797	0.499	0.309
<b>3-5 years (n=32)</b>	78.1%	43.8%	28.1%	43.8%
<b>p value</b>	0.001	0.110	0.056	0.068

284 Roux-en-Y gastric bypass (RYGB), sleeve gastrectomy (SG). Fisher's exact correlation, p values represent  
 285 comparison of given time interval versus previous data.  
 286  
 287  
 288  
 289  
 290  
 291  
 292



293  
294  
295  
296

**Table 4: Weight loss at different time points following RYGB and SG**

Time post-surgery	%WL in RYGB patients	%WL in SG patients
< 180 days	18.8 ± 2.2	17.0 ± 1.0
180-364 days	24.6 ± 1.4	23.1 ± 1.1
1-2 years	27.4 ± 1.9	26.2 ± 1.9
2-3 years	26.1 ± 1.9	19.9 ± 3.3
3-5 years	26.3 ± 1.8	20.3 ± 2.1

297 Roux-en-Y gastric bypass (RYGB), sleeve gastrectomy (SG), percentage weight loss (%WL)

298

**Relationship between subjective changes in appetite, taste, smell, food aversion and %WL**

300 We investigated the relationship between post-operative changes in appetite, taste, smell,  
301 food aversions and %WL. In the RYGB group there was significant association between a  
302 change in taste and higher %WL (27.8 ± 1.0 vs. 23.1 ± 1.6, p=0.036). In the SG group a  
303 significantly higher %WL was detected in patients with a change in appetite (21.9 ± 0.8 vs.  
304 13.4 ± 3.1, p=0.006) and food aversions (22.6 ± 1 vs. 19.2 ± 1.3, p=0.032). However, there was  
305 no association between subjective taste changes and %WL.

306

**Table 5: %WL in patients with and without subjective changes in appetite, taste, smell and food aversion post-RYGB**

307  
308  
309

	Appetite	Taste	Smell	Food aversion
<b>Change</b>	25.8 ± 0.9 n=89	27.8 ± 1.0 n=63	26.3 ± 1.5 n=40	26.5 ± 1.1 n=61
<b>No change</b>	23.8 ± 3.2 n=9	23.1 ± 1.6 n=35	25.2 ± 1.1 n=58	24.2 ± 1.2 n=37
<b>p value</b>	0.435	0.036	0.595	0.111

310 Roux-en-Y gastric bypass (RYGB), percentage weight loss (%WL)

311

312

**Table 6: %WL in patients with and without subjective changes in appetite, taste, smell and food aversion post-SG**

313  
314  
315

	Appetite	Taste	Smell	Food aversion
<b>Change</b>	21.9 ± 0.8 n=141	21.9 ± 1.0 n=92	21.7 ± 1.8 n=43	22.6 ± 1.0 n=92
<b>No change</b>	13.4 ± 3.1 n=14	20.3 ± 1.3 n=63	21.1 ± 0.9 n=112	19.2 ± 1.3 n=63
<b>p value</b>	0.006	0.520	0.772	0.032

316 Sleeve gastrectomy (SG), percentage weight loss (%WL)

317

**Subjective changes in taste towards sweet and salty foods**

319 Patients were asked if they had experienced a change in their taste towards sweet and salty  
320 foods. They were asked separately if they had experienced an increase or decrease in the

321 taste of sweet or salty tasting foods (Appendix: Questions 9, 10, 12 and 13). 87.8% of RYGB  
 322 patients responded that their taste for sweet foods had either increased or decreased  
 323 compared to 65.2% of SG patients (p=0.001). Changes in taste towards salty tasting foods  
 324 were also significantly more common following RYGB (% of patients reporting change in taste  
 325 towards salty tasting foods: RYGB =56.1, SG= 40.6%, p=0.020, Figure 2).

326  
 327 Following RYGB, there were no gender differences in subjective taste change towards sweet  
 328 foods (females 89% and males 84%, p=0.685). However, post-SG a change in taste towards  
 329 sweet foods, either an increase or decrease, was significantly more common in females  
 330 compared to males (females 70% and males 45%, p=0.009).

331  
 332 **(Figure 2 here)**

333  
 334 **Influence of T2D**

335 T2D was present in 92 patients (53 RYGB and 39 SG). No differences were seen in the  
 336 prevalence of subjective change in appetite, taste, smell or food aversion between individuals  
 337 with and without T2D post-RYGB (appetite change: no T2D=93.3%, T2D=88.7%; taste change:  
 338 no T2D=73.3%, T2D=56.6%, smell change: no T2D=40%, T2D=41.5%; food aversions: no  
 339 T2D=62.2%, T2D=62.3%; all p > 0.05) or post-SG (appetite change: no T2D=89.7%, T2D=94.9%;  
 340 taste change: no T2D=62.1%, T2D=51.3%, smell change: no T2D=26.7%, T2D=30.8%; food  
 341 aversions: no T2D=62.1%, T2D=51.3%; all p > 0.05). However, within the SG group, taste  
 342 changes were significantly less common in males with T2D compared to females with T2D (%  
 343 of patients reporting change in taste: T2D females=61.3%, T2D males=12.5%, p=0.020.  
 344 Furthermore, taste changes were less common in males with T2D following SG than post-  
 345 RYGB (change in taste prevalence: T2D men RYGB=69.2%, T2D men SG=12.5%, p=0.024).  
 346 Moreover, %WL was significantly lower in male T2D patients following SG compared to post-  
 347 RYGB (%WL: SG=14.6 ± 2.1, RYGB=27.5 ± 2.7, p=0.003).

348 **Table 7: Frequency of reported appetite, taste, smell changes and food aversions by gender**  
 349 **in patients with T2D following RYGB and SG.**

350

	RYGB			SG		
	Females	Males	p value	Females	Males	p value
<b>Appetite</b>	87.5%	92.3%	0.370	96.8%	87.5%	0.372
<b>Taste</b>	52.5%	69.2%	1.0	61.3%	12.5%	0.019
<b>Smell</b>	42.5%	38.5%	1.0	35.5%	12.5%	0.393
<b>Food aversions</b>	65.0%	53.4%	0.522	58.1%	25%	0.123

351 Roux-en-Y gastric bypass (RYGB), sleeve gastrectomy (SG)

352  
 353 **Predictors of weight loss**

354 Linear regression analyses were performed in order to correct for characteristics within  
 355 groups and identify predictors of %WL. The models for RYGB and SG patients are illustrated  
 356 in Table 8. The basic model (M1) adjusted for gender and days since surgery. Subjective taste  
 357 changes following surgery were associated with greater %WL post-RYGB, even after adjusting  
 358 for gender, age, duration since surgery and the presence of T2D. In contrast, post-SG,

359 subjective taste change was not related to %WL. However, in patients post-SG subjective  
 360 change in appetite strongly associated with greater %WL after adjusting for gender, age,  
 361 duration since surgery and T2D, which is not observed in RYGB.

362

363 **Table 8: Linear regression model testing %WL by appetite, taste, smell and aversions**

364

RYGB				
Model	Appetite	Taste	Smell	Aversions
	Coefficient (95% CI)	Coefficient (95% CI)	Coefficient (95% CI)	Coefficient (95% CI)
Basic Model (M1)	2.4 (-3.5; 8.4)	4.0 (0.5; 7.5)*	1.3 (-2.2; 4.9)	1.1 (-2.6; 4.8)
M1 + age	2.5 (-3.4; 8.4)	3.6 (0.7; 7.1)*	1.1 (-2.4; 4.5)	0.9 (-2.7; 4.5)
M1 + T2D	2.3 (-3.7; 8.3)	3.8 (0.2; 7.4)*	1.4 (-2.1; 4.9)	1.3 (-2.4; 4.7)
SG				
Model	Appetite	Taste	Smell	Aversions
Basic Model (M1)	8.1 (2.5; 13.7)**	0.9 (-2.5; 4.3)	0.1 (-3.5; 3.7)	3.1 (-0.1; 6.4)
M1 + age	8.7 (3.2; 14.1)**	0.6 (-2.7; 3.9)	0.4 (-3.2; 3.9)	2.8 (-0.5; 5.9)
M1 + T2D	8.9 (3.4; 14.5)**	0.5 (-2.8; 3.9)	0.3 (-3.2; 3.8)	2.8 (-0.4; 6.0)

\* p<0.05, \*\* p<0.01; Basic model adjusts for gender and days since surgery

365 Roux-en-Y gastric bypass (RYGB), sleeve gastrectomy (SG), Type 2 diabetes (T2D), M1 (Model 1)

366

367

## 368 Discussion

369

370 In this cross-sectional study we examined the prevalence of food aversions and subjective  
 371 changes in appetite, taste and smell in a cohort of patients 6 months to 5 years following  
 372 primary RYGB or SG. Our cohort included the largest SG group examined to date with the  
 373 longest duration post-surgery. Our data show that whilst subjective changes in appetite, taste,  
 374 smell and food aversions are common following both RYGB and SG they differ in their impact  
 375 on subjective changes in smell and taste, durability of changes, influence of gender and  
 376 relationship with %WL.

377

378 In the RYGB group the prevalence of changes in appetite, taste, smell and food aversions was  
 379 similar across all periods examined, with 86% of patients reporting appetite changes and 64%  
 380 taste changes 3-5 years following RYGB. We observed no influence of gender or T2D status  
 381 upon changes in appetite, taste or smell. %WL was similar at 1-2, 2-3 and 3-5 years post-  
 382 surgery. Linear regression analysis showed that subjective taste changes associated with and  
 383 predicted %WL.

384

385 After SG, smell changes were less common than after RYGB, as were also changes in taste  
 386 towards sweet and salty tasting foods. The prevalence of changes in appetite significantly  
 387 decreased with time post-surgery; the prevalence of changes in taste, smell and food aversion  
 388 showed a trend to decrease with time post-surgery. Importantly, following SG the prevalence  
 389 of changes in taste and smell was lower in males compared to females particularly in male  
 390 patients with T2D. While %WL was comparable between RYGB and SG during the first post-

391 operative 2-year period, between 2-5 years %WL was significantly greater in the RYGB group  
392 at 2-5 years post-surgery compared to the SG group. Subjective appetite changes associated  
393 with and predicted %WL in the SG group.

394

395 Gender differences in taste sensitivity have been reported in obesity (Doty & Cameron, 2009;  
396 Hwang et al., 2016). Similarly, T2D, which is known to impact upon gut hormone profiles, has  
397 also been suggested to affect taste sensitivity, particularly toward sweet tasting foods  
398 (Bustos-Saldana et al., 2009). However, until now, no data existed with regards to effects of  
399 gender or T2D on appetite, taste or smell in patients following bariatric surgery. While no  
400 significant gender differences were seen within our RYGB group, there was a marked gender  
401 effect following SG. In particular, following SG the prevalence of taste changes was  
402 significantly lower in males with T2D compared to females with T2D. Moreover, the  
403 prevalence of taste changes and %WL were significantly lower in males with T2D following SG  
404 compared to RYGB. Our findings highlight the need for further research into the underlying  
405 physiology of these gender differences.

406

407 The role of weight loss in mediating subjective changes in taste and smell remains to be  
408 clarified. Interestingly, reduced subjective taste and smell following GI surgery were first  
409 reported in normal weight patients as a transient complication following gastrectomy and  
410 oesophagectomy for GI malignancies (Harris & Griffin, 2003), suggesting that these changes  
411 are not restricted to people with obesity. In our cohort, the prevalence of smell changes was  
412 greater in the RYGB group compared to the SG despite comparable %WL during the first two  
413 post-operative years, suggesting weight independent processes may be involved. The  
414 perception of flavour is thought to be predominantly mediated through smell (Yeomans,  
415 2006), thus our findings of much higher prevalence of taste changes compared to smell  
416 changes following RYGB and SG is somewhat surprising. However, recent studies have  
417 identified gut hormone receptors on taste buds, gut hormones within saliva and postulated a  
418 role for saliva gut hormones in taste modulation (Acosta et al., 2011; Y. K. Shin et al., 2008).  
419 Circulating levels of gut hormones, in particular ghrelin, GLP-1 and PYY, change post-  
420 operatively (Yousseif et al., 2014). Thus, these changes offer not only a plausible biological  
421 explanation for appetite changes post-surgery but also for the higher prevalence of taste  
422 compared to smell changes. In addition, the differential gut hormone pattern observed  
423 following RYGB and SG may underlie the procedural differences in appetite, taste and smell  
424 post-surgery (Cummings, 2015; Fabian et al., 2015). Longitudinal studies examining hormone  
425 levels in blood and saliva coupled with subjective and objective assessments of appetite, taste  
426 and smell in large cohorts of males and females undergoing RYGB and SG are now needed.

427

428 Our study does have limitations. The study was cross-sectional, which resulted in a large  
429 variation in time post-surgery. Reported changes in appetite, taste and smell were subjective  
430 and no control group was included. In addition, since no data was collected pre-operatively,  
431 our data could be subject to recall bias, which is likely to increase with longer duration from  
432 surgery. Furthermore, the number of male participants in our study cohort was small.  
433 However, this reflects the higher number of females undergoing bariatric surgery in the UK  
434 compared to males (Wellbourn R, 2014). Our findings of appetite, taste, smell changes and  
435 food aversions include the largest cohort of SG patients to date. Our SG sample size was larger  
436 compared to the RYGB group, while the post-RYGB patients had a significantly longer duration  
437 since surgery, which reflects the increasing popularity of SG over recent years and the trend

438 to perform a higher number SG (Angrisani et al., 2015). The higher %WL in the RYGB group is  
439 in concordance with previous findings from studies that have compared post-operative  
440 outcomes between RYGB and SG (Schauer et al., 2014; Szczepaniak et al., 2015). Similarly,  
441 weight loss post-RYGB was also higher compared to SG in the study conducted by Zerrweck  
442 *et al* (Zerrweck et al., 2015). The higher percentage of T2D in the RYGB population reflects  
443 that RYGB remains the preferred procedure in T2D, given the better T2D outcomes post-RYGB  
444 in the literature (Schauer et al., 2014). Graham *et al.* and Zerrweck *et al.* reported associations  
445 with food aversions and weight loss (Graham et al., 2014; Zerrweck et al., 2015). This effect  
446 was observed only following SG in our cohort. However, the wording of the question relating  
447 to food aversions did not allow for a clear distinction between true food aversions, and post-  
448 ingestive phenomena such as dumping syndrome. This ambiguity needs to be borne in mind  
449 when interpreting these results, as food tolerance and GI quality of life may play a role as  
450 aversive drivers of eating behaviour (Overs, Freeman, Zarshenas, Walton, & Jorgensen, 2012).  
451 Furthermore, feedback from our participants also highlighted the ambiguity of some of the  
452 questions, particularly relating to “increase” or “loss” of taste.

453  
454 Nevertheless, compared to previous studies, our study design has several strengths and  
455 provides novel findings. We obtained consent in person and completed data collection in one  
456 visit. Our exclusion criteria allowed for elimination of patients with additional factors that  
457 could impact on %WL, including low B12 and zinc. Furthermore, in contrast to previous  
458 studies we have used %WL as our outcome for weight loss, in order to avoid for confounding  
459 outcomes by pre-operative BMI. Our analysis identified procedure dependent differences in  
460 appetite, taste, smell changes and their impact upon %WL. In addition, we investigated the  
461 effect of gender and found marked gender differences within the SG group that are  
462 exacerbated by the presence of T2D. Finally, we performed linear regression analyses and  
463 thereby, for the first time, identified appetite and taste changes as predictors of %WL for SG  
464 and RYGB respectively.

465  
466 In conclusion, whilst subjective changes in appetite, taste, smell and food aversions are  
467 common following both RYGB and SG, marked differences between them exist in relation to  
468 the prevalence of changes in smell, taste, durability of changes, influence of gender and %WL.  
469 The presence of gender differences highlights the need to study comparable numbers of  
470 males and females. Furthermore, correlating gut hormone profiles with appetite and taste  
471 changes, will not only aid to further our understanding of the biological mediators for weight  
472 loss post-bariatric surgery allowing for personalised procedure allocation but may also lead  
473 to novel therapies, such as taste modulation approaches.

474  
475  
476  
477  
478

## 479 **Acknowledgements**

480  
481 We would like to thank all members, past and present, of the University College London  
482 Hospitals Bariatric Group and all our patients. This work was supported by the Rosetrees Trust,  
483 Stoneygate Trust and the National Institute for Health Research.

484 **Figure Legends**

485

486 **Figure 1: Frequency of reported appetite, taste, smell changes and food aversions following**  
487 **Roux-en-Y gastric bypass (RYGB) and sleeve gastrectomy (SG)**

488

489 A: Appetite changes (Q1) B: Taste changes (Q2) C: Smell changes (Q3) D: Food aversions (Q4)

490

491 **Figure 2: Change in taste toward sweet and salty foods**

492

493 Reported frequency of increased and decreased taste toward sweet (A) and salty (B) tasting  
494 foods in Roux-en-Y gastric bypass (RYGB) and sleeve gastrectomy (SG) respectively.

495

496

497 **References**

498

499 Abu-Jaish, W., & Rosenthal, R. J. (2010). Sleeve gastrectomy: a new surgical approach for  
500 morbid obesity. *Expert Rev Gastroenterol Hepatol*, 4(1), 101-119.

501 doi:10.1586/egh.09.68

502 Acosta, A., Hurtado, M. D., Gorbatyuk, O., La Sala, M., Duncan, D., Aslanidi, G., . . .

503 Zolotukhin, S. (2011). Salivary PYY: a putative bypass to satiety. *PLoS One*, 6(10),

504 e26137. doi:10.1371/journal.pone.0026137

505 Altun, H., Hanci, D., Altun, H., Batman, B., Serin, R. K., Karip, A. B., & Akyuz, U. (2016).

506 Improved Gustatory Sensitivity in Morbidly Obese Patients After Laparoscopic Sleeve

507 Gastrectomy. *Ann Otol Rhinol Laryngol*. doi:10.1177/0003489416629162

508 Angrisani, L., Santonicola, A., Iovino, P., Formisano, G., Buchwald, H., & Scopinaro, N. (2015).

509 Bariatric Surgery Worldwide 2013. *Obes Surg*, 25(10), 1822-1832.

510 doi:10.1007/s11695-015-1657-z

511 Atalayer, D., Pantazatos, S. P., Gibson, C. D., McOuatt, H., Puma, L., Astbury, N. M., &

512 Geliebter, A. (2014). Sexually dimorphic functional connectivity in response to high

513 vs. low energy-dense food cues in obese humans: an fMRI study. *Neuroimage*, 100,

514 405-413. doi:10.1016/j.neuroimage.2014.05.054

515 Berthoud, H. R. (2011). Metabolic and hedonic drives in the neural control of appetite: who

516 is the boss? *Curr Opin Neurobiol*, 21(6), 888-896. doi:10.1016/j.conb.2011.09.004

517 Bustos-Saldana, R., Alfaro-Rodriguez, M., Solis-Ruiz Mde, L., Trujillo-Hernandez, B., Pacheco-

518 Carrasco, M., Vazquez-Jimenez, C., & Celis-de la Rosa Ade, J. (2009). [Taste sensitivity

519 diminution in hyperglycemic type 2 diabetics patients]. *Rev Med Inst Mex Seguro*

520 *Soc*, 47(5), 483-488.

521 Coluzzi, I., Raparelli, L., Guarnacci, L., Paone, E., Del Genio, G., le Roux, C. W., & Silecchia, G.

522 (2016). Food Intake and Changes in Eating Behavior After Laparoscopic Sleeve

523 Gastrectomy. *Obes Surg*. doi:10.1007/s11695-015-2043-6

524 Courcoulas, A. P., Christian, N. J., O'Rourke, R. W., Dakin, G., Patchen Dellinger, E., Flum, D.

525 R., . . . Belle, S. H. (2015). Preoperative factors and 3-year weight change in the

526 Longitudinal Assessment of Bariatric Surgery (LABS) consortium. *Surgery for Obesity*

527 *and Related Diseases*, 11(5), 1109-1118. doi:10.1016/j.soard.2015.01.011

528 Cummings, D. E. (2015). Taste and the regulation of food intake: it's not just about flavor.

529 *Am J Clin Nutr*, 102(4), 717-718. doi:10.3945/ajcn.115.120667

530 Deglaire, A., Mejean, C., Castetbon, K., Kesse-Guyot, E., Hercberg, S., & Schlich, P. (2015).

531 Associations between weight status and liking scores for sweet, salt and fat

532 according to the gender in adults (The Nutrinet-Sante study). *Eur J Clin Nutr*, 69(1),

533 40-46. doi:10.1038/ejcn.2014.139

534 Dirksen, C., Damgaard, M., Bojsen-Moller, K. N., Jorgensen, N. B., Kielgast, U., Jacobsen, S.

535 H., . . . Madsen, J. L. (2013). Fast pouch emptying, delayed small intestinal transit,

536 and exaggerated gut hormone responses after Roux-en-Y gastric bypass.

537 *Neurogastroenterol Motil*, 25(4), 346-e255. doi:10.1111/nmo.12087

538 Dirksen, C., Jorgensen, N. B., Bojsen-Moller, K. N., Kielgast, U., Jacobsen, S. H., Clausen, T.

539 R., . . . Hansen, D. L. (2013). Gut hormones, early dumping and resting energy

540 expenditure in patients with good and poor weight loss response after Roux-en-Y

541 gastric bypass. *Int J Obes (Lond)*, 37(11), 1452-1459. doi:10.1038/ijo.2013.15

542 Doty, R. L., & Cameron, E. L. (2009). Sex differences and reproductive hormone influences  
543 on human odor perception. *Physiol Behav*, *97*(2), 213-228.  
544 doi:10.1016/j.physbeh.2009.02.032

545 Fabian, T. K., Beck, A., Fejerdy, P., Hermann, P., & Fabian, G. (2015). Molecular mechanisms  
546 of taste recognition: considerations about the role of saliva. *Int J Mol Sci*, *16*(3),  
547 5945-5974. doi:10.3390/ijms16035945

548 Graham, L., Murty, G., & Bowrey, D. J. (2014). Taste, smell and appetite change after Roux-  
549 en-Y gastric bypass surgery. *Obes Surg*, *24*(9), 1463-1468. doi:10.1007/s11695-014-  
550 1221-2

551 Harris, A. M., & Griffin, S. M. (2003). Postoperative taste and smell deficit after upper  
552 gastrointestinal cancer surgery--an unreported complication. *J Surg Oncol*, *82*(3),  
553 147-150; discussion 150-142. doi:10.1002/jso.10199

554 Holinski, F., Menenakos, C., Haber, G., Olze, H., & Ordemann, J. (2015). Olfactory and  
555 Gustatory Function After Bariatric Surgery. *Obes Surg*. doi:10.1007/s11695-015-  
556 1683-x

557 Hwang, S. H., Kang, J. M., Seo, J. H., Han, K. D., & Joo, Y. H. (2016). Gender Difference in the  
558 Epidemiological Association between Metabolic Syndrome and Olfactory  
559 Dysfunction: The Korea National Health and Nutrition Examination Survey. *PLoS One*,  
560 *11*(2), e0148813. doi:10.1371/journal.pone.0148813

561 Jurowich, C. F., Seyfried, F., Miras, A. D., Bueter, M., Deckelmann, J., Fassnacht, M., Germer,  
562 CT, Thalheimer, A. (2014). Does bariatric surgery change olfactory perception?  
563 Results of the early postoperative course. *Int J Colorectal Dis*, *29*(2), 253-260.  
564 doi:10.1007/s00384-013-1795-5

565 le Roux, C. W., Welbourn, R., Werling, M., Osborne, A., Kokkinos, A., Laurenus, A., . . .  
566 Olbers, T. (2007). Gut hormones as mediators of appetite and weight loss after Roux-  
567 en-Y gastric bypass. *Ann Surg*, *246*(5), 780-785. doi:10.1097/SLA.0b013e3180caa3e3

568 Loch, D., Breer, H., & Strotmann, J. (2015). Endocrine Modulation of Olfactory  
569 Responsiveness: Effects of the Orexigenic Hormone Ghrelin. *Chem Senses*, *40*(7),  
570 469-479. doi:10.1093/chemse/bjv028

571 Manning, S., Pucci, A., & Batterham, R. L. (2015). Roux-en-Y gastric bypass: effects on  
572 feeding behavior and underlying mechanisms. *J Clin Invest*, *125*(3), 939-948.  
573 doi:10.1172/jci76305

574 Miras, A. D., & le Roux, C. W. (2010). Bariatric surgery and taste: novel mechanisms of  
575 weight loss. *Curr Opin Gastroenterol*, *26*(2), 140-145.  
576 doi:10.1097/MOG.0b013e328333e94a

577 National Institute for Health and Care Excellence. (2014). Obesity: identification, assessment  
578 and management. *NICE guidelines [CG189]*. Retrieved from  
579 <https://www.nice.org.uk/guidance/cg189>

580 Ochner, C. N., Kwok, Y., Conceicao, E., Pantazatos, S. P., Puma, L. M., Carnell, S., Taxeira J,  
581 Hirsh J, Geliebter, A. (2011). Selective reduction in neural responses to high calorie  
582 foods following gastric bypass surgery. *Ann Surg*, *253*(3), 502-507.  
583 doi:10.1097/SLA.0b013e318203a289

584 Ochner, C. N., Stice, E., Hutchins, E., Afifi, L., Geliebter, A., Hirsch, J., & Teixeira, J. (2012).  
585 Relation between changes in neural responsivity and reductions in desire to eat high-  
586 calorie foods following gastric bypass surgery. *Neuroscience*, *209*, 128-135.  
587 doi:10.1016/j.neuroscience.2012.02.030



588 Olbers, T., Lonroth, H., Fagevik-Olsen, M., & Lundell, L. (2003). Laparoscopic gastric bypass:  
589 development of technique, respiratory function, and long-term outcome. *Obes Surg*,  
590 13(3), 364-370. doi:10.1381/096089203765887679

591 Overs, S. E., Freeman, R. A., Zarshenas, N., Walton, K. L., & Jorgensen, J. O. (2012). Food  
592 tolerance and gastrointestinal quality of life following three bariatric procedures:  
593 adjustable gastric banding, Roux-en-Y gastric bypass, and sleeve gastrectomy. *Obes*  
594 *Surg*, 22(4), 536-543. doi:10.1007/s11695-011-0573-0

595 Patel, Z. M., DelGaudio, J. M., & Wise, S. K. (2015). Higher Body Mass Index Is Associated  
596 with Subjective Olfactory Dysfunction. *Behav Neurol*, 2015, 675635.  
597 doi:10.1155/2015/675635

598 Pepino, M. Y., Bradley, D., Eagon, J. C., Sullivan, S., Abumrad, N. A., & Klein, S. (2014).  
599 Changes in taste perception and eating behavior after bariatric surgery-induced  
600 weight loss in women. *Obesity (Silver Spring)*, 22(5), E13-20. doi:10.1002/oby.20649

601 Rissanen, A., Hakala, P., Lissner, L., Mattlar, C. E., Koskenvuo, M., & Ronnema, T. (2002).  
602 Acquired preference especially for dietary fat and obesity: a study of weight-  
603 discordant monozygotic twin pairs. *Int J Obes Relat Metab Disord*, 26(7), 973-977.  
604 doi:10.1038/sj.ijo.0802014

605 Schauer, P. R., Bhatt, D. L., Kirwan, J. P., Wolski, K., Brethauer, S. A., Navaneethan, S. D., . . .  
606 Kashyap, S. R. (2014). Bariatric surgery versus intensive medical therapy for diabetes-  
607 -3-year outcomes. *N Engl J Med*, 370(21), 2002-2013. doi:10.1056/NEJMoa1401329

608 Scholtz, S., Miras, A. D., Chhina, N., Prechtel, C. G., Sleeth, M. L., Daud, N. M., . . . Goldstone,  
609 A. P. (2014). Obese patients after gastric bypass surgery have lower brain-hedonic  
610 responses to food than after gastric banding. *Gut*, 63(6), 891-902.  
611 doi:10.1136/gutjnl-2013-305008

612 Scott, W. R., & Batterham, R. L. (2011). Roux-en-Y gastric bypass and laparoscopic sleeve  
613 gastrectomy: understanding weight loss and improvements in type 2 diabetes after  
614 bariatric surgery. *Am J Physiol Regul Integr Comp Physiol*, 301(1), R15-27.  
615 doi:10.1152/ajpregu.00038.2011

616 Sczepaniak, J. P., Owens, M. L., Shukla, H., Perlegos, J., & Garner, W. (2015). Comparability  
617 of weight loss reporting after gastric bypass and sleeve gastrectomy using BOLD data  
618 2008-2011. *Obes Surg*, 25(5), 788-795. doi:10.1007/s11695-014-1496-3

619 Shin, A. C., Townsend, R. L., Patterson, L. M., & Berthoud, H. R. (2011). "Liking" and  
620 "wanting" of sweet and oily food stimuli as affected by high-fat diet-induced obesity,  
621 weight loss, leptin, and genetic predisposition. *Am J Physiol Regul Integr Comp*  
622 *Physiol*, 301(5), R1267-1280. doi:10.1152/ajpregu.00314.2011

623 Shin, Y. K., Martin, B., Golden, E., Dotson, C. D., Maudsley, S., Kim, W., . . . Munger, S. D.  
624 (2008). Modulation of taste sensitivity by GLP-1 signaling. *J Neurochem*, 106(1), 455-  
625 463. doi:10.1111/j.1471-4159.2008.05397.x

626 Sjostrom, L. (2013). Review of the key results from the Swedish Obese Subjects (SOS) trial - a  
627 prospective controlled intervention study of bariatric surgery. *J Intern Med*, 273(3),  
628 219-234. doi:10.1111/joim.12012

629 Tichansky, D. S., Boughter, J. D., Jr., & Madan, A. K. (2006). Taste change after laparoscopic  
630 Roux-en-Y gastric bypass and laparoscopic adjustable gastric banding. *Surg Obes*  
631 *Relat Dis*, 2(4), 440-444. doi:10.1016/j.soard.2006.02.014

632 Wellbourn R, S. P., Finlay I, Sareela A, Somers S, Mahawar K, et al. (2014). *The United*  
633 *Kingdom National Bariatric Surgery Registry: second registry report*. Retrieved from  
634 London:

635 Westenhoefer, J., & Pudel, V. (1993). Pleasure from food: importance for food choice and  
636 consequences of deliberate restriction. *Appetite*, *20*(3), 246-249.  
637 doi:10.1006/appe.1993.1029

638 Yeomans, M. R. (2006). Olfactory influences on appetite and satiety in humans. *Physiol*  
639 *Behav*, *89*(1), 10-14. doi:10.1016/j.physbeh.2006.04.010

640 Yousseif, A., Emmanuel, J., Karra, E., Millet, Q., Elkalaawy, M., Jenkinson, A. D., . . .  
641 Batterham, R. L. (2014). Differential effects of laparoscopic sleeve gastrectomy and  
642 laparoscopic gastric bypass on appetite, circulating acyl-ghrelin, peptide YY3-36 and  
643 active GLP-1 levels in non-diabetic humans. *Obes Surg*, *24*(2), 241-252.  
644 doi:10.1007/s11695-013-1066-0

645 Zerrweck, C., Zurita, L., Alvarez, G., Maydon, H. G., Sepulveda, E. M., Campos, F., Caviedes  
646 A, Guilbert, L. (2015). Taste and Olfactory Changes Following Laparoscopic Gastric  
647 Bypass and Sleeve Gastrectomy. *Obes Surg*. doi:10.1007/s11695-015-1944-8  
648  
649

650 **Appendix**

651

652 **Taste and smell questionnaire**

653

654 **Please circle the most appropriate answer or circle the relevant number on the scale (1-**  
655 **10)**

656

657 Q1. Have you noticed any change in your appetite since your weight loss surgery?

658

659 **YES/ NO**

660

661 Q2. Have you noticed any change in the taste of food or drink since your weight loss  
662 surgery?

663

664 **YES/ NO**

665

666 Q3. Have you noticed any change in your sense of smell since your weight loss surgery?

667

668 **YES/ NO**

669

670 Q4. Have you experienced an overall loss of taste since your weight loss surgery?

671

672 **YES/ NO**

673

674

675 **If you answered YES to the above questions please continue. If you answered NO to all of**  
676 **the above questions, do not proceed any further. Please return the questionnaire.**

677

678 Q5. If you have had a loss of taste, is that loss:

679

680 **No loss**

**Partial**

**Complete**

681

682 **1 2 3 4 5 6 7 8 9 10**

683

684 Q6. Are there any foods that are repulsive or intolerable to you since your weight loss  
685 surgery?

686

687 **YES/ NO**

688

689 **If you answered yes then please state what:**

690

691

692 Q7. Are there any foods that taste different to you since your weight loss surgery?

693

694 **YES/ NO**

695 **If yes, please state what food:**

696

697  
698  
699  
700  
701  
702  
703  
704  
705  
706  
707  
708  
709  
710  
711  
712  
713  
714  
715  
716  
717  
718  
719  
720  
721  
722  
723  
724  
725  
726  
727  
728  
729  
730  
731  
732  
733  
734  
735  
736  
737  
738  
739  
740  
741  
742  
743

Q8. Has your overall taste increased in intensity since your weight loss surgery?

**YES/ NO**

Q9. Have you experienced an increase in taste for sweet foods?

**YES/ NO**

Q10. Have you experienced a decrease in taste for sweet foods?

**YES/ NO**

Q11. If you have had a loss of sweet taste, is that loss:

<b>No loss</b>				<b>Partial</b>				<b>Complete</b>	
<b>1</b>	<b>2</b>	<b>3</b>	<b>4</b>	<b>5</b>	<b>6</b>	<b>7</b>	<b>8</b>	<b>9</b>	<b>10</b>

Q12. Have you experienced an increase in taste for salty foods?

**YES/ NO**

Q13. Have you experienced a decrease in taste for salty foods?

**YES/ NO**

Q14. If you have had a loss of salty taste, is that loss:

<b>No loss</b>				<b>Partial</b>				<b>Complete</b>	
<b>1</b>	<b>2</b>	<b>3</b>	<b>4</b>	<b>5</b>	<b>6</b>	<b>7</b>	<b>8</b>	<b>9</b>	<b>10</b>

Q15. Have you experienced an increase in taste for sour foods?

**YES/ NO**

744 Q16. Have you experienced a decrease in taste for sour foods?

745

746 **YES/ NO**

747

748

749 Q17. If you have had a loss of sour taste, is that loss:

750

751 **No loss**

**Partial**

**Complete**

752

753 **1 2 3 4 5 6 7 8 9 10**

754

755

756 Q18. Has this change in taste affected how much you eat?

757

758 **YES/ NO**

759

760

761 Q19. In your opinion, has your change in taste affected your weight loss?

762

763 **YES/ NO**

764

765 Q20. Overall, do you feel that your taste has increased or decreased in intensity  
766 since your weight loss surgery?

767

768 **Increased/ Decreased**

769

770

771 Q21. In your opinion, does loss of taste lead to better weight loss?

772

773 **YES/ NO**

774

775

776 Q22. Is your postoperative change in taste greater or less than what you expected  
777 preoperatively?

778

779 **Greater/ Less**

780

781

782 Q23. How important is taste to the enjoyment of food?

783

784 **Important/ Not important**

785

786

787 Q24. Have you experienced an overall loss in your sense of smell since your weight loss  
788 surgery?

789

790 **YES/ NO**

791  
792  
793  
794  
795  
796  
797  
798  
799  
800  
801  
802  
803  
804  
805  
806  
807  
808  
809  
810  
811  
812  
813  
814  
815  
816  
817  
818  
819  
820  
821  
822  
823  
824  
825  
826  
827  
828  
829  
830  
831  
832  
833  
834  
835  
836  
837

Q25. If you have had a loss of smell, is that loss:

<b>No loss</b>		<b>Partial</b>						<b>Complete</b>	
<b>1</b>	<b>2</b>	<b>3</b>	<b>4</b>	<b>5</b>	<b>6</b>	<b>7</b>	<b>8</b>	<b>9</b>	<b>10</b>

Q26. Are there any foods that smell differently to you since your weight loss surgery?

**YES/ NO**

**If yes, please state what food?**

Q27. In your opinion, has your change in smell affected your weight loss?

**YES/ NO**

Q28. Overall, do you feel that your smell has increased or decreased in intensity since your weight loss surgery?

**Increased/ Decreased**

Q29. Is your postoperative change in smell greater or less than what you expected preoperatively?

**Greater/ Less**

Q30. How important is smell to the enjoyment of food?

**Important/ Not important**

Q31. Do you eat less food because it does not taste or smell good?

**YES/ NO**

Q32. Do you eat less food because you are simply not hungry?

**YES/ NO**

838

839 Q33. Is your postoperative change in appetite greater or less than what you expected  
840 preoperatively?

841

842 **Greater/ Less**

843

844

845