

ARTICLE

Received 22 Dec 2015 | Accepted 23 May 2016 | Published 7 Jul 2016

DOI: [10.1038/ncomms12051](https://doi.org/10.1038/ncomms12051)

OPEN

Microcanonical and resource-theoretic derivations of the thermal state of a quantum system with noncommuting charges

Nicole Yunger Halpern¹, Philippe Faist², Jonathan Oppenheim³ & Andreas Winter^{4,5}

The grand canonical ensemble lies at the core of quantum and classical statistical mechanics. A small system thermalizes to this ensemble while exchanging heat and particles with a bath. A quantum system may exchange quantities represented by operators that fail to commute. Whether such a system thermalizes and what form the thermal state has are questions about truly quantum thermodynamics. Here we investigate this thermal state from three perspectives. First, we introduce an approximate microcanonical ensemble. If this ensemble characterizes the system-and-bath composite, tracing out the bath yields the system's thermal state. This state is expected to be the equilibrium point, we argue, of typical dynamics. Finally, we define a resource-theory model for thermodynamic exchanges of noncommuting observables. Complete passivity—the inability to extract work from equilibrium states—implies the thermal state's form, too. Our work opens new avenues into equilibrium in the presence of quantum noncommutation.

¹Institute for Quantum Information and Matter, Division of Physics, Mathematics, and Astronomy, California Institute of Technology, Pasadena, California 91125, USA. ²Institute for Theoretical Physics, Department of Physics, ETH Zürich, 8093 Zürich, Switzerland. ³Department of Physics and Astronomy, University College London, Gower Street, London WC1E 6BT, UK. ⁴Departament de Física, Grup d'Informació Quàntica, Universitat Autònoma de Barcelona, ES-08193 Barcelona, Spain. ⁵ICREA, Pg. Lluís Companys 23, 08010 Barcelona, Spain. Correspondence and requests for materials should be addressed to N.Y.H. (email: nicoleyh@caltech.edu) or to P.F. (email: pfaist@phys.ethz.ch) or to A.W. (email: andreas.winter@uab.cat).

Recently reignited interest in quantum thermodynamics has prompted information-theoretic approaches to fundamental questions^{1–4}. The role of entanglement, for example, has been clarified with canonical typicality^{5–8}. Equilibrium-like behaviours have been predicted^{9–12} and experimentally observed in integrable quantum gases^{13,14}.

Thermodynamic resource theories offer a powerful tool for analysing fundamental properties of the thermodynamics of quantum systems. Heat exchanges with a bath are modelled with ‘free states’ and ‘free operations’^{15–18}. These resource theories have been extended to model exchanges of additional physical quantities, such as particles and angular momentum^{18–22}.

A central concept in thermodynamics and statistical mechanics is the thermal state. The thermal state has several important properties. First, typical dynamics evolve the system towards the thermal state. The thermal state is the equilibrium state. Second, consider casting statistical mechanics as an inference problem. The thermal state is the state that maximizes the entropy under constraints on physical quantities^{23,24}. Third, consider the system as interacting with a large bath. The system-and-bath composite occupies a microcanonical state. Physical observables of the composite, such as the total energy and total particle number, have sharply defined values. The system’s reduced state is the thermal state. Finally, in a resource theory, the thermal state is the only completely passive state. No work can be extracted from any number of copies of the thermal state^{25,26}.

If a small system exchanges heat and particles with a large environment, the system’s thermal state is a grand canonical ensemble: $e^{-\beta(H-\mu N)}/Z$. The system’s Hamiltonian and particle number are represented by H and N . β and μ denote the environment’s inverse temperature and chemical potential. The partition function Z normalizes the state. The system-and-bath dynamics conserves the total energy and total particle number. More generally, subsystems exchange conserved quantities, or ‘charges,’ Q_j , $j=1, 2, \dots, c$. To these charges correspond generalized chemical potentials μ_j . The μ_j ’s characterize the bath.

We address the following question. Suppose that the charges fail to commute with each other: $[Q_j, Q_k] \neq 0$. What form does the thermal state have? We call this state ‘the non-Abelian thermal state’ (NATS). Jaynes applied the principle of maximum entropy to this question²⁴. He associated fixed values v_j with the charges’ expectation values. He calculated the state that, on satisfying these constraints, maximizes an entropy. This thermal state has a generalized Gibbs form:

$$\gamma_v = \frac{1}{Z} e^{-\sum_{j=0}^c \mu_j Q_j} \quad (1)$$

wherein the v_j ’s determine the μ_j ’s.

Our contribution is a mathematical, physically justified derivation of the thermal state’s form for systems whose dynamics conserve noncommuting observables. We recover the state (1) via several approaches, demonstrating its physical importance. We address puzzles raised in refs 21,27 about how to formulate a resource theory in which thermodynamic charges fail to commute. Closely related, independent work was performed by Guryanova *et al.*²⁸ We focus primarily on the nature of passive states. Guryanova *et al.*, meanwhile, focus more on the resource theory for multiple charges and on trade-offs amongst types of charge extractions.

In this paper, we derive the NATS’s form from a microcanonical argument. A simultaneous eigenspace of all the noncommuting physical charges might not exist. Hence, we introduce the notion of an approximate microcanonical subspace. This subspace consists of the states in which the charges have sharply defined values. We derive conditions under which this subspace exists. We show that a small subsystem’s reduced state

lies, on average, close to γ_v . Second, we invoke canonical typicality^{7,8}. If the system-and-bath composite occupies a random state in the approximate microcanonical subspace, we argue, a small subsystem’s state likely lies close to the NATS. Typical dynamics are therefore expected to evolve a well-behaved system’s state towards the NATS. Third, we define a resource theory for thermodynamic exchanges of noncommuting conserved charges. We extend existing resource theories to model the exchange of noncommuting quantities. We show that the NATS is the only possible free state that renders the theory nontrivial: Work cannot be extracted from any number of copies of γ_v . We show also that the NATS is the only state preserved by free operations. From this preservation, we derive ‘second laws’ that govern state transformations. This work provides a well-rounded, and novelly physical, perspective on equilibrium in the presence of quantum noncommutation. This perspective opens truly quantum avenues in thermodynamics.

Results

Overview. We derive the NATS’s form via three routes: from a microcanonical argument; from a dynamical argument built on canonical typicality; and from complete passivity in a resource theory. Details appear in Supplementary Notes 1–3.

Microcanonical derivation. In statistical mechanics, the form $e^{-\beta(H-\mu N)}/Z$ of the grand canonical ensemble is well known to be derivable as follows. The system of interest is assumed to be part of a larger system. Observables of the composite have fixed values v_j . For example, the energy equals E_0 , and the particle number equals N_0 . The microcanonical ensemble is the whole-system state spread uniformly across these observables’ simultaneous eigenspace. Tracing out the environmental degrees of freedom yields the state $e^{-\beta(H-\mu N)}/Z$.

We derive the NATS’s form similarly. Crucially, however, we adapt the above strategy to allow for noncommuting observables. Observables might not have well-defined values v_j simultaneously. Hence a microcanonical ensemble as discussed above, suitable for commuting observables, may not exist. We overcome this obstacle by introducing an approximate microcanonical ensemble Ω . We show that, for every state satisfying the conditions of an approximate microcanonical ensemble, tracing out most of the larger system yields, on average, a state close to the NATS. We exhibit conditions under which an approximate microcanonical ensemble exists. The conditions can be satisfied when the larger system consists of many noninteracting replicas of the system. An important step in the proof consists of reducing the noncommuting case to the commuting one. This reduction relies on a result by Ogata (ref. 29, Theorem 1.1). A summary appears in Fig. 1.

Set-up. Let \mathcal{S} denote a system associated with a Hilbert space \mathcal{H} , with a Hamiltonian $H \equiv Q_0$, and with observables (which we call ‘charges’) Q_1, Q_2, \dots, Q_c . The charges do not necessarily commute with each other: $[Q_j, Q_k] \neq 0$.

Consider N replicas of \mathcal{S} , associated with the composite-system Hilbert space $\mathcal{H}^{\otimes N}$. We average each charge Q_j over the N copies:

$$\bar{Q}_j := \frac{1}{N} \sum_{\ell=0}^{N-1} \mathbb{I}^{\otimes \ell} \otimes Q_j \otimes \mathbb{I}^{\otimes (N-1-\ell)} \quad (2)$$

The basic idea is that, as N grows, the averaged operators \bar{Q}_j come increasingly to commute. Indeed, there exist operators \bar{Y}_j that commute with each other and that approximate the averages (ref. 29, Theorem 1.1). An illustration appears in Fig. 2.

Derivation. Since the \bar{Y}_j ’s commute mutually, they can be measured simultaneously. More importantly, the joint Hilbert space $\mathcal{H}^{\otimes n}$ contains a subspace on which each \bar{Q}_j has prescribed values close to v_j . Let \mathcal{M} denote the subspace. Perhaps




Figure 1 | Non-Abelian thermal state. We derive the form of the thermal state of a system that has charges that might not commute with each other. Example charges include the components J_j of the spin \mathbf{J} . We derive the thermal state’s form by introducing an approximate microcanonical state. An ordinary microcanonical ensemble could lead to the thermal state’s form if the charges commuted: suppose, for example, that the charges were a Hamiltonian H and a particle number N that satisfied $[H, N] = 0$. Consider many copies of the system. The composite system could have a well-defined energy E_{tot} and particle number N_{tot} simultaneously. E_{tot} and N_{tot} would correspond to some eigensubspace $\mathcal{H}_{E_{\text{tot}}, N_{\text{tot}}}$ shared by the total Hamiltonian and the total-particle-number operator. The (normalized) projector onto $\mathcal{H}_{E_{\text{tot}}, N_{\text{tot}}}$ would represent the composite system’s microcanonical state. Tracing out the bath would yield the system’s thermal state. But the charges J_j under consideration might not commute. The charges might share no eigensubspace. Quantum noncommutation demands a modification of the ordinary microcanonical argument. We define an approximate microcanonical subspace \mathcal{M} . Each state in \mathcal{M} simultaneously has almost-well-defined values of noncommuting whole-system charges: measuring any such whole-system charge has a high probability of outputting a value close to an ‘expected’ value analogous to E_{tot} and N_{tot} . We derive conditions under which the approximate microcanonical subspace \mathcal{M} exists. The (normalized) projector onto \mathcal{M} represents the whole-system state. Tracing out most of the composite system yields the reduced state of the system of interest. We show that the reduced state is, on average, close to the NATS. This microcanonical derivation of the NATS’s form strengthens the link between Jaynes’s information-theoretic derivation and physics.

unsurprisingly, because the \bar{Y}_j ’s approximate the \bar{Q}_j ’s, each state in \mathcal{M} has a nearly well-defined value of \bar{Q}_j near v_j . If \bar{Q}_j is measured, the distribution is sharply peaked around v_j . We can also show the opposite: every state with nearly well-defined values v_j of all \bar{Q}_j ’s has most of its probability weight in \mathcal{M} .

These two properties show that \mathcal{M} is an approximate microcanonical subspace for the \bar{Q}_j ’s with values v_j . The notion of the approximate microcanonical subspace is the first major contribution of our work. It captures the idea that for large N we can approximately fix the values of the noncommuting charges Q_j . An approximate microcanonical subspace \mathcal{M} is any subspace consisting of the whole-system states whose average observables \bar{Q}_j have nearly well-defined values v_j . More precisely, a measurement of any \bar{Q}_j has a high probability of yielding a value near v_j if and only if most of the state’s probability weight lies in \mathcal{M} .

Normalizing the projector onto \mathcal{M} yields an approximate microcanonical ensemble, Ω . Tracing out every copy of S but the ℓ th yields the reduced state Ω_ℓ . The distance between Ω_ℓ and the NATS γ_v can be quantified by the relative entropy

$$D(\Omega_\ell || \gamma_v) := -S(\Omega_\ell) - \text{Tr}(\Omega_\ell \log \gamma_v). \quad (3)$$




Figure 2 | Noncommuting charges. We consider a thermodynamic system S that has conserved charges Q_j . These Q_j ’s might not commute with each other. The system occupies a thermal state whose form we derive. The derivation involves an approximate microcanonical state of a large system that contains the system of interest. Consider a block of n copies of S . Most copies act, jointly, similarly to a bath for the copy of interest. We define \bar{Q}_j as the average of the Q_j ’s of the copies in the block. Applying results from Ogata²⁹, we find operators \bar{Y}_j that are close to the \bar{Q}_j ’s and that commute with each other. Next, we consider m such blocks. This set of m blocks contains $N = mn$ copies of S . Averaging the \bar{Q}_j ’s over the blocks, for a fixed j -value, yields a global observable \bar{Q}_j . The \bar{Q}_j ’s are approximated by \bar{Y}_j ’s. The \bar{Y}_j ’s are the corresponding averages of the \bar{Y}_j ’s. The approximating global charges \bar{Y}_j commute with each other. The commuting \bar{Y}_j ’s enable us to extend the concept of a microcanonical ensemble from the well-known context in which all charges commute to truly quantum systems whose charges do not necessarily commute.

Here $S(\Omega_\ell) := -\text{Tr}(\Omega_\ell \log \Omega_\ell)$ is the von Neumann entropy. The relative entropy D is bounded by the trace norm $\|\cdot\|_1$, which quantifies the distinguishability of Ω_ℓ and γ_v (ref. 30):

$$D(\Omega_\ell || \gamma_v) \geq \frac{1}{2} \|\Omega_\ell - \gamma_v\|_1^2. \quad (4)$$

Our second main result is that if Ω is an approximate microcanonical ensemble, then the average, over systems ℓ , of the relative entropy D between Ω_ℓ and γ_v is small:

$$\frac{1}{N} \sum_{\ell=0}^{N-1} D(\Omega_\ell || \gamma_v) \leq \theta + \theta'. \quad (5)$$

The parameter $\theta = (\text{const.})/\sqrt{N}$ vanishes in the many-copy limit. θ' depends on the number c of charges, on the approximate expectation values v_j , on the eigenvalues of the charges Q_j and on the (small) parameters in terms of which \mathcal{M} approximates a microcanonical subspace.

Inequality (5) capstones the derivation. The inequality follows from bounding each term in equation (3), the definition of the relative entropy D . The entropy $S(\Omega_\ell)$ is bounded with θ . This bound relies on Schumacher’s Theorem, which quantifies the size of a high-probability subspace like \mathcal{M} with an entropy $S(\gamma_v)$ ³¹. We bound the second term in the D definition with θ' . This bound relies on the definition of \mathcal{M} : outcomes of measurements of the \bar{Q}_j ’s are predictable up to parameters on which θ' depends.

Finally, we present conditions under which the approximate microcanonical subspace \mathcal{M} exists. Several parameters quantify the approximation. The parameters are shown to be interrelated and to approach zero simultaneously as N grows. In particular, the approximate microcanonical subspace \mathcal{M} exists if N is great enough.

This microcanonical derivation offers a physical counterpoint to Jaynes’s maximum-entropy derivation of the NATS’s form. We relate the NATS to the physical picture of a small subsystem in a vast universe that occupies an approximate microcanonical state. This vast universe allows the correspondence principle to underpin our argument. In the many-copy limit as $N \rightarrow \infty$, the principle implies that quantum behaviours

should vanish, as the averages of the noncommuting charges Q_j come to be approximated by commuting \bar{Y}_j 's. Drawing on Ogata's Theorem 1.1 (ref. 29), we link thermality in the presence of noncommutation to the physical correspondence principle.

Dynamical considerations. The microcanonical and maximum-entropy arguments rely on kinematics and information theory. But we wish to associate the NATS with the fixed point of dynamics. The microcanonical argument, combined with canonical typicality, suggests that the NATS is the equilibrium state of typical dynamics. Canonical typicality enables us to model the universe's state with a pure state in the approximate microcanonical subspace \mathcal{M} . If a large system occupies a randomly chosen pure state, the reduced state of a small subsystem is close to thermal^{5–8}.

Consider, as in the previous section, N copies of the system \mathcal{S} . By Ω , we denoted the composite system's approximately microcanonical state. We denoted by Ω_ℓ the reduced state of the ℓ th copy, formed by tracing out most copies from Ω . Imagine that the whole system occupies a pure state $|\psi\rangle \in \mathcal{M}$. Denote by ρ_ℓ the reduced state of the ℓ th copy. ρ_ℓ is close to Ω_ℓ , on average, by canonical typicality⁷:

$$\langle \|\rho_\ell - \Omega_\ell\|_1 \rangle \leq \frac{d}{\sqrt{D_M}}. \quad (6)$$

The average $\langle \cdot \rangle$ is over pure states $|\psi\rangle \in \mathcal{M}$. The trace norm is denoted by $\|\cdot\|_1$; $d := \dim(\mathcal{H})$ denotes the dimensionality of the Hilbert space \mathcal{H} of one copy of \mathcal{S} ; and $D_M := \dim(\mathcal{M})$ denotes the dimensionality of the approximate microcanonical subspace \mathcal{M} .

We have bounded, using canonical typicality, the average trace norm between ρ_ℓ and Ω_ℓ . We can bound the average trace norm between Ω_ℓ and the NATS γ_v , using our microcanonical argument. (Supplementary equation (10) bounds the average relative entropy D between Ω_ℓ and γ_v . Pinsker's inequality, Ineq. (4), lower bounds D in terms of the trace norm.) Combining these two trace-norm bounds via the triangle inequality, we bound the average distance between ρ_ℓ and γ_v :

$$\left\langle \frac{1}{N} \sum_{\ell=0}^{N-1} \|\rho_\ell - \gamma_v\|_1 \right\rangle \leq \frac{d}{\sqrt{D}} + \sqrt{2(\theta + \theta')}. \quad (7)$$

If the whole system occupies a random pure state $|\psi\rangle$ in \mathcal{M} , the reduced state ρ_ℓ of a subsystem is, on average, close to the NATS γ_v .

Sufficiently ergodic dynamics is expected to evolve the whole-system state to a $|\psi\rangle$ that satisfies Ineq. (7): suppose that the whole system begins in a pure state $|\psi(t=0)\rangle \in \mathcal{M}$. Suppose that the system's Hamiltonian commutes with the charges: $[H, Q_j] = 0$ for all $j = 1, \dots, c$. The dynamics conserves the charges. Hence, most of the amplitude of $|\psi(t)\rangle$ remains in \mathcal{M} for appreciable times. Over sufficient times, ergodic dynamics yields a state $|\psi(t)\rangle$ that can be regarded as random. Hence the reduced state is expected to be close to $\Omega_\ell \approx \gamma_v$ for most long-enough times t .

Exploring how the dynamics depends on the number of copies of the system offers promise for interesting future research.

Resource theory. A thermodynamic resource theory is an explicit characterization of a thermodynamic system's resources, free states and free operations with a rigorous mathematical model. The resource theory specifies what an experimenter considers valuable (for example, work) and what is considered plentiful, or free (for example, thermal states). To define a resource theory, we specify allowed operations and which states can be accessed for free. We use this framework to quantify the resources needed to transform one state into another.

The first resource theory was entanglement theory³². The theory's free operations are local operations and classical communication. The free states are the states that can be easily prepared with local operations and classical communication, the separable states. Entangled states constitute valuable resources. One can quantify entanglement using this resource theory.

We present a resource theory for thermodynamic systems that have noncommuting conserved charges Q_j . The theory is defined by its set of free operations, which we call 'non-Abelian thermal operations' (NATO). NATO generalize thermal operations^{15,18}. How to extend thermodynamic resource theories to conserved quantities other than energy was noted in refs 18,20,21. The NATO theory is related to the resource theory in ref. 27.

We supplement these earlier approaches with two additions. First, a battery has a work payoff function dependent on chemical potentials. We use this payoff function to define chemical work. Second, we consider a reference system for a non-Abelian group. The reference system is needed to resolve the difficulty encountered in refs 21,27: There might be no nontrivial operations which respect all the conservation laws. The laws of physics require that any operation performed by an experimenter commutes with all the charges. If the charges fail to commute with each other, there might be no non-trivial unitaries that commute with all of them. In practice, one is not limited by such a stringent constraint. The reason is that an experimenter has access to a reference frame^{33–35}.

A reference frame is a system W prepared in a state such that for any unitary on a system S which does not commute with the charges of S , some global unitary on WS conserves the total charges and approximates the unitary on S to arbitrary precision. The reference frame relaxes the strong constraint on the unitaries. The reference frame can be merged with the battery, in which the agent stores the ability to perform work. We refer to the composite as 'the battery'. We denote its state by ρ_W . The battery has a Hamiltonian H_W and charges $Q_{j,W}$, described below.

Within this resource theory, the NATS emerges in two ways:

1. The NATS is the unique state from which work cannot be extracted, even if arbitrarily many copies are available. That is, the NATS is completely passive.
2. The NATS is the only state of S that remains invariant under the free operations during which no work is performed on S .

On proving the latter condition, we prove second laws for thermodynamics with noncommuting charges. These laws provide necessary conditions for a transition to be possible. In some cases, we show, the laws are sufficient. These second laws govern state transitions of a system ρ_S , governed by a Hamiltonian H_S , whose charges $Q_{j,S}$ can be exchanged with the surroundings. We allow the experimenter to couple ρ_S to free states ρ_R . The form of ρ_R is determined by the Hamiltonian H_R and the charges $Q_{j,R}$ attributable to the free system. We will show that these free states have the form of the NATS. As noted above, no other state could be free. If other states were free, an arbitrarily large amount of work could be extracted from them.

Before presenting the second laws, we must define 'work'. In textbook examples about gases, one defines work as $\delta W = p dV$, because a change in volume at a fixed pressure can be translated into the ordinary notion of mechanical work. If a polymer is stretched, then $\delta W = F dx$, wherein x denotes the polymer's linear displacement and F denotes the restoring force. If B denotes a magnetic field and M denotes a medium's magnetization, $\delta W = B dM$. The definition of 'work' can depend on one's ability to transform changes in thermodynamic variables into a standard notion of 'work,' such as mechanical or electrical work.

Our approach is to define a notion of chemical work. We could do so by modelling explicitly how the change in some quantity Q_j can be used to extract $\mu_j \delta Q_j$ work. Explicit modelling would involve adding a term to the battery Hamiltonian H_W . Rather than considering a specific work Hamiltonian or model of chemical work, however, we consider a work payoff function,

$$\mathcal{W} = \sum_{j=0}^c \mu_j Q_{jw} \quad (8)$$

The physical situation could determine the form of this \mathcal{W} . For example, the μ_j 's could denote the battery's chemical potentials. In such a case, \mathcal{W} would denote the battery's total Hamiltonian, which would depend on those potentials.

We choose a route conceptually simpler than considering an explicit Hamiltonian and battery system, however. We consider equation (8) as a payoff function that defines the linear combination of charges that interests us. We define the (chemical) work expended or distilled during a transformation as the change in the quantum expectation value $\langle \mathcal{W} \rangle$.

The form of \mathcal{W} is implicitly determined by the battery's structure and by how charges can be converted into work. For our purposes, however, the origin of the form of \mathcal{W} need not be known. \mathcal{W} will uniquely determine the μ_j 's in the NATS. Alternatively, we could first imagine that the agent could access, for free, a particular NATS. This NATS's form would determine the work function's form. If the charges commute, the corresponding Gibbs state is known to be the unique state that is completely passive with respect to the observable (8).

In Supplementary Note 3, we specify the resource theory for noncommuting charges in more detail. We show how to construct allowable operations, using the reference frame and battery. From the allowable operations, we derive a zeroth law of thermodynamics.

Complete passivity and zeroth law. This zeroth law relates to the principle of complete passivity, discussed in refs 25,26. A state is complete passive if an agent cannot extract work from arbitrarily many copies of the state. In the resource theory for heat exchanges, completely passive states can be free. They do not render the theory trivial because no work can be drawn from them¹⁷.

In the NATO resource theory, we show, the only reasonable free states have the NATS's form. The free states' chemical potentials equal the μ_j 's in the payoff function \mathcal{W} , at some common fixed temperature. Any other state would render the resource theory trivial: from copies of any other state, arbitrarily much work could be extracted for free. Then, we show that the NATS is preserved by NATO, the operations that perform no work on the system.

The free states form an equivalence class. They lead to notions of temperature and chemical potentials μ_j . This derivation of the free state's form extends complete passivity and the zeroth law from ref. 17 to noncommuting conserved charges. The derivation further solidifies the role of the NATS in thermodynamics.

Second laws. The free operations preserve the NATS. We therefore focus on contractive measures of states' distances from the NATS. Contractive functions decrease monotonically under the free operations. Monotones feature in 'second laws' that signal whether NATO can implement a state transformation. For example, the α -Rényi relative entropies between a state and the NATS cannot increase.

Monotonicity allows us to define generalized free energies as

$$F_\alpha(\rho_S, \gamma_S) := k_B TD_\alpha(\rho_S || \gamma_S) - k_B T \log Z \quad (9)$$

wherein $\beta \equiv 1/(k_B T)$ and k_B denotes Boltzmann's constant. γ_S denotes the NATS with respect to the Hamiltonian H_S and the

charges Q_{js} of the system S . The partition function is denoted by Z . Various classical and quantum definitions of the Rényi relative entropies D_α are known to be contractive^{17,36–39}. The free energies F_α decrease monotonically if no work is performed on the system. Hence the F_α 's characterize natural second laws that govern achievable transitions.

For example, the classical Rényi divergences $D_\alpha(\rho_S || \gamma_S)$ are defined as

$$D_\alpha(\rho_S || \gamma_S) := \frac{\text{sgn}(\alpha)}{\alpha - 1} \log \left(\sum_k p_k^\alpha q_k^{1-\alpha} \right), \quad (10)$$

wherein p_k and q_k denote the probabilities of ρ_S and of γ_S in the \mathcal{W} basis. The D_α 's lead to second laws that hold even in the absence of a reference frame and even outside the context of the average work.

The F_α 's reduce to the standard free energy when averages are taken over large numbers. Consider the asymptotic ('thermodynamic') limit in which many copies $(\rho_S)^{\otimes n}$ of ρ_S are transformed. Suppose that the agent has some arbitrarily small probability ε of failing to implement the desired transition. ε can be incorporated into the free energies via a technique called 'smoothing'¹⁷. The average, over copies of the state, of every smoothed F_α^ε approaches F_1 (ref. 17):

$$\lim_{n \rightarrow \infty} \frac{1}{n} F_\alpha^\varepsilon((\rho_S)^{\otimes n}, (\gamma_S)^{\otimes n}) = F_1 \quad (11)$$

$$= k_B TD(\rho_S || \gamma_S) - k_B T \log(Z) \quad (12)$$

$$= \langle H_S \rangle_{\rho_S} - k_B TS(\rho_S) + \sum_{j=1}^c \mu_j \langle Q_{js} \rangle. \quad (13)$$

We have invoked the relative entropy's definition

$$D(\rho_S || \gamma_S) := \text{Tr}(\rho_S \log(\rho_S)) - \text{Tr}(\rho_S \log(\gamma_S)). \quad (14)$$

Note the similarity between the many-copy average F_1 in equation (13) and the ordinary free energy, $F = E - TS + \sum_j \mu_j N_j$. The monotonic decrease of F_1 constitutes a necessary and sufficient condition for a state transition to be possible in the presence of a reference system in the asymptotic limit.

In terms of the generalized free energies, we formulate second laws.

Proposition 1: in the presence of a heat bath of inverse temperature β and chemical potentials μ_j , the free energies $F_\alpha(\rho_S, \gamma_S)$ decrease monotonically:

$$F_\alpha(\rho_S, \gamma_S) \geq F_\alpha(\rho'_S, \gamma'_S) \quad \forall \alpha \geq 0 \quad (15)$$

wherein ρ_S and ρ'_S denote the system's initial and final states. The system's Hamiltonian and charges may transform from H_S and Q_{js} to H'_S and Q'_{js} . The NATSs associated with the same Hamiltonians and charges are denoted by γ_S and γ'_S . If

$$[\mathcal{W}, \rho'_S] = 0 \quad \text{and} \quad F_\alpha(\rho_S, \gamma_S) \geq F_\alpha(\rho'_S, \gamma'_S) \quad \forall \alpha \geq 0 \quad (16)$$

some NATO maps ρ_S to ρ'_S .

As in ref. 17, additional laws can be defined in terms of quantum Rényi divergences^{36–39}. This amounts to choosing, in Proposition 1, a definition of the Rényi divergence which accounts for the possibility that ρ_S and ρ'_S have coherences relative to the \mathcal{W}_S eigenbasis. Several measures are known to be contractive^{36–39}. They, too, provide a new set of second laws.

Extractable work: in terms of the free energies F_α , we can bound the work extractable from a resource state via NATO. We consider the battery W separately from the system S of interest. We assume that W and S occupy a product state.

(This assumption is unnecessary if we focus on average work.) Let ρ_W and ρ'_W denote the battery's initial and final states.

For all α ,

$$F_\alpha(\rho_S \otimes \rho_W, \gamma_{SW}) \geq F_\alpha(\rho'_S \otimes \rho'_W, \gamma_{SW}). \quad (17)$$

Since $F_\alpha(\rho_S \otimes \rho_W, \gamma_{SW}) = F_\alpha(\rho_S, \gamma_S) + F_\alpha(\rho_W, \gamma_W)$,

$$F_\alpha(\rho'_W, \gamma_W) - F_\alpha(\rho_W, \gamma_W) \leq F_\alpha(\rho_S, \gamma_S) - F_\alpha(\rho'_S, \gamma_S). \quad (18)$$

The left-hand side of Ineq. (18) represents the work extractable during one implementation of $\rho_S \rightarrow \rho'_S$. Hence the right-hand side bounds the work extractable during the transition.

Consider extracting work from many copies of ρ_S (that is, extracting work from $\rho_S^{\otimes n}$) in each of many trials. Consider the average-over-trials extracted work, defined as $\text{Tr}(\mathcal{W}[\rho'_W - \rho_W])$. The average-over-trials work extracted per copy of ρ_S is $\frac{1}{n} \text{Tr}(\mathcal{W}[\rho'_W - \rho_W])$. This average work per copy has a high probability of lying close to the change in the expectation value of the system's work function, $\frac{1}{n} \text{Tr}(\mathcal{W}[\rho'_W - \rho_W]) \approx \text{Tr}(\mathcal{W}[\rho'_S - \rho_S])$, if n is large.

Averaging over the left-hand side of Ineq. (18) yields the average work $\delta\langle W \rangle$ extracted per instance of the transformation. The average over the right-hand side approaches the change in F_1 (equation (13)):

$$\delta\langle W \rangle \leq \delta\langle H_S \rangle_{\rho_S} - T \delta S(\rho_S) + \sum_{j=1}^c \mu_j \langle \delta_j \rangle \quad (19)$$

This bound is achievable with a reference system, as shown in refs 40,41.

We have focused on the extraction of work defined by \mathcal{W} . One can extract, instead, an individual charge Q_j . The second laws do not restrict single-charge extraction. But extracting much of one charge Q_j precludes the extraction of much of another charge, Q_k . In Supplementary Note 3, we discuss the trade-offs amongst extractions of different charges Q_j .

Discussion

We have derived, via multiple routes, the form of the thermal state of a system that has noncommuting conserved charges. First, we regarded the system as part of a vast composite that occupied an approximate microcanonical state. Tracing out the environment yields a reduced state that lies, on average, close to a thermal state of the expected form. This microcanonical argument, with canonical typicality, suggests that the NATS is the fixed point of typical dynamics. Defining a resource theory, we showed that the NATS is the only completely passive state and is the only state preserved by free operations. These physical derivations buttress Jaynes's information-theoretic derivation from the principle of maximum entropy.

Our derivations also establish tools applicable to quantum noncommutation in thermodynamics. In the microcanonical argument, we introduced an approximate microcanonical state Ω . This Ω resembles the microcanonical ensemble associated with a fixed energy, a fixed particle number and so on but accommodates noncommuting charges. Our complete-passivity argument relies on a little-explored resource theory for thermodynamics, in which free unitaries conserve noncommuting charges.

We expect that the equilibrium behaviours predicted here may be observed in experiments. Quantum gases have recently demonstrated equilibrium-like predictions about integrable quantum systems^{11,13}.

From a conceptual perspective, our work shows that notions previously considered relevant only to commuting charges—for example, the microcanonical subspace—extend to

noncommuting charges. This work opens fully quantum thermodynamics to analysis with familiar, but suitably adapted, technical tools.

Data availability. Data sharing is not applicable to this article, as no data sets were generated or analysed during this study.

References

- Gemmer, J., Michel, M., Michel, M. & Mahler, G. *Quantum Thermodynamics: Emergence of Thermodynamic Behavior within Composite Quantum Systems* (Springer, 2009).
- Gogolin, C. & Eisert, J. Equilibration, thermalisation, and the emergence of statistical mechanics in closed quantum systems. *Rep. Prog. Phys.* **79**, 056001 (2016).
- Goold, J., Huber, M., Riera, A., del Rio, L. & Skrzypczyk, P. The role of quantum information in thermodynamics—a topical review. *J. Phys. A: Math. Theor.* **49**, 143001 (2016).
- Vinjanampathy, S. & Anders, J. Quantum thermodynamics. Preprint at <http://arxiv.org/abs/1508.06099> (2015).
- Goldstein, S., Lebowitz, J. L., Tumulka, R. & Zanghi, N. Canonical typicality. *Phys. Rev. Lett.* **96**, 050403 (2006).
- Gemmer, J., Michel, M. & Mahler, G. *18 Equilibrium Properties of Model Systems* (Springer, 2004).
- Popescu, S., Short, A. J. & Winter, A. The foundations of statistical mechanics from entanglement: individual states versus averages. *Nat. Phys.* **2**, 754–758 (2006).
- Linden, N., Popescu, S., Short, A. J. & Winter, A. Quantum mechanical evolution towards thermal equilibrium. *Phys. Rev. E* **79**, 061103 (2009).
- Fermi, E., Pasta, J. & Ulam, S. *Studies of Nonlinear Problems I*. Los Alamos Report LA-1940 (Los Alamos Scientific Laboratory, 1955).
- Kinoshita, T., Wenger, T. & Weiss, D. S. A quantum Newton's cradle. *Nature* **440**, 900–903 (2006).
- Rigol, M., Dunjko, V., Yurovsky, V. & Olshanii, M. Relaxation in a completely integrable many-body quantum system: an *Ab Initio* study of the dynamics of the highly excited states of 1d lattice hard-core bosons. *Phys. Rev. Lett.* **98**, 050405 (2007).
- Polkovnikov, A., Sengupta, K., Silva, A. & Vengalattore, M. *Colloquium: nonequilibrium dynamics of closed interacting quantum systems*. *Rev. Mod. Phys.* **83**, 863–883 (2011).
- Langen, T. *et al.* Experimental observation of a generalized Gibbs ensemble. *Science* **348**, 207–211 (2015).
- Langen, T., Geiger, R. & Schmiedmayer, J. Ultracold atoms out of equilibrium. *Annu. Rev. Condens. Matter Phys.* **6**, 201–217 (2015).
- Janzing, D., Wocjan, P., Zeier, R., Geiss, R. & Beth, T. Thermodynamic cost of reliability and low temperatures: tightening Landauer's principle and the second law. *Int. J. Theor. Phys.* **39**, 2717–2753 (2000).
- Brandão, F. G. S. L., Horodecki, M., Oppenheim, J., Renes, J. M. & Spekkens, R. W. Resource theory of quantum states out of thermal equilibrium. *Phys. Rev. Lett.* **111**, 250404 (2013).
- Brandao, F. G. S. L., Horodecki, M., Ng, N. H. Y., Oppenheim, J. & Wehner, S. The second laws of quantum thermodynamics. *Proc. Natl Acad. Sci. USA* **112**, 3275–3279 (2015).
- Horodecki, M. & Oppenheim, J. Fundamental limitations for quantum and nano thermodynamics. *Nat. Commun.* **4**, 2059 (2013).
- Vaccaro, J. A. & Barnett, S. M. Information erasure without an energy cost. *Proc. R. Soc. A* **467**, 1770–1778 (2011).
- Yunger Halpern, N. & Renes, J. M. Beyond heat baths: generalized resource theories for small-scale thermodynamics. *Phys. Rev. E* **93**, 022126 (2016).
- Yunger Halpern, N. Beyond heat baths II: Framework for generalized thermodynamic resource theories. Preprint at <http://arxiv.org/abs/1409.7845> (2014).
- Weilenmann, M., Krämer, L., Faist, P. & Renner, R. Axiomatic relation between thermodynamic and information-theoretic entropies. Preprint at <http://arxiv.org/abs/1501.06920> (2015).
- Jaynes, E. T. Information theory and statistical mechanics I. *Phys. Rev.* **106**, 620–630 (1957).
- Jaynes, E. T. Information theory and statistical mechanics II. *Phys. Rev.* **108**, 171–190 (1957).
- Pusz, W. & Woronowicz, S. L. Passive states and KMS states for general quantum systems. *Comm. Math. Phys.* **58**, 273–290 (1978).
- Lenard, A. Thermodynamical proof of the Gibbs formula for elementary quantum systems. *J. Stat. Phys.* **19**, 575 (1978).
- Lostaglio, M., Jennings, D. & Rudolph, T. Thermodynamic resource theories, non-commutativity and maximum entropy principles. Preprint at <http://arxiv.org/abs/1511.04420> (2015).

28. Guryanova, Y., Popescu, S., Short, A. J., Silva, R. & Skrzypczyk, P. Thermodynamics of quantum systems with multiple conserved quantities. *Nat. Commun.* **7**, 12049 (2016).
29. Ogata, Y. Approximating macroscopic observables in quantum spin systems with commuting matrices. *J. Funct. Anal.* **264**, 2005–2033 (2013).
30. Hiai, F., Ohya, M. & Tsukada, M. Sufficiency, KMS condition and relative entropy in von Neumann algebras. *Pacific J. Math.* **96**, 99–109 (1981).
31. Schumacher, B. Quantum coding. *Phys. Rev. A* **51**, 2738–2747 (1995).
32. Horodecki, R., Horodecki, P., Horodecki, M. & Horodecki, K. Quantum entanglement. *Rev. Mod. Phys.* **81**, 865–942 (2009).
33. Aharonov, Y. & Susskind, L. Charge superselection rule. *Phys. Rev.* **155**, 1428–1431 (1967).
34. Kitaev, A., Mayers, D. & Preskill, J. Superselection rules and quantum protocols. *Phys. Rev. A* **69**, 052326 (2004).
35. Bartlett, S. D., Rudolph, T. & Spekkens, R. W. Reference frames, superselection rules, and quantum information. *Rev. Mod. Phys.* **79**, 555–609 (2007).
36. Hiai, F., Mosonyi, M., Petz, D. & Bény, C. Quantum f -divergences and error correction. *Rev. Math. Phys.* **23**, 691–747 (2011).
37. Müller-Lennert, M., Dupuis, F., Szehr, O., Fehr, S. & Tomamichel, M. On quantum Rényi entropies: a new generalization and some properties. *J. Math. Phys.* **54**, 122203 (2013).
38. Wilde, M. M., Winter, A. & Yang, D. Strong converse for the classical capacity of entanglement-breaking and Hadamard channels via a sandwiched Rényi relative entropy. *Commun. Math. Phys.* **331**, 593–622 (2014).
39. Jaksic, V., Ogata, Y., Pautrat, Y. & Pillet, C.-A. in *Quantum Theory from Small to Large Scales: Lecture Notes of the Les Houches Summer School*. Vol. **95** (eds Fröhlich, J., Manfred, S., Vieri, M., De Roeck, W. & Cugliandolo, L. F.) (Oxford Univ. Press, 2012).
40. Åberg, J. Catalytic coherence. *Phys. Rev. Lett.* **113**, 150402 (2014).
41. Korzekwa, K., Lostaglio, M., Oppenheim, J. & Jennings, D. The extraction of work from quantum coherence. *N. J. Phys.* **18**, 023045 (2016).

Acknowledgements

We thank David Jennings, Tim Langen, Elliott Lieb (who pointed us to ref. 29), Matteo Lostaglio, Shelly Moram, Joseph M. Renes and Terry Rudolph for interesting conversations. We thank the authors of refs 27,28 for their community spirit. Much of this paper was developed at ‘Beyond i.i.d. in Information Theory 2015,’ hosted by BIRS. A.W.’s work was supported by the EU (STREP ‘RAQUEL’), the ERC

(AdG ‘IRQUAT’), the Spanish MINECO (grant FIS2013-40627-P) with the support of FEDER funds, as well as by the Generalitat de Catalunya CIRIT, project 2014-SGR-966. J.O. is supported by an EPSRC Established Career Fellowship, the Royal Society and FQXi. N.Y.H. was supported by an IQIM Fellowship and NSF grant PHY-0803371. The Institute for Quantum Information and Matter (IQIM) is an NSF Physics Frontiers Center supported by the Gordon and Betty Moore Foundation. P.F. acknowledges support from the European Research Council (ERC) via grant No. 258932, from the Swiss National Science Foundation through the National Centre of Competence in Research ‘Quantum Science and Technology’ (QSIT), and by the European Commission via the project ‘RAQUEL.’ This work was partially supported by the COST Action MP1209.

Author contributions

All authors contributed to all aspects of this work.

Additional information

Supplementary Information accompanies this paper at <http://www.nature.com/naturecommunications>

Competing financial interests: The authors declare no competing financial interests.

Reprints and permission information is available online at <http://npg.nature.com/reprintsandpermissions/>

How to cite this article: Yunger Halpern, N. *et al.* Microcanonical and resource-theoretic derivations of the thermal state of a quantum system with noncommuting charges. *Nat. Commun.* **7**:12051 doi: 10.1038/ncomms12051 (2016).



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article’s Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>

© The Author(s) 2016