

Prevalence of Late Life Anxiety Disorders

TITLE: PREVALENCE OF ANXIETY DISORDERS IN COMMUNITY LIVING
OLDER ADULTS IN HONG KONG

Ms Ada, Wai Tung, Fung, BA, MSc ¹

Prof. Linda, Chiu Wa, Lam, MD, FRCPsych, FHKCPsych, FHKAM (Psychiatry) ¹

Dr Wai-Chi Chan, MRCPsych, FHKCPsych, FHKAM (Psychiatry). ²

Ms Corine Sau-Man Wong, MSocSc ²

Prof. Eric Yu-Hai Chen, MA, MD, FRCPsych, FHKCPsych, FHKAM (Psychiatry) ²

Dr Roger Man-Kin Ng, FRCPsych, FHKAM (Psychiatry) ³

Dr Edwin Ho-Ming Lee, MRCPsych, FHKCPsych, FHKAM (Psychiatry) ².

Dr Wing-Chung Chang, MRCPsych, FHKCPsych, FHKAM (Psychiatry) ².

Dr Se-Fong Hung, FRCPsych, FHKCPsych, FHKAM (Psychiatry) ⁴

Dr Eric Fuk-Chi Cheung, FRCPsych, FHKCPsych, FHKAM (Psychiatry) ⁵

Dr Pak-Chung Sham, BA, MSc, PhD, MRPsych ²

Dr Helen Fung-Kum Chiu, FRCPsych, FHKCPsych, FHKAM (Psychiatry) ¹

Dr Ming Lam, MRCPsych, FHKCPsych, FHKAM (Psychiatry) ⁵

Dr Tin-Po Chiang, MRCPsych, FHKCPsych, FHKAM (Psychiatry) ⁵

Prof. Jim van Os, MD, PhD ⁶

Prof. Joseph Tak-Fai Lau, BSSc, MA, PhD, FFPH ⁷

Prof. Glyn Lewis, PhD, FRCPsych ⁸

Prof. Paul Bebbington, PhD, FRCPsych ⁹

The Hong Kong Mental Morbidity Survey Team ¹

Corresponding Author

Ada, Wai Tung, Fung

E-mail address: ada_fung@cuhk.edu.hk

Tel.no: 852-26076045

Fax.no: 852-26675464

Department of Psychiatry

The Chinese University of Hong Kong

Shatin, New Territories

Hong Kong SAR

China

¹ Department of Psychiatry, The Chinese University of Hong Kong, Hong Kong SAR, China

² Department of Psychiatry, The University of Hong Kong, Hong Kong SAR, China

³ Kowloon Hospital, Hong Kong SAR, China

⁴ Kwai Chung Hospital, Hong Kong SAR, China.

⁵ Castle Peak Hospital, Hong Kong SAR, China

⁶ Department of Psychiatry and Psychology, Maastricht University Medical Centre, Maastricht, The Netherlands

⁷ Jockey Club School of Public Health and Primary Care, The Chinese University of Hong Kong, Hong Kong SAR, China

⁸ Academic Unit of Psychiatry, University of Bristol, Bristol, England

⁹ Department of Mental Health Sciences, University College London, London, England

Abstract

Background: Anxiety disorders are prevalent yet under-recognized in late life. We examined the prevalence of anxiety disorders on a representative sample of community living older adults in Hong Kong.

Method: Data on 1158 non-demented respondents aged 60 to 75 years were extracted from the Hong Kong Mental Morbidity survey (HKMMS). Anxiety was assessed with the revised Clinical Interview Schedule (CIS-R).

Result: 137 respondents (11.9%, 95% CI=10-13.7%) had common mental disorders with a CIS-R score of 12 or above. 8% (95% CI=6.5 – 9.6%) had anxiety, 2.2% (95% CI=1.3 – 3%) had comorbid mood disorders, and 1.7% (95% CI=1 – 2.5%) had depression. Anxious individuals were more likely to be females ($\chi^2=25.3$, $p<0.001$), had higher chronic physical burden ($t=-9.3$, $p<0.001$), lower SF-12 physical functioning score ($t=9.2$, $p<0.001$), and poorer delayed recall ($t=2.3$, $p=0.022$). Two risk factors were identified: females (OR 2.9, 95% C.I. 1.2 – 6.7, $p<0.05$) and poor physical health (OR 1.4, 95% C.I. 1.3 – 1.6, $p<.001$). The risk of anxiety disorders increased in those with disorders of cardiovascular (OR 1.3, 95% C.I. 1.1– 1.6, $p=0.01$), musculoskeletal (OR 2.0, 95% C.I. 1.5– 2.7, $p<0.001$), and genitourinary system (OR 1.8, 95% C.I. 1.3– 2.7, $p=0.001$).

Conclusions: The prevalence of anxiety disorders in Hong Kong older population was 8%. Female gender and those with poor physical health were at greater risk of developing

anxiety disorders. Our findings also suggested potential risk for early sign of memory impairment in cognitively healthy individuals with anxiety disorders.

KEYWORDS: prevalence, anxiety disorders, community, older adults, memory, cognitive function, early sign, mental health

Introduction

Previous studies on mental illness in old age have focused heavily on depression and dementia. Yet, interests in anxiety disorders have increased in recent years, but the range of prevalence estimates of anxiety disorders in old age varied markedly. The prevalence of any anxiety disorders ranges from 3.2% to 15.3% in community based studies (Forsell & Winblad, 1997; Kessler *et al.*, 2005). Most of the studies of the prevalence of anxiety disorders were based on the DSM-IV diagnostic criteria (DSM-4 Ed.). It requires that excessive anxiety and worry occur for at least 6 months, the worry is difficult to control, and 3 or more anxiety related somatic symptoms are present: restlessness, fatigue, difficulty concentrating, irritability, muscle tension, sleep disturbance (DSM-4 Ed.). Despite this well delineated description, there are still about 15% to 52.3% of older adults experiencing subclinical level of anxiety symptoms that are also disturbing to everyday life (Bryant *et al.*, 2008). This variability could be due to methodological differences, such as the sampling frame recruited and diagnostic criteria in use (Bryant *et al.*, 2008).

Anxiety disorders appear to decline in prevalence with advancing age, and it is more common than depressive disorders (Schaub & Linden, 2000). Studies from western countries estimated that the 12 months prevalence rate of late-life depression in community dwelling older people ranges between 3% and 4.5% (Aziz & Steffens, 2013), whereas late-life anxiety disorders range between 6 to 10% (DSM-4 Ed.). Furthermore, the LASA study indicated a 6-month prevalence of depression to be 2% compared to a prevalence of 7.3% for anxiety disorder in the same cohort. These evidences seem to

suggest that anxiety disorders are more common than depression in the community older population.

It is also important to highlight that anxiety disorders are an under-recognized condition in the older population. Although treatments of anxiety disorders are available, only 17% to 22% of older people identified in an epidemiological survey received proper treatment (Himmelfarb & Murrell, 1984). Furthermore, anxiety disorders are also associated with disability and suffering that is not limited to the elderly people who affected by the condition alone, but also includes burdens for family members and the society. Recently, research has focused on the relationship between anxiety disorders and cognitive functions, but the results are still inconclusive. Therefore, the aim of this study is to extend available knowledge on the prevalence and correlates of anxiety disorders in the older population.

In Hong Kong, the prevalence of anxiety disorders in late life has never been systematically investigated. Although figures from other developed countries are available, we could not directly generalize the findings to reflect the situation in Hong Kong, due to cultural differences and different demographic makeup. Besides, most of the systematic epidemiological surveys on the prevalence of late life anxiety disorders were conducted a decade ago (Kessler *et al.*, 2005; Regier *et al.*, 1988). Therefore, a local study on current prevalence estimates of anxiety disorders in this target population is both timely and relevant.

Methodology

Study Design and participants

The sample was drawn from the dataset of Hong Kong Mental Morbidity Survey (HKMMS) that conducted between November 2010 and May 2013. The HKMMS sample was recruited through a multi-layered random sampling of community living adults in Hong Kong. A detailed description of the population survey has been provided elsewhere (Lam *et al.*, 2015). Face to face interviews were completed with a sample of 5,719 community-living Chinese adults aged 16 to 75 years in Hong Kong, with a participation rate of 68%.

Random address localities were generated from the Census and Statistics Department of the Government of Hong Kong SAR. The addresses were stratified by geographical locations and nature of premises (private versus public housing). Samples of geographical areas representative of the residential distribution in Hong Kong were selected, and followed by subsequent stages of geographical sampling (e.g. estates within districts, blocks within estates, households with blocks).

For each address identified, an advance letter was sent to introduce the survey and stated that an interviewer would be calling to seek permission for interview. At least three advance letters and/or one home visit were conducted within three months of first posting to ascertain the availability of eligible participants. For households with either no eligible participant, failure to gain access, non-response or unknown eligibility, a replacement address quarter with similar geographical and housing characteristics was generated. No substitution was carried out if the original household resident refused to participate. For each eligible household, a trained lay interviewer conducted the assessment. A household member with closest birthday to the day of interview was selected for assessment in order to avoid response bias.

The inclusion criteria of the HKMMS are: (i) aged between 16 to 75 years; (ii) Ethnically Chinese; (iii) birthday closest to the date of first scheduled interview; and (iv) consent for participation. Proxy information from first degree relatives was obtained if the subjects were unavailable or unable to provide full information (e.g. intellectual disability or sensory impairments).

For the present analyses, participants aged 60 years or older were considered (N=1,228) from the HKMMS dataset. Individuals with a Cantonese Version of Mini Mental State Examination (CMMSE) score of 18 or below were excluded from the analysis.

Measurements

Diagnosis of anxiety disorders

Ascertainment of diagnosis of common mental disorders was based on the use of the revised Clinical Interview Schedule (CIS-R) (Lewis *et al.*, 1992). CIS-R consists of 14 sections of psychological symptoms: somatic symptoms, fatigue, concentration and forgetfulness, sleep problems, irritability, worry about physical health, depression, depressive ideas, worry, anxiety, phobia, panic, compulsions and obsessions. Each section starts with two screening questions to establish if a particular symptom is present in the past month. Following positive responses, follow up questions on the symptoms in past week are asked. Scores for each section range from 0-4, except for depressive ideas with a maximum score of 5, with higher score indicating a higher level of symptom severity. Symptom scores are calculated for each section. A total score is generated by summation of the section scores. This indicates the overall severity of the psychological

symptoms. The general cut-off point of threshold for adults indicating for significant psychological symptoms is 11/12, which could yield a sensitivity of 69.3% and a good specificity of 92.7%. The cut-off point of threshold indicating the need for clinical attention is 17/18, which could yield a sensitivity of 70.2% and a specificity of 95.2% (Lam *et al.*, 2015). The CIS-R generated diagnosis of common mental disorders, including depressive disorder (DEP); generalized anxiety disorders (GAD); mixed anxiety and depressive disorder (MADD); other anxiety disorders (OAD); and comorbid mood disorders; according to the Tenth Revision of the International Classification of Diseases of the World Health Organization (ICD-10) (Lewis *et al.*, 1992). The 1-week ICD-10 diagnoses of common mental disorders were generated from CIS-R data.

Basic demographic, psychosocial functioning, chronic medical conditions, and global cognitive functioning were also collected. Physical health burden was measured with the Cumulative Illness Rating Scale (CIRS) (Parmelee *et al.*, 1995). Impairment of each medical condition was rated on a 0 to 4 grading scale. The total score is the sum of all physical system scores. The short-Form 12 (SF-12) health survey was used to assess the quality of life related functioning of participants (Lam *et al.*, 2005). It generates two summary scores on physical and mental component functioning. The scores are on a scale from 0 to 100, with 100 representing the highest level of functioning. Global cognitive functioning was evaluated by the use of CMMSE (Chiu *et al.*, 1994). It is a 30-points scale that examines different cognitive domains, including orientation to time, orientation to place, registration, attention and calculation, delayed recall, language, ability to follow commands, and visuospatial perception. Basic socio-demographic data on age, gender, years of completed education, marital status and occupational status were also obtained.

Data Analysis

This is a post hoc analysis of HKMMS that evaluate the prevalence of common mental disorders in a representative sample aged 16 to 75 years old in Hong Kong. We extracted respondents aged 60 to 75 years old from the HKMMS to describe the prevalence of anxiety disorders in the Chinese older population in Hong Kong. Before we estimated the prevalence of late life anxiety disorders in the community living older adults in Hong Kong, we first compared the HKMMS data to the Hong Kong Census data 2011 to check for the similarity on the distribution of basic demographic characteristics. The post stratification weights constructed were then entered into the SPSS and applied when calculating weight prevalence of anxiety disorders. Weights were applied to adjust for differences in education and type of housing between the general population and the HKMMS sample, because their effect size is larger than 0.2.

Prevalence of all categories on common mental disorder was calculated following the CIS-R algorithms. The CIS-R algorithms allow categorization according to ICD-10 diagnostic criteria for depression, anxiety disorders (phobias, generalized anxiety disorder (GAD), panic disorder and obsessive-compulsive disorder) and mixed anxiety and depressive disorder (MADD). For the present study, we have grouped all anxiety disorders and MADD into one category as any anxiety disorders without depressive symptoms. Anxiety disorders co-occurred with depressive episode is defined as comorbid mood disorders.

Logistic regression was then used to explore any association between anxiety disorders and psychosocial factors by comparing respondents with anxiety disorders to a

mentally healthy group. Data analysis was performed using PASW 20.0 for Windows. Statistical significance was set as $p < 0.05$.

Results

Sample Characteristics

One thousand one hundred and fifty eight subjects aged 60 to 75 years were assessed. The demographics of the participants were compared with the Hong Kong population in 2011 for the same age range (HKC&SD, 2012). As shown in Table 1, the sample was comparable to the 2011 census data for Hong Kong on gender. The goodness-of-fit test indicated a small effect size for gender (Table 1). The mean age of the HKMMS participants was 65.9 (SD=4.4) years, and 52.8% were female. The mean years of education was 8.9 (SD=5.3) years. The sample had a CIRS scores of 3.8 (SD=2.2). The mean CIS-R score was 4.7 (SD=6.8).

Prevalence of Anxiety Disorders

After adjusted the age, gender, and type of housing, 138 participants (11.8%, 95% CI=10-13.7%) have a CIS-R score of 12 or above. According to the ICD-10 diagnosis generated by the CIS-R algorithms, the weighted prevalence of anxiety disorders occurring was 8% (95% CI=6.5 – 9.6%), followed by 2.1% (95% CI=1.3 – 3%) with comorbid mood disorders, and 1.7% (95% CI=1 – 2.5%) with depression. Table 2 depicted the unweighted and weighted prevalence of all categories on common mental disorders.

Symptoms Cluster of Anxiety Disorders

Symptom presentation of anxiety group is predominant of physical symptoms, including fatigue, forgetfulness and sleeping problems, following by psychological symptoms, such as irritability, worry about health, feeling anxious, and having depressive ideas. Table 3 displayed the proportion of symptoms endorsed by participants with and without anxiety disorders.

Correlates of Anxiety Disorders

Of the 101 participants with anxiety disorders, the mean age was 65.7 (SD=4.5) years, and 76.2% of the sample are female. The mean years of education was 8.0 (SD=5.4) years. The sample had a mean chronic illness burden score of 5.4 (SD=2.3). The mean CIS-R score was 17.2 (SD=4.4). Compared with those without mental disorders (N=1006), individuals with anxiety disorders are more likely to be women ($X^2=25.3$, $p<0.001$), had more chronic illness burden ($t=-9.3$, $p<0.001$), had lower scores in SF12 physical component summary ($t=9.2$, $p<0.001$), CMMSE delayed recall ($t=2.3$, $p=0.022$), and CMMSE total score ($t=2.3$, $p=0.023$). Age and years of completed education were not different from the group without depression or anxiety. Table 4 showed the socio-demographic information of participants with and without anxiety disorders.

Variables that differed significantly between groups were subjected to a logistic regression analysis to determine which factors were associated with anxiety in the elderly people. These variables included gender, CIRS score, and CMMSE score, but SF12 physical component summary (PCS) score. PCS was excluded from logistic regression analysis, because it is strongly influenced by the existing clinical conditions and is highly correlated with the CIRS score. As a result, it is more worthwhile to examine potential

association between the presence of medical comorbidity and anxiety disorders in older adults.

Two significant factors for anxiety disorders were identified by binary logistic regression, including being women (OR 2.9, 95% C.I. 1.2 – 6.7, $p < 0.05$) and high CIRS score (OR 1.4, 95% C.I. 1.3 – 1.6, $p < .001$). We also assessed the association of different physical illnesses against the presence of anxiety disorders. Anxiety disorders was found to be associated with cardiovascular disease (CVD) (OR 1.3, 95% C.I. 1.1– 1.6, $p = 0.01$), musculoskeletal diseases (OR 2.0, 95% C.I. 1.5– 2.7, $p < 0.001$), and genitourinary diseases (OR 1.8, 95% C.I. 1.3– 2.7, $p = 0.001$).

Discussion

The objective of the present thesis was to comprehensively evaluate the prevalence of late life anxiety disorders in the Chinese community population. It was based on a random sample of person aged 60 to 75. Since most previous studies on anxiety in old age have focused on individuals aged 55 or above and also included elderly people with depression, the symptomatology of anxiety disorders in the older population is not yet fully developed. Therefore, we have chosen an age cut-off at 60 years old, and specifically looked only at those with pure anxiety disorder and independent of depression.

Several major findings were obtained in this study. First, our results indicated that anxiety disorders are prevalent condition in the Chinese older community. The Hong Kong Mental Morbidity Survey (HKMMS) is the first population-based survey on common mental disorders conducted locally. The current findings gave an overview on

the pattern of anxiety disorders in the older Hong Kong population. Based on the use of ICD-10 diagnostic criteria generated by CIS-R, anxiety disorders are common in our community sample with a one week prevalence of 8%. Our finding on prevalence estimates was much higher than that reported in other studies of similar survey length ranging from 3.2 % in Sweden (Forsell & Winblad, 1997), 4.5% in Berlin(Schaub & Linden, 2000), and 5.5% in the United States (Regier et al, 1988).

Second, the study showed that the effect of gender was consistently found as with other studies, with higher risk of anxiety disorders in females than in males. Women are more likely to report anxiety symptoms to their health care professionals than men. For men, they would not seek help from doctors until the conditions get worse and have serious impact on their daily activities and health. Indeed, findings from a recent study have already found association of increased rate of mortality in older men with anxiety disorders (van Hout *et al.*, 2004). The risk of mortality in anxious older men is as high as 87% over a 7 years period. The excess subsequent mortality figure reflected that very few older men with anxiety disorders received treatment and led to worsening of existing physical illnesses, particularly cardiovascular disorders. It is important to explore on how to increase utilization of medical services in older men in the community, particularly those with comorbid affective disorders.

Third, elderly with anxiety disorders were more likely to have poor physical health. Yet, due to our cross-sectional nature, it was not sure whether these chronic conditions are a cause or a consequence of anxiety disorders in the elderly population. Still, a possible explanation could be that an increase in chronic diseases in anxious older individuals may exacerbate the deterioration in functioning as well as wellbeing. Actually,

the NCS-R study also reported similar results. They found that the presence of anxiety disorders is associated with having three or more chronic illnesses (Gum *et al.*, 2009). In addition, a Netherland study involved a sample of 659 participants aged 55 to 85 years old found that individuals with either anxiety symptoms or disorders reported more functional limitations, chronic physical diseases, and decreased well-being in terms of loneliness, satisfaction with life and self-perceived health (de Beurs *et al.*, 1999). In particular, we found association between cardiovascular, genitourinary and musculoskeletal diseases and increased risk of anxiety disorders. The relationship of anxiety disorders with certain cardiovascular conditions, such as hypertension and coronary heart disease, are already well documented (Kubzansky *et al.*, 1997). Anxiety disorders were also found to be a predictor of recurrent cardiac events in those who have suffered from myocardial infarction (Strik *et al.*, 2003; Huffman *et al.*, 2008).

Forth, it is of interests to note that the association between early cognitive impairment and anxiety disorders in older people without cognitive impairment. We found that individuals with anxiety disorders were poorer in global cognition and performed worse in CMMSE delayed recall when compared to the group without mood symptoms. In fact, there is some evidence that supports a linear relationship between increased level of anxiety and decreased cognitive performance in older adults. A clinical study compared cognitive performance among patients with GAD, patients with MADD and normal older individuals. They found that older adults with GAD performed worse on the California Verbal Learning Test (CVLT) than those with no psychiatric illness (Mantella *et al.*, 2007). The results on reduced episodic memory in anxious older adults

reflected the possibility of anxious mood affecting retrieval process and hence delayed recall.

To put this forward, the findings seemed to suggest that anxiety symptoms or disorders were an early sign of cognitive decline, especially memory decline, in the non-demented community older population. This interpretation has gained concrete supports from a new longitudinal study on the interplay among amyloid plaque, anxiety symptoms and cognitive decline. The study has found association between anxiety and cognitive decline in older adults with elevated levels of amyloid-beta. Furthermore, individuals with higher level of anxiety have greater rate of decline in global cognition, verbal memory, language and executive function than those with lower level of anxiety (Pietrzak *et al.*, 2015).

Strength and Limitations

The symptomatology of late life anxiety disorders has not been clearly defined. We have taken the chance to group individuals with different types of anxiety disorders, including generalized anxiety disorders, phobias, panic disorders and mixed anxiety and depressive disorders, into one category. This broad definition of anxiety can allow us to consider both clinical and subclinical level of anxiety disorders. At the same time, the definition can offer promise of obtaining a reliable estimate of anxiety disorders in old age for mental health services planning.

A number of limitations were noted. We measured anxiety symptoms with CIS-R in this study, but it is not originally designed specifically for assessing anxiety symptoms in the older population. It is possible to have missed some elderly individuals who

exhibited anxiety in a way different from the diagnostic criteria developed for the younger population. However, CIS-R enabled us to transform self-report symptoms directly into a classification of common mental disorders on the basis of the disease algorithms. As therefore, it has enabled us to estimate the proportion of anxiety symptomatology in a large community sample. Second, this study is cross-sectional in nature, direction of associations with the correlates of anxiety disorders could not be determined. Longitudinal studies in the future are needed to get more inference on causation, especially on the relationship between prolonged anxiety disorders and subsequent development of cognitive impairment. Third, the three items delayed recall in CMMSE used in the survey is only a screening tool for assessing the memory of our sample. In order to better understand how anxiety disorders affect cognition, more comprehensive cognitive assessments, should be adopted to investigate the impact of anxiety disorders on different cognitive domains and memory process in future studies.

Conclusion

Clearly, our study has demonstrated that anxiety disorders are prevalent among the community dwelling older adults in Hong Kong. The next important question is how many of those left untreated in the community? As pointed out in previous literatures, anxiety disorders are a chronic mental illness that can last for many years. Elderly people lived with anxiety disorders frequently found to have greater medical comorbidity that may affect their quality of life and independency. Our findings also suggested a potential risk for early sign of memory impairment in clinically non-demented individuals with anxiety disorders. It is imperative to increase elderly's awareness of early identification

of anxiety disorders and its potential consequences for future physical and cognitive impairment.

Conflict of interest

None

Description of authors' roles

Ada Fung analyzed the data and wrote the paper. Corine Wong collected the data and supervised the data collection. All other authors designed the study and supervised the study progress.

Acknowledgement

The Hong Kong Mental Morbidity Survey is a commissioned project supported by Health and Health Services Research Fund (Ref: 09101601), Food and Health Bureau, Hong Kong SAR Government.

Reference

Aziz, R. and Steffens, D.C. (2013). What are the causes of late-life depression? The *Psychiatric Clinics of North American*, 36, 497-516. doi: 10.1016/j.psc.2013.08.001

Bryant, C., Jackson, H. and Ames, D. (2008). The prevalence of anxiety in older adults: methodological issues and a review of the literature. *Journal of Affective Disorders*, 109, 233–250. doi:10.1016/j.jad.2007.11.008

Chiu, H.F., Lee, H.C., Chung, W.S. and Kwong, P.K. (1994). Reliability and validity of the Cantonese version of the Mini-Mental State Examination — a preliminary study. *Journal of Hong Kong College Psychiatry*, 4, 25-28. doi:10.1017/S1041610209008588.

de Beurs, E., Beekman, A.T., van Balkom, A.J., Deeg, D.J., van Dyck, R. and van Tilburg W. (1999). Consequences of anxiety in older persons: its effect on disability, well-being and use of health services. *Psychological Medicine*, 29(3), 583-593. doi: 10.1017/S0033291799008351

Diagnostic and statistical manual of mental disorders, 4th ed, text rev. (2000). Washington, DC: American Psychiatric Association.

Forsell, Y. and Winblad, B. (1997). Anxiety disorders in non-demented and demented elderly patients: prevalence and correlates. *Journal of Neurology*, 62, 294–295. doi:10.1016/j.cpr.2008.02.008

Gum, A.M., King-Kallimanis, B. and Kohn, R. (2009). Prevalence of mood, anxiety, and substance—abuse disorders for older Americans in the National Comorbidity Survey Replication. *The American Journal of Geriatric Psychiatry*, 17, 769–781. doi: 10.1097/JGP.0b013e3181ad4f5a

Himmelfarb, S. and Murrell, S.A. (1984). The prevalence and correlates of anxiety symptoms in older adults. *The Journal of Psychology*, 116, 159-167. doi: 10.1097/JGP.0b013e318202bc62.

Hong Kong Census and Statistic Department. (2012). Demographic trends in Hong Kong 1981-2011.

Huffman, J.C., Smith, F.A., Blais, M.A., Januzzi, J.L. and Fricchione G.L. (2008). Anxiety, independent of depressive symptoms, is associated with in hospital cardiac complications after acute myocardial infarction. *Journal of Psychosomatic Research*, 65, 557–563. doi: 10.1016/j.jpsychores.2008.08.001

Kessler, R.C., Berglund, P., Demler, O., Jin, R., Merikangas, K.R. and Walters, E.E. (2005). Lifetime prevalence and age-of-onset distributions of DSM-IV disorders in the national comorbidity survey replication. *Archives of General Psychiatry*, 62, 593–602. doi:10.1001/archpsyc.62.6.593.

Kubzansky, L.D., Kawachi, I., Spiro III, A., Weiss, S.T., Vokonas, P.S. and Sparrow, D. (1997). Is Worrying Bad for Your Heart? A Prospective Study of Worry and Coronary Heart Disease in the Normative Aging Study. *Circulation*, 95, 818-824. doi: 10.1161/01

Lam, C.L., Tse, E.Y. and Gandek, B. (2005). Is the standard SF-12 health survey valid and equivalent for a Chinese population? *Quality of Life Research*, 14, 539-547. doi: <http://dx.doi.org/10.1007/s11136-004-0704-3>

Lam, L.C., et al. (2015). Prevalence, psychosocial correlates and service utilization of depressive and anxiety disorders in Hong Kong: the Hong Kong Mental Morbidity Survey (HKMMS). *Social psychiatry and psychiatric epidemiology*, 50(9), 1379-1388. doi: 10.1007/s00127/s00127-015-1014-5. Epub 2015 Feb 8.

Lewis, G., Pelosi, A. J., Araya, R. and Dunn, G. (1992). Measuring psychiatric disorder in the community: a standardized assessment for use by lay interviewers. *Psychological Medicine*, 22, 465–486. doi:10.1016/j.jad.2015.06.001

Mantella, R.C., et al. (2007). Cognitive impairment in late-life generalized anxiety disorder. *Am J Geriatr Psychiatry*, 15(8), 673-679.

Murphy, L.B., Sacks, J.J., Brady, T.J., Hootman, J.M. and Chapman, D.P. (2012). Anxiety is more common than depression among US adults with arthritis. *Arthritis Care and Research*, 64(7), 968–976. doi: 10.1002/acr.21685.

Parmelee, P.A., Thuras, P.D., Katz, I.R. and Lawton, M.P. (1995). Validation of the Cumulative Illness Rating Scale in a geriatric residential population. *Journal of the American Geriatrics Society*, 43(2), 130-137. doi: 10.1111/j.1532-5415.1995.tb06377.x.

Pietrzak, R.H., et al. (2015). Amyloid- β , Anxiety, and Cognitive Decline in Preclinical Alzheimer Disease A Multicenter, Prospective Cohort Study. *JAMA Psychiatry*, Published online. doi:10.1001/jamapsychiatry.2014.2476.

Regier, D.A., et al. (1988). One-month prevalence of mental disorders in the United States: based on five epidemiologic catchment area sites. *Archives of General Psychiatry*, 45, 977–986. doi: 10.1111/j.1600-0447.1993.tb03411.x

Schaub, R.T. and Linden, M. (2000). Anxiety and anxiety disorders in the old and very old—results from the Berlin Aging Study (BASE). *Comprehensive Psychiatry*, 41, 48–54. doi: 10.1016/S0010-440X(00)80008-5.

Strik, J.L.M.H., Denollet, J., Lousberg, R. and Honig, A. (2003). Comparing symptoms of depression and anxiety as predictors of cardiac events and increased health care consumption after myocardial infarction. *Journal of American College of Cardiology*, 42, 1801–1807. doi:10.1016/j.jacc.2003.07.007

van Hout, H.P., et al. (2004). Anxiety and the risk of death in older men and women. *The British Journal of Psychiatry*, 185, 399–404. doi: 10.1192/bjp.185.5.399

Table 1 Comparisons on gender, education level, and type of housing of the participants interviewed (N=1,158) and corresponding population (aged 60-75) in Hong Kong.

		HKMMS (60-75 years old)		2011 HK Census (60-75 years old)	Sig.	Effect size
		N	%	%		
Gender	Males	546	47.2%	50.5%	<0.05	0.07
	Females	612	52.8%	49.5%		
Education	None	81	7.0%	13.3%	<0.001	0.26
	Primary	388	33.5%	39%		
	Secondary	522	45.1%	37.2%		
	Tertiary	167	14.4%	10.4%		
Housing	Public	552	47.7%	35.8%	<0.001	0.25
	Private	606	52.3%	64.2%		

*HKMMS = Hong Kong Mental Morbidity Survey

Table 2 Unweighted and weighted prevalence of late life common mental disorders by CIS-R score and by ICD-10 diagnosis in Study I.

	Unweighted (N=1156*)		Weighted (N=1156*)	
	N	%	N	%
CIS-R score				
CISR = 0-5	830	71.7	853	73.7
CISR = 6-11	176	15.2	165	14.3
CISR \geq 12	150	13	137	11.9
ICD-10 Diagnosis				
Anxiety Disorders	101	8.7	93	8
Depression	23	2	20	1.7
Comorbid mood disorders	26	2.2	25	2.1

*2 cases with missing data were excluded from analysis

Table 3 Symptom endorsement of participants with and without anxiety disorders.

CISR Symptoms	ICD-10 Diagnosis	
	None (N=1,006)	Anxiety (N=101)
Somatic (Y, %)	3.98	49.50
Fatigue (Y, %)	24.35	87.13
Forgetfulness (Y, %)	21.37	84.16
Sleeping problems (Y, %)	33.40	91.09
Irritability (Y, %)	12.92	66.34
Worry about health (Y, %)	11.13	67.33
Depression (Y, %)	3.18	49.50
Depressive Idea (Y, %)	7.36	64.36
Worry (Y, %)	12.33	54.46
Anxiety (Y, %)	7.75	67.33
Phobia (Y, %)	1.69	14.85
Panic (Y, %)	0.10	4.95
Compulsions (Y, %)	0.89	6.93
Obsessions (Y, %)	1.79	16.83

Table 4 Sociodemographic information of participants with and without anxiety disorders.

	None (N=1006)		Anxiety (N=101)		t-test
	Mean	SD	Mean	SD	p-value
Age	65.9	4.4	65.7	4.5	0.591
Edu years	9.0	5.3	8.0	5.4	0.061
Female (%)	50	-	76.2	-	<0.001
CIRS	3.5	2.0	5.4	2.3	<0.001
SF12 (PCS)	49.3	7.6	41.7	9.8	<0.001
SF12 (MCS)	54.8	5.4	45.2	7.4	<0.001
CIS-R	2.5	2.9	17.2	4.4	<0.001
CMMSE	27.8	2.1	27.2	2.5	0.023
CMMSE delayed recall	2.5	0.7	2.3	0.8	0.022