

Staging fibrosis and excluding advanced fibrosis in patients with NAFLD: comparison of non-invasive markers in an interim analysis from a prospective multicenter study

Peter J. Eddowes¹, Quentin Anstee², Indra Neil Guha³, David A. Sheridan⁴, Emmanouil Tsochatzis⁵, Jeremy Cobbold⁶, Michael E. Allison⁷, Valerie Paradis⁸, Pierre Bedossa⁸, Philip N. Newsome¹

1. NIHR Liver Biomedical Research Unit and Centre for Liver Research, University of Birmingham, Birmingham, United Kingdom

2. Institute of cellular medicine - Faculty of Medical Sciences, Newcastle University, Newcastle upon Tyne, United Kingdom

3. NIHR Nottingham Digestive Diseases Biomedical Research Unit, NHS Trust and University of Nottingham, Nottingham, United Kingdom

4. Institute of Translational and Stratified Medicine, Plymouth University, Plymouth, United Kingdom

5. UCL Institute for Liver and Digestive Health, Royal Free Hospital, London, United Kingdom

6. Department of Gastroenterology, Oxford University Hospitals NHS Trust, John Radcliffe Hospital, Oxford, United Kingdom

7. Department of Hepatology, Addenbrookes Hospital, Cambridge, United Kingdom;

8. Pathology department, Hôpital Beaujon, APHP, Clichy, France

Background & Aims: Hepatic fibrosis is a major determinant of clinical outcomes in non-alcoholic fatty liver disease (NAFLD) and there remains a clear need to establish the accuracy of non-invasive markers of fibrosis. This study aims to prospectively compare the diagnostic performance and ability to exclude advanced fibrosis of the

following non-invasive tests in NAFLD: FibroScan, FibroMeter V, FibroMeter NAFLD, FibroMeter VCTE, NAFLD Fibrosis score (NFS), Fib4, APRI, BARD and AST/ALT ratio.

Methods: Patients with suspected NAFLD prospectively underwent FibroScan examination and blood sampling within 2 weeks of a standard of care liver biopsy (LB) between March 2014 and January 2016 at seven UK centres. LB were staged in a blinded manner by two expert pathologists according to the NASH CRN system. Diagnostic performance was assessed in terms of area under the ROC curves (AUC). Ability to exclude advanced fibrosis was assessed using published cut-offs except for FibroMeter (FM), for which cut-offs have not yet been published. Cut-offs for FM were determined that maximized the Youden index.

Results: 155 patients (57% male, median age 54 [IQR 20] years, median BMI 33.2 [8.1] kg/m²) had a complete dataset for analysis. Fibrosis distribution was: F0: 23%, F1: 25%, F2: 21%, F3: 25%, F4: 6%. 43% of the patients had a NAS score \geq 5. Performance summary of the tests is presented below in the table.

Conclusion: FibroMeter VCTE, which combines biochemical parameters with liver stiffness measured by FibroScan, has the highest performance characteristics with positive and negative predictive values of 67 and 93% respectively at confirming or excluding \geq F3 fibrosis.