

Daniel Barrera Salazar · Lucio Guerberoff 

On critical values of L -functions of potentially automorphic motives

Received: 10 May 2016 / Accepted: 28 October 2016

Abstract. In this paper we prove a version of Deligne's conjecture for potentially automorphic motives, twisted by certain algebraic Hecke characters. The Hecke characters are chosen in such a way that we can use automorphic methods in the context of totally definite unitary groups.

1. Introduction

The goal of this paper is to prove results on Deligne's conjecture for potentially automorphic motives, twisted by certain algebraic Hecke characters. Let K be a totally real number field and L/K be a CM extension. We refer the reader to Sect. 2 for an overview of motives and realizations. For a realization M over K and an algebraic Hecke character χ of L , we let $M(\chi)$ denote the tensor product $M \otimes \text{Res}_{L/K}[\chi]$, where $[\chi]$ is the CM motive over L attached to χ . A realization M is automorphic of unitary type if it looks like the conjectural motive attached to a self-dual, cohomological, cuspidal automorphic representation Π of $\text{GL}_n(\mathbb{A}_K)$; for more details, see Sect. 4.1. A realization M is potentially automorphic of unitary type if there exists a finite, Galois, totally real extension K'/K such that $M_{K'}$ is automorphic. Our main result (see Theorem 4.4.1) is the following. In the statement, \tilde{L} is a certain finite Galois extension of L , which equals the Galois closure of L if M is automorphic. We refer the reader to the main text for more details and for any unexplained notation.

Theorem. *Let M be a potentially automorphic realization of unitary type over K , with coefficients in E , of rank n , satisfying condition (3) in Theorem 4.4.1. Let ψ be a critical algebraic Hecke character of L of infinity type $(m_\tau)_{\tau \in \text{Hom}(L, \mathbb{C})}$ and weight w . Assume that:*

D. Barrera Salazar: Departament de Matemàtiques, Universitat Politècnica de Catalunya, Planta 3, 330. Campus Nord, Edificio Omega, c/Jordi Girona, 1-3, 08034 Barcelona, Spain. e-mail: danielbarreras@hotmail.com

L. Guerberoff (✉): Department of Mathematics, University College London, 25 Gordon Street, London WC1H 0AY, UK. e-mail: l.guerberoff@ucl.ac.uk

Mathematics Subject Classification: 11F67 (Primary) · 11S40 · 11F66 · 11G40 · 11G18 · 11R39 (Secondary)

- (1) either n is even, or n is odd and Hypothesis 4.3.1 is satisfied for M , and
 (2) $|m_\tau - m_{\bar{\tau}}| > \max\{n - p_n(\sigma, 1)\}_{\sigma \in \text{Hom}(K, \mathbb{C})}$ for any $\tau \in \text{Hom}(L, \mathbb{C})$.

Then for all critical integers $k > w + n$ of $M(\chi)$, we have

$$\frac{L(M(\chi), k)_1}{c^+(M(\chi)(k))_1} \in (E\mathbb{Q}(\psi)\tilde{L})^\times.$$

Here $p_1(\sigma, 1) > \dots > p_n(\sigma, 1)$ are the Hodge numbers of M (see Sect. 2.2) and χ is an algebraic Hecke character constructed from ψ as follows: there exists a finite order character ψ_0 of $\mathbb{A}_K^\times/K^\times$ such that $\psi|_{\mathbb{A}_K^\times} = \psi_0 \|\cdot\|^{-w}$, and we put $\chi = \psi^2(\psi_0 \circ N_{L/K})^{-1}$. We stress that, given M , there always exists algebraic Hecke characters and integers $k > w + n$ critical for $M(\chi)$ as in the statement.

The proof of our theorem works as follows. First, suppose that M is automorphic. Hypothesis (2) in the theorem allows us, using the results proved by one of the authors [14] generalizing earlier results of Harris [17] to totally real fields, to write the value of the L -function of $M(\chi)$ at k in terms of a CM period attached to χ . The way this works is by using descent to a totally definite unitary group G and expressing this L -function as that of a cohomological automorphic representation on G . The corresponding critical value is expressed in terms of a CM period and a Petersson norm, which in our case turns out to be algebraic. On the other hand, the motivic computations of [14] and Hypothesis (2) allow us to express the Deligne period of $M(\chi)(k)$ in terms of a CM period, which turns out to match the previous one. Thus, we prove in this way that the value $L(M(\chi), k)_1$ at any critical integer k of $M(\chi)$ which satisfies $k > w + n$ equals the Deligne period $c^+(M(\chi)(k))_1$, up to multiplication by an element of $E\mathbb{Q}(\psi)\tilde{L}$. The requirement that $k > w + n$ is necessary for the automorphic methods to work. In the same vein, automorphic methods only allow us to consider multiples by $E\mathbb{Q}(\psi)\tilde{L}$. This consideration is included in Conjecture 2.2.1, which is a weak form of Deligne's conjecture.

When M is a potentially automorphic realization over K , we use Brauer's induction and solvable base change for GL_n , as developed in the theory of Arthur–Clozel [1]. We prove the theorem for M by using the previous automorphic case and further compatibilities between the CM periods that appear, which are a consequence of Deligne's conjecture for algebraic Hecke characters, proved in this case by Blasius [4].

Let us say a few words about the hypotheses of the theorem. As we mentioned before, a potentially automorphic realization of unitary type M is one such that it becomes automorphic of unitary type after extension of scalars to a totally real, Galois extension K'/K . A number of techniques is available to prove that certain motives, or Galois representations, are potentially automorphic. We refer the reader to [2] for some general results in this direction. From now on, for simplicity, assume that M is already automorphic. In the statement of the theorem, we require ψ to be critical. This means that $m_\tau \neq m_{\bar{\tau}}$ for any $\tau \in \text{Hom}(L, \mathbb{C})$. This is needed for the motivic computations in [14]. Regarding Hypothesis (1) in the theorem, it is needed to express the full period $\delta(M)$ in a convenient way. This period appears in the motivic computations that give rise to the expression of $c^+(M(\chi)(k))$. Automorphic realizations of unitary type are endowed with a polarization, which can be used to

show that $\delta(M)$ can be replaced by $(2\pi i)^{-[K:\mathbb{Q}]n(n-1)/2}$ when n is even. Hypothesis (1) says that when n is odd, we can also replace $\delta(M)$ by the same power of $2\pi i$. This can be shown to be true if one assumes the much stronger Tate conjecture. Hypothesis (2), as we explained above, is made so that $M(\chi)$ has critical values and so that they can be expressed essentially as a CM period (in the terminology of [17], χ belongs to the n -th critical interval of M). It is essential to the method, and a modification of it, still assuming that $M(\chi)$ has critical values, would entail the appearance of additional quadratic periods in the formulas, which are harder to relate to critical values of L -functions. Hypothesis (3) says that Π has a descent to a totally definite unitary group satisfying a number of conditions. This is expected to hold in our setting, and many, if not most, cases have been already proved [22–24]. Combined with Hypothesis (2), this implies that we can express the L -function as that of a cohomological automorphic representation on a totally definite unitary group.

While doing the motivic computations behind the expression of $c^+(M(\chi)(k))$, we draw some consequences of the general formula proved in [14] for the case of arbitrary critical intervals. More precisely, if M is a regular, polarized realization, then after fixing embeddings of the coefficient fields into \mathbb{C} , we construct certain algebraic Hecke characters χ with the property that $M(\chi)$ has critical values which can be expressed in terms of CM periods and additional quadratic periods $Q_j(M)$ attached to M . The set of quadratic periods appearing in the expression depends on the critical interval of χ , which can be arbitrarily prescribed. Combining the formulas for different characters χ gives an expression of the quadratic periods in terms of quotients of Deligne periods of various twists $M(\chi)$ and CM periods (see Proposition 2.4.1 and Corollary 2.4.1). Assuming Deligne’s conjecture for these motives, we can then express the quadratic periods in terms of critical values of L -functions and certain Gauss sums (Proposition 5.1.1). These expressions should be helpful in certain applications, and we plan to exploit this in a future work in relation to p -adic interpolation and p -adic L -functions.

To finish this introduction we would like to say a few words about the background and motivation for this work. The study of values of L -functions at integers is a subject with a long history starting with Euler. Based on experimental results, in 1979 Deligne proposed a general conjecture relating the values of motivic L -functions at certain integer points to periods of integrals. Arguably the most useful way to approach this and other conjectures on critical values of L -functions is by using automorphic methods. In this direction, let us mention Blasius’s result [4], which proves the conjecture for the motives attached to algebraic Hecke characters of CM fields (see also [15]). Let us also mention Shimura’s works on critical values of L -functions in the case of modular forms and Hilbert modular forms (see for instance [28] and [29]), based on the classical Rankin–Selberg theory of L -functions and an analysis of the arithmetic properties of ratios of Petersson inner products. Following these works, Harris in [17] treated the case of the polarized regular motives over \mathbb{Q} coming from automorphic representations of $\mathrm{GL}_n(\mathbb{A}_{\mathbb{Q}})$. One of the main ingredients in this work is the doubling method. Finally, in [14], the author generalized Harris’s results over arbitrary totally real number fields. The present work is based on [14].

1.1. Organization of the paper

Here is the outline of the paper. Section 2 is devoted to the motivic considerations of this work. The main result of this section is a formula for the Deligne period of $M(\chi)(k)$ in terms of CM periods and powers of $2\pi i$, based on [14] (Proposition 2.5.1). As mentioned above, with an eye towards future applications, we also include a formula for the quadratic periods of a regular, polarized realization in terms of quotients of various twists of M and CM periods (Proposition 2.4.1 and its corollary). In Sect. 3 we specialize to the case of totally definite unitary groups the results of [14] on critical values of automorphic L -functions (Proposition 3.3.1). In Sect. 4 we combine the previous sections and prove the main result of this paper (Theorems 4.3.1 and 4.4.1 for automorphic and potentially automorphic realizations respectively). Finally, in Sect. 5, which is more speculative, we write down an expression of the quadratic periods in terms of critical values of L -functions (Proposition 5.1.1), which follows from assuming Deligne's conjecture for the corresponding motives.

Notation and conventions. We fix an algebraic closure \mathbb{C} of \mathbb{R} , a choice of $i = \sqrt{-1}$, and we let $\bar{\mathbb{Q}}$ denote the algebraic closure of \mathbb{Q} in \mathbb{C} . We let $c \in \text{Gal}(\mathbb{C}/\mathbb{R})$ denote complex conjugation on \mathbb{C} , and we use the same letter to denote its restriction to $\bar{\mathbb{Q}}$. We write $c(z) = \bar{z}$ for $z \in \mathbb{C}$.

For a number field K , we let \mathbb{A}_K and $\mathbb{A}_{K,f}$ denote the rings of adèles and finite adèles of K respectively. When $K = \mathbb{Q}$, we write $\mathbb{A} = \mathbb{A}_{\mathbb{Q}}$ and $\mathbb{A}_f = \mathbb{A}_{\mathbb{Q},f}$. After fixing an algebraic closure \bar{K} of K , we denote $\Gamma_K = \text{Gal}(\bar{K}/K)$.

A CM field L is a totally imaginary quadratic extension of a totally real field K . A CM type Φ for L/K is a subset $\Phi \subset \text{Hom}(L, \mathbb{C})$ such that $\text{Hom}(L, \mathbb{C}) = \Phi \amalg c\Phi$ (equivalently, a choice of one of the two possible extensions to L of each embedding of K in \mathbb{C}).

All vector spaces over fields will be finite-dimensional except otherwise stated. A tensor product without a subscript between \mathbb{Q} -vector spaces will always mean tensor product over \mathbb{Q} . For any number field K , we denote by $J_K = \text{Hom}(K, \mathbb{C})$. For $\sigma \in J_K$, we let $\bar{\sigma} = c\sigma$. Let E and K be number fields, and $\sigma \in J_K$. If $\alpha, \beta \in E \otimes \mathbb{C}$, we write $\alpha \sim_{E \otimes K, \sigma} \beta$ if either $\beta = 0$ or if $\beta \in (E \otimes \mathbb{C})^\times$ and $\alpha/\beta \in (E \otimes \sigma(K))^\times$. There is a natural isomorphism $E \otimes \mathbb{C} \simeq \prod_{\varphi \in J_E} \mathbb{C}$ given by $e \otimes z \mapsto (\varphi(e)z)_\varphi$ for $e \in E$ and $z \in \mathbb{C}$. Under this identification, we denote an element $\alpha \in E \otimes \mathbb{C}$ by $(\alpha_\varphi)_{\varphi \in J_E}$. When K is given from the context as a subfield of \mathbb{C} , we write $\sim_{E \otimes K}$ for $\sim_{E \otimes K, 1}$, where $1 : K \hookrightarrow \mathbb{C}$ is the given embedding.

Suppose that $\mathbf{r} = (r_\varphi)_{\varphi \in J_E}$ is a tuple of nonnegative integers. Given Q_1, \dots, Q_n in $E \otimes \mathbb{C}$ (with $n \geq r_\varphi$ for all φ), we denote by

$$\prod_{j=1}^{\mathbf{r}} Q_j \in E \otimes \mathbb{C}$$

the element whose φ -th coordinate is $\prod_{j=1}^{r_\varphi} Q_{j,\varphi}$

2. Motives and periods

2.1. Algebraic Hecke characters

Let L be a number field. For a place v of L , we denote by L_v the corresponding completion, and by $(L_v^\times)^+$ the connected component of the identity in L_v^\times . An algebraic Hecke character of L is a continuous character $\chi : L^\times \backslash \mathbb{A}_L^\times \rightarrow \mathbb{C}^\times$, with the property that for each embedding $\tau \in J_L$, there exists an integer n_τ such that if v is the archimedean place of L induced by τ , then for every $x \in (L_v^\times)^+$,

$$\chi(x) = \begin{cases} \tau(x)^{-n_\tau} & \text{if } v \text{ is real,} \\ \tau(x)^{-n_\tau} \bar{\tau}(x)^{-n_{\bar{\tau}}} & \text{if } v \text{ is complex.} \end{cases}$$

The integer $n_\tau + n_{\bar{\tau}}$ is independent of τ , and it's called the weight $w(\chi)$ of χ . The tuple $(n_\tau)_{\tau \in J_L}$ is called the infinity type of χ . Let $\mathbb{Q}(\chi) \subset \mathbb{C}$ be the field generated by the values of χ on the finite idèles $\mathbb{A}_{L,f}^\times$. Then $\mathbb{Q}(\chi)$ is either \mathbb{Q} or a CM field.

From now on, assume that L is a CM field, which is the case that we will be interested in. Let K denote the maximal totally real subfield of L . The restriction of χ to K must necessarily be of the form

$$\chi|_{\mathbb{A}_K^\times} = \chi_0 \| \cdot \|^{-w(\chi)},$$

where $\| \cdot \|$ is the idèlic norm on \mathbb{A}_K^\times , and χ_0 is of finite order (see [27], Chapter 0).

Let Φ be a CM type for L/K , that is, $\Phi \subset J_L$ consists of a choice of one of the two possible extensions of each $\sigma \in J_K$ to L .

Proposition 2.1.1. *Let $(a_\tau)_{\tau \in \Phi}$ be a tuple of integers such that*

$$a_\tau \equiv a_{\tau'}(2)$$

for every $\tau, \tau' \in \Phi$. Then there exists an algebraic Hecke character χ of L of infinity type $(n_\tau)_{\tau \in J_L}$ such that

$$n_\tau - n_{\bar{\tau}} = a_\tau \quad (\tau \in \Phi).$$

Moreover, if $w_0 \in \mathbb{Z}$ satisfies that $w_0 \equiv a_\tau(2)$ for one (or every) $\tau \in J_L$, then χ can be taken to have weight w_0 .

Proof. As in [27], Chaper 0, any tuple of integers $(n_\tau)_{\tau \in J_L}$ with the property that $n_\tau + n_{\bar{\tau}}$ is independent of τ is the infinity type of an algebraic Hecke character χ of L . To arrive at the conditions of the proposition, choose an arbitrary w_0 , with the same parity as the a_τ , and take the tuple $(n_\tau)_{\tau \in \Phi}$ as

$$\begin{aligned} n_\tau &= \frac{w_0 + a_\tau}{2}, & \tau \in \Phi, \\ n_\tau &= \frac{w_0 - a_{\bar{\tau}}}{2}, & \tau \notin \Phi. \end{aligned}$$

□

2.2. Polarized regular motives

In this subsection, we recall the main result of [14], Sect. 2. Let K be a totally real number field, L/K a CM extension and E any number field. We are fixing throughout an algebraic closure \bar{K} of K , and we take L inside \bar{K} . Let M be a realization over K of rank n with coefficients in E , pure of weight $w(M)$. We refer to [14] for a basic overview of realizations. Here we stress that M consists of a collection of vector spaces and comparison isomorphisms, the interesting cases being given by collections of realizations coming from motives for absolute Hodge cycles over K , as in Deligne's article [12]. For each $\sigma \in J_K$, we can define the period $\delta_\sigma(M)$. These are elements of $(E \otimes \mathbb{C})^\times$, well defined up to multiplication by an element of $(E \otimes \sigma(K))^\times$. We let $\delta(M) = \delta_1(\text{Res}_{K/\mathbb{Q}} M)$, with $1 \in J_{\mathbb{Q}}$ being the unique embedding of \mathbb{Q} .

We say that M is *special* if, for every $\sigma \in J_K$, the action of the Frobenius automorphism F_σ on the Hodge component $M_\sigma^{w(M)/2, w(M)/2}$ is given by a scalar $\varepsilon = \pm 1$ (independent of σ). Here the E -vector space M_σ is the Betti realization of M attached to the embedding σ . Under the condition that M is special, we can define the Deligne σ -periods $c_\sigma^\pm(M)$, again elements of $(E \otimes \mathbb{C})^\times$ well defined modulo $(E \otimes \sigma(K))^\times$. We let $c^\pm(M) = c^\pm(\text{Res}_{K/\mathbb{Q}} M)$. We define $n^\pm = \dim_E M_\sigma^\pm$, where

$$M_\sigma^\pm = \{x \in M_\sigma : F_\sigma(x) = \pm x\}.$$

This is independent of σ . The following factorization formula is proved in [32] or [25] (we also include a similar formula for the δ 's):

$$c^\pm(M) \sim_E D_K^{n^\pm/2} \prod_\sigma c_\sigma^\pm(M), \quad (2.2.1)$$

$$\delta(M) \sim_E D_K^{n/2} \prod_\sigma \delta_\sigma(M). \quad (2.2.2)$$

Here D_K is the discriminant of K . Note that D_K is a positive integer such that $D_K^{1/2} \in K^{\text{Gal}}$, where $K^{\text{Gal}} \subset \bar{\mathbb{Q}}$ is the Galois closure of K in $\bar{\mathbb{Q}}$.

Under the assumption that the system of λ -adic representations $(M_\lambda)_\lambda$ is strictly compatible, we can define the L -function of M , $L^*(M, s) = (L(M, s)_\varphi)_{\varphi \in J_E} \in E \otimes \mathbb{C}$ (see [12]). We will always assume this to be the case. We also refer the reader to *op. cit.* for the definition of critical integers. Deligne's conjecture is the statement saying that if M is critical, meaning that 0 is a critical integer, then

$$\frac{L^*(M, 0)}{c^+(M)} \in E.$$

We will later prove, for certain realizations M , a weaker version of this conjecture which we state as follows. Note that if k is a critical integer for M , then the k -th Tate twist $M(k)$ is critical, and $L^*(M(k), 0) = L^*(M, k)$. Also, $c^+(M(k)) \sim (2\pi i)^{[K:\mathbb{Q}]n^\pm k} c^\pm(M)$, where $\pm = (-1)^k$.

Conjecture 2.2.1. *Let $F \subset \mathbb{C}$ be a number field and $\varphi \in J_E$. Let k be a critical integer of M . Then*

$$\frac{L(M, k)_\varphi}{c^+(M(k))_\varphi} \in F\varphi(E).$$

We will usually refer to the above as Conjecture 2.2.1 for M and the critical integer k , over F (for the embedding φ).

We say that the realization M is *regular* if, for every $\sigma \in J_K$ and $\varphi \in J_E$, the spaces $M_\sigma^{p,q}(\varphi)$ have dimension at most 1 over \mathbb{C} . Associated to the pair (σ, φ) , there is a sequence of integers

$$p_1(\sigma, \varphi) > \cdots > p_n(\sigma, \varphi)$$

with the property that $M_\sigma^{p,q}(\varphi) \neq 0$ if and only if $p = p_i(\sigma, \varphi)$ for some $i = 1, \dots, n$. We let $q_i(\sigma, \varphi) = w(M) - p_i(\sigma, \varphi)$, which in fact equals $p_{n+1-i}(\sigma, \varphi)$. We let $p_0(\sigma, \varphi) = +\infty$ and $p_{n+1}(\sigma, \varphi) = -\infty$. Note that if $n = 2k - 1$ is odd, then $p_k(\sigma, \varphi) = p_{n+1-k}(\sigma, \varphi)$, which implies that $w(M)$ is even. In particular, $w(M)n$ is even in all cases.

Let $\chi : L^\times \backslash \mathbb{A}_L^\times \rightarrow \mathbb{C}^\times$ be an algebraic Hecke character of L of infinity type $(n_\tau)_{\tau \in J_L}$. Attached to χ is a CM motive over L with coefficients in $\mathbb{Q}(\chi)$, which we denote by $[\chi]$ (this is denoted by $M(\chi)$ in [14]). See [27] for the construction of $[\chi]$. We let $\text{Res}_{L/K}[\chi]$ be the motive over K obtained by restriction of scalars from L to K of $[\chi]$. We say that χ is *critical* if $n_\tau \neq n_{\bar{\tau}}$ for every $\tau \in J_L$. In this case, $\text{Res}_{L/K}[\chi]$, a realization of rank 2, is regular in the sense defined above, and we denote the corresponding Hodge numbers by

$$p_1^\chi(\sigma, \rho) > p_2^\chi(\sigma, \rho),$$

where $\rho \in J_{\mathbb{Q}(\chi)}$. If we let $n(\tau, \rho) = n_{\tilde{\rho}^{-1}\tau}$, where $\tilde{\rho} : \mathbb{C} \rightarrow \mathbb{C}$ is an extension of ρ to \mathbb{C} , then

$$\{p_1^\chi(\sigma, \rho), p_2^\chi(\sigma, \rho)\} = \{n(\tau, \rho), n(\bar{\tau}, \rho)\},$$

where τ and $\bar{\tau}$ are the two embeddings of L extending σ . For χ critical, we define

$$t_{\sigma, \rho}(\chi) = p_1^\chi(\sigma, \rho) - p_2^\chi(\sigma, \rho).$$

We let $c_\sigma^\pm(\chi) = c_\sigma^\pm(\text{Res}_{L/K}[\chi])$. For any $\tau \in J_L$, let $e_\tau = (e_{\tau, \rho})_{\rho \in J_{\mathbb{Q}(\chi)}} \in (\mathbb{Q}(\chi) \otimes \mathbb{C})^\times$ be the element whose ρ -coordinate is $e_{\tau, \rho} = 1$ if $n(\tau, \rho) > n(\bar{\tau}, \rho)$ and $e_{\tau, \rho} = -1$ if $n(\tau, \rho) < n(\bar{\tau}, \rho)$. Recall that the restriction of χ to \mathbb{A}_K^\times can be written as

$$\chi|_{\mathbb{A}_K^\times} = \chi_0 \| \cdot \|^{-w(\chi)},$$

where χ_0 is a finite order character. We let ε_L denote the finite order character of $K^\times \backslash \mathbb{A}_K^\times$ corresponding under class field theory to the quadratic character of $\text{Gal}(\bar{K}/K)$ associated with the extension L/K . We let $[\chi_0 \varepsilon_L]$ denote the Artin motive of rank 1 over K , with coefficients in $\mathbb{Q}(\chi)$, attached to the finite order character $\chi_0 \varepsilon_L$. For each $\sigma \in J_K$, we let

$$G_\sigma(\chi) = \delta_\sigma[\chi_0 \varepsilon_L].$$

Remark 2.2.1. When $K = \mathbb{Q}$, such elements are classically given by Gauss sums (see [12], 6.4). There is no such simple relation with Gauss sums when $K \neq \mathbb{Q}$ (see [27], 3.4).

We let $M(\chi) = M \otimes \text{Res}_{L/K}[\chi]$ be the tensor product of the realizations M and $\text{Res}_{L/K}[\chi]$. This tensor product is over \mathbb{Q} , meaning that it has coefficients in $E(\chi) = E \otimes \mathbb{Q}(\chi)$. To be more precise, $E(\chi)$ is a product of number fields, and $M(\chi)$ is a collection of realizations with coefficients in each of these fields. For simplicity in most of what follows, we will simply assume that $E(\chi)$ is a field. Suppose that M is regular and χ is critical. It's shown in [14] (Proposition 2.5.1) that $M(\chi)$ has critical values if and only if, for every $\sigma \in J_K$ and every $\rho \in J_{\mathbb{Q}(\chi)}$,

$$t_{\sigma, \rho}(\chi) \neq w(M) - 2p_i(\sigma, \varphi)$$

for any $i = 1, \dots, n$ and any $\varphi \in J_E$. Assuming this is the case, we can find integers $r_{\sigma, \varphi, \rho}(\chi) \in \{0, \dots, n\}$ such that

$$w(M) - 2p_{r_{\sigma, \varphi, \rho}(\chi)}(\sigma, \varphi) < t_{\sigma, \rho}(\chi) < w(M) - 2p_{r_{\sigma, \varphi, \rho}(\chi)+1}(\sigma, \varphi).$$

We stress here that if $K \neq \mathbb{Q}$, the numbers $p_i(\sigma, \varphi)$ depend in general on the choice of the embeddings. Moreover, the integers $r_{\sigma, \varphi, \rho}(\chi)$ also depend on the choice of the three embeddings σ , φ and ρ .

We say that M is *polarized* if there exists a non-degenerate morphism of realizations

$$\langle \cdot, \cdot \rangle : M \otimes_E M \rightarrow E(-w(M)).$$

We also impose the condition that $\langle \cdot, \cdot \rangle$ is symmetric if $w(M)$ is even, and alternated if $w(M)$ is odd. We refer to Sect. 2.3 of [14] for the definition of the quadratic periods $Q_{j, \sigma} \in (E \otimes \mathbb{C})^\times$ attached to a polarized realization M . Assume from now on that M is a regular, polarized, special realization, pure of weight $w(M)$ and rank n .

Let χ be an critical algebraic Hecke character of infinity type $(n_\tau)_{\tau \in J_L}$ and weight $w(\chi)$. We suppose that $M(\chi)$ has critical values, and we let $s_{\sigma, \varphi, \rho}(\chi) = n - r_{\sigma, \varphi, \rho}(\chi)$, $\mathbf{r}_\sigma = (r_{\sigma, \varphi, \rho}(\chi))_{\varphi, \rho}$ and $\mathbf{s}_\sigma = (s_{\sigma, \varphi, \rho}(\chi))_{\varphi, \rho}$. We let

$$a_\sigma^\pm(\chi) = (2\pi i)^{w(\chi)} G_\sigma(\chi)^{-1} c_\sigma^\mp(\chi)$$

and

$$Q_\sigma(\chi) = (2\pi i)^{w(\chi)} G_\sigma(\chi)^{-2} e_\tau c_\sigma^+(\chi)^2.$$

Here τ is any of the two embeddings of L extending σ . Since $e_\tau = -e_{\bar{\tau}}$, this definition makes sense in $(\mathbb{Q}(\chi) \otimes \mathbb{C})^\times$ modulo $(\mathbb{Q}(\chi) \otimes \sigma(K))^\times$.

Remark 2.2.2. These quantities are defined more conceptually in [14]. The formulas above are obtained (up to multiples in $(\mathbb{Q}(\chi) \otimes \sigma(K))^\times$) in (2.4.3) and Proposition 2.5.2 of *op. cit.*

Theorem 2.5.1 of [14] (see also Proposition 1.7.6 of [17] when $K = \mathbb{Q}$, and [33]) says that

$$c_{\sigma}^{+}(M(\chi)) \sim (2\pi i)^{-\lceil n/2 \rceil w(\chi)} G_{\sigma}(\chi)^{r_{\sigma}} \delta_{\sigma}(M) a_{\sigma}^{*}(\chi) Q_{\sigma}(\chi)^{r_{\sigma} - \lceil n/2 \rceil} \prod_{j=1}^{s_{\sigma}} Q_{j,\sigma}, \quad (2.2.3)$$

where $a_{\sigma}^{*}(\chi) = 1$ if n is even, and $a_{\sigma}^{*}(\chi) = a_{\sigma}^{\pm}(\chi)$ if n is odd, with $\pm = -$ if $n^{+} > n^{-}$ and $\pm = +$ if $n^{-} > n^{+}$. In the formula, \sim means $\sim_{E(\chi) \otimes K, \sigma}$. In particular, if we look at the coordinates of the elements corresponding to an embedding $\varphi \in J_E$ and the embedding $1 \in J_{\mathbb{Q}(\chi)}$, we get

$$c_{\sigma}^{+}(M(\chi))_{\varphi,1} \sim (2\pi i)^{-\lceil n/2 \rceil w(\chi)} G_{\sigma}(\chi)_{1}^{r_{\sigma,\varphi,1}(\chi)} \delta_{\sigma}(M)_{\varphi} a_{\sigma}^{*}(\chi)_{1} Q_{\sigma}(\chi)_{1}^{r_{\sigma,\varphi,1}(\chi) - \lceil n/2 \rceil} \prod_{j=1}^{s_{\sigma,\varphi,1}(\chi)} Q_{j,\sigma,\varphi}, \quad (2.2.4)$$

where now both sides are complex numbers and \sim stands for $\sim_{\varphi(E)\mathbb{Q}(\chi)\sigma(K)}$. When E is given as a subfield of \mathbb{C} with a given embedding $1 \in J_E$, we write $c^{+}(M(\chi))_1$ for $c^{+}(M(\chi))_{1,1}$, and similarly for any other element of $E(\chi) \otimes \mathbb{C}$.

2.3. Twists with prescribed critical intervals

In this subsection, we consider twists of M by certain algebraic Hecke characters with prescribed r . Let M be a polarized, regular, special realization over K with coefficients in E . We let n be the rank of M , and assume that M is pure of weight $w(M)$. Fix a CM type Φ for L/K , $\varphi \in J_E$ and $r \in \{0, \dots, n\}$. If $r > 0$, let $(a_{\tau}^{(r,\varphi)})_{\tau \in \Phi}$ be the tuple of integers defined by

$$a_{\tau}^{(r,\varphi)} = w(M) - 2p_r(\sigma, \varphi) + 1, \quad (2.3.1)$$

where σ is the restriction of τ to K . For $r = 0$, let $a^{(0)}$ be any integer such that

$$\begin{aligned} a^{(0)} &< \min\{w(M) - 2p_1(\sigma, \varphi)\}_{\sigma \in J_K, \varphi \in J_E}, \\ a^{(0)} &\equiv w(M) + 1(2). \end{aligned} \quad (2.3.2)$$

For any τ and φ , we let $a_{\tau}^{(0,\varphi)} = a^{(0)}$. Note that $w(M) - 2p_1(\sigma, \varphi) = p_n(\sigma, \varphi) - p_1(\sigma, \varphi) \leq 0$, and hence $a_{\tau}^{(0,\varphi)} < 0$. In all cases, we have that

$$a_{\tau}^{(r,\varphi)} \equiv a_{\tau'}^{(r,\varphi)}(2) \quad (\tau, \tau' \in \Phi),$$

and thus by Proposition 2.1.1, there exists an algebraic Hecke character $\chi^{(r,\varphi)}$ of L , of infinity type $(n_{\tau}^{(r,\varphi)})_{\tau \in \Phi}$, such that

$$n_{\tau}^{(r,\varphi)} - n_{\tau'}^{(r,\varphi)} = a_{\tau}^{(r,\varphi)} \quad (\tau \in \Phi).$$

- Lemma 2.3.1.** (i) *The algebraic Hecke characters $\chi^{(r,\varphi)}$ are all critical, except for $r = n/2$ when n is even. In this case, $\chi^{(n/2,\varphi)}$ is critical if and only if $p_{\frac{n}{2}}(\sigma, \varphi) \neq p_{\frac{n}{2}+1}(\sigma, \varphi) + 1$ for any $\sigma \in J_K$.*
- (ii) *The infinity type satisfies that $n_{\tau}^{(r,\varphi)} > n_{\bar{\tau}}^{(r,\varphi)}$ for one (or every) $\tau \in \Phi$ if and only if $r \in \{\lfloor \frac{n}{2} \rfloor + 1, \dots, n\}$.*

Proof. If $r = 0$, then $a_{\tau}^{(0,\varphi)} < 0$, which implies that $n_{\tau}^{(0,\varphi)} \neq n_{\bar{\tau}}^{(0,\varphi)}$ for every $\tau \in \Phi$, so $\chi^{(0,\varphi)}$ is critical. From now on assume that $r > 0$. Then $n_{\tau}^{(r,\varphi)} - n_{\bar{\tau}}^{(r,\varphi)} = w(M) - 2p_r(\sigma, \varphi) + 1$ for $\tau \in \Phi$. Suppose that $\chi^{(r,\varphi)}$ is not critical, so that $n_{\tau}^{(r,\varphi)} = n_{\bar{\tau}}^{(r,\varphi)}$ for some $\tau \in \Phi$. Then

$$w(M) = 2p_r(\sigma, \varphi) - 1.$$

Since $w(M) = p_{n+1-r}(\sigma, \varphi) + p_r(\sigma, \varphi)$, this amounts to say that

$$p_{n+1-r}(\sigma, \varphi) = p_r(\sigma, \varphi) - 1. \quad (2.3.3)$$

This necessarily implies that $r \neq n$ and

$$p_{n+1-r}(\sigma, \varphi) = p_{r+1}(\sigma, \varphi).$$

This in turn implies that $n + 1 - r = r + 1$, so that n is even and $r = n/2$. Then (2.3.3) implies that $p_{\frac{n}{2}}(\sigma, \varphi) = p_{\frac{n}{2}+1}(\sigma, \varphi) + 1$. This proves part (i).

For part (ii), the condition is equivalent to $a_{\tau}^{(r,\varphi)} > 0$ for every $\tau \in \Phi$, which means that $r \neq 0$ and

$$p_{n+1-r}(\sigma, \varphi) - p_r(\sigma, \varphi) \geq 0$$

for every $\sigma \in J_K$. This means that $r \geq \frac{n+1}{2}$, or, what is the same, $r \geq \lfloor \frac{n}{2} \rfloor + 1$. \square

Lemma 2.3.2. *Let $r \in \{0, \dots, n\}$. If n is even and $r = n/2$, assume that $p_{\frac{n}{2}}(\sigma, \varphi) \neq p_{\frac{n}{2}+1}(\sigma, \varphi) + 1$. Then the realization $M(\chi^{(r,\varphi)})$ has critical values, and for every $\sigma \in J_K$,*

$$r_{\sigma,\varphi,1}(\chi^{(r,\varphi)}) = \begin{cases} n - r & \text{if } 0 \leq r \leq \lfloor \frac{n}{2} \rfloor \\ r & \text{if } \lfloor \frac{n}{2} \rfloor < r \leq n, \end{cases}$$

where $1 \in J_{\mathbb{Q}(\chi^{(r,\varphi)})}$ denotes the given embedding of $\mathbb{Q}(\chi^{(r,\varphi)})$.

Proof. First we need to check the condition that guarantees that $M(\chi^{(r,\varphi)})$ has critical values. Thus, we need to see that for any $\sigma \in J_K$, $\psi \in J_E$, $\rho \in J_{\mathbb{Q}(\chi^{(r,\varphi)})}$ and $i = 1, \dots, n$, $t_{\sigma,\rho}(\chi^{(r,\varphi)}) \neq w(M) - 2p_i(\sigma, \psi)$. But

$$t_{\sigma,\rho}(\chi^{(r,\varphi)}) = \pm(n^{(r,\varphi)}(\tau, \rho) - n^{(r,\varphi)}(\bar{\tau}, \rho)),$$

where $\tau \in \Phi$ extends σ . Suppose first that $r \neq 0$. Then

$$n^{(r,\varphi)}(\tau, \rho) - n^{(r,\varphi)}(\bar{\tau}, \rho) = n_{\rho^{-1}\tau}^{(r,\varphi)} - n_{\rho^{-1}\bar{\tau}}^{(r,\varphi)} = w(M) - 2p_r(\sigma', \varphi) + 1,$$

where $\sigma' \in J_K$ is the restriction of $\tilde{\rho}^{-1}\sigma$ to K . Thus, using that $p_r(\sigma', \varphi) + p_{n+1-r}(\sigma', \varphi) = w(M)$,

$$t_{\sigma, \rho}(\chi^{(r, \varphi)}) = \pm(w(M) - 2p_r(\sigma', \varphi) + 1) = w(M) - 2p_{r'}(\sigma', \varphi) \pm 1,$$

where r' is either r or $n + 1 - r$, according to whether the sign is $+$ or $-$. This differs from $w(M)$ by an odd integer, so it can never be equal to $w(M) - 2p_i(\sigma, \psi)$.

If $r = 0$, then

$$t_{\sigma, \rho}(\chi^{(r, \varphi)}) = -a^{(0)},$$

which by our choices is never equal to any of the $w(M) - 2p_i(\sigma, \varphi)$. This shows that $M(\chi^{(r, \varphi)})$ has critical values.

Now, if $\tau \in \Phi$ extends σ then, by part (ii) of Lemma 2.3.1,

$$t_{\sigma, 1}(\chi) = \begin{cases} n_{\tau}^{(r, \varphi)} - n_{\tau}^{(r, \varphi)} & \text{if } \lfloor \frac{n}{2} \rfloor < r \leq n \\ n_{\tau}^{(r, \varphi)} - n_{\tau}^{(r, \varphi)} & \text{if } 0 \leq r \leq \lfloor \frac{n}{2} \rfloor. \end{cases}$$

If $\lfloor \frac{n}{2} \rfloor < r \leq n$, then $t_{\sigma, 1}(\chi^{(r, \varphi)}) = w(M) - 2p_r(\sigma, \varphi) + 1$. Since $p_{r+1}(\sigma, \varphi) + \frac{1}{2} < p_r(\sigma, \varphi)$, it follows that

$$w(M) - 2p_r(\sigma, \varphi) < t_{\sigma, 1}(\chi^{(r, \varphi)}) < w(M) - 2p_{r+1}(\sigma, \varphi),$$

so that $r_{\sigma, \varphi, 1}(\chi^{(r, \varphi)}) = r$. If $0 \leq r \leq \lfloor \frac{n}{2} \rfloor$, then $t_{\sigma, 1}(\chi^{(r, \varphi)}) = -w(M) + 2p_r(\sigma, \varphi) - 1 = w(M) - 2p_{n+1-r}(\sigma, \varphi) - 1$, and by a similar reasoning we get that

$$w(M) - 2p_{n-r}(\sigma, \varphi) < t_{\sigma, 1}(\chi^{(r, \varphi)}) < w(M) - 2p_{n-r+1}(\sigma, \varphi),$$

so that $r_{\sigma, \varphi, 1}(\chi^{(r, \varphi)}) = n - r$. □

2.4. Formulas for quadratic periods

In this subsection, we combine the results above to obtain a formula for the quadratic periods $Q_{j, \sigma}$ in terms of quotients of Deligne σ -periods of various twists $M(\chi)$, with an eye towards future applications. We first introduce some notation. For any $r = 0, \dots, n$, and $\varphi \in J_E$, we fix $\chi^{(r, \varphi)}$ as above. Given our choices in (2.3.1) and (2.3.2), we can assume by Proposition 2.1.1 that all the characters $\chi^{(r, \varphi)}$ have the same weight $w_0 \equiv w(M) + 1(2)$. We let

$$P_{\sigma}(\chi^{(r, \varphi)}) = (2\pi i)^{-\lceil \frac{n}{2} \rceil w_0} G_{\sigma}(\chi^{(r, \varphi)})^r a_{\sigma}^*(\chi^{(r, \varphi)}) Q_{\sigma}(\chi^{(r, \varphi)})^{r - \lceil \frac{n}{2} \rceil}. \quad (2.4.1)$$

Implicit here is the integer n and its decomposition as $n = n^+ + n^-$, depending on M . This is an element of $(\mathbb{Q}(\chi) \otimes \mathbb{C})^{\times}$, well defined up to multiples in $(\mathbb{Q}(\chi) \otimes \sigma(K))^{\times}$. Note that if $1 \leq j < \lceil \frac{n}{2} \rceil$, then $\lfloor \frac{n}{2} \rfloor < n - j \leq n - 1$.

Proposition 2.4.1. *Let $\varphi \in J_E$ and $j \in \mathbb{Z}$ be an integer with $1 \leq j < \lfloor \frac{n}{2} \rfloor$. Then, for any $\sigma \in J_K$, we have that*

$$Q_{j,\sigma,\varphi} \sim_{\varphi(E)\mathbb{Q}(\chi)\sigma(K)} \begin{cases} \frac{c_\sigma^+(M(\chi^{(n-1,\varphi)}))_{\varphi,1}}{P_\sigma(\chi^{(n-1,\varphi)})_1} \delta_\sigma(M)_\varphi^{-1} & \text{if } j = 1 \\ \frac{c_\sigma^+(M(\chi^{(n-j,\varphi)}))_{\varphi,1}}{c_\sigma^+(M(\chi^{(n-j+1,\varphi)}))_{\varphi,1}} \frac{P_\sigma(\chi^{(n-j+1,\varphi)})_1}{P_\sigma(\chi^{(n-j,\varphi)})_1} & \text{if } j \neq 1. \end{cases}$$

Proof. Let $r = n - j \in \{\lfloor \frac{n}{2} \rfloor + 1, \dots, n - 1\}$. We apply the formula (2.2.4) to $\chi^{(r,\varphi)}$. By Lemma 2.3.2, the formula in this case, looking at the coordinate given by the embeddings $\varphi \in J_E$ and $1 \in J_{\mathbb{Q}(\chi)}$, says that

$$c_\sigma^+(M(\chi^{(r,\varphi)}))_{\varphi,1} \sim \delta_\sigma(M)_\varphi P_\sigma(\chi^{(r,\varphi)})_1 \prod_{i=1}^j Q_{i,\sigma,\varphi}. \quad (2.4.2)$$

Thus the formula for $j = 1$ in the proposition is clear. If $j \geq 2$, then we also have

$$c_\sigma^+(M(\chi^{(r+1,\varphi)}))_{\varphi,1} \sim \delta_\sigma(M)_\varphi P_\sigma(\chi^{(r+1,\varphi)})_1 \prod_{i=1}^{j-1} Q_{i,\sigma,\varphi}. \quad (2.4.3)$$

Hence, the result follows by dividing (2.4.2) by (2.4.3) (note that all the relevant elements belong to \mathbb{C}^\times). \square

We define

$$Q_j = \prod_{\sigma \in J_K} Q_{j,\sigma} \in (E \otimes \mathbb{C})^\times / (E \otimes K^{\text{Gal}})^\times$$

and

$$P(\chi^{(r,\varphi)}) = \prod_{\sigma \in J_K} P_\sigma(\chi^{(r,\varphi)}) \in (\mathbb{Q}(\chi) \otimes \mathbb{C})^\times / (\mathbb{Q}(\chi) \otimes K^{\text{Gal}})^\times. \quad (2.4.4)$$

Using formulas (2.2.1) and (2.2.2), we obtain the following corollary of Proposition 2.4.1.

Corollary 2.4.1. *Let the notation and assumptions be as above. Then, for each $\varphi \in J_E$, we have that*

$$Q_{j,\varphi} \sim_{\varphi(E)\mathbb{Q}(\chi)K^{\text{Gal}}} \begin{cases} \frac{c^+(M(\chi^{(n-1,\varphi)}))_{\varphi,1}}{P(\chi^{(n-1,\varphi)})_1} \delta(M)_\varphi^{-1} & \text{if } j = 1 \\ \frac{c^+(M(\chi^{(n-j,\varphi)}))_{\varphi,1}}{c^+(M(\chi^{(n-j+1,\varphi)}))_{\varphi,1}} \frac{P(\chi^{(n-j+1,\varphi)})_1}{P(\chi^{(n-j,\varphi)})_1} & \text{if } j \neq 1. \end{cases}$$

Fix $\varphi \in J_E$ and $r \in \{0, \dots, n\}$. Let $s = n - r$. For the rest of this subsection, we let $\chi = \chi^{(r,\varphi)}$. Recall that w_0 is the weight of χ . The period $P(\chi)$ of (2.4.4) can be interpreted in terms of CM periods, as in Sect. 2.6 of [14]. We are only interested in the case $r > \lfloor \frac{n}{2} \rfloor$, and we assume that this is the case from now. In particular, $n_\tau > n_{\bar{\tau}}$ for $\tau \in \Phi$. Moreover, we will only write down the formulas after fixing

the embedding $1 \in J_{\mathbb{Q}(\chi)}$, since $P(\chi)_1$ is what appears in Corollary 2.4.1. In [21] (see also [16]), a family of CM periods attached to χ is defined. As a particular case, there is a period

$$p(\chi; \Phi) \in \mathbb{C}^\times,$$

well defined up to multiples in $\mathbb{Q}(\chi)^\times$. For each embedding $\rho \in J_{\mathbb{Q}(\chi)}$, let $\tilde{\rho}$ be an extension of ρ to \mathbb{C} . We can define an algebraic Hecke character χ^ρ , of infinity type $(n_{\tilde{\rho}^{-1}\tau})_{\tau \in J_L}$, obtained by applying ρ to the values of χ on $\mathbb{A}_{L,f}^\times$. We can also define a CM type $\tilde{\rho}\Phi = \{\tilde{\rho}\tau : \tau \in \Phi\}$. All of these are independent of the extension $\tilde{\rho}$, and thus we get a well defined CM period $p(\chi^\rho; \rho\Phi) \in \mathbb{C}^\times$. We let

$$\mathbf{p}(\chi; \Phi) = (p(\chi^\rho; \rho\Phi))_{\rho \in J_{\mathbb{Q}(\chi)}} \in (\mathbb{Q}(\chi) \otimes \mathbb{C})^\times.$$

The following formula is a theorem of Blasius, and we use the formulation that appears as Proposition 1.8.1 of [16], corrected as in the Introduction to [17]. Combined with Deligne's conjecture for the motive $[\chi]$, proved by Blasius ([4]), we get that if m is a critical integer for $[\chi]$, then

$$c^+([\chi](m)) \sim_{\mathbb{Q}(\chi)} D_K^{1/2} (2\pi i)^{[K:\mathbb{Q}]m} \mathbf{p}(\check{\chi}; \Phi), \quad (2.4.5)$$

where $\check{\chi} = \chi^{\iota, -1}$. Here $\iota \in \text{Gal}(L/K)$ is the non-trivial element.

The following lemma allows us to relate the quadratic periods $Q_{j,\varphi}$ to quotients of Deligne periods and CM periods, via Corollary 2.4.1. Let

$$G(\chi) = \prod_{\sigma \in J_K} G_\sigma(\chi) \in (\mathbb{Q}(\chi) \otimes \mathbb{C})^\times.$$

Lemma 2.4.1. *Let the notation and assumptions be as above. Let $t = 0$ if there exist even critical integers m for $[\chi]$, and let $t = 1$ otherwise. Then*

$$P(\chi) \sim_{\mathbb{Q}(\chi) \otimes K^{\text{Gal}}} (2\pi i)^{-[K:\mathbb{Q}]w_0 s} G(\chi)^s \mathbf{p}(\check{\chi}; \Phi)^{r-s} \left(\prod_{\tau \in \Phi} e_\tau^{r-n^++t} \right).$$

In particular,

$$P(\chi)_1 \sim_{\mathbb{Q}(\chi)K^{\text{Gal}}} (2\pi i)^{-[K:\mathbb{Q}]w_0 s} G(\chi)_1^s p(\check{\chi}; \Phi)^{r-s}.$$

Proof. Throughout, we write \sim for $\sim_{\mathbb{Q}(\chi) \otimes K^{\text{Gal}}}$. By definition, we have that

$$P(\chi) \sim (2\pi i)^{-\lceil \frac{n}{2} \rceil w_0 [K:\mathbb{Q}]} G(\chi)^r \left(\prod_{\sigma \in J_K} a_\sigma^*(\chi) \right) \left(\prod_{\sigma \in J_K} Q_\sigma(\chi)^{r - \lceil \frac{n}{2} \rceil} \right), \quad (2.4.6)$$

where $a_\sigma^*(\chi) = 1$ if n is even and $a_\sigma^*(\chi) = a_\sigma^\pm(\chi)$ if n is odd, where $\pm = -$ if $n^+ > n^-$ and $\pm = +$ if $n^+ < n^-$.

By Lemma 2.4.1 of [14], we have that

$$c_\sigma^-(\chi) \sim_{\mathbb{Q}(\chi) \otimes K, \sigma} e_\tau c_\sigma^+(\chi), \quad (2.4.7)$$

for τ any extension of σ . It follows from (2.2.1) that

$$\prod_{\sigma \in J_K} a_{\sigma}^{\pm}(\chi) \sim (2\pi i)^{[K:\mathbb{Q}]w_0} D_K^{-1/2} G(\chi)^{-1} c^+(\chi) \left(\prod_{\tau \in \Phi} e_{\tau}^? \right), \quad (2.4.8)$$

where $? = 1$ if $\pm = +$, and $? = 0$ if $\pm = -$. The factor $D_K^{-1/2}$ belongs to K^{Gal} , so it can be ignored in the formula. Similarly,

$$\prod_{\sigma \in J_K} Q_{\sigma}(\chi) \sim (2\pi i)^{[K:\mathbb{Q}]w_0} G(\chi)^{-2} c^+(\chi)^2 \left(\prod_{\tau \in \Phi} e_{\tau} \right). \quad (2.4.9)$$

Since $c^+([\chi](m)) \sim (2\pi i)^{[K:\mathbb{Q}]m} c^{(-1)^m}(\chi)$, we get from (2.4.5) and (2.4.7) that

$$c^+(\chi) \sim_{\mathbb{Q}(\chi)} D_K^{1/2} \mathbf{p}(\check{\chi}; \Phi) \prod_{\tau \in \Phi} e_{\tau}^t. \quad (2.4.10)$$

The lemma follows by combining (2.4.6), (2.4.7), (2.4.8), (2.4.9) and (2.4.10). \square

Remark 2.4.1. The last lemma is true for any critical χ such that $n_{\tau} > n_{\bar{\tau}}$ for $\tau \in \Phi$, as long as the definition (2.4.1) of $P(\chi)$ uses the same r, s and n^+ .

2.5. Deligne periods when $r = n$

In this subsection, we obtain a formula for the Deligne period of $M(\chi)(k)$ whenever χ has $r = n$ and k is a critical integer. We can use the above formulas for $\chi^{(n, \varphi)}$, but it will be more useful to allow more general characters. Thus, suppose that χ is an algebraic Hecke character, critical of infinity type $(n_{\tau})_{\tau \in J_L}$. Suppose that the CM type Φ and the infinity type of χ are related by the condition

$$n_{\tau} > n_{\bar{\tau}} \quad (\tau \in \Phi).$$

Moreover, fix $\varphi \in J_E$, and suppose that

$$n_{\tau} - n_{\bar{\tau}} > \max\{w(M) - 2p_n(\sigma, \varphi)\}_{\sigma \in J_K} \quad (2.5.1)$$

for all $\tau \in \Phi$.

Remark 2.5.1. If (2.5.1) holds for the embedding $\varphi \in J_E$, then it holds for any other $\psi \in J_E$ as well. This follows from the following more general fact. Let N be a realization over K , with coefficients in E , pure of weight $w(N)$. For $\varphi \in J_E$, let

$$T(\varphi) = \bigcup_{\sigma \in J_K} \{p \in \mathbb{Z} : N_{\sigma}^{pq}(\varphi) \neq 0\}.$$

Then $T(\varphi) = T(\psi)$ for $\varphi, \psi \in J_E$. Indeed, let $p \in T(\varphi)$. Then $\text{gr}^p(N_{\text{dR}}) \otimes_{E \otimes K, \varphi \otimes \sigma} \mathbb{C} \neq 0$. There exists an element $h \in \text{Aut}(\mathbb{C})$ such that $\psi = h\varphi$, and then

$$0 \neq \text{gr}^p(N_{\text{dR}}) \otimes_{E \otimes K, \varphi \otimes \sigma} \mathbb{C} \otimes_{\mathbb{C}, h} \mathbb{C} = \text{gr}^p(N_{\text{dR}}) \otimes_{E \otimes K, \psi \otimes h\sigma},$$

so $p \in T(\psi)$. Then, (2.5.1) can be stated as

$$n_{\tau} - n_{\bar{\tau}} > \max\{w(M) - 2p\}_{p \in T(\varphi)},$$

which is independent of φ .

We then have, just as in the proof of Lemma 2.3.2, that $M(\chi)$ has critical values and

$$r_{\sigma, \varphi, 1}(\chi) = n \quad (2.5.2)$$

for all $\sigma \in J_K$ and $\varphi \in J_E$. As in (2.4.1) and (2.4.4), we let

$$P_\sigma(\chi) = (2\pi i)^{-\lceil \frac{n}{2} \rceil w(\chi)} G_\sigma(\chi)^n a_\sigma^*(\chi) Q_\sigma(\chi)^{\lfloor \frac{n}{2} \rfloor}$$

and

$$P(\chi) = \prod_{\sigma \in J_K} P_\sigma(\chi),$$

where again $n = n^+ + n^-$ is implicit in the notation.

Taking into account (2.5.2), formula (2.2.3) applied to the case of χ says that

$$c_\sigma^+(M(\chi))_{\varphi, 1} \sim_{\varphi(E)\mathbb{Q}(\chi)\sigma(K)} \delta_\sigma(M)_\varphi P_\sigma(\chi)_1.$$

Using formulas (2.2.1) and (2.2.2), we get

$$c^+(M(\chi))_{\varphi, 1} \sim_{\varphi(E)\mathbb{Q}(\chi)K^{\text{Gal}}} \delta(M)_\varphi P(\chi)_1. \quad (2.5.3)$$

The set of critical integers for $M(\chi)$ is computed in (2.5.2) of [14]. In this case, this set consist of those integers k such that

$$p_1(\sigma, \varphi) + n_{\bar{\tau}} < k \leq p_n(\sigma, \varphi) + n_\tau \quad (2.5.4)$$

for every $\tau \in \Phi$ (with $\sigma = \tau|_K$) and a fixed $\varphi \in J_E$.

Remark 2.5.2. By Remark 2.5.1, these inequalities are independent of the chosen φ . For this we need to assume, as we do, that $E(\chi)$ is a field. In the general case, $M(\chi)$ is a collection of realizations with coefficients in the fields appearing in $E(\chi)$, and a critical integer is defined to be one which is critical for each of these realizations.

Let k be an integer satisfying (2.5.4). By Lemma 2.4.1 of [14],

$$c_\sigma^+(M(\chi)(k)) \sim_{E(\chi) \otimes K, \sigma} (2\pi i)^{kn} c_\sigma^+(M(\chi)) e_\tau^{(-1)^k}$$

with $\tau \in \Phi$ extending σ . Using (2.2.1), we get that

$$c^+(M(\chi)(k)) \sim_{E(\chi) \otimes K^{\text{Gal}}} (2\pi i)^{[K:\mathbb{Q}]kn} c^+(M(\chi)) \prod_{\tau \in \Phi} e_\tau^{(-1)^k}.$$

Combining this with (2.5.3), we obtain that

$$c^+(M(\chi)(k))_{\varphi, 1} \sim_{\varphi(E)\mathbb{Q}(\chi)K^{\text{Gal}}} (2\pi i)^{[K:\mathbb{Q}]kn} \delta(M)_\varphi P(\chi)_1.$$

Finally, the following result follows from this and Lemma 2.4.1 (see also (2.6.2) of [14]). For clarity, we recall all the relevant hypotheses.

Proposition 2.5.1. *Let M be a regular, special, polarized realization over K with coefficients in E , pure of weight $w(M)$ and rank n . Let Φ be a CM type for L/K , and let χ be a critical algebraic Hecke character of L of infinity type $(n_\tau)_{\tau \in J_L}$. Suppose that $n_\tau > n_{\bar{\tau}}$ for $\tau \in \Phi$, and that (2.5.1) holds. Let k be an integer satisfying (2.5.4). Then*

$$c^+(M(\chi)(k))_{\varphi,1} \sim_{\varphi(E)\mathbb{Q}(\chi)K^{\text{Gal}}} (2\pi i)^{[K:\mathbb{Q}]kn} \delta(M)_\varphi p(\check{\chi}; \Phi)^n$$

for any $\varphi \in J_E$.

Remark 2.5.3. The factor $\delta(M)_\varphi$ in the previous proposition needs to be dealt with. It can easily be replaced with a suitable power of $2\pi i$ if $w(M)$ is odd (and hence n is even and the polarization is alternated). More precisely, it can be shown (see Lemma 1.4.12 of [17] or Remark 2.3.1 of [14]) that $\delta_\sigma(M) \sim_{E \otimes K, \sigma} (2\pi i)^{-w(M)n/2}$. Thus, (2.2.2) implies that

$$\delta(M) \sim_{E \otimes K^{\text{Gal}}} (2\pi i)^{-w(M)[K:\mathbb{Q}]n/2}$$

in this case. When $w(M)$ is even, there is apparently no simple way to obtain such an expression. When comparing with automorphic motives in the following sections, we will deal with this case assuming an extra conjecture.

3. Critical values of automorphic L -functions

In this section we recall the main results of [14] regarding the critical values of L -functions of cohomological automorphic forms on unitary groups. In this paper, we will only care about totally definite unitary groups.

3.1. Totally definite unitary groups

Let L/K be a CM extension and Φ a CM type for L/K . Let V be a finite-dimensional L -vector space, and $h : V \times V \rightarrow L$ be a non-degenerate hermitian form, relative to the non-trivial automorphism $\iota \in \text{Gal}(L/K)$. Let $n = \dim_L V$. We let G be the similitude unitary group, with similitude factors in \mathbb{Q} , attached to (V, h) . Thus, for a \mathbb{Q} -algebra R , the points of G with values in R are given by

$$G(R) = \{g \in \text{Aut}_{L \otimes R}(V \otimes R) : h_R(gu, gv) = v(g)h_R(u, v) \quad \forall u, v \in V \otimes R\},$$

where $v(g) \in R^\times$. Here $h_R : V \otimes R \times V \otimes R \rightarrow L \otimes R$ is given by $h_R(u \otimes a, v \otimes b) = h(u, v) \otimes ab$. For each $\tau \in J_L$, let $V_\tau = V \otimes_{L, \tau} \mathbb{C}$. This is equipped with a hermitian form h_τ relative to complex conjugation on \mathbb{C}/\mathbb{R} . In particular, there is a well-defined signature (r_τ, s_τ) . We will assume throughout the paper that V is totally definite. This means that for any $\tau \in \Phi$, the signature is $(r_\tau, s_\tau) = (n, 0)$. We also fix an L -basis $\beta = \{v_1, \dots, v_n\}$ of V , orthogonal for h . As in (3.1.1) of [14], we can write

$$G_{\mathbb{R}} \cong \left(\prod_{\tau \in \Phi} GU(n, 0) \right)', \quad G_{\mathbb{C}} \cong \left(\prod_{\tau \in \Phi} \text{GL}_{n, \mathbb{C}} \right) \times \text{GL}_{1, \mathbb{C}},$$

where the symbol $'$ means that we are looking at tuples where all the elements have the same multiplier ν . Here, the group $GU(n, 0)$ is the usual similitude unitary group over \mathbb{R} of the identity matrix I_n . There is a maximal torus $T \subset G$, namely the subgroup of automorphism which are diagonal with respect to the basis β , such that $T_{\mathbb{C}}$ corresponds to the subgroup of diagonal matrices under the second isomorphism above. We let $B \subset G_{\mathbb{C}}$ be the Borel subgroup corresponding to $(\prod_{\tau \in \Phi} B_{n, \mathbb{C}}) \times GL_{1, \mathbb{C}}$, where $B_{n, \mathbb{C}}$ is the group of upper triangular matrices in $GL_{n, \mathbb{C}}$.

We use the notation of Sect. 3.3 of [14] regarding roots and weights. In particular, we identify the group $\Lambda = X^*(T)$ with the group of tuples

$$\mu = ((a_{\tau, 1}, \dots, a_{\tau, n})_{\tau \in \Phi}; a_0) \in \left(\prod_{\tau \in \Phi} \mathbb{Z}^n \right) \times \mathbb{Z}.$$

The set of dominant weights Λ^+ , with respect to the Borel subgroup B , are those μ for which $a_{\tau, 1} \geq \dots \geq a_{\tau, n}$ for all $\tau \in \Phi$. In the notation of *op. cit.*, the group K_x is the whole group $G_{\mathbb{R}}$, $\Lambda_{x, c}^+ = \Lambda^+$ and $w_0^1 = 1$.

3.2. Automorphic forms

A Shimura datum (G, X) is constructed in Sect. 3.2 of [14], assuming that V is not totally definite. In our case, we can still define zero-dimensional varieties S_U for a compact open subgroup $U \subset G(\mathbb{A}_f)$. These are algebraic varieties over the reflex field E , which is the field generated over \mathbb{Q} by the elements $\sum_{\tau \in \Phi} \tau(b)$, for $b \in L$. In particular, $E \subset L^{\text{Gal}}$, the Galois closure of L in $\bar{\mathbb{Q}}$. The set of complex points of S_U is the finite set

$$S_U(\mathbb{C}) = G(\mathbb{Q}) \backslash G(\mathbb{A}_f) / U.$$

Most of what is contained in Sect. 3 of [14] also applies to these zero-dimensional varieties. We denote by S the projective limit of the S_U .

From now on, fix $\mu \in \Lambda^+$ such that the corresponding representation $W = W_{\mu}$ of $G_{\mathbb{C}}$ is defined over \mathbb{Q} . This implies that

$$a_{\tau, i} = -a_{\tau, n+1-i}$$

for every $\tau \in \Phi$ and $i = 1, \dots, n$. Let $\xi = 2a_0$. We let $\text{Coh}_{G, \mu}$ be the set of cuspidal automorphic representations π of $G(\mathbb{A})$ which are essentially tempered and cohomological of type μ . The last condition means that

$$(\pi_{\infty} \otimes_{\mathbb{C}} W_{\mu})^{G(\mathbb{R})} \neq 0.$$

Let $\pi \in \text{Coh}_{G, \mu}$. The motivic normalization of the standard L -function is given by

$$L^{\text{mot}, S}(s, \pi, \text{St}) = L^S \left(s - \frac{n-1}{2}, \pi, \text{St} \right),$$

where St stands for the L -function corresponding to the standard representation of the L -group of G , and S is a big enough finite set of places of L , included to ensure that the local base change from G (rather, the unitary group) to GL_n is defined at places outside S . We let $E(\pi)$ be a CM field containing L^{Gal} over which π_f can be realized. Such a CM field always exist (see [5], Theorem 4.4.1, and [17], 2.6). We let $\pi_{f,0}$ be a model of π_f over $E(\pi)$. We will assume from now on the following list of hypotheses.

Hypotheses 3.2.1. The representation $\pi \in \text{Coh}_{G,\mu}$ satisfies:

1. $\pi^\vee \cong \pi \otimes \|\nu\|^\xi$,
2. for any $\sigma \in J_{E(\pi)}$, $\pi_f^\sigma = \pi_{f,0} \otimes_{E(\pi),\sigma} \mathbb{C}$ is essentially tempered, and
3. $\dim_{\mathbb{C}} \text{Hom}_{\mathbb{C}[G(\mathbb{A}_f)]}(\pi_f^\sigma, H^0(S_{\mathbb{C}}, \mathcal{E}_\mu)) \leq 1$.

Hypothesis 3.2.1, (1), is assumed for simplifying purposes, and it will be satisfied in the applications of Sect. 4. In (3), \mathcal{E}_μ is the automorphic vector bundle over the Shimura variety S defined by the representation W_μ .

Remark 3.2.1. Hypotheses 3.2.1, (2) and (3) are expected to be satisfied in most cases in our applications. Conjugation of automorphic representations as in (2) is discussed in several sources. See for instance [5, 13] and [18]. Hypothesis (3) is part of Arthur’s multiplicity conjectures, a proof of which is expected to appear soon (see [22] and their forthcoming sequels).

Automorphic quadratic periods for π are defined in Sect. 3.10 of [14]. Under our running hypotheses, we can define a holomorphic quadratic period

$$Q^{\text{hol}}(\pi) \in E(\pi) \otimes \mathbb{C}.$$

We don’t need to recall the precise definition of it, but rather its interpretation as a Petersson norm. As in Remarks 3.9.2 and 3.10.1 of [14],

$$Q^{\text{hol}}(\pi) \sim_{E(\pi) \otimes L^{\text{Gal}}} \int_{G(\mathbb{Q})Z(\mathbb{A}) \backslash G(\mathbb{A})} f(g) \bar{f}(g) \|\nu(g)\|^\xi dg,$$

where f is an automorphic form on $G(\mathbb{A})$, contributing to π rationally in the sense of the canonical model of \mathcal{E}_μ over L^{Gal} .

3.3. The main formula for critical values

Here we recall the main result (Theorem 4.5.1) of [14] in the case of totally definite unitary groups. As above, we let $\pi \in \text{Coh}_{G,\mu}$, with W_μ defined over \mathbb{Q} , be an automorphic representation satisfying Hypotheses 3.2.1. Let ψ be an algebraic Hecke character of L , of infinity type $(m_\tau)_{\tau \in J_L}$, and let $m > n$ be an integer satisfying

$$m \leq a_{\tau,n} + m_\tau - m_{\bar{\tau}} \tag{3.3.1}$$

for every $\tau \in \Phi$ (this is inequality (4.2.1) of [14]). Note that n needs to satisfy the same inequality for such an integer m to exist, which poses the condition that

$m_\tau - m_{\bar{\tau}} > n - a_{\tau,n}$ for every $\tau \in \Phi$. For the time being, let $m > n$ be any integer satisfying (3.3.1). The representation $\pi \otimes (\psi \circ \det)$ will be denoted by $\pi \otimes \psi$. We let

$$\tilde{\psi} = \frac{\psi}{\psi^t}.$$

In the context of the present paper (totally definite unitary groups) Theorem 4.5.1 of [14] is basically the following result.

Proposition 3.3.1. *Let the notation and assumptions be as above. Then*

$$L^{S,\text{mot}}(m, \pi \otimes \psi, \text{St}) \sim_{E(\pi)\mathbb{Q}(\psi)} (2\pi i)^{[K:\mathbb{Q}](mn - \frac{n(n-1)}{2}) - \xi} p(\tilde{\psi}; \Phi)^n.$$

Proof. We apply the statement of the main theorem of [14] given as formula (4.5.2) of *op. cit.*, and ignoring $D_K^{n/2}$, since it already belongs to $K^{\text{Gal}} \subset L^{\text{Gal}} \subset E(\pi)$. We also fix the embeddings $1 \in J_{E(\pi)}$ and $1 \in J_{\mathbb{Q}(\psi)}$. The formula then is

$$L^{S,\text{mot}}(m, \pi \otimes \psi, \text{St}) \sim_{E(\pi)\mathbb{Q}(\psi)} (2\pi i)^{[K:\mathbb{Q}](mn - \frac{n(n-1)}{2}) - \xi} Q^{\text{hol}}(\pi)_1 p(\psi; h) p(\psi^{-1}, \bar{h}). \quad (3.3.2)$$

Here $\mathbb{S} = \text{Res}_{\mathbb{C}/\mathbb{R}} \mathbb{G}_{m,\mathbb{C}}$, where \mathbb{G}_m is the multiplicative group, and $h : \mathbb{S} \rightarrow (\text{Res}_{L/\mathbb{Q}} \mathbb{G}_{m,L})_{\mathbb{R}} \cong \prod_{\tau \in \Phi} \mathbb{S}$ is given as $h = (h_\tau)_{\tau \in \Phi}$, with $h_\tau : \mathbb{S} \rightarrow \mathbb{S}$ defined by $h_\tau(z) = z^n$ (see [14], 3.10). The map \bar{h} is given by $\bar{h}(z) = h(\bar{z})$. The elements $p(\psi; h)$ and $p(\psi^{-1}; \bar{h})$ are CM periods satisfying

$$p(\psi; h) \sim_{\mathbb{Q}(\psi)L^{\text{Gal}}} \prod_{\tau \in \Phi} p(\psi^n; \{\tau\})$$

and

$$p(\psi^{-1}; \bar{h}) \sim_{\mathbb{Q}(\psi)L^{\text{Gal}}} \prod_{\tau \in \Phi} p(\psi^{t,-n}; \{\tau\})$$

(see *op. cit.*, (3.10.3)). By Proposition 1.4 and Corollary 1.5 of [16], we can write

$$p(\psi; h) p(\psi^{-1}; \bar{h}) \sim_{\mathbb{Q}(\psi)L^{\text{Gal}}} p(\tilde{\psi}; \Phi)^n. \quad (3.3.3)$$

Finally, since the hermitian space V is totally definite, the quadratic period $Q^{\text{hol}}(\pi)$ can be taken to be in $E(\pi)$ (see for instance [20], Section 5). Thus, it can be taken to be 1 in (3.3.2), which, together with (3.3.3), proves the formula in the statement of the theorem. \square

4. The main theorems

In this section, we prove a version of Deligne’s conjecture for certain twists $M(\chi)$ of realizations M which are potentially automorphic of unitary type, in the sense that, after extending the scalars to a totally real Galois extension K'/K , they look like the motives (conjecturally) attached to self-dual automorphic representations of $\mathrm{GL}_n(\mathbb{A}_K)$. We relate this, via base change and descent, to automorphic representations of unitary groups, and apply the results of Sect. 3 to express the critical values of the corresponding L -functions. We then compare this expression with the one obtained in Sect. 2 to deduce Deligne’s conjecture for automorphic realizations (Theorem 4.3.1), after working over the Galois closure L^{Gal} and fixing embeddings of the coefficient fields. We then prove the theorem for potentially automorphic realizations (Theorem 4.4.1) by means of Brauer’s induction and the previous case.

4.1. Automorphic representations of $\mathrm{GL}_n(\mathbb{A}_K)$

Fix a totally real field K . Let Π be a cuspidal automorphic representation of $\mathrm{GL}_n(\mathbb{A}_K)$, satisfying the following properties:

- $\Pi^\vee \cong \Pi$ (self-duality), and
- Π is cohomological.

The second condition can be expressed by saying that Π_∞ has the same infinitesimal character as an irreducible representation of $(\mathrm{Res}_{K/\mathbb{Q}} \mathrm{GL}_{n,K})_{\mathbb{C}} \simeq \prod_{\sigma \in J_K} \mathrm{GL}_{n,\mathbb{C}}$. Such an irreducible representation can be parametrized, in the standard way, by a collection of integers $(a_{\sigma,1}, \dots, a_{\sigma,n})_{\sigma \in J_K}$, called the weight of Π , with $a_{\sigma,1} \geq \dots \geq a_{\sigma,n}$ for every $\sigma \in J_K$.

We let $\mathbb{Q}(\Pi_f)$ be the field of definition of Π_f . By Theorem 3.13 of [8], $\mathbb{Q}(\Pi_f)$ is a number field, and Π_f can be defined over $\mathbb{Q}(\Pi_f)$. We expect the existence of a motive $M = M(\Pi)$ over K , with coefficients in a finite extension $E(\Pi) \subset \mathbb{Q}$ of $\mathbb{Q}(\Pi_f)$, attached to Π (the reason we need to allow non-trivial extensions of $\mathbb{Q}(\Pi_f)$ is that the associated Galois representations may not be defined over the λ -adic completions of $\mathbb{Q}(\Pi_f)$; see 1.1 of [19]). The motive M should have rank n , weight $w(M) = n - 1$, and it should have the property that, for v outside a finite set of places S ,

$$L_v \left(s - \frac{n-1}{2}, \Pi^\varphi \right) = L_v(M, s)_\varphi, \quad (4.1.1)$$

where $\varphi \in J_{E(\Pi)}$. The λ -adic realizations of M have been constructed by a number of people ([6, 10, 30, 31]). Moreover, M should be polarized (see for instance [3] regarding the sign of the polarization) and regular, and the Hodge numbers are recovered from the weight of Π by the following recipe. We fix $1 \in J_{E(\Pi)}$, and we have that

$$p_i(\sigma, 1) = a_{\sigma,i} + n - i \quad (4.1.2)$$

for every $\sigma \in J_K, i = 1, \dots, n$. The other Hodge numbers $p_i(\sigma, \varphi)$ are obtained by a similar recipe by conjugating the weight of Π . We stress that the motives $M(\Pi)$ are conjectural.

Definition 4.1.1. Let M be a realization of rank n over K with coefficients in E . We say that M is *automorphic of unitary type* if it is regular, polarized, pure of weight $n - 1$, and there exists a self-dual, cohomological, cuspidal automorphic representation Π of $\mathrm{GL}_n(\mathbb{A}_K)$ such that (4.1.1) holds for $\varphi = 1$ and v outside a finite set of places S , and (4.1.2) holds. In this case we also say that M is associated with Π .

We say that M is *potentially automorphic of unitary type* if it is polarized and there exists a finite, totally real Galois extension K'/K such that $M_{K'} = M \times_K K'$ is automorphic of unitary type.

Remark 4.1.1. There exist cohomological cuspidal representations of $\mathrm{GL}_n(\mathbb{A}_K)$ which are not self-dual, and not even essentially self-dual (see [26]). The general philosophy of the Langlands program predicts the existence of motives attached to such representations as well. These motives do not fit into the framework of this paper since they are not of unitary type, in the sense that their automorphic representations do not come from unitary groups as in the self-dual case.

4.2. Transfer

Let Π be a self-dual, cohomological, cuspidal automorphic representation of $\mathrm{GL}_n(\mathbb{A}_K)$. For a totally imaginary quadratic extension L/K , let Π_L denote the base change of Π from $\mathrm{GL}_n(\mathbb{A}_K)$ to $\mathrm{GL}_n(\mathbb{A}_L)$ (see Theorems 4.2 and 5.1 of [1]). If $\Pi \not\cong \Pi \otimes \varepsilon_{L/K}$, then Π_L is cuspidal. This is always the case if n is odd, for instance. In any case, there always exists a totally imaginary quadratic extension L of the form $L = KF$ for a quadratic imaginary field F , such that Π_L is cuspidal (see Section 1 of [9]). From now on, we will fix an L such that Π_L is cuspidal. We also have that Π_L is a cohomological and $\Pi_L^\vee \cong \Pi_L^t \cong \Pi_L$.

Let G be a unitary group attached to an n -dimensional totally definite hermitian space V over L/K , as in Sect. 3. We expect the existence of a descent π of Π_L , from $\mathrm{GL}_n(\mathbb{A}_L)$ to $G(\mathbb{A})$. Actually, Π_L should descent to an L -packet of representations of $G(\mathbb{A})$, but for our purposes, we will just pick one of its members. In a significant number of cases, this has already been proved ([23]; see also [24] and [22]). For any $\tau \in J_L$, let

$$a_{\tau,i} = a_{\sigma,i},$$

where $\sigma = \tau|_K$, and let

$$\mu = ((a_{\tau,1}, \dots, a_{\tau,n})_{\tau \in \Phi}; 0).$$

We say that π is a *good descent* of Π_L (or of Π) if it is cuspidal, cohomological of type μ , and satisfies Hypotheses 3.2.1 (with W_μ defined over \mathbb{Q}). The first hypothesis, 3.2.1 (1), is easy to verify in this case (see [14], Remark 5.2.1). The

other two hypotheses are expected to hold, so that good descents are expected to exist. The condition that W_μ is defined over \mathbb{Q} is included in [14] for simplicity of notation, and it shouldn't be hard to remove.

Let ψ be an algebraic Hecke character of L , of infinity type $(m_\tau)_{\tau \in J_L}$ and weight $w = w(\psi)$. Write

$$\psi|_{\mathbb{A}_K^\times} = \psi_0 \|\cdot\|^{-w}$$

as before, with ψ_0 of finite order. Define

$$\chi = \psi^2(\psi_0 \circ N_{L/K})^{-1}.$$

Suppose that π is a descent of Π to G , and that M is an automorphic realization of unitary type associated with Π . Then, for a certain finite set of places S , we have that

$$L^{\text{mot}, S}(s - w, \pi \otimes \psi, \text{St}) = L^S(M(\chi), s)_1,$$

where $1 \in J_E$ stands for the given embedding of $E = E(\Pi)$ into \mathbb{C} (see [17], (3.5.2)).

4.3. Deligne's conjecture: the automorphic case

Keep the assumptions and notation of the last subsections. In particular, M is an automorphic realization of unitary type associated with Π . Assume that ψ is critical, and let Φ be the CM type defined by the condition

$$m_\tau > m_{\bar{\tau}} \quad (\tau \in \Phi).$$

Furthermore, assume that

$$m_\tau - m_{\bar{\tau}} > \max\{n - p_n(\sigma, 1)\}_{\sigma \in J_K} \tag{4.3.1}$$

for any $\tau \in \Phi$. Let $(n_\tau)_{\tau \in \Phi}$ be the infinity type of χ . Then $n_\tau = 2m_\tau$, and thus it satisfies Eq. (2.5.1) for M . Then, $M(\chi)$ has critical values, and the critical integers are determined by the inequalities (2.5.4), which become

$$a_{\sigma, 1} + n - 1 + 2m_{\bar{\tau}} < k \leq a_{\sigma, n} + 2m_\tau. \tag{4.3.2}$$

Note that the condition (4.3.1) implies that there always exists at least one k satisfying (4.3.2) and $k > w + n$.

At some point we need to deal with the factor $\delta(M)$. This is relatively easy to do when n is even (see Remark 2.5.3), but for the moment we need the conclusion of this remark as a hypothesis when n is odd. It can be proved assuming a much stronger hypothesis, namely Tate's conjecture for the realization M (see [14], 5.4).

Hypothesis 4.3.1. If n is odd, then $\delta(M)_1 \sim_{EK^{\text{Gal}}} (2\pi i)^{-[K:\mathbb{Q}] \frac{n(n-1)}{2}}$.

The following theorem is our main result in the case of automorphic realizations. We recall all the relevant hypotheses. We stress that assumption (3) below is expected to be satisfied in general, and most of what it involves is already proved in many cases (see [22] and its sequels). We also stress that given M , there always exist algebraic Hecke characters ψ and integers k as in the statement of the theorem.

Theorem 4.3.1. *Let M be an automorphic realization of unitary type. Let ψ be a critical algebraic Hecke character of L , of infinity type $(m_\tau)_{\tau \in J_L}$ and weight w , let Φ be the CM type defined by the condition $m_\tau > m_{\bar{\tau}}$ for $\tau \in \Phi$, and let*

$$\chi = \psi^2(\psi_0 \circ N_{L/K})^{-1}.$$

Assume that

1. either n is even, or n is odd and Hypothesis 4.3.1 is satisfied,
2. $m_\tau - m_{\bar{\tau}} > \max\{n - p_n(\sigma, 1)\}_{\sigma \in J_K}$ for any $\tau \in \Phi$, and
3. the automorphic representation Π giving rise to M has a good descent to a totally definite unitary group over L/K .

Then Conjecture 2.2.1 is true for $M(\chi)$ and all critical integers $k > w + n$, over $\mathbb{Q}(\psi)L^{\text{Gal}}$ (for the embedding $1 \in J_{E(\chi)}$). That is, for such integers k , we have

$$\frac{L(M(\chi), k)_1}{c^+(M(\chi)(k))_1} \in (E\mathbb{Q}(\psi)L^{\text{Gal}})^\times.$$

Proof. Recall that we defined

$$\check{\chi} = \chi^{\iota, -1}, \quad \check{\psi} = \frac{\psi}{\psi^\iota},$$

where $\iota \in \text{Gal}(L/K)$ is the non-trivial element. Let G be a totally definite unitary group as in the hypotheses, and let π be a good descent of Π to G , so that

$$L^{\text{mot}, S}(s - w, \pi \otimes \psi, \text{St}) = L^S(M(\chi), s)_1,$$

where $w = m_\tau + m_{\bar{\tau}}$. The remaining (finite) Euler factors, evaluated at $k \in \mathbb{Z}$, only affect this equation up to a multiple in the compositum $(E\mathbb{Q}(\chi))^\times \subset (E\mathbb{Q}(\psi))^\times$, so we may write

$$L(M(\chi), k)_1 \sim_{E\mathbb{Q}(\psi)} L^{\text{mot}, S}(m, \pi \otimes \psi, \text{St}),$$

where $m = k - w$. Our hypotheses imply that $m > n$ and

$$m \leq a_{\tau, n} + m_\tau - m_{\bar{\tau}}$$

for any $\tau \in \Phi$. This follows directly from the fact that k is a critical integer of $M(\chi)$ and $w = m_\tau + m_{\bar{\tau}}$, so that

$$k \leq a_{\sigma, n} + 2m_\tau$$

by (4.3.2). Thus, all the hypotheses of Proposition 3.3.1 are satisfied with the integer m , and we can write

$$L^{\text{mot}, S}(m, \pi \otimes \psi, \text{St}) \sim_{E\mathbb{Q}(\psi)L^{\text{Gal}}} (2\pi i)^{[K:\mathbb{Q}]\left((k-w)n - \frac{n(n-1)}{2}\right)} p(\tilde{\psi}; \Phi)^n \quad (4.3.3)$$

(note that $a_0 = 0$ and thus $\xi = 2a_0 = 0$ in this situation, and $E(\pi)$ was taken to contain L^{Gal} in Proposition 3.3.1).

Now, note that (2.5.1) is satisfied for χ , and hence, by Proposition 2.5.1, we have that

$$c^+(M(\chi)(k))_1 \sim_{E\mathbb{Q}(\chi)K^{\text{Gal}}} (2\pi i)^{[K:\mathbb{Q}]kn} \delta(M)_1 p(\check{\chi}; \Phi)^n. \quad (4.3.4)$$

By Remark 2.5.3 in the case n even, or by Hypothesis 4.3.1 in the case n odd, we can write

$$\delta(M)_1 \sim_{EK^{\text{Gal}}} (2\pi i)^{-[K:\mathbb{Q}]\frac{n(n-1)}{2}}. \quad (4.3.5)$$

Now, note that $\check{\chi} = \tilde{\psi} \parallel \cdot \parallel^w$. Using Proposition 1.4 and Lemma 1.8.3 of [16], we obtain that

$$p(\check{\chi}; \Phi) \sim_{\mathbb{Q}(\psi)} p(\tilde{\psi}; \Phi) (2\pi i)^{-[K:\mathbb{Q}]w}. \quad (4.3.6)$$

It follows by combining (4.3.4), (4.3.5) and (4.3.6), that

$$c^+(M(\chi)(k))_1 \sim_{E\mathbb{Q}(\psi)K^{\text{Gal}}} (2\pi i)^{[K:\mathbb{Q}]\left(nk-nw - \frac{n(n-1)}{2}\right)} p(\tilde{\psi}; \Phi)^n. \quad (4.3.7)$$

This is exactly the right-hand side of (4.3.3), which proves the theorem. \square

4.4. Deligne's conjecture: the potentially automorphic case

Suppose that M is a realization over K , which becomes automorphic of unitary type over K' , where K'/K is a Galois, totally real extension contained in \bar{K} . Let $M' = M_{K'}$, and let $p_i^{M'}(\sigma', \varphi)$ for $\sigma' \in J_{K'}$, $\varphi \in J_E$ and $i = 1, \dots, n$, be the Hodge numbers of M' . Then

$$p_i^{M'}(\sigma', \varphi) = p_i(\sigma'|_K, \varphi). \quad (4.4.1)$$

Let Π' be the automorphic representation of $\text{GL}_n(\mathbb{A}_{K'})$ such that M' is associated with Π' . Then Π' is cohomological of weight $(a_{\sigma', 1}, \dots, a_{\sigma', n})_{\sigma' \in J_{K'}}$, where

$$a_{\sigma', i} = p_i(\sigma|_K, 1) + i - n.$$

This follows from (4.1.2) and (4.4.1).

By Brauer-Salomon's Theorem (see [11], 15.10), there exists a finite family of intermediate fields $K \subset K_j \subset K'$ and integers $n_j \in \mathbb{Z}$ such that

- each $\text{Gal}(K'/K_j)$ is solvable, and

- we have an isomorphism

$$1_{\text{Gal}(K'/K)} \simeq \bigoplus_j n_j \text{Ind}_{\text{Gal}(K'/K_j)}^{\text{Gal}(K'/K)} 1_{\text{Gal}(K'/K_j)}. \quad (4.4.2)$$

Let $M_j = M_{K_j}$. Using the Arthur–Clozel theory of base change developed in [1], we can show that M_j is automorphic of unitary type, associated with a certain cuspidal automorphic representation Π_j of $\text{GL}_n(\mathbb{A}_{K_j})$. The proof of this fact uses cyclic base change, strong multiplicity one, and an argument of Harris (a nice explanation of this is given in [7], 1). The representation Π_j is a descent of Π' , meaning that its base change $\Pi_{j,K'}$ to K' is isomorphic to Π' . Moreover, Π_j is cohomological of weight $(a_{\sigma_j,1}^{\Pi_j}, \dots, a_{\sigma_j,n}^{\Pi_j})_{\sigma_j \in J_{K_j}}$, where

$$a_{\sigma_j,i}^{\Pi_j} = p_i(\sigma_j|_K, 1) + i - n.$$

For L/K a totally imaginary quadratic extension, we let $L_j = LK_j$ and $L' = LK'$. Then each of the extensions L_j/K_j and L'/K' is a CM extension. We fix from now on L with the property that $\Pi'_{L'}$ is cuspidal. We let \tilde{L} be the compositum of the Galois closures L_j^{Gal} . We claim that each base change Π_{j,L_j} is also cuspidal. Indeed, we can suppose that K'/K is cyclic of prime degree. If Π_{j,L_j} is not cuspidal, then $\Pi_j \cong \Pi_j \otimes \varepsilon_{L_j/K_j}$, by Theorem 4.2 of [1]. Since the base change of Π_j (resp. ε_{L_j/K_j}) to K' is Π' (resp. $\varepsilon_{L'/K'}$), this would imply that $\Pi' \cong \Pi' \otimes \varepsilon_{L'/K'}$, which would in turn imply by the same theorem that $\Pi'_{L'}$ is not cuspidal.

We now come to the main result of this paper. Hypothesis (3) below is assumed in order to apply Theorem 4.3.1 to M_j . We again stress that this is expected to hold, and it has been proved in many cases (see [22] and its sequels).

Theorem 4.4.1. *Let M be a potentially automorphic realization of unitary type over K . Let ψ be a critical algebraic Hecke character of L of infinity type $(m_\tau)_{\tau \in J_L}$, let Φ be the CM type defined by $m_\tau > m_{\bar{\tau}}$ for $\tau \in \Phi$, and let*

$$\chi = \psi^2(\psi_0 \circ N_{L/K})^{-1}.$$

Assume that:

1. either n is even, or n is odd and Hypothesis 4.3.1 is satisfied for M ,
2. $m_\tau - m_{\bar{\tau}} > \max\{n - p_n(\sigma, 1)\}_{\sigma \in J_K}$ for any $\tau \in \Phi$, and
3. for each j the automorphic representation Π_j has a good descent to a totally definite unitary group over L_j/K_j .

Then Conjecture 2.2.1 is true for $M(\chi)$ and all critical integers $k > w + n$, over $\mathbb{Q}(\psi)\tilde{L}$ (for the embedding $1 \in J_{E(\chi)}$). That is, for such integers k , we have

$$\frac{L(M(\chi), k)_1}{c^+(M(\chi)(k))_1} \in (E\mathbb{Q}(\psi)\tilde{L})^\times.$$

Proof. From (4.4.2) we deduce a formal equality between the λ -realizations of $M(\chi)$ and $\text{Res}_{K_j/K}(M(\chi)_{K_j})$, which implies the following equality of L -functions:

$$L(M(\chi), s)_1 = \prod_j L(M(\chi)_{K_j}, s)_1^{n_j}. \quad (4.4.3)$$

We denote by ψ_j the Hecke character of L_j obtained from ψ by base change. Thus,

$$\psi_j = \psi \circ N_{L_j/L}.$$

We define χ_j in the same way as χ was constructed from ψ . Then $\chi_j = \chi \circ N_{L_j/L}$. Before continuing with the proof of the theorem, we need a lemma. Recall that we use the notation $M(\chi) = M \otimes \text{Res}_{L/K}[\chi]$, $M_j = M_{K_j}$, and $M_j(\chi_j) = M_j \otimes \text{Res}_{L_j/K_j}[\chi_j]$.

Lemma 4.4.1. *We have an equality of L -functions*

$$L(M(\chi)_{K_j}, s)_1 = L(M_j(\chi_j), s)_1.$$

Proof. First note that $M(\chi)_{K_j} = M_j \otimes (\text{Res}_{L/K}[\chi])_{K_j}$. Then it is enough to verify that $(\text{Res}_{L/K}[\chi])_{K_j}$ and $\text{Res}_{L_j/K_j}[\chi_j]$ have the same λ -adic realizations for each finite place λ of $\mathbb{Q}(\chi)$. Denote by χ_λ and $\chi_{j,\lambda}$ the λ -adic realizations of χ and χ_j respectively. By definition,

$$((\text{Res}_{L/K}[\chi])_{K_j})_\lambda = \text{Ind}_{\Gamma_L}^{\Gamma_K}(\chi_\lambda) |_{\Gamma_{K_j}}$$

and

$$(\text{Res}_{L_j/K_j}[\chi_j])_\lambda = \text{Ind}_{\Gamma_{L_j}}^{\Gamma_{K_j}}(\chi_{j,\lambda}).$$

The proof of the lemma finishes by considering the following Γ_{K_j} -equivariant isomorphism:

$$\text{Ind}_{\Gamma_L}^{\Gamma_K}(\chi_\lambda) |_{\Gamma_{K_j}} \longrightarrow \text{Ind}_{\Gamma_{L_j}}^{\Gamma_{K_j}}(\chi_{j,\lambda}), \quad f \longmapsto f |_{\Gamma_{K_j}}.$$

□

Returning to the proof of the theorem, we claim now that the hypotheses of Theorem 4.3.1 are satisfied for the motive M_j , associated with Π_j , the CM extension L_j/K_j , the character ψ_j and the integer k .

Note that ψ_j has infinity type $(m_\tau^{\psi_j})_{\tau \in J_{L_j}}$, where $m_\tau^{\psi_j} = m_{\tau|_L}$. In particular, ψ_j is critical. If we let Φ_j be the CM type of L_j/K_j determined by $m_\tau^{\psi_j} > m_{\bar{\tau}}^{\psi_j}$ for $\tau \in \Phi_j$, then Φ_j consists of those embeddings $\tau \in J_{L_j}$ such that $\tau|_L \in \Phi$.

Concerning hypothesis (1) of Theorem 4.3.1, note that $\text{Res}_{K_j/K} M_j \cong M^{[K_j:K]}$, as can be easily checked. It then follows that

$$\delta(M_j)_1 \sim \delta(M)_1^{[K_j:K]}.$$

Thus, hypothesis (1) is satisfied for M_j if n is odd.

For hypothesis (2), note that if $\tau \in \Phi_j$, then $m_\tau^{\psi_j} - m_{\bar{\tau}}^{\psi_j} = m_{\tau|_L} - m_{\bar{\tau}|_L}$. Since $\tau|_L$, assumption (2) for M and ψ says that this is strictly larger than $\max\{n - p_n(\sigma, 1)\}_{\sigma \in J_K}$. Then, using relation (4.4.1) with K_j instead of K' , we have that

$$m_\tau^{\psi_j} - m_{\bar{\tau}}^{\psi_j} > \max\{n - p_n^{M_j}(\sigma, 1)\}_{\sigma \in J_{K_j}}, \quad (\tau \in \Phi_j),$$

where we are denoting the Hodge numbers of M_j by $p_i^{M_j}(\sigma, 1)$.

Hypothesis (3) of Theorem 4.3.1 for Π_j is already within our assumptions. Finally, suppose that $k > w + n$ is a critical integer for $M(\chi)$. Recall from (2.5.4) that being critical means that

$$p_1(\sigma, 1) + 2m_{\bar{\tau}} < k \leq p_n(\sigma, 1) + 2m_\tau$$

for every $\tau \in \Phi$. Then, by (4.4.1) and the fact that $m_\tau^{\psi_j} = m_{\tau|_L}$, we have that

$$p_1^{M_j}(\sigma, 1) + 2m_{\bar{\tau}}^{\psi_j} < k \leq p_n^{M_j}(\sigma, 1) + 2m_\tau^{\psi_j}$$

for every $\tau \in \Phi_j$. Thus, we are under the conditions of Theorem 4.3.1, and using Lemma 4.4.1, we obtain that

$$L(M(\chi)_{K_j}, k)_1 = L(M_j(\chi_j), k)_1 \sim_{E\mathbb{Q}(\psi)_L^{\text{Gal}}} c^+(M_j(\chi_j)(k))_1. \quad (4.4.4)$$

Now, the triple (M_j, χ_j, k) satisfies the hypotheses of Proposition 2.5.1. Thus,

$$c^+(M_j(\chi_j)(k))_1 \sim_{E\mathbb{Q}(\psi)_{K_j}^{\text{Gal}}} (2\pi i)^{[K_j:\mathbb{Q}]kn} \delta(M_j)_1 p(\check{\chi}_j; \Phi_j)^n. \quad (4.4.5)$$

From (4.4.3), (4.4.4) and (4.4.5) we obtain:

$$L(M(\chi), k)_1 \sim_{E\mathbb{Q}(\psi)\tilde{L}} (2\pi i)^{\sum_j n_j [K_j:\mathbb{Q}]kn} \prod_j \delta(M_j)_1^{n_j} \left(\prod_j p(\check{\chi}_j; \Phi_j)^{n_j} \right)^n. \quad (4.4.6)$$

From (4.4.2) we deduce that $[K:\mathbb{Q}] = \sum_j n_j [K_j:\mathbb{Q}]$, so we obtain that

$$(2\pi i)^{\sum_j n_j [K_j:\mathbb{Q}]kn} = (2\pi i)^{[K:\mathbb{Q}]kn} \quad (4.4.7)$$

As we noted above, $\delta(M_j)_1 \sim \delta(M)_1^{[K_j:K]}$. Then, from the formula $1 = \sum_j n_j [K_j:K]$, we deduce that

$$\prod_j \delta(M_j)_1^{n_j} \sim_{E\tilde{L}} \delta(M)_1. \quad (4.4.8)$$

Finally, we claim that

$$\left(\prod_j p(\check{\chi}_j; \Phi_j)^{n_j} \right) \sim_{\mathbb{Q}(\chi)\tilde{L}} p(\check{\chi}; \Phi). \quad (4.4.9)$$

Indeed, all the characters χ_j and χ are critical, and the set of critical integers is the same for all (this follows for example from (2.5.4) taking $M = \mathbb{Q}(0)$). Fix $m \in \mathbb{Z}$ such an integer. Then the results of Blasius ([4]) ((2.4.5) and Deligne's conjecture for the motives $[\chi]$ and $[\chi_j]$) imply that

$$L(\chi, m)_1 \sim_{\mathbb{Q}(\psi)\tilde{L}} (2\pi i)^{[K:\mathbb{Q}]m} p(\check{\chi}; \Phi)$$

and

$$L(\chi_j, m)_1 \sim_{\mathbb{Q}(\psi)\tilde{L}} (2\pi i)^{[K_j:\mathbb{Q}]m} p(\check{\chi}_j; \Phi_j).$$

Then (4.4.9) follows from formula (4.4.3) and Lemma 4.4.1 for $M = \mathbb{Q}(0)$.

From (4.4.6), (4.4.7), (4.4.8), (4.4.9) and Proposition 2.5.1 for the triple (M, χ, k) , we obtain that

$$L(M(\chi), k)_1 \sim_{E\mathbb{Q}(\psi)\tilde{L}} (2\pi i)^{[K:\mathbb{Q}]kn} \delta(M)_1 p(\check{\chi}; \Phi) \sim_{E\mathbb{Q}(\psi)L'} c^+(M(\chi)(k))_1,$$

which ends the proof of the theorem. \square

5. Remarks about quadratic periods

5.1. Quadratic periods and critical values

In this subsection, we draw some final remarks about quadratic periods. We plan to apply these ideas in a future project involving p -adic interpolation and p -adic L -functions. Let M be an automorphic realization of unitary type of rank n over K with coefficients in E . In this section we suppose that n is even. We will use the same notation as in Sect. 4. Let $r \in \{\frac{n}{2} + 1, \dots, n - 1\}$ and suppose that:

$$a_{\sigma,r} \equiv a_{\sigma',r}(2), \quad \text{for all } \sigma, \sigma' \in J_K. \quad (5.1.1)$$

This hypothesis is equivalent to $p_r(\sigma, 1) \equiv p_r(\sigma', 1)(2)$ for all $\sigma, \sigma' \in J_K$. As n is even and the weight of M is $n - 1$, using Proposition 2.1.1 we can construct an algebraic Hecke character $\psi^{(r,1)}$ of L of infinity type $(m_\tau)_{\tau \in J_L}$, such that

$$m_\tau - m_{\bar{\tau}} = \frac{n}{2} - p_r(\sigma, 1) = \frac{n}{2} - a_{\sigma,r} - s \quad (\tau \in \Phi).$$

As before, we let

$$\chi^{(r,1)} = (\psi^{(r,1)})^2 (\psi_0^{(r,1)} \circ N_{L/K}),$$

so that its infinity type is $(n_\tau)_{\tau \in J_L}$, with $n_\tau = 2m_\tau$ and

$$n_\tau - n_{\bar{\tau}} = n - 2p_r(\sigma, 1) = n - 2s - 2a_{\sigma,r} \quad (\tau \in \Phi).$$

Let $\sigma \in J_K$ and suppose that

$$a_{\sigma,r+1} \equiv a_{\sigma,r} + 1(2), \quad \text{for all } r \in \left\{ \frac{n}{2} + 1, \dots, n - 1 \right\}. \quad (5.1.2)$$

This condition allows us to choose $\psi^{(\frac{n}{2}+1,1)}, \dots, \psi^{(n-1,1)}$ such that each $\chi^{(r,1)}$, for $r = \frac{n}{2} + 1, \dots, n - 1$, has weight w_0 , independent of r , and satisfying:

- (a) $w_0 \equiv n(2)$,
- (b) $2r - n$ is a divisor of w_0 for any $r \in \{\frac{n}{2} + 1, \dots, n - 1\}$, and
- (c) 4 is a divisor of $w_0 + n$.

Lemma 5.1.1. *Let $r \in \{\frac{n}{2} + 1, \dots, n - 1\}$. Then the only integer $m \in \mathbb{Z}$ which is critical for $M(\chi^{(r,1)})$ is $m = \frac{n+w_0}{2}$.*

Proof. Recall that in this case we have $p_r(\sigma, 1) = a_{\sigma,r} + n - r$ for each $\sigma \in J_K$. Now we apply the formulas obtained in Sect. 5.3 of [14] which determine the set of critical integers, considering the following facts: (i) $a_{\sigma,1} \geq \dots \geq a_{\sigma,n}$, (ii) $a_{\sigma,r} + a_{\sigma,n-r+1} = 0$ and (iii) $m_{\tau}^{(r,1)} - m_{\bar{\tau}}^{(r,1)} = \frac{n-2p_r(\sigma,1)}{2}$. \square

Let F be the number field generated by the image of $\chi^{(r,1)}$ on the finite adèles $\mathbb{A}_{L,f}^{\times}$ for $r = \frac{n}{2} + 1, \dots, n - 1$ (that is, we take the compositum of the $\mathbb{Q}(\chi^{(r,1)})$ for $r = \frac{n}{2} + 1, \dots, n - 1$).

Proposition 5.1.1. *Let $j \in \{2, \dots, \frac{n}{2} - 1\}$. Suppose that Conjecture 2.2.1 is true for $M(\chi^{(n-j,1)})$ and $M(\chi^{(n-j+1,1)})$ for the critical integer $\frac{n+w_0}{2}$, over FK^{Gal} (for the embedding 1). Then*

$$\frac{L(M(\chi^{(n-j,1)}), \frac{n+w_0}{2})_1}{L(M(\chi^{(n-j+1,1)}), \frac{n+w_0}{2})_1} \stackrel{Q_{j,1} \sim_{EFK^{\text{Gal}}}}{=} \frac{L(\chi^{(n-j+1,1)}, -\frac{w_0(j-1)}{n-2(j-1)})^{n-2(j-1)}}{L(\chi^{(n-j,1)}, -\frac{w_0 j}{n-2j})^{n-2j}} \frac{G(\chi^{(n-j+1,1)})_1^{j-1}}{G(\chi^{(n-j,1)})_1^j}$$

Proof. First note that for each $r = 0, \dots, n$, by Deligne's conjecture for algebraic Hecke characters (proved by Blasius in [4]) and Lemma 2.4.1, we obtain that

$$P(\chi^{(r,1)})_1 \sim_{\mathbb{Q}(\chi^{(r,1)})L^{\text{Gal}}} L\left(\chi^{(r,1)}, -\frac{w_0(n-r)}{2r-n}\right)^{2r-n} G(\chi)_1^{n-r}.$$

Then the proof finishes by using this formula, Corollary 2.4.1 and the hypothesis. \square

Remark 5.1.1. When $j = 1$, we deduce an analogous formula to that of Proposition 5.1.1.

Remark 5.1.2. In Sect. 3.10 of [14], the author defines *automorphic quadratic periods*. A particular case of this is the period denoted by $Q^{\text{hol}}(\pi)$ in Sect. 3 above, but in arbitrary signatures they can be defined for automorphic forms contributing in coherent cohomology to other non-holomorphic degrees. Proposition 5.1.1, combined with the comparisons in 5.4 of *op. cit.*, suggests an expression for these periods in terms of critical values of automorphic L -functions. This formula should be useful, for example, in questions on p -adic interpolation related with automorphic quadratic periods.

Acknowledgements. The second author wants to thank Adrian Iovita and the Centre de Recherches Mathématiques in Montréal for supporting his visit in Spring 2016, during which this work was carried out. The authors would also like to thank the referee for helpful comments and suggestions.

Open Access This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

References

- [1] Arthur, J., Clozel, L.: Simple Algebras, Base Change, and the Advanced Theory of the Trace Formula, *Annals of Mathematics Studies*, vol. 120. Princeton University Press, Princeton (1989)
- [2] Barnet-Lamb, T., Gee, T., Geraghty, D., Taylor, R.: Potential automorphy and change of weight. *Ann. Math.* **179**(2), 501–609 (2014)
- [3] Bellaïche, J., Chenevier, G.: The sign of Galois representations attached to automorphic forms for unitary groups. *Compos. Math.* **147**(5), 1337–1352 (2011)
- [4] Blasius, D.: On the critical values of Hecke L -series. *Ann. Math.* **124**(1), 23–63 (1986)
- [5] Blasius, D., Harris, M., Ramakrishnan, D.: Coherent cohomology, limits of discrete series, and Galois conjugation. *Duke Math. J.* **73**(3), 647–685 (1994)
- [6] Chenevier, G., Harris, M.: Construction of automorphic Galois representations, II. *Camb. J. Math.* **1**(1), 53–73 (2013)
- [7] Clozel, L.: The Sato-Tate Conjecture, *Current Developments in Mathematics*, pp. 1–34. Int. Press, Somerville (2008)
- [8] Clozel, L.: Motifs et formes automorphes: applications du principe de fonctorialité, *Automorphic forms, Shimura varieties, and L -functions*, Vol. I (Ann Arbor, 1988) *Perspect. Math.*, vol. 10. Academic Press, Boston, pp. 77–159 (1990)
- [9] Clozel, L.: Purity reigns supreme. *Int. Math. Res. Not. IMRN* **2013**(2), 328–346 (2013)
- [10] Clozel, L., Harris, M., Labesse, J.-P.: Construction of Automorphic Galois Representations, I, On the Stabilization of the Trace Formula, *Stab. Trace Formula Shimura Var. Arith. Appl.*, vol. 1, Int. Press, Somerville, MA, pp. 497–527 (2011)
- [11] Curtis, C.W., Reiner, I.: *Methods of Representation Theory*, vol. I. Wiley, New York (1981). (With applications to finite groups and orders. A Wiley-Interscience Publication, Pure and Applied Mathematics)
- [12] Deligne, P.: Valeurs de fonctions L et périodes d'intégrales, *Automorphic forms, representations and L -functions*, Part 2, *Proc. Sympos. Pure Math.*, XXXIII, Amer. Math. Soc., Providence, pp. 313–346 (1979)
- [13] Gan, W.T., Raghuram, A.: Arithmeticity for periods of automorphic forms, *Automorphic representations and L -functions*, *Tata Inst. Fundam. Res. Stud. Math.*, vol. 22, Tata Inst. Fund. Res., Mumbai, pp. 187–229 (2013)
- [14] Guerberoff, L.: Period relations for automorphic forms on unitary groups and critical values of L -functions, to appear in *Documenta Mathematica*
- [15] Harder, G., Schappacher, N.: Special values of Hecke L -functions and abelian integrals. *Workshop Bonn 1984* (Bonn, 1984), *Lecture Notes in Math.*, vol. 1111, Springer, Berlin, pp. 17–49 (1985)
- [16] Harris, M.: L -functions of 2×2 unitary groups and factorization of periods of Hilbert modular forms. *J. Amer. Math. Soc.* **6**(3), 637–719 (1993)
- [17] Harris, M.: L -functions and periods of polarized regular motives. *J. Reine Angew. Math.* **483**, 75–161 (1997)
- [18] Harris, M.: Beilinson-Bernstein localization over \mathbb{Q} and periods of automorphic forms. *Int. Math. Res. Not. IMRN* **2013**(9), 2000–2053 (2013) (MR3053412)

- [19] Harris, M.: L -functions and periods of adjoint motives. *Algeb. Number Theory* **7**(1), 117–155 (2013)
- [20] Harris, M.: p -adic and analytic properties of period integrals and values of L -functions. *Annales mathématiques du Québec* **40**(2), 435–452 (2016)
- [21] Harris, M., Kudla, S.S.: The central critical value of a triple product L -function. *Ann. Math.* **133**(3), 605–672 (1991)
- [22] Kaletha, T., Mínguez, A., Shin, S.W., White, P.-J.: Endoscopic classification of representations: inner forms of unitary groups. [arXiv:1409.3731v3](https://arxiv.org/abs/1409.3731v3) [math.NT]
- [23] Labesse, J.-P.: Changement de base CM et séries discrètes, On the stabilization of the Trace Formula, *Stab. Trace Formula Shimura Var. Arith. Appl.*, vol. 1, Int. Press, Somerville, pp. 429–470 (2011)
- [24] Mok, C.P.: Endoscopic classification of representations of quasi-split unitary groups. *Mem. Amer. Math. Soc.* **235**(1108), vi+248 (2015)
- [25] Alexei, A.: Panchishkin, motives over totally real fields and p -adic L -functions. *Ann. Inst. Fourier (Grenoble)* **44**(4), 989–1023 (1994)
- [26] Ramakrishnan, D., Wang, S.: A cuspidality criterion for the functorial product on $GL(2) \times GL(3)$ with a cohomological application. *Int. Math. Res. Not. IMRN* **2004**(27), 1355–1394 (2004)
- [27] Schappacher, N.: Periods of Hecke characters. *Lecture Notes in Math*, vol. 1301. Springer-Verlag, Berlin (1988)
- [28] Shimura, G.: The special values of the zeta functions associated with cusp forms. *Comm. Pure Appl. Math.* **29**(6), 783–804 (1976)
- [29] Shimura, G.: The special values of the zeta functions associated with Hilbert modular forms. *Duke Math. J.* **45**(3), 637–679 (1978)
- [30] Shin, S.W.: Galois representations arising from some compact Shimura varieties. *Ann. Math.* **173**(3), 1645–1741 (2011)
- [31] Sorensen, C.M.: A patching lemma. manuscript available at <http://www.imj-prg.fr/fa/spip.php?article10>
- [32] Yoshida, H.: On the zeta functions of Shimura varieties and periods of Hilbert modular forms. *Duke Math. J.* **75**(1), 121–191 (1994)
- [33] Yoshida, H.: Motives and Siegel modular forms. *Amer. J. Math.* **123**(6), 1171–1197 (2001)