

**HIV MEDICINE**British HIV Association  
**BHIVA**

## Pregnancies in older women living with HIV in the United Kingdom and Ireland

Journal:	<i>HIV Medicine</i>
Manuscript ID	HIV-SR-05-2016-3464.R1
Manuscript Type:	Short communication
Date Submitted by the Author:	n/a
Complete List of Authors:	Townsend, Claire; University College London, Institute of Child Health de Ruitter, Annemiek; Guy's and St Thomas' NHS Foundation Trust, Department of GUM and HIV; Peters, Helen; University College London, National Study of HIV in Pregnancy and Childhood Nelson-Piercy, Catherine; Guy's and Saint Thomas' NHS Foundation Trust Tookey, Pat; UCL Institute of Child Health, PPP Programme Thorne, Claire; UCL Institute of Child Health, University College London, UK, MRC Centre of Epidemiology for Child Health,
Keywords:	HIV, maternal age, stillbirth, surveillance, pregnancy outcomes

SCHOLARONE™  
Manuscripts

Review

1  
2  
3 **Pregnancies in older women living with HIV in the United Kingdom and Ireland**  
4  
5  
6

7 Claire Louise Townsend PhD<sup>1</sup>, Annemiek de Ruiter FRCP<sup>2</sup>, Helen Peters MSc<sup>1</sup>, Cathy Nelson-Piercy MA

8  
9 FRCP FRCOG<sup>2</sup>, Pat Tookey PhD<sup>1</sup>, Claire Thorne PhD<sup>1</sup>  
10

11 <sup>1</sup> Population, Policy and Practice Programme, UCL Institute of Child Health, University College

12  
13 London, London, United Kingdom  
14

15 <sup>2</sup> Guys & St Thomas' NHS Foundation Trust, London, United Kingdom  
16  
17

18  
19  
20 **Corresponding Author:** Dr Claire Townsend, Population, Policy, and Practice Programme, UCL  
21

22 Institute of Child Health, 30 Guilford Street, London WC1N 1EH, United Kingdom  
23

24 Email: [c.townsend@ucl.ac.uk](mailto:c.townsend@ucl.ac.uk); [ctownsend156@gmail.com](mailto:ctownsend156@gmail.com), Tel: +33 770 31 95 66  
25  
26

27 **Alternative contact:** Dr Claire Thorne, Population, Policy, and Practice Programme, UCL Institute of  
28

29 Child Health, 30 Guilford Street, London WC1N 1EH, United Kingdom  
30

31 Email: [claire.thorne@ucl.ac.uk](mailto:claire.thorne@ucl.ac.uk), Tel: +44 20 7905 2105  
32  
33

34  
35 **Word count:** 2157  
36

37 **Running head:** Pregnancies in older women living with HIV  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Abstract**

Objectives: To compare maternal characteristics and pregnancy outcomes in women aged <40 years and ≥40 years in a large unselected population of HIV-positive women delivering in the UK and Ireland between 2000 and 2014.

Methods: Comprehensive population-based surveillance data on all HIV-positive pregnant women and their children seen for care in the UK and Ireland are collected through the National Study of HIV in Pregnancy and Childhood. All singleton and multiple pregnancies reported by the end of June 2015 resulting in live birth or stillbirth to women diagnosed before delivery and delivering in 2000-2014 were included. Logistic regression models were fitted in analyses examining the association between older maternal age and specific outcomes (preterm delivery and stillbirth).

Results: Among 15,501 pregnancies in HIV-positive women, the proportion in older women (≥40 years) increased from 2.1% (73/3419) in 2000-2004 to 8.9% (510/5748) in 2010-2014 ( $p<0.001$ ). Compared with pregnancies in younger women, those in older women were more likely to result in multiple birth (3.0% versus 1.9%,  $p=0.03$ ), stillbirth (adjusted odds ratio 2.39,  $p=0.004$ ) or an infant with a chromosomal abnormality (1.6% versus 0.2%,  $p<0.001$ ). However, there was no increased risk of preterm delivery, low birth weight or mother-to-child HIV transmission among older mothers.

Conclusions: There has been a significant increase over time in the proportion of deliveries to women living with HIV aged 40 years or more, which has implications for pregnancy management, given their increased risk of multiple births, stillbirth and chromosomal anomalies, as also apparent in the general population.

**Keywords:** HIV; pregnancy outcomes; maternal age; stillbirth; surveillance

## Introduction

Pregnancy incidence among women living with HIV has increased in recent years (1, 2) alongside increases in life expectancy (3) and a decline in mother-to-child HIV transmission (MTCT) rates to less than 1% in some regions (4, 5). In many countries in North America and Europe, including the United Kingdom (UK), maternal age has been increasing both in the general obstetric population (6) and among HIV-positive women (7). Advanced maternal age in pregnancy (over 35 or 40 years) is associated with an increased risk of obstetric complications including pre-eclampsia, gestational diabetes, preterm birth and stillbirth (8, 9). In addition, both HIV infection and antiretroviral therapy in pregnancy are themselves risk factors for adverse pregnancy outcomes such as preterm birth (10). There is to date little research on the small but increasing group of women with both HIV- and age-related risks during pregnancy (11, 12). One small study in the United States found a significantly higher rate of preterm delivery among HIV-positive (41%) compared with HIV-negative (17%) women aged over 35 years (12). High rates of illicit drug use were also reported in the study population (28% and 14% respectively), although the association between HIV infection status and preterm delivery remained after adjusting for this. A larger study in Italy (11) found higher rates of preterm delivery and low birth weight in HIV-positive mothers  $\geq 35$  years of age compared with mothers  $< 35$  years, but the association was not significant in multivariable analysis. The authors reported no differences by maternal age in rates of congenital abnormalities or other pregnancy complications (e.g. hypertension, diabetes or anaemia) or delivery complications (11).

Our objective was to compare maternal characteristics and pregnancy outcomes in women aged  $< 40$  years and  $\geq 40$  years in a large unselected population of HIV-positive women delivering in the UK and Ireland between 2000 and 2014.

## Methods

Comprehensive population-based surveillance data on all HIV-positive pregnant women and their children seen for care in the UK and Ireland are collected through the National Study of HIV in Pregnancy and Childhood (NSHPC) (4, 7). Information on pregnancy and delivery are collected from obstetric respondents in each maternity unit, and HIV-exposed infants are followed up through their paediatricians to establish infection status. Full methodological details are described elsewhere (4, 7). All singleton and multiple pregnancies reported by the end of June 2015 resulting in live birth or stillbirth to HIV-positive women diagnosed before delivery and delivering between 2000 and 2014 were included.

UK guidelines recommend that pregnant women should book for antenatal care by around 10-13 weeks gestation and that all women should routinely be offered ultrasound screening for fetal anomalies at 18-20 weeks gestation. Screening policies for Down's Syndrome have varied over time; in 2003 it was recommended that all women should be offered screening, and since 2008, that this should be performed by the end of the first trimester.

Maternal age was categorised as <40 ("younger women") or  $\geq 40$  years ("older women") at the time of delivery, as numbers were sufficiently large (over 850 women  $\geq 40$  years) to focus on this higher risk group. Year of delivery was grouped into five-year periods: 2000-2004, 2005-2009, and 2010-2014. Mode of delivery was classified as elective caesarean section (pre-labour, pre-rupture of membranes), emergency caesarean section (after rupture of membranes and/or onset of labour, or for other emergency obstetric indications), or vaginal delivery. Information on operative vaginal delivery was only collected from July 2008 onwards. Delivery at <37 completed weeks was classified as preterm. Maternal HIV viral load was measured closest to delivery (30 days before or up to seven days postpartum) and categorised as <50 (undetectable), 50-399, 400-999, 1000-9999, and  $\geq 10,000$  copies/ml. Maternal CD4 cell count closest to delivery was classified as <200, 200-349, 350-499 and  $\geq 500$  cells/mm<sup>3</sup>. Congenital abnormalities were classified according to the World Health

1  
2  
3 Organization International Classification of Diseases version 10 (13). Infant HIV infection status was  
4 established based on PCR tests carried out after one month of age (4)

5  
6  
7 Data were managed in Access 2010 (Microsoft Corp., Redmond, Washington, USA) and analysed  
8 using Stata version 12.1 (Stata Corp. LP, College Station, Texas, USA). Means were compared using *t*-  
9 tests and medians using Wilcoxon tests. Trends in medians were assessed using Cuzick's  
10 nonparametric test for trend across ordered groups (function 'nptrend' in Stata). Logistic regression  
11 models were fitted to obtain odds ratios (ORs) and 95% confidence intervals (CIs) in analyses  
12 examining the association between older maternal age and a) preterm delivery and b) stillbirth. In  
13 multivariable analyses, variables were included in the model if they were considered potential  
14 confounders and significantly improved the model's goodness of fit (significance level  $p < 0.10$ ). To  
15 allow for repeat pregnancies in the same woman in preterm delivery analyses only (as <5 women  
16 experienced more than one stillbirth), generalized linear mixed effects were used to fit logistic  
17 regression models accounting for random effects attributed to the mother (14).

18  
19  
20 The National Study of HIV in Pregnancy and Childhood has London Multi-Centre Research Ethics  
21 Committee approval (MREC/04/2/009).

## 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 **Results**

40 Overall 15,501 pregnancies in 10,997 women delivering in 2000-2014 were included, 856 (5.5%) of  
41 which were in women aged  $\geq 40$  years at delivery. Most pregnancies (79%) were in women from sub-  
42 Saharan Africa, and only 2% in women who acquired HIV through injecting drug use (Table 1). First  
43 antenatal appointment (collected since 2008) occurred at a median of 12.4 weeks gestation  
44 (interquartile range (IQR) 10.1, 16.0,  $n=7154$  pregnancies). Median maternal age was 31.5 years (IQR  
45 27.6, 35.4, range 14.5, 50.6), increasing over time from 29.6 years in 2000-2004, to 31.0 years in  
46 2005-2009, and 33.2 years in 2010-2014 (trend  $p < 0.001$ ). The proportion of pregnancies in older  
47 women ( $\geq 40$  years at delivery) also increased significantly over time, from 2.1% (73/3419) in 2000-  
48 2004, to 4.3% (273/6334) in 2005-2009, and 8.9% (510/5748) in 2010-2014 (trend  $p < 0.001$ ).

1  
2  
3 Similar patterns were observed among pregnancies in primiparous women ( $n=4408$ ), with an  
4  
5 increase in those  $\geq 40$  years from 0.9% (10/1112) in 2000-2004, to 2.0% (39/1923) in 2005-2009, and  
6  
7 5.2% (71/1373) in 2010-2014 (trend  $p<0.001$ ) (2.7% overall, 2000-2014). Median maternal age  
8  
9 among pregnancies in primiparous women was 28.8 years, increasing from 27.2 years in 2000-2004  
10  
11 to 28.5 years in 2005-2009 and 30.6 years in 2010-2014 (trend  $p<0.001$ ).

12  
13 Compared with pregnancies in younger women, those in older women were more likely to be in  
14  
15 women with three or more previous births, and in those diagnosed with HIV before their pregnancy  
16  
17 (Table 1). There were no differences by age group in maternal region of birth, HIV risk factor  
18  
19 (injecting drug use or other/unknown), or type of combination antiretroviral therapy (cART) in  
20  
21 pregnancy (Table 1).  
22  
23

24  
25 However, treatment in pregnancy, and at conception, was more likely for pregnancies in older  
26  
27 women, and cART was started slightly earlier compared with pregnancies in younger women  
28  
29 (median week of cART initiation 22.8 versus 23.5 weeks,  $p=0.02$ ), even though there were no  
30  
31 differences in timing of first antenatal appointment (median 14 weeks for both groups,  $p=0.88$ ).

32  
33 Pregnancies in older women were more commonly associated with suppressed viral load at delivery  
34  
35 than those in younger women, and with low CD4 count ( $<200$  cells/ $\mu\text{l}$ ) (Table 1), although median  
36  
37 CD4 cell count did not differ by maternal age group: 431 cells/ $\mu\text{l}$  among younger women versus 445  
38  
39 cells/ $\mu\text{l}$  among older women ( $p=0.19$ ). The viral load difference was restricted to women starting  
40  
41 treatment in pregnancy (viral load was suppressed in 74.9% (143/191) of pregnancies in older versus  
42  
43 63.3% (3506/5536) of those in younger women), and was not apparent where women were on  
44  
45 treatment at conception ( $p=0.71$ ).  
46  
47

48  
49 There was no evidence of an increased rate of emergency caesarean section or operative vaginal  
50  
51 delivery among pregnancies in older compared with younger women, but the risk of multiple births  
52  
53 and an infant with a chromosomal abnormality was higher (Table 2). There was no increased risk of  
54  
55 preterm delivery ( $<37$ ,  $<34$  or  $<32$  weeks) or low birth weight among deliveries to older compared  
56  
57 with younger women (Table 2); there was a suggestion that infants of older women were more likely  
58  
59  
60

1  
2  
3 to have very low (<1.5 kg) or high ( $\geq 4$  kg) birth weight but neither association reached statistical  
4  
5 significance. In multivariable analysis of singleton pregnancies adjusting for time period, injecting  
6  
7 drug use (reported as the route of maternal HIV acquisition) and type of antiretroviral therapy (cART  
8  
9 versus 1-2 antiretroviral drugs, excluding untreated pregnancies), maternal age  $\geq 40$  years was not  
10  
11 significantly associated with preterm delivery (<37 weeks) (adjusted OR (AOR) 1.21, 95% CI 0.92,  
12  
13 1.60,  $p=0.18$ ). All other variables included in the model were significantly associated with the  
14  
15 outcome ( $p<0.05$ ).  
16

17  
18 More pregnancies in older mothers ended in stillbirth (1.6% versus 1.0%), although this did not reach  
19  
20 statistical significance in univariable analysis (OR 1.70, 95% CI: 0.99-3.01,  $p=0.05$ ). After adjusting for  
21  
22 time period, parity (0, 1+ previous births) and type of antiretroviral therapy (as above), the strength  
23  
24 of this association increased and was statistically significant (AOR 2.39, 95% CI 1.32, 4.32,  $p=0.004$ );  
25  
26 in this model, the only other variable significantly associated with the outcome was time period.  
27  
28 There was no difference in the MTCT rate according to maternal age (Table 2). Thirteen maternal  
29  
30 deaths were reported, but none were in older mothers.  
31  
32  
33  
34

### 35 Discussion

36  
37 In this population-based surveillance study of HIV-positive pregnant women, we observed a  
38  
39 significant increase over time in median maternal age, and in the proportion of deliveries to women  
40  
41 aged over 40 years, such that in the most recent time period, 9% of all deliveries were to women  
42  
43 aged  $\geq 40$  years. Compared with pregnancies in younger HIV-positive women (<40 years), those in  
44  
45 older women were significantly more likely to result in multiple births and an infant with a  
46  
47 chromosomal abnormality. Pregnancies in older women were also more likely to result in stillbirth  
48  
49 (1.6% versus 1.0%); although not statistically significant in univariable analysis, it was significant  
50  
51 after adjusting for time period, parity and cART. There was no evidence that older maternal age was  
52  
53 associated with an increased risk of preterm delivery, low birth weight or MTCT in this population.  
54  
55  
56  
57  
58  
59  
60



1  
2  
3 Our findings are consistent with studies in the general population showing an association between  
4 advanced maternal age and outcomes such as multiple pregnancy (15), chromosomal abnormalities  
5 (16), and stillbirth (8, 17). Data on multiple maternity rates in England and Wales for 2014 are  
6 consistent with our results, with a rate of 2.9% for women aged 40-44 years (compared with 3.0%  
7 here) and ranging from 0.96% for 20-24 year olds to 2.3% for 35-39 year olds (compared with 1.9%  
8 here for <40 year olds) (18). Higher twinning rates in older women may reflect a biological  
9 propensity for multiple births as well as higher rates of fertility treatment; we were unable to  
10 explore the contribution of fertility treatment to the multiple birth rate here as we do not collect this  
11 information.  
12

13 The doubling of risk of stillbirth associated with maternal age over 40 years is of a similar magnitude  
14 to that estimated through meta-analysis of five studies in the general obstetric population in high  
15 income countries (8). The stillbirth rates in HIV-positive women were higher than in the general  
16 population (0.47% overall, 0.76% for 40-44 year olds and 0.95% for  $\geq 45$  year olds), with the greatest  
17 difference seen for younger women (18). However, unlike other studies (17, 19) we did not observe  
18 an increased risk of preterm delivery, although the rate was higher overall than in the general  
19 population (13.2% versus 6.2%) (20). It was also reassuring that rates of emergency caesarean  
20 section and operative vaginal deliveries were not significantly increased in older mothers in this  
21 study.  
22

23 Overall, most pregnancies were in women already aware of their HIV status from before conception,  
24 and the greater proportion of older women in this group likely reflects greater opportunity for HIV  
25 testing, particularly in previous pregnancies, given their higher parity. As a result, a higher  
26 proportion of pregnancies in older women were conceived on cART, whilst those in which cART was  
27 started in pregnancy were more likely to have an undetectable viral load at delivery; this latter  
28 finding is consistent with other studies and may reflect better adherence in older people (21) as well  
29 as more effective and less toxic antiretroviral drugs in later calendar years, when most pregnancies  
30 in older women occurred.  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 The NSHPC is a well-established, ongoing surveillance study which aims to include all pregnancies in  
4 HIV-positive women in the UK and Ireland and has high response rates. However, limited data are  
5 collected on background characteristics and clinical measures in pregnancy, and we had no  
6 information on factors such as smoking, or history of complications such as hypertension or delivery  
7 complications. Our findings are likely generalizable to other Western European HIV pregnancy  
8 cohorts which have similarly high proportions of sub-Saharan African migrants (22,23).  
9  
10 This report highlights an increase over time in pregnancies among HIV-positive women aged 40 years  
11 or more. These findings have implications for pregnancy management of older HIV-positive women,  
12 given the increased risk of multiple births in this group, as well as pre-existing co-morbidities and  
13 adverse outcomes such as stillbirth and chromosomal anomalies, as has been reported in the HIV-  
14 negative population.  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Acknowledgements**

National surveillance of obstetric and paediatric HIV is undertaken through the National Study of HIV in Pregnancy and Childhood (NSHPC) in collaboration with the Health Protection Agency Centre for Infections, now Public Health England, and Health Protection Scotland. The authors gratefully acknowledge the contribution of the midwives, obstetricians, genitourinary physicians, paediatricians, clinical nurse specialists, and all other colleagues who report to the NSHPC through the British Paediatric Surveillance Unit of the Royal College of Paediatrics and Child Health, and the obstetric reporting scheme run under the auspices of the Royal College of Obstetricians and Gynaecologists. We also thank Icina Shakes (former Study Assistant), Kate Francis, Rebecca Sconza and Anna Horn (Study Assistants/Research Assistants) for their essential contributions to the NSHPC.

The National Study of HIV in Pregnancy and Childhood receives core funding from Public Health England, with additional funding from the Public Health England National Health Service Infectious Diseases in Pregnancy Screening Programme. The UCL Institute of Child Health receives a proportion of funding from the Department of Health's National Institute for Health Research Biomedical Research Centres funding scheme.

**Contribution to authorship**

CLT carried out the statistical analyses. CLT, AdR and CT drafted the paper, and all authors contributed to interpreting the results, and critically revising the article. CT is responsible for the NSHPC and is the guarantor.

## Reference List

1. French CE, Cortina-Borja M, Thorne C, *et al.* Incidence, patterns, and predictors of repeat pregnancies among HIV-infected women in the United Kingdom and Ireland, 1990-2009. *J Acquir Immune Defic Syndr.* 2012; **59**:287-293.
2. Huntington SE, Thorne C, Bansi LK, *et al.* Predictors of pregnancy and changes in pregnancy incidence among HIV-positive women accessing HIV clinical care. *AIDS.* 2013; **27**:95-103.
3. Samji H, Cescon A, Hogg RS, *et al.* Closing the gap: increases in life expectancy among treated HIV-positive individuals in the United States and Canada. *PLoS One.* 2013; **8**:e81355.
4. Townsend CL, Byrne L, Cortina-Borja M, *et al.* Earlier initiation of ART and further decline in mother-to-child HIV transmission rates, 2000-2011. *AIDS.* 2014; **28**:1049-1057.
5. Nesheim S, Harris LF, Lampe M. Elimination of perinatal HIV infection in the USA and other high-income countries: achievements and challenges. *Curr Opin HIV AIDS.* 2013; **8**:446-455.
6. Office for National Statistics. *Characteristics of Mother 1, England and Wales, 2013.* 2014. Available at: <http://www.ons.gov.uk/ons/rel/vsob1/characteristics-of-Mother-1--england-and-wales/2013/stb-characteristics-of-mother-1--2013.html>.
7. Townsend CL, Cortina-Borja M, Peckham CS, *et al.* Trends in management and outcome of pregnancies in HIV-infected women in the UK and Ireland, 1990-2006. *BJOG.* 2008;**115**:1078-1086.

- 1  
2  
3 8. Flenady V, Koopmans L, Middleton P, *et al.* Major risk factors for stillbirth in high-income  
4  
5 countries: a systematic review and meta-analysis. *Lancet*. 2011;**377**:1331-1340.  
6  
7
- 8  
9 9. Luke B, Brown MB. Elevated risks of pregnancy complications and adverse outcomes with  
10  
11 increasing maternal age. *Hum Reprod*. 2007;**22**:1264-1272.  
12  
13
- 14  
15 10. Short CE, Taylor GP. Antiretroviral therapy and preterm birth in HIV-infected women. *Expert*  
16  
17 *Rev Anti Infect Ther*. 2014;**12**:293-306.  
18  
19
- 20  
21 11. Liuzzi G, Pinnetti C, Floridia M, *et al.* Pregnancy outcomes in HIV-infected women of  
22  
23 advanced maternal age. *HIV Clin Trials*. 2013;**14**:110-119.  
24  
25
- 26  
27 12. Brown K, Holland B, Mosquera C, *et al.* Human immunodeficiency virus infection in advanced  
28  
29 maternal age gravidas. *AIDS Res Hum Retroviruses*. 2012;**28**:265-269.  
30  
31
- 32  
33 13. World Health Organization. *International Statistical Classification of Diseases and Related*  
34  
35 *Health Problems*. Geneva: WHO; 1992.  
36  
37
- 38  
39 14. Rabe-Hesketh S, Skrondal A, Pickles A. Reliable estimation of generalized linear mixed  
40  
41 models using adaptive quadrature. *The Stata Journal*. 2002;**2**:1-21.  
42  
43
- 44  
45 15. Bortolus R, Parazzini F, Chatenoud L, *et al.* The epidemiology of multiple births. *Hum Reprod*  
46  
47 *Update*. 1999; **5**:179-187.  
48  
49
- 50  
51 16. Hook EB, Cross PK, Schreinemachers DM. Chromosomal abnormality rates at amniocentesis  
52  
53 and in live-born infants. *JAMA*. 1983; **249**:2034-2038.  
54  
55  
56  
57  
58  
59  
60

- 1  
2  
3 17. Kenny LC, Lavender T, McNamee R, *et al.* Advanced maternal age and adverse pregnancy  
4  
5 outcome: evidence from a large contemporary cohort. *PLoS One*. 2013;**8**:e56583.  
6  
7  
8  
9 18. Office for National Statistics. Statistical Bulletin: Birth characteristics in England and Wales:  
10  
11 2014.  
12  
13 <http://www.ons.gov.uk/peoplepopulationandcommunity/birthsdeathsandmarriages/livebirths/bulletins/birthcharacteristicsinenglandandwales/2015-10-08>  
14  
15  
16  
17  
18  
19 19. Laopaiboon M, Lumbiganon P, Intarut N, *et al.* Advanced maternal age and pregnancy  
20  
21 outcomes: a multicountry assessment. *BJOG*. 2014;**121** Suppl 1:49-56.  
22  
23  
24  
25  
26 20. Moser K, Stanfield KM, Leon DA. Birthweight and gestational age by ethnic group, England  
27  
28 and Wales 2005: introducing new data on births. *Health Stat Q*. 2008(**39**):22-55.  
29  
30  
31  
32 21. Sherr L, Lampe FC, Clucas C, *et al.* Self-reported non-adherence to ART and virological  
33  
34 outcome in a multiclinic UK study. *AIDS Care*. 2010;**22**:939-945.  
35  
36  
37  
38  
39 22. Mandelbrot L, Tubiana R, Le Chenadec J, *et al.* No perinatal HIV-1 transmission from women  
40  
41 with effective antiretroviral therapy starting before conception. *Clin Infect Dis*. 2015;  
42  
43 **61**:1715-25.  
44  
45  
46  
47 23. Reitter A, Stücker AU, Linde R, *et al.* Pregnancy complications in HIV-positive women: 11-  
48  
49 year data from the Frankfurt HIV Cohort. *HIV Med*. 2014;**15**:525-536.  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Table 1. Baseline maternal characteristics for pregnancies in HIV-positive pregnant women according to age at delivery (<40 years or ≥40 years at delivery)**

	Maternal age at delivery				<i>p</i> -value ( $\chi^2$ )*
	<40 years		≥40 years		
	<i>n</i>	%	<i>n</i>	%	
<b>Maternal region of birth (n=15375)</b>					
UK/Ireland	2061	14.2	100	11.8	
Africa	11093	76.4	670	79.2	
Elsewhere	1375	9.5	76	9.0	0.1
<b>HIV risk factor (n=15501)</b>					
Other risk	14343	97.9	839	98.0	
IDU	302	2.1	17	2.0	0.9
<b>Timing of diagnosis (n=15477)</b>					
Before this pregnancy	9385	64.2	715	83.6	
During this pregnancy	5237	35.8	140	16.4	<0.001
<b>Parity (number of previous births) (n=14497)</b>					
None	4288	31.3	120	14.8	
One	4953	36.2	213	26.3	
Two	2870	21.0	210	25.9	
Three or more	1575	11.5	268	33.0	<0.001
<b>Type of ART (n=15329)</b>					
Untreated	248	1.7	6	0.7	0.03*
ZDV monotherapy	860	5.9	19	2.2	
Dual therapy	134	0.9	7	0.8	
cART (3 or more drugs)	13239	91.4	816	96.2	<0.001

**cART drug class (n=14055)**

NRTI + NNRTI	4234	32.0	286	35.1	
NRTI + PI	8075	61.0	474	58.1	
NRTI + PI + NNRTI	665	5.0	36	4.4	
Other combinations*	265	2.0	20	2.5	0.2

**cART at conception (n=13882)**

No	7984	61.1	268	33.2	
Yes	5091	38.9	539	66.8	<0.001

**HIV RNA viral load closest to delivery (copies/ml) (n=9181)\*\***

<50	6226	72.0	457	85.4	
50-999	1758	20.3	53	9.9	
1000-9999	415	4.8	13	2.4	
≥10,000	247	2.9	12	2.2	<0.001

**CD4 cell count (cells/μl) (n=14036)**

≥500	5042	38.1	318	39.6	
350-499	3780	28.6	242	30.1	
200-349	3143	23.8	192	23.9	
<200	1268	9.6	51	6.4	0.02

---

ART, antiretroviral therapy; cART, combination antiretroviral therapy; IDU, injecting drug use; NRTI, nucleoside reverse transcriptase inhibitor; NNRTI, non-nucleoside reverse transcriptase inhibitor; PI, protease inhibitor; UK, United Kingdom; ZDV, zidovudine. \* Includes triple NRTIs and regimens containing integrase inhibitors. \*\* From 30 days before to 7 days after delivery.



Table 2. Pregnancy outcomes according to maternal age at delivery (<40 years or ≥40 years at delivery)

	Maternal age at delivery				p-value ( $\chi^2$ )
	<40 years		≥40 years		
	%	(n/N)	%	(n/N)	
<b>Emergency caesarean section (n=14567)</b>	23.9	(3487/14567)	25.7	(218/849)	0.2
<b>Operative vaginal delivery (n=6203*)</b>	4.6	(265/5712)	3.1	(15/491)	0.1
<b>Stillbirth (n=15501)</b>	1.0	(146/14645)	1.6	(14/856)	0.07
<b>Multiple birth (n=15501)</b>	1.9	(285/14645)	3.0	(26/856)	0.03
<b>Preterm delivery (n=15464)</b>					
<37 weeks	13.1	(1919/14609)	14.3	(122/855)	0.3
<34 weeks	5.2	(753/14609)	6.1	(52/855)	0.2
<32 weeks	3.2	(469/14609)	4.2	(36/855)	0.1
<b>Low birth weight (n=14399)</b>					
<2.5 kg	14.1	(1910/13585)	14.4	(117/814)	0.8
<1.5 kg	2.8	(376/13585)	3.8	(31/814)	0.08
<b>Birth weight ≥4 kg (n=14399)</b>	3.4	(469/14399)	4.7	(38/814)	0.07
<b>Congenital Abnormality (n=15041)</b>	2.8	(403/14212)	4.2	(35/829)	0.02
- Chromosomal	0.2	(31/14212)	1.6	(13/829)	<0.001
- Structural	2.6	(372/14212)	2.7	(22/829)	0.9
<b>MTCT (n=13031)</b>	0.8	(105/12381)	0.6	(4/650)	0.5

MTCT, mother-to-child HIV transmission. \* Information on operative vaginal delivery was only collected from July 2008 onward.