

Clinical commentary

Cardiac stunning as first manifestation of multiple sclerosis: A case report reminding us not to overlook cardiovascular autonomic dysfunction in MS

Max J. Hilz

Department of Neurology, University of Erlangen-Nuremberg, Germany

Corresponding author: Max J. Hilz

Department of Neurology, University of Erlangen-Nuremberg

Schwabachanlage 6, D-91054 Erlangen, Germany

Phone: +49-9131-853-4444

Fax: +49-9131-853-4328

E-mail: max.hilz@uk-erlangen.de

Abstract:

Autonomic dysfunction is common but frequently overlooked in multiple sclerosis (MS) patients. The case of a Tako-Tsubo cardiomyopathy on which this commentary is based shows that centrally triggered autonomic dysfunction may be the first, life-threatening manifestation of MS.

Key words: Multiple sclerosis, cardiovascular autonomic dysfunction, Tako-Tsubo Cardiomyopathy

Autonomic dysfunction is known to contribute to disability in multiple sclerosis (MS) patients.¹

Up to 84% of MS patients experience dysfunction of autonomic organ regulation.¹ Common

complaints include bowel and bladder disturbances, sexual dysfunction, or sweating abnormalities.^{1,2}

Cardiovascular autonomic dysfunction may include baroreflex dysfunction with orthostatic intolerance

and orthostatic hypotension, postural tachycardia syndrome or decreased heart rate variability³ which is associated with an increased risk of cardiovascular complications and reduced life expectancy.^{3,4} Although autonomic dysfunction frequently impairs the patient's quality of life,³ autonomic dysregulation and its prognostic impact seem to be poorly addressed in the clinical evaluation of MS patients. The most common scales evaluating MS severity, the Expanded Disability Status Scale (EDSS)⁵ and Multiple Sclerosis Functional Composite (MSFC)⁶ largely ignore the clinical and prognostic relevance of autonomic dysfunction.

This is remarkable since the autonomic nervous system is not only afflicted by the neuro-inflammatory process but may even contribute to maintaining inflammation and disease progression.⁷ MS induced changes in sympathetic or parasympathetic modulation may in turn alter immune responses.⁷

Particularly during the first years after disease manifestation, MS patients seem to have an increased risk of cardiovascular diseases⁸ which is very likely to be due to a central up-regulation of sympathetic outflow to the cardiovascular system.⁷

Still, acute cardiovascular emergencies are rarely recognized as initial manifestation of MS, probably because cardiologists and neurologists are unaware of this etiology.

However, sympathetic activity may increase significantly during early stages of MS while later relapses may lead to progressive noradrenergic failure and adrenergic receptor dysfunction.⁷ In stroke patients, neurologists are increasingly aware of the risk of abundant cardiovascular autonomic disorders^{9,10} In contrast, acute cardiologic emergencies might not be recognized as centrally mediated in patients who experience such an emergency, and, supposedly by coincidence, also happen to have MS.

In this issue of the *MSJ*, Madaglia et al. present the case of a previously healthy young man who was admitted to internal medicine because of severe headache and diaphoresis and a first-degree atrioventricular block.¹¹ Two weeks later, symptoms reoccurred but were more severe, with additional tachycardia- and dyspnoea, arterial hypertension, and electrocardiographic changes that are typical for stress-cardiomyopathy,¹² also called Tako-Tsubo cardiomyopathy (TTC). TTC may be life threatening and is characterized by apical ballooning, left ventricular dysfunction with ventricular wall-motion

abnormalities.^{12, 13} Although the pathogenesis of TTC is not yet fully understood,¹² there is a dysfunction in the brain-heart axis, most likely with increased sympathetic activity and catecholamine surges.¹³ The patient had no myocarditis or acute myocardial infarction but turned out to have MS with several active lesions, some of which involved areas of the central autonomic network e.g. the medulla oblongata. Methylprednisolone treatment quickly reversed the myocardial stunning.

The case alerts neurologists to the fact that MS may manifest as cardiovascular emergency³. Although it seems to be the first report of TTC as the initial symptom of MS, cardiovascular complications of MS are frequent and range from subclinical to life-threatening events, and may cause fatalities that have been coined as “SUDIMS”, Sudden Unexpected Death in MS,³ in analogy to “SUDEP”, the Sudden Unexpected Death in Epilepsy.¹⁴

In summary, the case reminds us to pay closer attention to centrally mediated autonomic dysfunction in MS patients and to better investigate the association between neuro-inflammation and cardiovascular autonomic dysfunction.

Conflict of interest

The author declares that there is no conflict of interest with respect to the research, authorship and/or publication of this article.

Funding

The author received no financial support for the research, authorship and/or publication of this article.

References

1. Racosta JM, Kimpinski K, Morrow SA, et al. Autonomic dysfunction in multiple sclerosis. *Autonomic neuroscience : basic & clinical*. 2015;193:1-6.
2. Merkelbach S, Haensch CA, Hemmer B, et al. Multiple sclerosis and the autonomic nervous system. *Journal of neurology*. 2006;253 Suppl 1:I21-25.
3. Kaplan TB, Berkowitz AL, Samuels MA. Cardiovascular dysfunction in multiple sclerosis. *The neurologist*. 2015;20:108-114.

4. Scafari A, Knappertz V, Cutter G, et al. Mortality in patients with multiple sclerosis. *Neurology*. 2013;81:184-192.
5. Kurtzke JF. Rating neurologic impairment in multiple sclerosis: An expanded disability status scale (edss). *Neurology*. 1983;33:1444-1452.
6. Cutter GR, Baier ML, Rudick RA, et al. Development of a multiple sclerosis functional composite as a clinical trial outcome measure. *Brain : a journal of neurology*. 1999;122 (Pt 5):871-882.
7. Racosta JM, Kimpinski K. Autonomic dysfunction, immune regulation, and multiple sclerosis. *Clinical autonomic research : official journal of the Clinical Autonomic Research Society*. 2016;26:23-31.
8. Christiansen CF. Risk of vascular disease in patients with multiple sclerosis: A review. *Neurological research*. 2012;34:746-753.
9. Hilz MJ, Schwab S. Stroke-induced sudden-autonomic death: Areas of fatality beyond the insula. *Stroke; a journal of cerebral circulation*. 2008;39:2421-2422.
10. Soros P, Hachinski V. Cardiovascular and neurological causes of sudden death after ischaemic stroke. *The Lancet. Neurology*. 2012;11:179-188.
11. Midaglia L, Juega Mariñob JM, Sastre-Garrigaa J, et al. An uncommon first manifestation of multiple sclerosis: Tako-tsubo cardiomyopathy. *Mult Scler*. 2016
12. Templin C, Ghadri JR, Diekmann J, et al. Clinical features and outcomes of takotsubo (stress) cardiomyopathy. *The New England journal of medicine*. 2015;373:929-938.
13. Prasad A, Lerman A, Rihal CS. Apical ballooning syndrome (tako-tsubo or stress cardiomyopathy): A mimic of acute myocardial infarction. *American heart journal*. 2008;155:408-417.
14. Matthias Dütsch, Hilz MJ. Epilepsy and autonomic regulation. In: David Robertson, Italo Biaggioni, Geoffrey Burnstock, Phillip A. Low, Paton JFR, eds. *Primer on the autonomic nervous system*. San Diego, USA: Academic Press, Elsevier; 2012:549-552.