

# Large-volume lava flows fed by a deep magmatic reservoir at Ağrı Dağı (Ararat) volcano, Eastern Turkey

Özgür Karaoğlu<sup>1,2,\*</sup>, Abdelsalam Elshaafi<sup>2</sup>, Mohamed K. Salah<sup>3</sup>, John Browning<sup>4</sup>,

Agust Gudmundsson<sup>2</sup>

<sup>1</sup>Eskişehir Osmangazi University, Department of Geological Engineering, 26040 Eskişehir, Turkey

<sup>2</sup>Department of Earth Sciences, Royal Holloway University of London, Egham, TW20 0EX, UK

<sup>3</sup>Department of Geology, American University of Beirut, Riad El Solh 1107 2020, Beirut, Lebanon

<sup>4</sup>Department of Earth Sciences, University College London, Gower Street, London WC1E 6BT, UK

\* Correspondence to: Ö. Karaoğlu, [ozgur.karaoglu@deu.edu.tr](mailto:ozgur.karaoglu@deu.edu.tr); [cougarforever@gmail.com](mailto:cougarforever@gmail.com)

## Abstract

1 Ağrı Dağı (Ararat), whilst being the tallest volcano in Turkey, is largely understudied.  
2 Two predominant peaks, Greater and Lesser Ağrı, make up the main edifice which has been  
3 built during four main phases. The most recent phase consisted of two volcanic eruptions. The  
4 respective surface area and volume of the first volcanic eruption were estimated at 96 km<sup>2</sup> and  
5 3.2 km<sup>3</sup>, whereas those of second eruption were much smaller with the surface area and volume  
6 estimated at 25 km<sup>2</sup> and 0.6 km<sup>3</sup>. It is unusual for stratovolcanoes to produce basaltic eruptions  
7 of over 3 km<sup>3</sup>, although these and larger volumes are not uncommon in flood basalt-type  
8 eruptions. Large basaltic eruptions from stratovolcanoes normally require volcano-tectonic  
9 forcing (e.g. subsidence of collapse caldera and graben). However, there is no evidence for  
10 such volcano-tectonic forcing, during the most recent eruptions at Ağrı Dağı (Ararat), and  
11 therefore their comparatively large volume basaltic lavas need to be explained in a different  
12 way. Here we present an analytical method for calculating the source volume needed to supply  
13 magma to the eruptions at Ağrı Dağı. We find that the lava flow of 3.2 km<sup>3</sup> was likely fed by  
14 a very large magma reservoir (~13,000 km<sup>3</sup>) while the second flow of 0.6 km<sup>3</sup> was fed by a

15 reservoir of a much smaller effective size, or  $\sim 2000 \text{ km}^3$ ). ‘Effective size’ depends on what  
16 fraction of the reservoir participates in the eruption. We propose that entire reservoir supplied  
17 magma to the larger eruption, but only one of its compartments (about 1/5 of the total volume  
18 of the reservoir) supplied magma to the smaller eruption. Although seismic tomography  
19 indicates a magma reservoir at great depths ( $>20\text{-}30 \text{ km}$ ) below the Ağrı Dağı volcano,  
20 geochemical constraints on some of the later-formed rocks suggest an interaction between a  
21 shallow chamber (at 8-10 km depth) and the deep reservoir approximately 0.5 Ma. We provide  
22 numerical models whose results indicate that dykes injected from the lateral margins of the  
23 deep-seated reservoir are more likely to reach the surface directly rather than replenish the  
24 shallow magma chamber, suggesting also that the compartment for the second eruption was at  
25 the margin of the reservoir.

Keywords: large eruptions, magma chambers, magma reservoirs, volcano-tectonic forcing,  
crustal stresses, numerical models

## 26 **1. Introduction**

27 Magma or melt transport in the mantle is somewhat different from magma transport in the  
28 upper crust. Magma in the mantle, and partly in the lower crust, ascends by porous flow ([Scott  
29 and Stevenson 1986](#)). At shallower crustal levels, magma ascent is primarily through magma-  
30 driven fractures, that is, dykes. Dyke initiation and propagation is known to be partly controlled  
31 by regional stress fields, particularly those induced by crustal extension (e.g. [Gudmundsson  
32 1990, 2006](#); [Daniels et al. 2012](#); [Le Corvec et al. 2013](#); [Maccaferri et al. 2014](#); [Tibaldi 2015](#)).  
33 Reservoirs which are underlying the shallow magma chamber may directly supply magma to  
34 areas outside of the stratovolcano ([Gudmundsson 2006](#)). Thus, less evolved magmas can erupt  
35 at the margins of stratovolcanoes while more evolved magmas erupt within the central parts  
36 of the stratovolcano.

37 Long-lived ( $>1 \text{ Ma}$ ) major volcanic edifices, such as a stratovolcano, a caldera volcano, or  
38 a large shield volcano (basaltic edifice), are commonly supplied with magma from a  
39 comparatively shallow crustal magma chamber ([Browning et al. 2015](#); [Gudmundsson 2016](#);  
40 [Karaoğlu et al. 2016](#)). While active, a shallow magma chamber acts as a sink for magma from  
41 a deeper magma source (or reservoir) ([Gudmundsson 2012](#); [Le Corvec et al. 2013](#)). If new  
42 magma is injected from a deeper source during an eruption, that magma is likely to be of high  
43 density and may accumulate at the floor of the magma chamber ([Coppola et al. 2009](#);

44 [Gudmundsson 2012](#)). For an eruption to occur, the necessary conditions are that the magma  
45 chamber or reservoir (deep-seated magma chamber) ruptures and a fluid-driven fracture is able  
46 to propagate from the chamber to the surface ([Gudmundsson 2012](#)). There is a close  
47 relationship between the excess chamber pressure ( $p_e$ ) and magma recharge volume. At the  
48 most active volcanoes, rupture probability based on increasing excess pressure within the  
49 shallow chamber allow forecasts of dyke formation to be made in real time during magma  
50 recharge events ([Browning et al. 2015](#)). Stratovolcanoes in Turkey or elsewhere are commonly  
51 fed by shallow crustal magma chambers with estimated volumes that commonly range from  
52 about 5 km<sup>3</sup> to 500 km<sup>3</sup> (e.g. [Chester 1993](#)). Lavas issued from stratovolcanoes commonly  
53 range in volume between 0.01 km<sup>3</sup> or less to 0.1 km<sup>3</sup>. Whilst these small eruption volumes can  
54 be considered ‘normal’, more voluminous eruptions are known to erupt at stratovolcanoes such  
55 as the 1981 lateral blast event at Mt Saint Helens, USA (2.5 km<sup>3</sup>), the Plinian eruption of  
56 Krakatoa, Indonesia in 1881 (18-21 km<sup>3</sup>), the 1991 dome collapse of Mt Unzen, Japan (1 km<sup>3</sup>),  
57 and the the Plinian eruption of Mt Nemrut, Eastern Turkey (2.5 km<sup>3</sup>) ([Karaoğlu et al. 2005](#)).  
58 Such events cannot be considered ‘normal’ as they are often associated with some degree of  
59 volcano-tectonic forcing, particularly graben or caldera formation or slip. By volcano-tectonic  
60 forcing we mean processes where the strain energy needed for displacement on a ring-fault of  
61 a caldera or the boundary faults of a graben is primarily of tectonic origin and the displacement  
62 cause reduction in volume, shrinkage, of the chamber/reservoir source. The volume reduction  
63 maintains the magmatic excess pressure in the source until the very end of the eruption, thereby  
64 squeezing out an exceptionally large fraction of the magma in the source and producing a large  
65 eruption ([Gudmundsson, 2015, 2016](#)). As said, we do not find evidence of volcano-tectonic  
66 forcing of this kind for these two eruptions and therefore seek alternative explanations for their  
67 sizes.

68 The type and composition of magma feeding an eruption can also influence the eruptive  
69 volume. For example, eruptions of felsic magmas commonly produce somewhat larger  
70 volumes than mafic ones, as exemplified by the eruption of Puyehue Cordon-Caulle which  
71 produced a rhyolitic lava flow of volume 0.5 km<sup>3</sup> ([Tuffen et al. 2013](#)). Nevertheless, large-  
72 volume basaltic lava flows are commonly associated with flood basalt events such as the  
73 Deccan Plateau and the Columbia Basalt Plateau ([Reidel et al. 2013](#)).

74 It is seemingly rare for stratovolcanoes to produce both normal-size eruptions and large  
75 volume effusive eruptions without an element of local volcano-tectonic forcing  
76 ([Gudmundsson, 2015, 2016](#)). The Ağrı Dağı volcano, however, seems to exhibit such rare

77 behaviour. Where most of the lavas that make up Ağrı Dağı were produced in relatively small  
78 eruptions ( $<0.1 \text{ km}^3$ ), two massive basaltic lava flows with total volume exceeding  $3.8 \text{ km}^3$ ,  
79 that formed roughly during the period between the peak activity of the greater and lesser Ağrı  
80 volcano. There is currently no explanation as to why such voluminous eruptions occurred  
81 during this time.

82 At the height of 5165 m, Ağrı Dağı (Ararat) is the tallest volcano in Turkey and is  
83 comprised of two main peaks: Greater and Lesser Ağrı (Fig. 1). The most recent eruption ( $<$   
84  $0.5 \text{ Ma}$ ) of Ağrı Dağı occurred at  $39^\circ 30' 20'' \text{ N} / 44^\circ 22' 23'' \text{ E}$  and produced two generations of  
85 basaltic lava flows. The former volcanic eruption occupies an area of about  $96 \text{ km}^2$  and a  
86 volume of around  $3.2 \text{ km}^3$  while the later volcanic eruption was much smaller with an area of  
87  $25 \text{ km}^2$  and a volume of  $0.6 \text{ km}^3$  (Fig. 1). The exact age difference between these lava flows,  
88 however, is unknown. The nearest major population centres (about 145,000 inhabitants) are  
89 only 6 km away from the volcano. Many of the stratovolcanoes in Eastern Turkey are poorly  
90 studied and understood, particularly in terms of their relationship to the current tectonics. This  
91 is an important issue because Ağrı Dağı and other neighbouring volcanoes are situated close to  
92 major strike-slip faults and areas of triple junction tectonics (Fig. 1).

93 The Ağrı Dağı volcano covers the largest area ( $\sim 1100 \text{ km}^2$ ) of any volcano in Turkey. The  
94 volcano has erupted some  $1150 \text{ km}^3$  of volcanic materials over its  $\sim 1.5 \text{ Ma}$  of activity (Yılmaz  
95 et al. 1998) (Fig. 2). There are no calderas or grabens dissecting the volcano, which is in  
96 contrast with the common calderas on most stratovolcanoes in Eastern Turkey, such as the  
97 Nemrut caldera (Karaoğlu et al. 2005). The orientations of the parasitic cones and main  
98 volcanic fissures indicate that the dominant direction of tension in the area is NW-SE (e.g.  
99 Karakhanian et al. 2002). Dextral faults are common and form several pull-apart structures,  
100 some of which may be linked to volcanic activity (Karakhanian et al. 2002).

101 Ağrı Dağı is a typical stratovolcano mostly built up by calc-alkaline volcanic rocks (Yılmaz  
102 et al. 1998, Fig. 2). Initial products (pre-cone phase) observed in the eastern part of the volcano  
103 are mainly intermediate (dacitic and andesitic in composition) pyroclastic rocks and lavas (e.g.  
104 Yılmaz et al. 1998). K-Ar radiometric age data show that the oldest lavas are basaltic and were  
105 erupted between 1.51 Ma and 1.09 Ma ago (Sanver 1968; Pearce 1990). Basaltic lava flows  
106 overlay the oldest volcanic rocks. Following the first eruptive stages, the main cone of the  
107 volcano was built up mostly by andesite and dacite lavas. The last stage (flank eruption phase)  
108 is represented by alternating andesitic and basaltic lava flows from the main cone and parasitic

109 scoria spatter cones on the flanks. During the last and most recent phase; basaltic lava flows  
110 were particularly dominant at the margin of the Ağrı Dağı volcano (Fig. 2).

111 One objective of this paper is to provide models that give insights into the magma  
112 storage systems feeding the Ağrı Dağı volcano and how their characteristics can account for  
113 the contrasting eruption volumes issued at the volcano. More specifically, we aim to find the  
114 feeding mechanism of the large-volume basaltic lava flows. Furthermore, in the absence of  
115 evidence of volcano-tectonic forcing contributing to the generation of the lava flows, we seek  
116 an alternative mechanism for their comparatively large sizes. In particular, we propose that the  
117 entire reservoir supplied magma during the eruption of the larger and more primitive lava flow.  
118 By contrast, we suggest that only a small compartment within the reservoir supplied magma  
119 during the eruption of the smaller and more evolved lava flow.

120 Geochemical constraints indicate that the Agri Dagi volcano was predominantly  
121 constructed from acidic to intermediate lavas and the later-formed rocks indicate an interaction  
122 between a shallow chamber (at 8-10 km depth) and the deep reservoir. As such, we have  
123 developed numerical models to study how of the magma systems of Ağrı Dağı volcano interact  
124 over time. These models are combined with approximate estimations as to the volume of the  
125 magma system underlying the Ağrı Dağı volcano in order to understand how and why such  
126 comparatively voluminous lavas can be erupted from stratovolcanoes such as Ağrı Dağı. The  
127 results provide information which is vital for understanding such large eruptions, particularly  
128 because they pose a significant threat to nearby population centres (e.g. [Small and Naumann](#)  
129 [2001](#)).

## 130 2. Tectonics, geology and geochemistry of the Ağrı Dağı volcano

131 The East Anatolian High Plateau (EAHP) displays a very complex volcano-tectonic  
132 history of continental collision. After the closure of the Neotethyan Ocean as a result of Africa-  
133 Eurasian convergence ([Barka 1992](#); [Okay and Tüysüz 1999](#); [Bozkurt 2001](#)), syn- and post-  
134 collisional magmatism dominate in the EAHP since the Middle Miocene (15 My, [Lebedev et](#)  
135 [al. 2010](#)). Four stages of Neogene-Quaternary volcanism have been identified: Middle Miocene  
136 (15.0–13.5 Ma), Late Miocene (10–9 Ma), Pliocene (5.8–3.7 Ma), and Quaternary (1.0–0.4  
137 Ma) ([Lebedev et al. 2010](#)). Quaternary stratovolcanoes (e.g. Nemrut, Suphan, Ağrı Dağı  
138 volcanoes, Fig. 1) and shield volcanoes (e.g. Tendürek) on the Eurasian Plate produce  
139 predominantly calc-alkaline type eruptive materials (e.g. [Pearce et al. 1990](#); [Keskin et al. 1998](#);  
140 [Yılmaz et al. 1998](#); [Keskin 2007](#), [Lustrino et al. 2010](#)), with minor alkaline igneous rocks (e.g.

141 [Innocenti et al. 1976, 1980](#); [Pearce et al. 1990](#); [Keskin et al. 1998](#); [Yılmaz et al. 1998](#); [Alici et](#)  
142 [al. 2001](#); [Keskin 2007](#)). [Lustrino et al. \(2010\)](#) proposed that extensive volcanic activity on the  
143 Arabian plate, such as Karacadağ shield volcano, surfaced on a 35–40 km thick crust mostly  
144 during the Late Miocene to Quaternary, with the production of large amounts of alkaline basic  
145 rocks ([Pearce et al. 1990](#); [Ercan et al. 1991](#); [Notsu et al. 1995](#)). The formation of Ağırı Dağı  
146 volcano has been tectonically linked to slab break-off and delamination in intraplate settings  
147 overlying hot asthenosphere through transtension ([Yılmaz et al. 1998](#); [Shabanian et al. 2012](#);  
148 [Sağlam-Selçuk et al. 2016](#)).

149 Recent seismic tomography studies have documented that the crust in Eastern Turkey has  
150 an average of thickness 65 km; it is thinner than average in the south, about 38 km (Arabian  
151 foreland ([Angus et al. 2006](#); [Ozacar et al. 2008](#); [Cakir et al. 2000](#); [Zor et al. 2003](#)) (Fig. 1).  
152 Many studies suggest that the lithospheric mantle may be either completely absent (e.g. [Al-](#)  
153 [Lazki et al. 2003](#)) or very thin (e.g. [Angus et al. 2006](#); [Ozacar et al. 2008](#)) beneath Eastern  
154 Turkey. Two controversial views have been expressed as to the origin of volcanism in Eastern  
155 Turkey; namely (i) that the region is reformed by melting and cooling of the asthenosphere and  
156 is as such an older lithospheric mantle ([Keskin 2007](#)), and (ii) that, on average, a 20 km thick  
157 lithosphere may have resulted from cooling of the asthenosphere from 15 Ma to 7 Ma ([Angus](#)  
158 [et al. 2006](#)). The crustal stress field has likely changed dramatically in the past 10 to 5 Ma  
159 ([Örgülü et al. 2003](#)). These seismic- and petrology-based studies indicate that the uppermost  
160 mantle is partially molten and that the asthenosphere is close to the base of the crust, consistent  
161 with the existence of volcanism in the region ([Örgülü et al. 2003](#)).

162 The volcano-stratigraphy of the Ağırı Dağı volcano was mapped by [Yılmaz et al. \(1998\)](#).  
163 Geological observations and published data ([Türkunal 1980](#); [Bingöl et al. 1989](#)) show that 14  
164 different types of geological units represent the stratigraphy of the region around the Ağırı Dağı  
165 volcano (Appendix 1). A cone-building phase produced mostly basaltic and rarely andesite  
166 rocks between 0.68 Ma and 0.5 Ma ([Sanver 1968](#); [Pearce 1990](#)). The final stages of activity  
167 resulted in flank eruptions between 0.3 Ma and 0.04 Ma ([Sanver 1968](#); [Pearce 1990](#); [Ercan et](#)  
168 [al. 1990](#); [Notsu et al. 1995](#)) and the most recent activity which occurred 20,000 years ago  
169 produced mostly andesitic lavas ([Nagao et al. 1989](#)). Since this period the volcano has been  
170 dormant, although there were unrest periods characterised by increased seismicity in 2500-  
171 2400 BC and 1840 AD ([Karakhanian et al. 2002](#)).

172 Here we consider the most recent basaltic lava flows erupted during the cone-building and  
173 flank eruption phases (< 0.5 Ma), which are located on the southern flank of the Greater Ağırı

174 Dağı volcano (lava flows I and II, Fig. 3). The flows are easily distinguishable from the older  
175 basaltic lava flows (lava flow III, Fig. 3) by colour and lack of both erosion and alteration.  
176 These most recent basaltic lava flows were erupted from a NW-SE aligned series of scoria  
177 cones dated at 0.5 Ma (Sanver 1968). The origin of those basaltic lava flows ~~are~~is poorly  
178 constrained in terms of petrology because previous sampling localities were not spatially  
179 defined (e.g. Pearce et al. 1990; Yılmaz et al. 1998). Generally, though, the volcanic rocks of  
180 the Ağrı Dağı volcano are classified through a wide compositional range from trachy-basalt,  
181 tephrite/basanite, basaltic andesite, andesite, dacite and rhyolite (Fig. 4a) (e.g. Pearce et al.  
182 1990; Nagao et al. 1989; Kheirkhah et al. 2009). A significant feature of the genesis and  
183 evolution of Quaternary magmas in Ağrı Dağı is the absence of basalt on the plot although the  
184 petrography shows them as basalt (Fig. 4). MORB-normalised trace element content of selected  
185 basaltic rocks are shown in Fig 4b. The basaltic lava flows at the main cone of the Ağrı Dağı  
186 are more enriched in LREE than the marginal lavas (Fig. 4b).

### 187 **3. Injected material and reservoir volume**

188 In order to estimate the relative contribution of a shallow magma chamber and the  
189 contribution of a deeper magma reservoir to the eruptions of Ağrı Dağı we calculated first the  
190 total volume of injected materials, that is, magma volume leaving the chamber/reservoir during  
191 the eruption, from the lava flows I and II. In this study we used ArcGIS 10.1 to calculate the  
192 geometry and area of the Quaternary basaltic lava flows I and II (Fig. 3).  
193 The maximum thickness for each flow was estimated based on the elevation difference with  
194 the surrounding area using a SRTM (Shuttle Radar Topography Mission) compiled digital  
195 elevation model (DEM). The thickness of each individual lava flow increases from the margins  
196 to the centre, and so the greatest thickness was recorded at the centre of each flow that appears  
197 to be similar to lava shield (Fig. 2).

198 We can make an approximation to the shape and emplacement style of a lava shield. The  
199 volume of a lava shield is generally computed by approximating its shape as a truncated cone  
200 for flat topped volcanoes or a pyramid for a volcano with a distinct peak (Hasenaka  
201 1994). Therefore, during this study the volume of each lava flow is calculated by  
202 approximating its shape to a cone, namely as:

$$203 \quad V_e = a(h/3) \quad (1)$$

204 where  $V_e$  is volume of the volcanic unit;  $a$  is area and  $h$  is the maximum thickness of the unit.  
205 The area of the base of each individual volcanic unit was calculated using ArcGIS. The volume  
206 of eruptive surface materials is somewhat uncertain because part of the flow may be partially  
207 submerged by younger thick lava sequences (Andrew and Gudmundsson 2007). The total  
208 volume of injected material is a combination of the volume of an individual lava flow on the  
209 surface and the volume of the feeder dyke that fed the eruption. There are no available data in  
210 the study area on dyke geometries, such as length (strike dimension), thickness and depth (dip  
211 dimension), to calculate the volume of feeder dykes. As such, we use rough estimations of the  
212 average volume of dykes in Eastern Turkey, where the volumes do not exceed 0,004  
213 km<sup>3</sup> (Karaoğlu et al. 2016). Therefore, the error produced in the total injected material due to  
214 neglected volume of feeder dyke is very small.

215 The total injected material or magma  $V_e$  from Eq. (1) for lava flow I is around 3.2 km<sup>3</sup>  
216 ( $\pm 0,1$  km<sup>3</sup>) while the total injected material for lava flow II is around 0.6 km<sup>3</sup> ( $\pm 0.02$  km<sup>3</sup>). Both  
217 volumes are quite similar to the sizes of monogenetic Holocene lava shields on the Reykjanes  
218 Peninsula, West Iceland, where the volume lava flow II is approximately the same size of the  
219 picrite lava-shields while the volume lava flow I is approximately the size of the olivine-  
220 tholeiite shields (Andrew and Gudmundsson 2007). The primary picrite or olivine basaltic  
221 magmas in Iceland are believed to come from deep magma reservoirs rather than crustal  
222 shallow magma chambers (Meyer et al. 1985).

223 It is known that magma can accumulate at the crust-mantle boundary, which is commonly  
224 the situation for deep-seated reservoirs. A deep reservoir may directly feed surface eruptions  
225 or form a shallow magma chamber in the upper or middle crust. Such shallow chambers can  
226 form due to abrupt changes in the mechanical properties of the crustal rocks, particularly  
227 changes in stiffness (Young's modulus) of those rocks (Barnett and Gudmundsson 2014). In  
228 areas of intense magmatism such as Iceland, the crust-mantle boundary is commonly referred  
229 to as the magma layer (Hermance 1981; Bjornsson 1983; Gudmundsson 1987). The porosity  
230 or melt fraction differs through a magma reservoir due to buoyancy and reduced potential  
231 energy such that magma tends to move towards the top (shallowest depth) of the reservoir.  
232 Therefore, the greatest melt fraction is normally in the uppermost compartments of the  
233 reservoir and gradually decreases with depth (Richter and McKenzie 1984). The average melt  
234 fraction throughout a reservoir is commonly assumed at 0.25 (e.g. Richter and McKenzie  
235 1984). The melt fraction of the lowest parts of a chamber may be higher if the reservoir is



236 continuously supplied with new primitive melt or magma from deeper sources in the mantle;  
 237 for example, from the upper parts of a mantle plume (Gudmundsson 1987). The mechanical  
 238 behaviour of a magma reservoir can be modelled to a first approximation as  
 239 a poroelastic material (Gudmundsson 1986, 2016; Tibaldi 2015). Hence, the volume of a  
 240 magma source during individual eruptions may be roughly estimated from Eq. 2, and is given  
 241 by (Gudmundsson 1987, 2016; Browning and Gudmundsson 2015):

$$242 \quad V_b = \frac{V_e}{p_e \phi \left( \beta_m + \frac{\beta_b}{\phi} \right)} \quad (2)$$

243 where  $V_e$  is the volume of injected material in a single eruption,  $\phi$  is fractional porosity of the  
 244 reservoir,  $p_e$  is the excess magmatic pressure in the reservoir,  $\beta_m$  and  $\beta_b$  are magma  
 245 compressibility and bulk compressibility of the reservoir, respectively.

246 Magmatic excess pressure in the reservoir can be considered nearly equal to the *in-situ*  
 247 tensile strength of the host rock at the time of rupture (Elshaafi and Gudmundsson 2016). The  
 248 average *in-situ* tensile strength of the upper crust in Eastern Turkey is around 3.5 MPa (Gurocak  
 249 et al. 2012). Compressibility is a measure of the relative volume change of a fluid or solid as a  
 250 response to change in stress. The static compressibility of basaltic magma  $\beta_m$  at 1100-1300 °C  
 251 is around  $1.25 \times 10^{-10} \text{ Pa}^{-1}$  (Murase and McBirney 1973). The Young's modulus for the  
 252 lowermost crust in Eastern Turkey is around 35 GPa at a depth of 20 km (e.g. Gurocak et al.  
 253 2012; Karaoğlu et al. 2016). The bulk modulus ( $K$ ) for this part of the crust can be calculated  
 254 from the relation:

$$255 \quad K = \frac{E}{3(1 - 2\nu)} \quad (3)$$

256 where  $K$  is the bulk modulus,  $E$  is the Young's modulus and  $\nu$  is the Poisson's ratio, whose  
 257 average value for most solid rocks is around 0.25 (Gudmundsson 2011). Hence the  
 258 compressibility of the crust in Eastern Turkey  $\left( \frac{1}{K} \right)$  is around  $4.28 \times 10^{-11} \text{ Pa}^{-1}$ .

259 From Eq. (2), if we assume the magma reservoir as partially melted with an average  
 260 porosity of 0.25 throughout the reservoir as previously mentioned, the volume of the reservoir  
 261 would be:

262  $V_b = 3858 \times V_e$  (4)

263 This equation can be applied to estimate the volume of magma within a reservoir supplying  
264 magma to individual eruptions. From Eq. (4), the volume of the magma reservoir during the  
265 first eruption (lava flow I) is around 12,345 km<sup>3</sup>. By contrast, the volume of the magma  
266 reservoir during the second eruption (lava flow II) is, at 2,403 km<sup>3</sup>, that is less by a factor of  
267 about 5. A much larger reservoir is thus needed to give rise to the first lava flow than the second  
268 lava flow, as expected, assuming the reservoir's elastic properties remained the same for both  
269 eruptions. To explain this difference in reservoir size and related aspects during these eruptions,  
270 we created a suite of numerical models which investigate the distribution of stresses around a  
271 deep magma reservoir, with some constraints from seismic tomography.

#### 272 **4. Seismic tomography models**

273 Low-velocity anomalies obtained from seismic tomography models can be used to detect  
274 magma chambers and reservoirs at depth. The seismic velocity model derived by [Salah et al.](#)  
275 [\(2011\)](#) is used to construct five vertical cross-sections of P-wave velocity across the area of  
276 Ağrı Dağı volcano. This model is constructed through the application of the seismic  
277 tomography method of [Zhao et al. \(1992, 1994\)](#) on P-wave (primary wave) arrival times in  
278 Eastern Anatolia. This method has been applied successfully on arrival times collected from  
279 seismic events occurring in different tectonic circumstances. The method is adaptable to a  
280 general velocity structure which includes several seismic velocity discontinuities of complex  
281 topography. Initially, a 3-D grid net is set in the model space to express the 3-D velocity  
282 variations, the seismic velocities are taken as unknown parameters. Velocity at any point in the  
283 model is calculated by linear interpolation of the velocity values at eight grid nodes surrounding  
284 that point. The method uses an efficient 3-D ray-tracing scheme which accurately calculates  
285 travel times and ray-paths. More details about the method can be found in [Zhao et al. \(1992,](#)  
286 [1994, 2012\)](#).

287 The Eastern Turkey data set comprises 31,730 P-wave arrival times generated by 7380  
288 seismic events which were recorded by 39 seismic stations distributed relatively uniformly in  
289 the study area. Analysis of ray-path coverage (both in plan and vertical views) and the results  
290 of a checkerboard resolution test, and the hit count rates all imply that the obtained velocity  
291 anomalies are reliable features down to a depth of 45 km ([Salah et al. 2011](#)). P-wave velocity  
292 along the selected five cross-sections are shown in Figure 5. The model shows that prominent

293 low P-wave velocity zones are visible at a depth range of 20-30 km beneath cross-sections 1-3  
294 which strike in a NW-SE direction. Cross-sections 4 and 5 run in an NE-SW direction and  
295 exhibit low P-wave velocities that extend to the base of the upper crust (Fig. 5). These low P-  
296 wave velocity zones most likely indicate the occurrence of partial melt which can be interpreted  
297 as magma reservoirs beneath Eastern Anatolia (Hearn 1999; Calvert et al. 2000; Zor et al.  
298 2003). These low-velocity zones seem to be consistent with previous seismological  
299 observations such as inefficient  $S_n$  propagation and low  $P_n$  velocity (Rodgers et al. 1997; Al-  
300 Lazki et al. 2004).

## 301 5. Numerical models

302 Whilst the seismic tomography data clearly indicates the presence of a deep reservoir  
303 there is little evidence in the tomography for a shallow magma chamber. However,  
304 geochemical constraints indicate that a shallow chamber was active approximately 5 Ma. As  
305 such, we built a suite of numerical models to test the stress conditions generated by different  
306 arrangements of magma chambers. The objective was to understand which conditions favour  
307 eruptions, and how could the relative size and location of those eruptions change due to the  
308 magma chamber arrangement.

309 The numerical models were built and solved using the finite element program  
310 COMSOL ([www.comsol.com](http://www.comsol.com); cf. Zienkiewicz 1979; Deb 2006). The models are based on the  
311 real geological setting of the Ağrı Dağı volcano as interpreted from field measurements,  
312 seismic wave profiles, and InSAR data (Cavalié and Jónsson 2014) (Fig. 6). All models are  
313 two dimensional where the magma chambers and reservoirs are modelled as cavities or holes  
314 with prescribed loads given at their boundaries to simulate overpressure (Gudmundsson 2011;  
315 Gerbault 2012) (Fig. 6).

### 316 5.1. Model set-up

317 The geometry of our 2-D models is based upon a simplified E-W striking profile  
318 through the Ağrı Dağı volcano (Fig. 6). The magma sources in our models are elliptical, which  
319 is likely a simplification of real magma chamber geometries (e.g. Gudmundsson 2012; Le  
320 Corvec et al. 2015; Karaoglu et al. 2016). Although it has been shown previously that  
321 topography can play a role in distributing near surface stresses, the primary focus of our  
322 investigation is on the stress differences resulting from different boundary conditions applied  
323 to the magma chamber itself, where the host-rock properties as well as the depth, shape and

324 size of the chamber are of main concern. Thus, we assume flat topography in all the models.  
325 The 14 different geological units as mechanical layers used in our models are based on direct  
326 geological observations and published literature (Yilmaz et al. 1998) (Fig. 6). The values used  
327 to calculate depth of the magma chamber encompass all of these mechanical layers.

328 The depths of shallow magma chambers are commonly located within a few kilometres  
329 of the ground surface (cf. Gudmundsson 1998). In this study we assume the magma chamber  
330 depth to be 8 km, although results are not sensitive to the shallow chamber depth. The depth of  
331 the deep seated magma reservoir is inferred from tomographic data at around 20 km. In Fig. 6  
332 we show only the model along an E-W strike. We performed two models, in order to investigate  
333 different eruption volumes i.e. (i) very large magma storage configuration for lava flow I (i.e.  
334  $\sim 13,000 \text{ km}^3$ ), and (ii) a smaller lava flow II (i.e.  $\sim 2,000 \text{ km}^3$ ). We assumed two magma storage  
335 regions: 1) a deeper and larger reservoir at a depth of 20 km (with a diameter of 40 km and a  
336 thickness of 7 km) a shallow magma chamber at 8 km depth (with a diameter of 16 km and a  
337 thickness of 5 km (Figs. 7a-b). The second model shows the same shallow magma chamber at  
338 8 km depth (with a diameter of 16 km and a thickness of 2 km) but with a much smaller volume  
339 deeper reservoir at 20 km depth (with a diameter of 30 km and thickness of 3 km (Figs. 7c-d).

340 In this model both magma chambers are residing within a heterogeneous, anisotropic  
341 elastic half space with Young's modulus (E) varying between individual layers from 50 GPa  
342 to 20 GPa, as shown in Appendix 1. The shallower magma chamber is modelled considering  
343 two criteria. First, that most stratovolcanoes are fed by shallow chambers and, second, that  
344 geological data (some magma mingling textures in the rocks) and geochemical records indicate  
345 the existence of a shallow magma chamber beneath the volcano. The deeper magma reservoir  
346 is modelled based on our seismic tomography data. The shallower magma chamber assumed  
347 that has a maximum diameter of 16 km to a first approximation (Figs. 7a-b), whereas the deeper  
348 chamber or reservoir has a maximum diameter of 40 km for the first volcanic eruption to  
349 correspond the shrinkage of the volume of reservoir with the time. Poisson's ratio ( $\nu$ ) does not  
350 vary significantly between individual layers; thus, in the models we use a constant typical value  
351 of 0.25 (Gudmundsson 2011). The E-W striking profile hosts predominantly horizontal layers  
352 where the layer thicknesses are taken from geological measurements (Fig. 2) and given in  
353 Appendix 1. All models are fixed at the corners, with boundary loads applied at the west and  
354 east edges and a free surface (a region free from shear stress) prescribed on the upper edge  
355 (Earth's surface).

356 In addition to boundary loads prescribed at the edge of the models, to simulate tectonic  
357 stressing, we also load the internal cavities to simulate excess magma pressure, which is 5 MPa  
358 in Figure 6. Magma-chamber rupture and dyke injection occur when the tensile stresses at any  
359 point at the boundary of the chamber/reservoir reach the tensile strength of the rock (0.5 to 9  
360 MPa) (Amadei and Stephenson 1997). Laboratory tensile strengths of rocks reach up to about  
361 30 MPa, but the *in-situ* tensile strengths are between 0.5 and 9 MPa, the most common values  
362 being 2-4 MPa (Gudmundsson 2011). By using excess pressure in the chamber/reservoir rather  
363 than total pressure, the effects of gravity are automatically considered (cf. Gudmundsson 2012).  
364 We use a triangular mesh with a maximum element size of 16 m and a minimum element size  
365 of 2 m. Our simplified models show that the most likely area of chamber rupture and surface  
366 eruption is fed by interconnected magma reservoirs, shallow and deeper magma chambers (Fig.  
367 7).

## 368 5.2. Results

369 To explore the potential magma propagation paths in the shallow crust beneath the Ağrı  
370 Dağı volcano, we constructed a numerical model (Fig. 7). It is first necessary to consider the  
371 stress required for magma chamber rupture. In the simplest terms, a magma chamber roof will  
372 rupture and inject a dyke (or an inclined sheet) when (Gudmundsson 1990, 2011):

$$373 \quad p_l + p_e = \sigma_3 + T_0 \quad (5)$$

374 where  $p_l$  is the lithostatic pressure and  $p_e$  is the excess pressure in the magma chamber,  $\sigma_3$   
375 is minimum principal compressive stress in the host rock, and  $T_0$  is the tensile strength of the  
376 host rock, which ranges from 0.5 to 9 MPa (Amadei and Stephenson 1997) and the average *in*  
377 *situ* tensile strength of the upper crust in East Turkey is around 3.5 MPa (in agreement with the  
378 common *in-situ* tensile strength range given above). When a chamber roof has failed in tension  
379 and a dyke is initiated then the magma follows the path or trajectories of maximum principal  
380 compressive stress,  $\sigma_1$  (Gudmundsson 2011). Here we present first the results on crustal stresses  
381 induced solely by magmatic excess pressure within each chamber, ignoring initially the effects  
382 of any regional tectonic loading. In Fig. 7 we show the magnitudes of the minimum principal  
383 compressive (maximum tensile) stress,  $\sigma_3$ , and von Mises shear stress,  $\tau$ .

384 In an E-W profile, the maximum tensile and shear stresses concentrate at the lateral  
385 margins of each magma chamber and at the Earth's surface above the magma chamber.

386 Complex stress patterns and interactions occur at depth due to the attitude and mechanical  
387 properties of the layers (Fig. 7a). There is a stress concentration zone or link between the deeper  
388 magma reservoir and the shallow chamber (Fig. 7b). Our model indicates that if magma  
389 propagates from the edge of the deeper reservoir it can reach the surface without interaction  
390 with the shallow chamber (Fig. 7a-b). However, this is partially dependant on the size and  
391 position of the deeper reservoir with respect to the shallow chamber. When the reservoir is  
392 smaller (Fig 7c) we find there is more likelihood of interaction with the shallow chamber. Here,  
393 the result show that the deeper magma reservoir has two options, so as to either 1) feed the  
394 volcanic edifice from the lateral margins or 2) replenish the shallow magma chamber. Dykes  
395 that propagate from the central part will not feed an eruption but instead charge the shallow  
396 magma chamber. These models indicate that most lava flows at the central part of the volcano  
397 will produce more evolved lavas compared to those lavas fed from the reservoir margins.

## 398 **6. Discussion**

### 399 *6.1. Magma discharge mechanism*

400 Field studies and stratigraphy of the volcano indicate three major andesitic and two basaltic  
401 lava flow eruption cycles, with tens of intermediate-composition lava stacks, from cone  
402 building to late stage of the Ağrı Dağı volcano (Fig. 2; [Yılmaz et al. 1998](#)). We focus on the  
403 latest basaltic lava flows (~0.5 Ma; [Sanver 1968](#)) which record a single magmatic pulse and  
404 path from chamber to the surface. The combined volume of lava flows I and II represents only  
405 0.06 % of the volume of the estimated magma reservoir.

406 The variety of volcanic products along Ağrı Dağı volcano range from contemporaneous  
407 intermediate (dacitic and andesitic) to basic (basaltic) eruptions, indicating that the magma in  
408 this volcanic edifice may be derived from double magma chambers rather than a single magma  
409 source. The more evolved intermediate volcanic rocks (e.g. dacite and andesite) are generally  
410 concentrated at the central part of the edifice while the less evolved basaltic rocks are  
411 distributed at the margin. The injection of dykes from the central part of the deeper magma  
412 chamber (magma reservoir) could feed the shallow magma chamber while dyke injection from  
413 the margin of the deeper magma reservoir can propagate directly to the surface to feed  
414 eruptions. Field observations and the numerical ~~model~~ models are consistent with this  
415 distribution, where less evolved magma can be observed around the periphery of the volcanic  
416 edifice whereas more evolved lava flows are present around the central part.

417 The results of the numerical model, supported by geochemical data, indicate that dyke  
418 injection from the central part of a deep magma reservoir could feed a shallow magma chamber.  
419 The magma arriving at the shallower depths could then begin a fractionation or differentiation  
420 process prior to the chamber rupture condition ( $p_e = T_0 \approx 5MPa$ ) being reached. Thus, we  
421 suggest that the shallow magma chamber produces more evolved magma (e.g. the young  
422 andesitic rocks of age 0.1-0.02 Ma; [Nagao et al. 1989](#)); whereas the deep-seated magma  
423 reservoir produces the older and less evolved lavas (e.g. 0.3-0.049 Ma basaltic rock).

424 The magma reservoir volume underneath Ağrı Dağı appears to have reduced considerably  
425 over a period of 0.5 Ma. Our models indicate a volume reduction from 12,345 km<sup>3</sup> for lava  
426 flow I to 2403 km<sup>3</sup> for lava flow II. Magma storage shrinkage has been interpreted at other  
427 volcanic provinces such as in Iceland ([Andrew and Gudmundsson 2007](#)) and at the Al Haruj  
428 Volcanic Province, central Libya ([Elshaafi and Gudmundsson 2016](#)) (Fig. 8).

429 The first volcanic eruptions may be envisaged as injection from the margins of the deeper  
430 part of reservoir, whereas the second volcanic units may be injected from the uppermost part  
431 of the reservoir where more fractionated (lighter) basaltic rocks form. These basaltic magmas  
432 tend to occupy the uppermost part of a reservoir due to buoyancy. This process might explain  
433 why the volume of the reservoir changed substantially through time.

434 The sizes and areas of individual volcanic eruptions are mainly dependent on the sizes of  
435 the source magma chambers. There are many examples around the world showing that  
436 individual volcanic eruptions can occur on the order of several hundred square kilometres and  
437 have volumes exceeding several cubic kilometres. In fact, the largest basaltic lava flows reach  
438 estimated volumes of thousands of cubic kilometres (Fig. 8). Much more commonly, however,  
439 lava flows cover only small areas and have volumes less than 0.5 km<sup>3</sup>. As an example of a large  
440 historical lava flow, the 1783 Laki lava in Iceland covers 565 km<sup>2</sup> and has a volume of about  
441 15 km<sup>3</sup> (Fig. 8). Also, some prehistorical (mainly 16-17 Ma old) individual lava flows of the  
442 Columbia River Plateau exceed volumes of 1000 km<sup>3</sup>. By contrast the lava flow erupted during  
443 the Krafla Fires in North Iceland, 1975-1984, covers an area of 0.3 km<sup>2</sup> and its volume is about  
444 0.17 km<sup>3</sup> ([Tryggvason 1984](#)), while Etna lava flow for the 1991-1993 eruption has an area of  
445 7.2 km<sup>2</sup> and an estimated volume between 0.022 and 0.072 km<sup>3</sup> ([Harris et al. 2000](#)) (Fig. 8).  
446 Many eruptions of 1-10 km<sup>3</sup> and even larger can be explained by local volcano-tectonic forcing  
447 (e.g. [Gudmundsson 2015, 2016](#)) or continuous supply from a large deeper reservoir to the

448 shallow chamber during the eruption (Gudmundsson 1987). By contrast ‘normal’ or small  
449 eruptions are usually less than 0.1 km<sup>3</sup> and commonly fed by crustal shallow magma chambers  
450 with little or no continuous magma replenishment from a large deeper reservoir during the  
451 eruption (Gudmundsson 1987, 2016). Thus, in the absence of evidence for local volcano-  
452 tectonic forcing, we assume that both lava flows I and II were emplaced from a deep reservoir  
453 in a normal eruption. This notion is supported by the chemistry of the lavas which indicates  
454 primitive magma, particularly of the larger lava flow.

455 To explain the difference in the volumes and chemistry of the lava flows, there are several  
456 possibilities. One possibility is that the size or volume of the entire reservoir decreased greatly  
457 between the two eruptions, in which case reduction in ‘effective size’ corresponds to reduction  
458 in true size. This possibility cannot be ruled out, but the reduction in size would then have to  
459 have happened within the time period of, at maximum, a few hundred thousand years (the lava  
460 flows are younger than 500,000 years). This is possible, but not very likely given that reservoir  
461 feeding volcanic systems, such as in Iceland, are commonly active for 0.5-1 Ma  
462 (Gudmundsson, 2006, 2012), and in many other areas similar reservoir are active for as long  
463 as millions of years. We therefore propose that the second and smaller eruption was supplied  
464 with magma from only a part of the reservoir, that is, for a compartment within the reservoir  
465 (see Gulen 1984 for discussion of the origin of the lavas). This suggestion is supported by the  
466 second and smaller lava flow being more evolved than the first and larger flow. It is clear  
467 particularly at the margin area of the Ađrı Dađı volcano (e.g. Kheirkhah et al. 2009). Thus, we  
468 suggest that only a fraction of the entire reservoir, a compartment (cf. Gudmundsson, 2012),  
469 contributed to the second eruption, thereby, partly at least, explaining their volume and  
470 chemical differences. Based on our calculations, the volume of that compartment is 2403 km<sup>3</sup>,  
471 or roughly 1/5 that of the entire reservoir. Formation and maintenance of compartments in  
472 magma sources is discussed by Gudmundsson (2012). Furthermore, based on our numerical  
473 studies, this compartment was most likely at one of the margins of the reservoir.

## 474 6.2. Tomography

475 The tomographic data ~~indieates~~indicate the presence of an active deep magma reservoir  
476 having low P-wave velocities that extend to the base of the upper crust (Fig. 5). The magma  
477 reservoir may extend between 20-30 km in depth and 35-45 km in width, showing a NW–SE-  
478 elongated tabular form (sill-like shape) in the crust (Figs. 5a-c). A diapiric-shaped dyke  
479 injection extending to the upper level of the crust in a NE–SW oriented profile (Fig. 5d) is



480 clearly observed. In all profiles, we note that diapiric-shaped dyke injection feeding the main  
481 vent of the Ağrı Dağı volcano are not aligned below the main volcanic centers (Figs. 5a-d). It  
482 seems there is no magma source below the Lesser Ağrı Dağı volcano. Greater Ağrı Dağı  
483 volcano is not situated directly over the centre of the large deeper reservoir. This suggests that  
484 the reservoir may have migrated laterally following constructing of the Ağrı Dağı volcano  
485 during the past 1.5 Ma. The shallow magma chamber may be fossilised as a plutonic body  
486 directly below the Ağrı Dağı volcano, which would not be possible to detect it with  
487 tomographic imaging.

488 At least 4 historical volcanic eruptions are known to have occurred from Ağrı Dağı  
489 volcano (Karakhanian et al. 2002): (i) pyroclastic flow in 1840 AD from Greater Ağrı Dağı  
490 volcano, (ii) unclear eruption type in 1450 AD from the SE slope of the Lesser Ağrı Dağı  
491 volcano, (iii) unclear eruption type in late 3<sup>rd</sup>-early 4<sup>th</sup> century AD from Greater Ağrı Dağı  
492 volcano, and (iv) explosive eruption-pyroclastic flow in 2500-2400 BC from the N-NE slope  
493 of Greater Ağrı Dağı volcano. Taking into account the huge magma reservoir below the  
494 volcano even a small future eruptive event coupled with volcano-flank instabilities could  
495 therefore pose a threat to the large populations living around Ağrı Dağı volcano, in Eastern  
496 Turkey and in the Armenian province.

### 497 *6.3. Numerical models in the geological context*

498 Our general numerical results provide insights into the mechanism of magma  
499 movement from a deep magma reservoir to the surface. Such a process can occur in two  
500 predominant ways: (i) the magma is fed directly to the surface from the lateral margins of the  
501 deep reservoir, or (ii) when the magma of deep origin is injected from the central part of the  
502 reservoir, the magma path (the dyke) connects with a shallow chamber which, in turn, ruptures  
503 and propagates a dyke to the surface. In the second case any erupting magma is then technically  
504 fed from the shallow chamber. Despite the tomography data which support an active deep  
505 magma reservoir (20-30 km in depth), the huge volume of intermediate and acidic lavas  
506 constructed at Ağrı Dağı stratovolcano (see Fig. 2) and other large stratovolcanoes most likely  
507 require the formation of a shallow magma chamber.

508 When taken together all of our results indicate that the bulk volume of the reservoir  
509 appears to be considerably reduced between the time of erupting Lava flow I and Lava flow II.  
510 The smaller size of the later magma reservoir increases the likelihood of interaction with the

511 shallow chamber, assuming it has not already solidified which seems to be the case in Ararat  
512 volcano. Regardless of the size of each individual chamber, the conditions for rupture remain  
513 the same, namely that the excess pressure must exceed the tensile strength of the wall rocks  
514 (Eq. 5). In both cases tested numerically (Fig. 7) we find that this failure is most likely at the  
515 margins of the chamber. Therefore more evolved basaltic magma ~~remains~~ inside reservoir  
516 during the quiescent time among two eruptions may be moved upward ~~compartment~~ due to  
517 buoyancy effects ~~of the reservoir to form compartment at the uppermost of the reservoir~~  
518 (Gudmundsson 2012), and then ~~reservoir it would be~~ ruptures ~~after while~~ when Eq. 5 becomes  
519 satisfied again (Fig. 7c).

520 Basaltic rocks generated in lava flow II are generally more fractionated than lava flow  
521 I which is exactly as expected. The lack of data concerning the petrogenesis and geochemistry  
522 for both volcanic flows makes further analysis challenging. We therefore encourage a  
523 systematic field survey which would greatly improve the understanding of Ağrı Dağı volcano.  
524 We hope that this paper encourages further research into this volcano.

## 525 7. Conclusions

526 1) We calculated the total injected materials  $V_e$  for two of the most recent basaltic eruptions at  
527 the Ağrı Dağı volcano. Lava flow I is around  $3.2 \text{ km}^3$  while the lava flow II is around  $0.6 \text{ km}^3$ .  
528 In addition, we present an approach for estimating the volume of the reservoir supplying each  
529 individual volcanic eruption. The effective reservoir volumes obtained were  $12,345 \text{ km}^3$  and  
530  $2403 \text{ km}^3$  for lava flows I and II, respectively.

531 32) Results of seismic tomography reveal a low-velocity zone at a depth of 20 to 30 km below  
532 the northwest part of the Ağrı Dağı volcano which interpret to be a deep magma reservoir. We  
533 do not find strong evidence of a shallow magma source from the present velocity models. This  
534 may indicate that the shallow magma chamber has already solidified.

535

536 23) We explore two scenarios to explain the difference in volume of these two flows. One is  
537 that the absolute reservoir volume decreased between the two eruptions. This is possible, but  
538 not very likely since the likely time between the eruptions is not very large in comparison with  
539 the lifetimes of large reservoirs. The other scenario involves reservoir compartments. In this  
540 scenario, while the less evolved lavas around the volcano was feeding only by deep reservoir,  
541 a comparatively small compartment within the reservoir contributed magma to the eruption,

542 which generated lava flows I and II at Ağrı Dağı volcano. In this view, calculated reservoir  
543 volume of 2403 km<sup>3</sup> for issuing lava flow II thus corresponds to that compartment and is about  
544 1/5 of the total volume of the reservoir.

545 ~~3) Results of seismic tomography reveal a low velocity zone at a depth of 20 to 30 km below~~  
546 ~~the northwest part of the Ağrı Dağı volcano which interpret to be a deep magma reservoir. We~~  
547 ~~do not find strong evidence of a shallow magma source from the present velocity models. This~~  
548 ~~may indicate that the shallow magma chamber has already solidified.~~

549 4) The combined results from our tomography models and analytical calculations were used to  
550 prepare a suite of numerical models. By simulating various crustal loading situations we show  
551 the most likely stress state that promoted feeder-dyke propagation to erupt lava flows I and II.  
552 Our data is useful in estimating the potential source of future eruptions at Ağrı Dağı volcano.  
553 The interpretation of our numerical models suggests that Ağrı Dağı basaltic volcanism has been  
554 fed by ~~either a shallow magma chamber located at about 8 km depth or lateral ends of~~ a deep-  
555 seated magma reservoir at 20-30 km depth which is supported by the geographical distribution  
556 of these basaltic lava flows.-

557 5) The basaltic magma feeding Ağrı Dağı stratovolcano is enriched in LILE elements which  
558 indicates an interaction between the shallow magma chamber and the deeper magma reservoir  
559 just below the volcano. However, lesser evolved basaltic volcanic rocks at the margin of the  
560 Ağrı Dağı volcano were presumably fed by a deeper magma reservoir with no interaction with  
561 the shallow chamber.

## 562 **Acknowledgments**

563 Özgür Karaoğlu is supported by The Scientific  
564 and Technological Research Council of Turkey (TUBITAK) International Postdoctoral  
565 Research Fellowship Programme. John Browning is supported by NERC project  
566 NE/N002938/1. We are grateful to the Editor Valerio Acocella and the reviewers, Alessandro  
567 Tibaldi and an anonymous reviewer, for comments which greatly improved this work.

568

## 569 **References**

- 570 Al-Lazki A, Seber D, Sandvol E, Turkelli N, Mohamad R, Barazangi M (2003)  
571 Tomographic Pn velocity and anisotropy structure beneath the Anatolian plateau  
572 (eastern Turkey) and the surrounding regions. *Geophys Res Lett* 30:24.  
573 doi:10.1029/2003GL017391.
- 574 Al-Lazki AI, Sandvol E, Seber D, Barazangi M, Turkelli N, Mohamad R (2004) Pn  
575 tomographic imaging of mantle lid velocity and anisotropy at the junction of the  
576 Arabian, Eurasian and African plates. *Geophys J Int* 158:1024–1040.  
577 doi:10.1111/j.1365-246X.2004.02355x.
- 578 Alici P, Temel A, Gourgaud A, Vidal P, Gundogdu MN (2001) Quaternary tholeiitic to  
579 alkaline volcanism in the Karasu Valley, Dead Sea rift zone, Southeast Turkey: Sr-  
580 Nd-Pb-O isotopic and trace-element approaches to crust-mantle interaction. *Int Geol*  
581 *Rev* 43 (2):120–138.
- 582 Amadei B, Stephenson O (1997) *Rock stress and its measurement*. Chapman and Hall,  
583 London.
- 584 Ambraseys N.N, Melville C.P (1982) *A History of Persian Earthquakes*. Cambridge Univ.  
585 Press, New York, p. 219.
- 586 Andrew RE, Gudmundsson A (2007) Distribution, structure, and formation of Holocene  
587 lava shields in Iceland. *J Volcanol Geoth Res* 168(1):137–154.
- 588 Angus DA, Wilson DC, Sandvol E, Ni JF (2006) Lithospheric structure of the Arabian and  
589 Eurasian collision zone in Eastern Turkey from S-wave receiver functions.  
590 *Geophys J Int* 166:1335–1346.
- 591 Barka AA (1992) The North Anatolian fault zone. *Ann Tecton* 6:164–195.
- 592 Barnett ZA, Gudmundsson A (2014) Numerical modelling of dykes deflected into sills to form  
593 a magma chamber. *J Volcanol Geoth Res* 281:1–11.
- 594 Bingöl E, Bal I, Can N (eds) (1989) *Geological map of Turkey, scale 1:2000000*, General  
595 Directorates of Mineral Research and Exploration, Ankara, Turkey.
- 596 Bjornsson H (1983) A natural calorimeter at Grimsvötn; an indicator of geothermal and  
597 volcanic activity. *Jökull* (33):13–18.
- 598 Bozkurt E (2001) Neotectonics of Turkey—a synthesis. *Geodin Acta* 14:3–30.

599 Browning J, Drymoni K, Gudmundsson A (2015) Forecasting magma-chamber rupture at  
600 Santorini Volcano, Greece. *Sci Rep* 5:15785. <http://doi.org/10.1038/srep15785>.

601 Cakir O, Erduran M, Cinar H, Yilmazturk A (2000) Forward modeling receiver functions  
602 for crustal structure beneath station TBZ (Trabzon, Turkey). *Geophys J Int* 140:341–  
603 356.

604 Calvert A, Sandvol E, Seber D, Barazangi M, Roecker S, Mourabit T, Vidal F, Alguacil G,  
605 Jabour N (2000) Geodynamic evolution of the lithosphere and upper mantle beneath  
606 the Alboran region of the western Mediterranean: constraints from travel time  
607 tomography. *J Geophys Res* 105(B5):10871–10898.

608 Cavalié O, Jónsson S (2014) Block-like plate movements in eastern Anatolia observed by  
609 InSAR. *Geophys Res Lett* 41:26–31.

610 Daniels KA, Kavanagh JL, Menand T, Stephen JSR (2012) The shapes of dikes: Evidence for  
611 the influence of cooling and inelastic deformation. *Geol Soc Am Bull* 124(7-  
612 8):1102–1112.

613 Coppola D, Piscopo D, Staudacher T, Cigolini C (2009) Lava discharge rate and effusive  
614 pattern at Piton de la Fournaise from MODIS data. *J Volcanol Geoth Res* 184(1):174–  
615 192.

616 Chester D (1993) *Volcanoes and Society*. Edward Arnold, London.

617 Deb D (2006) *Finite Element Method, Concepts and Applications in Geomechanics*. PHI  
618 Learning Private Limited, New Delhi.

619 Elshaafi A, Gudmundsson A (2016) Volcano-tectonics of the Al Haruh Volcanic Province,  
620 Central Libya. *J Volcanol Geoth Res* 325:189–202.

621 Ercan T, Fujitani T, Madsuda JI, Notsu K, Tokel S, Tadahide UI (1990) Dogu ve güneydogu  
622 Anadolu Neojen-Kuvaterner volkanitlerine iliskin yeni jeokimyasal, radyometrik ve  
623 izotopik verilerin yorumu: *M.T.A Dergisi* 110:143–164.

624 Gerbault M (2012) Pressure conditions for shear and tensile failure around a circular magma  
625 chamber; insight from elasto-plastic modelling. *Geol Soc (Lond.) Spec*  
626 *Publ* 367(1):111–130.

- 627 Gudmundsson A (1986) Mechanical aspects of postglacial volcanism and tectonics of the  
628 Reykjanes Peninsula, Southwest Iceland. *J Geophys Res-Solid* 91(B12):12711–  
629 12721.
- 630 Gudmundsson A (1987) Geometry, formation and development of tectonic fractures on the  
631 Reykjanes Peninsula, southwest Iceland. *Tectonophysics* 139(3-4):295–308.
- 632 Gudmundsson A (1990) Emplacement of dikes, sills and crustal magma chambers at  
633 divergent plate boundaries. *Tectonophysics* 176:257–275.
- 634 Gudmundsson A (2006) How local stresses control magma-chamber ruptures, dyke  
635 injections, and eruptions in composite volcanoes. *Earth-Sci Rev* 79:1–31.
- 636 Gudmundsson A (2011) *Rock Fractures in Geological Processes*. Cambridge University Press,  
637 Cambridge.
- 638 Gudmundsson A (2012) Magma chambers: formation, local stresses, excess pressures, and  
639 compartments. *J Volcanol Geoth Res* 237–238:19–41.
- 640 Gudmundsson A (2015) Collapse-driven large eruptions. *J Volcanol Geoth Res* 304:1–10.
- 641 Gudmundsson A (2016) The mechanics of large eruptions. *Earth Sci Rev.*  
642 <http://dx.doi.org/10.1016/j.earscirev.2016.10.003>.
- 643 Gurocak Z, Solanki P, Alemdag S, Zaman MM (2012) New considerations for empirical  
644 estimation of tensile strength of rocks. *Eng Geol* 145:1–8.
- 645 Gulen L (1984) Sr, Nd, Pb isotope and trace element geochemistry of calc-alkaline and  
646 alkaline volcanics, Eastern Turkey. Doctoral dissertation, Massachusetts Institute of  
647 Technology.
- 648 Hasenaka T (1994) Size, distribution, and magma output rate for shield volcanoes of the  
649 Michoac in-Guanajuato volcanic field, Central Mexico. *J Volcanol Geoth Res* 63:13–  
650 31.
- 651 Hearn TM (1999) Uppermost mantle velocities and anisotropy beneath Europe. *J Geophys Res*  
652 104:15123–15139.
- 653 Hermance JF (1981) Crustal genesis in Iceland: Geophysical constraints on crustal  
654 thickening with age. *Geophys Res Lett* 8(3): 203–206.

- 655 Innocenti F, Mazzuoli R, Pasquare G, Redicat di Brozolo F, Villari L (1976) Evolution of the  
656 volcanism in the area of interaction between the Arabian, Anatolian and Iranian plates  
657 (Lake Van, Eastern Turkey). *J Volcanol Geoth Res* 1:103–112.
- 658 Innocenti F, Mazzuoli R, Pasquare G, Serri G, Villari L (1980) Geology of the volcanic  
659 area north of Lake Van (Turkey). *Geol Rundsch* 69: 292–322.
- 660 Irvine TN, Baragar WRA (1971) A guide to the chemical classification of the common  
661 volcanic rocks. *Can J Earth Sci* 8:523–548.
- 662 Karakhanian A, Djrbashian R, Trifonov V, Philip H, Arakelian S, Avagian A (2002)  
663 Holocene–historical volcanism and active faults as natural risk factors for Armenia  
664 and adjacent countries. *J Volcanol Geoth Res* 113:319–344.
- 665 Karaoğlu Ö, Özdemir Y, Tolluoğlu AÜ, Karabıykoğlu M, Köse O, Froger JL (2005)  
666 Stratigraphy of the volcanic products around Nemrut Caldera: implications for  
667 reconstruction of the Caldera Formation. *Turk J Earth Sci* 14:123–143.
- 668 Karaoğlu Ö, Browning J, Bazargan M, Gudmundsson A (2016) Numerical modelling of  
669 triple-junction tectonics at Karlıova, Eastern Turkey, with implications for regional  
670 magma transport. *Earth Planet Sc Lett* 452:152–170.
- 671 Kervyn M, Ernst GG, Klaudius J, Keller J, Kervyn F, Mattsson HB, Belton F, Mbede E,  
672 Jacobs P (2008) Voluminous lava flows at Oldoinyo Lengai in 2006: chronology of  
673 events and insights into the shallow magmatic system. *B Volcanol* 70(9):1069–1086.
- 674 Keskin M (2007) Eastern Anatolia: a hot spot in a collision zone without a mantle plume. In:  
675 Foulger GR, Jurdy D(eds). *Plates, Plumes, and Planetary Processes*. *Geol Soc Am*  
676 *Spec Pap* 430:693722.
- 677 Keskin M, Pearce JA, Mitchell JG (1998) Volcano-stratigraphy and geochemistry of  
678 collision-related volcanism on the Erzurum-Kars Plateau, North Eastern Turkey. *J*  
679 *Volcanol Geoth Res* 85:355–404.
- 680 Kheirkhah M, Allen MB, Emami M (2009) Quaternary syn-collision magmatism from the  
681 Iran/Turkey borderlands. *J Volcanol Geoth Res* 182(1):1–12.
- 682 Le Corvec N, Menand T, Lindsay J (2013) Interaction of ascending magma with pre-  
683 existing crustal fractures in monogenetic basaltic volcanism: an experimental approach. *J*  
684 *Geophys Res-Sol Ea* 118(3):968–984.

- 685 Le Corvec N, McGovern PJ, Grosfils EB, Galgana G (2015) Effects of crustal-scale  
686 mechanical layering on magma chamber failure and magma propagation within the  
687 Venusian lithosphere. *J Geophys Res* 120(7):1279–1297.
- 688 Lebedev VA, Sharkov EV, Keskin M, Oyan V (2010) Erratum: Geochronology of Late  
689 Cenozoic volcanism in the area of Lake Van, Turkey: an example of developmental  
690 dynamics for magmatic processes. *Dokl Earth Sci* 433(2):1031–1037.
- 691 Lustrino M, Keskin M, Mattioli M, Lebedev V, Chugaev A, Sharkov E, Kavak O (2010).  
692 Early activity of the largest Cenozoic shield volcano in the circum- Mediterranean  
693 area: Mt. Karacadag SE Turkey. *Eur J Mineral* 22:343–362.
- 694 Maccaferri F, Rivalta E, Keir D, Acocella V (2014) Off-rift volcanism in rift zones determined  
695 by crustal unloading. *Nat Geosci* 7(4):297–300.
- 696 Murase T, McBirney AR (1973) Properties of some common igneous rocks and their melts at  
697 high temperatures. *Geol Soc Am Bull* 84(11):3563–3592.
- 698 Meyer PS, Sigurdsson H, Schilling JG (1985) Petrological and geochemical variations along  
699 Iceland's neovolcanic zones. *J Geophys Res-Solid* 90(B12): 10043–10072.
- 700 Nagao K, Matsuda JI, Kita I, Ercan T (1989) Noble gas and carbon isotopic  
701 compositions in Quaternary volcanic area in Turkey. *Bull Geomorphol* 17:101–110.
- 702 Notsu K, Fujitani T, Ui T, Matsuda J, Ercan T (1995) Geochemical features of collision-  
703 related volcanic rocks in central and eastern Anatolia, Turkey. *J Volcanol Geoth Res*  
704 64: 171–192. doi: 10.1016/0377-0273(94)00077-T.
- 705 Okay AI, Tüysüz O (1999) Tethyan sutures of northern Turkey. *Geol Soc Spec Publ* 156:475–  
706 515.
- 707 Ozacar AA, Gilbert H, Zandt G (2008) Upper mantle discontinuity structure beneath East  
708 Anatolian Plateau (Turkey) from receiver functions. *Earth Planet Sc Lett* 269:426–  
709 434.
- 710 Örgülü G, Aktar M, Türkelli N, Sandvol E, Barazangi M (2003) Contribution to the  
711 seismotectonics of Eastern Turkey from moderate and small size events. *Geophysc*  
712 *Res Lett* 30:(24).
- 713 Pearce JA, et al. (1990) Genesis of collision volcanism in Eastern Anatolia, Turkey. *J Volcanol*  
714 *Geotherm Res* 44:189–229.



- 715 Reidel SP, Camp VE, Tolan TL, Martin BS (2013) The Columbia River flood basalt  
716 province: stratigraphy, areal extent, volume, and physical volcanology. *Geol Soc Am*  
717 *Spec Pap* 497:1–43.
- 718 Reilinger R, et al (2006) GPS constraints on continental deformation in the Africa-Arabia-  
719 Eurasia continental collision zone and implications for the dynamics of plate  
720 interactions. *J Geophys Res* 111:B05411. doi:10.1029/2005JB004051.
- 721 Richter FM, McKenzie D (1984) Dynamical models for melt segregation from a deformable  
722 matrix. *J Geol* 92:729–740.
- 723 Riel B, Milillo P, Simons M, Lundgren P, Kanamori H, Samsonov S (2015) The collapse of  
724 Bárðarbunga caldera, Iceland. *Geophys J Int* 202(1): 446–453.
- 725 Rodgers AJ, Ni JF, Hearn TM (1997) Propagation characteristics of short-period Sn and Lg  
726 in the Middle East. *B Seismol Soc Am* 87:396–413.
- 727 Salah MK, Sahin S, Aydin U (2011) Seismic velocity and Poisson's ratio tomography of the  
728 crust beneath east Anatolia. *J Asian Earth Sci* 40:746–761.  
729 doi:10.1016/j.jseaes.2010.10.021.
- 730 Sanver M (1968) A palaeomagnetic study of Quaternary volcanic rocks from Turkey. *Phys*  
731 *Earth Planet Inter* 1:403–421.
- 732 Scott DR, Stevenson DJ (1986) Magma ascent by porous flow. *J Geophys Res-Sol*  
733 *Ea* 91(B9):9283–9296.
- 734 Selçuk AS, Erturaç MK, Nomade S (2016) Geology of the Çaldıran Fault, Eastern  
735 Turkey: Age, slip rate and implications on the characteristic slip  
736 behaviour. *Tectonophysics* 680:155–173.
- 737 Small C, Naumann T (2001) The global distribution of human population and recent  
738 volcanism. *Global Environmental Change Part B. Environmental Hazards* 3(3):93–  
739 109.
- 740 Şengör AMC, Özeren MS, Keskin M, Sakıncı M, Özbakır AD, Kayan I (2008) Eastern  
741 Turkish high plateau as a small Turkic-type orogen: implications for post-collisional  
742 crust-forming processes in Turkic-type orogens. *Earth Sci Rev* 90:1–48.
- 743 Shabanian E, Acocella V, Gioncada A, Habibollah G, Belier O (2012) Structural control on  
744 volcanism in intraplate post collisional settings: Late Cenozoic to Quaternary examples  
745 of Iran and eastern Turkey. *Tectonics* 31:TC3013.

- 746 Tezcan A (1995) Geothermal explorations and heat flow in Turkey. In: Gupta ML, Yamamo  
747 M (Eds.). *Terrestrial Heat Flow and Geothermal Energy in Asia*. Oxford and IBH  
748 Publishing Co. Pvt. Ltd., New Delhi, pp, 23–42.
- 749 Thordarson T, Self S (1993) The Laki (Skaftár Fires) and Grímsvötn eruptions in 1783–  
750 1785. *B Volcanol* 55(4):233–263.
- 751 Tibaldi A (2015) Structure of volcano plumbing systems: A review of multi-parametric  
752 effects. *J Volcanol Geotherm Res* 298:85–135.
- 753 Tryggvason E (1984) Widening of the Krafla fissure swarm during the 1975–1981 volcano-  
754 tectonic episode. *Bull Volcanol* 47:47–69.
- 755 Tuffen H, James MR, Castro JM, Schipper CI (2013) Exceptional mobility of an advancing  
756 rhyolitic obsidian flow at Cordón Caulle volcano in Chile. *Nature Communications* 4.
- 757 Türkünal S (1980) Doğu ve Güneydoğu Anadolu'nun jeolojisi. TMMOB Jeoloji  
758 Mühendisleri Odası.
- 759 Vernant P, Nilforoushan F, Chéry J, Bayer R, Djamour Y, Masson F, Nankali H, Ritz JF,  
760 Sedighi M, Tavakoli F (2004) Deciphering oblique shortening of central Alborz in  
761 Iran using geodetic data. *Earth Planet Sci Lett* 223, 177–185.  
762 doi:10.1016/j.epsl.2004.04.017.
- 763 White JDL, Houghton B (2000) Surtseyan and related phreatomagmatic  
764 eruptions. *Encyclopedia of volcanoes*. Academic, San Diego, pp. 495–513.
- 765 Yılmaz Y, Guner Y, Saroglu F (1998) Geology of the Quaternary volcanic centres of the  
766 East Anatolia. *J Volcanol Geoth Res* 85:173–210.
- 767 Zhao D, Hasegawa A, Horiuchi S (1992) Tomographic imaging of P- and S-wave velocity  
768 structure beneath northeastern Japan. *J Geophys Res* 97:19909–19928.
- 769 Zhao D, Hasegawa A, Kanamori H (1994) Deep structure of Japan subduction zone as derived  
770 from local, regional and teleseismic events. *J Geophys Res* 99:22313–22329.
- 771 Zhao D, Yanada T., Hasegawa A, Umino N, Wei W (2012) Imaging the subducting slabs and  
772 mantle upwelling under the Japan Islands. *Geophys J Int* 190:816–828.
- 773 Zienkiewicz OC (1979) *The Finite Element Method*. McGraw-Hill, New York, p. 787.

774 Zor E, Sandvol E, Gurbuz C, Turkelli N, Seber D, Barazangi M (2003) The crustal  
775 structure of East Anatolian plateau from receiver functions. Geophys Res Lett:30.  
776 doi:10.1029/2003GLO18192.

777

778

779

780

781

782

783

784

785

786

787

788

789

790

791

792

793

794

795

796

797

798

799

800

801 **Figure captions**

802 Figure 1. a) Map of the tectonic framework of Turkey; b) middle Miocene to recent volcanic  
803 centers in Eastern Turkey and location of population centres on a DEM-derived map. NAF:  
804 North Anatolian Fault, EAF: Eastern Anatolian Fault, KTJ: Karliova Triple Junction, VFZ:  
805 Varto Fault Zone.

806  
807 Figure 2. a) Ağrı Dağı volcano and surrounding region; b) volcano-stratigraphy of the Ağrı Dağı  
808 volcano; c) geological map of the Ağrı Dağı volcano; d) geological map of the last two basaltic  
809 lava eruptions, flows I and II.

810  
811 Figure 3. The last basaltic/most basaltic lava flows around Ağrı Dağı volcano. a-b) Google  
812 Earth Images of basaltic lava flows at around the Great and Lesser Ağrı Dağı volcanoes; c-d)  
813 Images of the most recent lava flows (I-II).

814  
815 Figure 4. a)  $K_2O+Na_2O-SiO_2$  (TAS) (Le Maitre 2002) diagram for the rock samples around  
816 Ağrı Dağı volcano (data taken from Gulen 1984; Pearce et al. 1990); alkaline-subalkaline line  
817 is according to Irvine and Baragar (1971); b) MORB-normalized multi-element diagrams for  
818 the volcanic rocks of the basaltic/most basaltic rock samples around Ağrı Dağı volcano (data  
819 taken from Gulen 1984; Pearce et al. 1990; Kheirkhah et al. 2009). Normalizing values are  
820 from Sun and McDonough (1989).

821  
822 Figure 5. (a-e) Five vertical cross-sections of P-wave velocity beneath the area of the Ağrı Dağı  
823 volcano (see Fig. 1 for the location of the cross-sections). Low velocities are shown in red,  
824 whereas high velocities are shown in blue. Large stars and small circles show, respectively, the  
825 location of moderate/large earthquakes ( $M \geq 5.0$ ) and the microseismic activity in a 30 km  
826 wide-zone around the profile. The perturbation scale ( $\pm 5\%$ ) is shown to the right. (f) The  
827 locations for these seismic profiles on the map.

828 Figure 6. 2-D numerical model setups. The 2-D example shown represents the geology of an  
829 E-W striking profile through Ağrı Dağı volcano. All 2-D models are layered  $E_{(1-14)}$  with each  
830 unit assigned a different value of Young's modulus. Magma chambers, represented as cavities,  
831 are given an excess pressure of 5 to 15 MPa.

832 Figure 7. Modelled stresses induced by excess magmatic pressure ( $p_e$ ) inside a shallow  
833 chamber of diameter 16 km and a deep reservoir of diameter 40 km. a) Magnitudes of the  
834 minimum principal compressive (maximum tensile) stress ( $\sigma_3$ ). b) Magnitude of von Mises  
835 shear stresses ( $\tau$ ) The excess magmatic pressure in each chamber is 5 MPa and is the only  
836 loading. Parts c) and d) show the same arrangement of shallow chamber and deep reservoir but  
837 in this case the reservoir is reduced in size with a diameter of 16 km, and 30 km in respectively.

838  
839 Figure 8. a) Area vs. volume constraints of some single lava discharge ratios for composite  
840 volcanoes and volcanic regions; b) Volume compilation for some historical eruptions. (1) This  
841 study; (2) Kervyn et al. (2008); (3) Haris et al. (2000); (4) Tryggvason (1984); (5) Thordarson

842 [and Self \(1993\)](#); (6) [Reidel et al. \(2013\)](#); (7) [White and Houghton \(2000\)](#). Arrows highlighting  
843 the last two basaltic eruptions of Ağrı Dağı volcano.