

PHYSICAL REVIEW C **71**, 055501 (2005)**Superaligned  $0^+ \rightarrow 0^+$  nuclear  $\beta$  decays: A critical survey with tests of the conserved vector current hypothesis and the standard model**J. C. Hardy<sup>1,\*</sup> and I. S. Towner<sup>1,2</sup><sup>1</sup>*Cyclotron Institute, Texas A&M University, College Station, Texas 77843*<sup>2</sup>*Physics Department, Queen's University, Kingston, Ontario K7L 3N6, Canada*

(Received 13 December 2004; published 19 May 2005)

A complete and critical survey is presented of all half-life, decay-energy, and branching-ratio measurements related to 20 superallowed  $0^+ \rightarrow 0^+$  decays; no measurements are ignored, although some are rejected for cause and others updated. A new calculation of the statistical rate function  $f$  is described and experimental  $ft$  values determined. The associated theoretical corrections needed to convert these results into “corrected”  $\mathcal{F}t$  values are discussed, and careful attention is paid to the origin and magnitude of their uncertainties. As an exacting confirmation of the conserved vector current hypothesis, the corrected  $\mathcal{F}t$  values are seen to be constant to three parts in  $10^4$ . These data are also used to set a new limit on any possible scalar interaction (assuming maximum parity violation) of  $C_S/C_V = -(0.0005 \pm 0.00130)$ . The average  $\mathcal{F}t$  value obtained from the survey, when combined with the muon lifetime, yields the up-down quark-mixing element of the Cabibbo-Kobayashi-Maskawa (CKM) matrix,  $V_{ud} = 0.9738 \pm 0.0004$ , and the unitarity test on the top row of the matrix becomes  $|V_{ud}|^2 + |V_{us}|^2 + |V_{ub}|^2 = 0.9966 \pm 0.0014$  using the Particle Data Group's currently recommended values for  $V_{us}$  and  $V_{ub}$ . If  $V_{us}$  comes instead from two recent results on  $K_{e3}$  decay, the unitarity sum becomes  $0.9996(11)$ . Either result can be expressed in terms of the possible existence of right-hand currents. Finally, we discuss the priorities for future theoretical and experimental work with the goal of making the CKM unitarity test more definitive.

DOI: 10.1103/PhysRevC.71.055501

PACS number(s): 23.40.Bw, 12.15.Hh, 12.60.-i

**I. INTRODUCTION**

Precise measurements of the  $\beta$  decay between nuclear analog states of spin,  $J^\pi = 0^+$ , and isospin,  $T = 1$ , provide demanding and fundamental tests of the properties of the electroweak interaction. Collectively, these transitions can sensitively probe the conservation of the vector weak current, set tight limits on the presence of scalar or right-hand currents and contribute to the most demanding available test of the unitarity of the Cabibbo-Kobayashi-Maskawa (CKM) matrix, a fundamental tenet of the electroweak standard model.

Eight transitions,  $^{14}\text{O}$ ,  $^{26}\text{Al}^m$ ,  $^{34}\text{Cl}$ ,  $^{38}\text{K}^m$ ,  $^{42}\text{Sc}$ ,  $^{46}\text{V}$ ,  $^{50}\text{Mn}$ , and  $^{54}\text{Co}$  are particularly amenable to experiment and, because of their significance to physics, have consequently received a good deal of attention over the past few decades. In each of these cases, the experimental  $ft$  value is known to better than 0.1%. In the 1990s,  $^{10}\text{C}$  was added to this list; its  $ft$  value is known to a precision of 0.15%. More recently, three more cases have been added:  $^{22}\text{Mg}$ ,  $^{34}\text{Ar}$ , and  $^{74}\text{Rb}$ , with  $ft$  value standard deviations ranging from from 0.24 to 0.40%. In the near future these uncertainties will undoubtedly be reduced and an additional eight cases could well be added to the list. Though improvements are still possible, with current data we can test the conserved vector current hypothesis at the level of three parts in  $10^4$  and the three-generation standard model at the level of its quantum corrections.

Over the past decade, it has become increasingly clear that the CKM unitarity test made possible by these measurements does not, in fact, quite agree with standard-model expectations.

The test involves the top row of the CKM matrix and requires that the sum of squares of the three experimentally determined elements,  $|V_{ud}|^2 + |V_{us}|^2 + |V_{ub}|^2$ , should equal 1. With results from superallowed  $\beta$  decay providing the input for  $V_{ud}$ , and values for  $V_{us}$  and  $V_{ub}$  taken from the Particle Data Group reviews, the sum falls short of unity by 0.3%, more than twice the quoted standard deviation [1]—a provocative but hardly definitive disagreement. Nevertheless, it has stimulated experimental activity not only on the nuclear decays used to determine  $V_{ud}$  but also on the  $K_{e3}$  branching ratio used for  $V_{us}$ . Strikingly, two new measurements, one of the  $K_{e3}^+$  branching ratio [2] and the other of the  $K_L$  branching fractions [3], have thrown the accepted value of  $V_{us}$  into doubt. Although the new results disagree significantly with previous measurements, they are consistent with one another and would, if taken by themselves, lead to a larger value for  $V_{us}$ : this would bring the CKM top-row sum into agreement with unity. At this time, the value of  $V_{us}$  remains controversial and there are a number of kaon-decay experiments currently underway, which should lead to a settled outcome within a very few years.

With all this activity in progress, and the likelihood that a new and reliable value of  $V_{us}$  will soon be forthcoming, this is an opportune time to produce a complete new survey of the nuclear data used to establish  $V_{ud}$ . This way, we will be able to view the value of  $V_{ud}$  with renewed confidence in anticipation of a revised result for  $V_{us}$ . ( $V_{ub}$  is very small and contributes a negligible 0.001% to the unitarity sum.) We have published four previous surveys, Refs. [4–7]: the most recent appeared 15 years ago and included only 8 superallowed transitions. In addition to bringing the results for these cases up to date, we are now incorporating data on 12 more transitions

\*Electronic address: [hardy@comp.tamu.edu](mailto:hardy@comp.tamu.edu)

and have continued the practice we began in 1984 [6] of updating all original data to take account of the most modern calibration standards. We have also made completely new calculations of the statistical rate function,  $f$ , and employed the most complete radiative and isospin-symmetry-breaking corrections in dealing with the  $ft$  values in the context of fundamental weak-interaction tests.

Superaligned Fermi  $\beta$  decay between  $0^+$  states depends uniquely on the vector part of the hadronic weak interaction. When it occurs between isospin  $T = 1$  analog states, the conserved vector current (CVC) hypothesis indicates that the  $ft$  values should be the same irrespective of the nucleus as given by:

$$ft = \frac{K}{G_V^2 |M_F|^2} = \text{const}, \quad (1)$$

where  $K/(\hbar c)^6 = 2\pi^3 \hbar \ln 2 / (m_e c^2)^5 = (8120.271 \pm 0.012) \times 10^{-10} \text{ GeV}^{-4} \text{ s}$ ,  $G_V$  is the vector coupling constant for semileptonic weak interactions, and  $M_F$  is the Fermi matrix element, which for  $T = 1$  states has the value  $M_F = \sqrt{2}$ . The CVC hypothesis asserts that the vector coupling constant,  $G_V$ , is a true constant and not renormalized to another value in the nuclear medium. A demonstration with the data assembled here that the  $ft$  values are indeed constant would provide a stringent test of the CVC hypothesis.

Unfortunately, Eq. (1) has to be amended slightly. First, there are radiative corrections because, for example, the emitted electron may emit a bremsstrahlung photon, which goes undetected in the experiment. Second, isospin is not an exact symmetry in nuclei so the nuclear matrix element,  $M_F$  is slightly reduced from its ideal value, leading us to write  $|M_F|^2 = 2(1 - \delta_C)$ . Thus, we define a ‘‘corrected’’  $ft$  value as follows:

$$\mathcal{F}t \equiv ft(1 + \delta_R)(1 - \delta_C) = \frac{K}{2G_V^2(1 + \Delta_R^V)} = \text{const}, \quad (2)$$

where  $\delta_C$  is the isospin-symmetry-breaking correction,  $\delta_R$  is the transition-dependent part of the radiative correction, and  $\Delta_R^V$  is the transition-independent part. Fortunately these corrections are all of the order of 1% but, even so, to maintain an accuracy criterion of 0.1% they must be calculated with an accuracy of 10% of their central value. This is a demanding request, especially for the nuclear-structure-dependent corrections.

To separate out those terms that are dependent on nuclear structure from those that are not, we split the transition-dependent radiative correction into two terms:

$$\delta_R = \delta'_R + \delta_{NS}, \quad (3)$$

of which the first,  $\delta'_R$ , is a function only of the electron’s energy and the charge of the daughter nucleus  $Z$ ; it therefore depends on the particular nuclear decay, but is *independent* of nuclear structure. The second term,  $\delta_{NS}$ , like  $\delta_C$ , depends in its evaluation on the details of nuclear structure. To emphasize the different sensitivities of the correction terms, we rewrite the expression for  $\mathcal{F}t$  as follows:

$$\mathcal{F}t \equiv ft(1 + \delta'_R)(1 + \delta_{NS} - \delta_C), \quad (4)$$

where the first correction in brackets is independent of nuclear structure, whereas the second incorporates the structure-dependent terms.

Our procedure is to examine all experimental data related to 20 superallowed transitions, comprising those that have been well studied, together with others that have only recently become accessible to precision measurement. The methods used and the data accepted are presented in Sec. II. The calculations and corrections required to extract final  $\mathcal{F}t$  values from these data are described and applied in Sec. III; in the same section, we use the resulting  $\mathcal{F}t$  values to test CVC. Finally, in Sec. IV we explore the impact of these results on a number of weak-interaction issues: CKM unitarity as well as the possible existence of scalar or right-hand currents.

## II. EXPERIMENTAL DATA

The  $ft$  value that characterizes any  $\beta$  transition depends on three measured quantities: the total transition energy,  $Q_{EC}$ , the half-life,  $t_{1/2}$ , of the parent state and the branching ratio,  $R$ , for the particular transition of interest. The  $Q_{EC}$  value is required to determine the statistical rate function,  $f$ , whereas the half-life and branching ratio combine to yield the partial half-life,  $t$ . In Tables I–VII we present the measured values of these three quantities and supporting information for a total of 20 superallowed transitions, incorporating the 8 cases we dealt with in our last complete survey [7] but now including 4 more cases that have been measured more recently with comparable precision and a further 8 that are likely to become accessible to precision measurements within the next few years.

### A. Evaluation principles

In our treatment of the data, we considered all measurements formally published before November 2004 and those we knew to be in an advanced state of preparation for publication by that date. We scrutinized all the original experimental reports in detail. Where necessary and possible, we used the information provided there to correct the results for calibration data that have improved since the measurement was made. If corrections were evidently required but insufficient information was provided to make them, the results were rejected. Of the surviving results, only those with (updated) uncertainties that are within a factor of 10 of the most precise measurement for each quantity were retained for averaging in the tables. Each datum appearing in the tables is attributed to its original journal reference *via* an alphanumeric code comprising the initial two letters of the first author’s name and the two last digits of the publication date. These alphanumeric codes are correlated with the actual reference numbers [8–153] in Table VIII.

The statistical procedures we have followed in analyzing the tabulated data are based on those used by the Particle Data Group in their periodic reviews of particle properties (e.g. Ref. [154]) and adopted by us in earlier surveys [5,7] of superallowed  $0^+ \rightarrow 0^+ \beta$  decay. In the tables and throughout this work, ‘‘error bars’’ and ‘‘uncertainties’’ always refer to plus/minus 1 standard deviation (68% confidence level). For a set of  $N$  uncoupled measurements,  $x_i \pm \delta x_i$ , of a particular

TABLE I. Decay energies,  $Q_{EC}$ , for superallowed  $\beta$ -decay branches. (See Table VIII for the correlation between the alphanumeric reference code used in this table and the actual reference numbers.)

Parent/daughter nuclei	property <sup>1</sup>	Measured energy, $Q_{EC}$ (keV)			Average value	
		1	2	3	Energy (keV)	Scale
$(T_z = -1)$						
<sup>10</sup> C → <sup>10</sup> B	$Q_{EC}(gs)$	3647.84 ± 0.34 [Ba84]	3647.95 ± 0.12 [Ba98]		3647.94 ± 0.11	1.0
	$E_x(d0^+)$	1740.15 ± 0.17 [Aj88]	1740.07 ± 0.02 <sup>2</sup>		1740.07 ± 0.02	1.0
	$Q_{EC}(sa)$				<b>1907.87 ± 0.11</b>	
<sup>14</sup> O → <sup>14</sup> N	$Q_{EC}(gs)$	5143.35 ± 0.60 [Bu61]	5145.09 ± 0.46 [Ba62]	5145.57 ± 0.48 [Ro70]		
		5142.71 ± 0.80 [Vo77]	5143.43 ± 0.37 [Wh77]	5144.34 ± 0.17 [To03]	5144.29 ± 0.28	2.1
	$E_x(d0^+)$	2312.798 ± 0.011 [Aj91]			2312.798 ± 0.011	
	$Q_{EC}(sa)$				<b>2831.18 ± 0.24<sup>3</sup></b>	<b>2.5</b>
<sup>18</sup> Ne → <sup>18</sup> F	$ME(p)$	5316.8 ± 1.5 [Ma94]	5317.63 ± 0.36 [Bl04b]		5317.58 ± 0.35	1.0
	$ME(d)$	873.31 ± 0.94 [Bo64]	875.5 ± 2.2 [Ho64]	876.5 ± 2.8 [Pr67]		
		877.2 ± 3.0 [Se73]	873.96 ± 0.61 [Ro75]		874.02 ± 0.48	1.0
	$Q_{EC}(gs)$	4438 ± 9 [Fr63]			4443.54 ± 0.60	1.0
	$E_x(d0^+)$	1041.55 ± 0.08 [Ti95]			1041.55 ± 0.08	
	$Q_{EC}(sa)$				<b>3401.99 ± 0.60</b>	
<sup>22</sup> Mg → <sup>22</sup> Na	$ME(p)$	-401.3 ± 3.0 [Ha74c]	-400.5 ± 1.3 <sup>4</sup>		-400.6 ± 1.2	1.0
	$ME(d)$	-5184.3 ± 1.5 [We68]	-5182.5 ± 0.5 [Be68]	-5181.3 ± 1.7 [An70]		
		-5183.2 ± 1.0 [Gi72]	-5181.56 ± 0.16 [Mu04]	-5181.08 ± 0.30 [Sa04]	-5181.58 ± 0.19	1.7
	$Q_{EC}(gs)$	4781.64 ± 0.28 [Mu04]	4781.40 ± 0.67 [Sa04]		4781.58 ± 0.25	1.0
	$E_x(d0^+)$	657.00 ± 0.14 [En98]			657.00 ± 0.14	
	$Q_{EC}(sa)$				<b>4124.58 ± 0.29</b>	
<sup>26</sup> Si → <sup>26</sup> Al	$ME(p)$	-7159 ± 18 [Mi67]	-7149 ± 30 [Mc67]	-7139 ± 30 [Ha68]		
		-7145.5 ± 3.0 [Ha74c]			-7145.8 ± 2.9	1.0
	$ME(d0^+)$	-11982.08 ± 0.19 <sup>5</sup>			-11982.08 ± 0.26	
	$Q_{EC}(sa)$	4850 ± 13 [Fr63]			<b>4836.9 ± 3.0</b>	<b>1.0</b>
<sup>30</sup> S → <sup>30</sup> P	$ME(p)$	-14060 ± 15 [Mi67]	-14054 ± 25 [Mc67]	-14068 ± 30 [Ha68]		
		-14063.4 ± 3.0 [Ha74c]			-14063.2 ± 2.9	1.0
	$ME(d)$	-20203 ± 3 [Ha67]	-20200.58 ± 0.40 [Re85]		-20200.62 ± 0.40	1.0
	$Q_{EC}(gs)$				6137.4 ± 2.9	
	$E_x(d0^+)$	677.29 ± 0.07 [En98]			677.29 ± 0.07	
	$Q_{EC}(sa)$	5437 ± 17 [Fr63]			<b>5459.5 ± 3.9</b>	<b>1.3</b>
<sup>34</sup> Ar → <sup>34</sup> Cl	$ME(p)$	-18380.2 ± 3.0 [Ha74c]	-18377.10 ± 0.41 [He02]		-18377.17 ± 0.40	1.0
	$ME(d)$	-24440.01 ± 0.23 <sup>5</sup>			-24440.01 ± 0.23	
	$Q_{EC}(sa)$				<b>6062.83 ± 0.46</b>	
<sup>38</sup> Ca → <sup>38</sup> K	$ME(p)$	-22056.0 ± 5.0 [Se74]			-22056.0 ± 5.0	
	$ME(d0^+)$	-28670.20 ± 0.32 <sup>5</sup>			-28670.20 ± 0.32	
	$Q_{EC}(sa)$				<b>6614.2 ± 5.0</b>	
<sup>42</sup> Ti → <sup>42</sup> Sc	$ME(p)$	-25121 ± 6 [Mi67]	-25086 ± 30 [Ha68]	-25124 ± 13 [Zi72]	-25120.7 ± 5.3	1.0
	$ME(d)$	-32121.55 ± 0.80 <sup>5</sup>			-32121.55 ± 0.80	
	$Q_{EC}(sa)$				<b>7000.9 ± 5.4</b>	
$(T_z = 0)$						
<sup>26</sup> Al <sup>m</sup> → <sup>26</sup> Mg	$Q_{EC}(gs)$	4004.79 ± 0.55 [De69]	4004.41 ± 0.10 <sup>6</sup>		4004.42 ± 0.10	1.0
	$E_x(p0^+)$	228.305 ± 0.013 [En98]			228.305 ± 0.013	
	$Q_{EC}(sa)$	4232.71 ± 0.60 [Vo77]	4232.19 ± 0.12 [Br94]		<b>4232.55 ± 0.17<sup>3</sup></b>	<b>2.7</b>

TABLE I. (Continued.)

Parent/daughter nuclei	property <sup>1</sup>	Measured energy, $Q_{EC}$ (keV)			Average value		
		1	2	3	Energy (keV)	Scale	
<sup>34</sup> Cl	<sup>34</sup> S	$Q_{EC}(sa)$	5490.3 ± 1.9 [Ry73a]	5491.6 ± 2.3 [Ha74d]	5491.71 ± 0.54 [Ba77c]	<b>5491.78 ± 0.20</b>	<b>1.0</b>
			5492.2 ± 0.4 [Vo77]	5491.65 ± 0.26 <sup>7</sup>			
<sup>38</sup> K <sup>m</sup>	<sup>38</sup> Ar	$Q_{EC}(gs)$	5914.76 ± 0.60 [Ja78]			5914.76 ± 0.60	
		$E_x(p0^+)$	130.4 ± 0.3 [En98]			130.4 ± 0.3	
		$Q_{EC}(sa)$	6044.6 ± 1.5 [Bu79]	6044.38 ± 0.12 [Ha98]		<b>6044.40 ± 0.11</b>	<b>1.0</b>
<sup>42</sup> Sc	<sup>42</sup> Ca	$Q_{EC}(sa)$	6423.71 ± 0.40 [Vo77]	6425.84 ± 0.17 <sup>8</sup>		<b>6425.63 ± 0.38<sup>3</sup></b>	<b>3.2</b>
<sup>46</sup> V	<sup>46</sup> Ti	$Q_{EC}(sa)$	7053.3 ± 1.8 [Sq76]	7050.41 ± 0.60 [Vo77]		<b>7050.71 ± 0.89</b>	<b>1.6</b>
<sup>50</sup> Mn	<sup>50</sup> Cr	$Q_{EC}(sa)$	7632.8 ± 2.8 [Ha74d]	7631.91 ± 0.40 [Vo77]		<b>7632.43 ± 0.23<sup>3</sup></b>	<b>1.0</b>
<sup>54</sup> Co	<sup>54</sup> Fe	$Q_{EC}(sa)$	8241.2 ± 1.8 [Ho74]	8245.6 ± 3.0 [Ha74d]	8241.61 ± 0.60 [Vo77]	<b>8242.60 ± 0.29<sup>3</sup></b>	<b>1.5</b>
<sup>62</sup> Ga	<sup>62</sup> Zn	$Q_{EC}(sa)$	9171 ± 26 [Da79]			<b>9171 ± 26</b>	
<sup>66</sup> As	<sup>66</sup> Ge	$Q_{EC}(sa)$	9550 ± 50 [Da80]			<b>9550 ± 50</b>	
<sup>70</sup> Br	<sup>70</sup> Se	$Q_{EC}(sa)$	9970 ± 170 [Da80]			<b>9970 ± 170</b>	
<sup>74</sup> Rb	<sup>74</sup> Kr	$ME(p)$	-51915.2 ± 4.0 [Ke04]			-51915.2 ± 4.0	
		$ME(d)$	-62332.0 ± 2.1 [Ke04]			-62332.0 ± 2.1	
		$Q_{EC}(sa)$				<b>10416.8 ± 4.5</b>	

<sup>1</sup>Abbreviations used in this column are as follows: “gs,” transition between ground states; “sa,” superallowed transition; “p,” parent; “d,” daughter; “ME,” mass excess; “ $E_x(0^+)$ ,” excitation energy of the  $0^+$  (analog) state. Thus, for example, “ $Q_{EC}(sa)$ ” signifies the  $Q_{EC}$  value for the superallowed transition, “ $ME(d)$ ,” the mass excess of the daughter nucleus; and “ $ME(d0^+)$ ,” the mass excess of the daughter’s  $0^+$  state.

<sup>2</sup>Result based on references [Ba88] and [Ba89].

<sup>3</sup>Average result includes the results of  $Q_{EC}$  pairs; see Table II.

<sup>4</sup>Result based on references [Bi03] and [Se05].

<sup>5</sup>Result obtained from data elsewhere in this table.

<sup>6</sup>Result based on references [Is80], [Al82], [Hu82], [Be85], [Pr90], [Ki91], and [Wa92].

<sup>7</sup>Result based on references [Wa83], [Ra83], and [Li94].

<sup>8</sup>Result based on references [Zi87] and [Ki89].

quantity, a Gaussian distribution is assumed, the weighted average being calculated according to the equation

$$\bar{x} \pm \delta\bar{x} = \frac{\sum_i w_i x_i}{\sum_i w_i} \pm \left( \frac{\sum_i w_i}{\sum_i w_i} \right)^{-1/2}, \quad (5)$$

where

$$w_i = 1/(\delta x_i)^2$$

and the sums extend over all  $N$  measurements. For each average, the  $\chi^2$  is also calculated and a scale factor,  $S$ , determined from

$$S = [\chi^2/(N - 1)]^{1/2}. \quad (6)$$

This factor is then used to establish the quoted uncertainty. If  $S \leq 1$ , the value of  $\delta\bar{x}$  from Eq. (5) is left unchanged. If  $S > 1$  and the input  $\delta x_i$  are all about the same size, then we increase  $\delta\bar{x}$  by the factor  $S$ , which is equivalent to assuming that all the experimental errors were underestimated by the same factor. Finally, if  $S > 1$  but the  $\delta x_i$  are of widely varying magnitudes,  $S$  is recalculated with only those results for which  $\delta x_i \leq 3N^{1/2}\delta\bar{x}$  being retained; the recalculated scale factor is then applied in the usual way. In all three cases,

TABLE II.  $Q_{EC}$ -value differences for superallowed  $\beta$ -decay branches. These data are also used as input to determine some of the average  $Q_{EC}$  values listed in Table I. (See Table VIII for the correlation between the alphabetical reference code used in this table and the actual reference numbers.)

Parent nucleus 1	Parent nucleus 2	$Q_{EC2} - Q_{EC1}$ (keV)	
		Measurement	Average <sup>a</sup>
<sup>14</sup> O	<sup>26</sup> Al <sup>m</sup>	1401.68 ± 0.13 [Ko87]	1401.37 ± 0.29
<sup>26</sup> Al <sup>m</sup>	<sup>42</sup> Sc	2193.5 ± 0.2 [Ko87]	2193.09 ± 0.42
<sup>42</sup> Sc	<sup>50</sup> Mn	1207.6 ± 2.3 [Ha74d]	1206.79 ± 0.44
<sup>42</sup> Sc	<sup>54</sup> Co	1817.2 ± 0.2 [Ko87]	1816.97 ± 0.48
<sup>50</sup> Mn	<sup>54</sup> Co	610.09 ± 0.17 <sup>[Ko87]</sup> <sub>[Ko97b]</sub>	610.18 ± 0.37

<sup>a</sup>Average values include the results of direct  $Q_{EC}$ -value measurements; see Table I.

no change is made to the original average  $\bar{x}$  calculated with Eq. (5).

The data for  $Q_{EC}$  include measurements of both individual  $Q_{EC}$  values and the differences between pairs of  $Q_{EC}$  values. This required a two-step analysis procedure. We first treated the individual  $Q_{EC}$ -value measurements for each particular

TABLE III. Half-lives,  $t_{1/2}$ , of superallowed  $\beta$  emitters. (See Table VIII for the correlation between the alphabetical reference code used in this table and the actual reference numbers.)

Parent nucleus	Measured half-lives, $t_{1/2}$ (ms)				Average value	
	1	2	3	4	$t_{1/2}$ (ms)	Scale
$\langle T_z = -1 \rangle$						
$^{10}\text{C}$	19280 $\pm$ 20 [Az74]	19295 $\pm$ 15 [Ba90]			19290 $\pm$ 12	1.0
$^{14}\text{O}$	70480 $\pm$ 150 [Al72]	70588 $\pm$ 28 [Cl73]	70430 $\pm$ 180 [Az74]	70684 $\pm$ 77 [Be78]		
	70613 $\pm$ 25 [Wi78]	70560 $\pm$ 49 [Ga01]	70641 $\pm$ 20 [Ba04]		70616 $\pm$ 14	1.1
$^{18}\text{Ne}$	1690 $\pm$ 40 [As70]	1670 $\pm$ 20 [Al70]	1669 $\pm$ 4 [Al75]	1687 $\pm$ 9 [Ha75]	1672.1 $\pm$ 4.6	1.3
$^{22}\text{Mg}$	3857 $\pm$ 9 [Ha75]	3875.5 $\pm$ 1.2 [Ha03]			3875.2 $\pm$ 2.4	2.0
$^{26}\text{Si}$	2210 $\pm$ 21 [Ha75]	2240 $\pm$ 10 [Wi80]			2234 $\pm$ 12	1.3
$^{30}\text{S}$	1180 $\pm$ 40 [Ba67]	1220 $\pm$ 30 [Mo71]	1178.3 $\pm$ 4.8 [Wi80]		1179.4 $\pm$ 4.7	1.0
$^{34}\text{Ar}$	844.5 $\pm$ 3.4 [Ha74a]	847.0 $\pm$ 3.7 [Ia03]			845.6 $\pm$ 2.5	1.0
$^{38}\text{Ca}$	470 $\pm$ 20 [Ka68]	439 $\pm$ 12 [Ga69]	450 $\pm$ 70 [Zi72]	430 $\pm$ 12 [Wi80]	440.0 $\pm$ 7.8	1.2
$^{42}\text{Ti}$	200 $\pm$ 20 [Ni69]	202 $\pm$ 5 [Ga69]	173 $\pm$ 14 [Al69]		198.8 $\pm$ 6.3	1.4
$\langle T_z = 0 \rangle$						
$^{26}\text{Al}^m$	6346 $\pm$ 5 [Fr69a]	6346 $\pm$ 5 [Az75]	6339.5 $\pm$ 4.5 [Al77]	6346.2 $\pm$ 2.6 [Ko83]		
	6345 $\pm$ 14 [Sc05]				6345.0 $\pm$ 1.9	1.0
$^{34}\text{Cl}$	1526 $\pm$ 2 [Ry73a]	1525.2 $\pm$ 1.1 [Wi76]	1527.7 $\pm$ 2.2 [Ko83]	1527.1 $\pm$ 0.5 [Ia03]	1526.77 $\pm$ 0.44	1.0
$^{38}\text{K}^m$	925.6 $\pm$ 0.7 [Sq75]	922.3 $\pm$ 1.1 [Wi76]	921.71 $\pm$ 0.65 [Wi78]	924.15 $\pm$ 0.31 [Ko83]		
	924.4 $\pm$ 0.6 [Ba00]	924.46 $\pm$ 0.14 [Ba05]			924.33 $\pm$ 0.27	2.3
$^{42}\text{Sc}$	680.98 $\pm$ 0.62 [Wi76]	680.67 $\pm$ 0.28 [Ko97a]			680.72 $\pm$ 0.26	1.0
$^{46}\text{V}$	422.47 $\pm$ 0.39 [Al77]	422.28 $\pm$ 0.23 [Ba77a]	422.57 $\pm$ 0.13 [Ko97a]		422.50 $\pm$ 0.11	1.0
$^{50}\text{Mn}$	284.0 $\pm$ 0.4 [Ha74b]	282.8 $\pm$ 0.3 [Fr75]	282.72 $\pm$ 0.26 [Wi76]	283.29 $\pm$ 0.08 [Ko97a]	283.24 $\pm$ 0.13	1.8
$^{54}\text{Co}$	193.4 $\pm$ 0.4 [Ha74b]	193.0 $\pm$ 0.3 [Ho74]	193.28 $\pm$ 0.18 [Al77]	193.28 $\pm$ 0.07 [Ko97a]	193.271 $\pm$ 0.063	1.0
$^{62}\text{Ga}$	115.95 $\pm$ 0.30 [Al78]	116.34 $\pm$ 0.35 [Da79]	115.84 $\pm$ 0.25 [Hy03]	116.19 $\pm$ 0.04 [Bl04a]		
	116.09 $\pm$ 0.17 [Ca05]				116.175 $\pm$ 0.038	1.0
$^{66}\text{As}$	95.78 $\pm$ 0.39 [Al78]	95.77 $\pm$ 0.28 [Bu88]			95.77 $\pm$ 0.23	1.0
$^{70}\text{Br}$	80.2 $\pm$ 0.8 [Al78]	78.54 $\pm$ 0.59 [Bu88]			79.12 $\pm$ 0.79	1.7
$^{74}\text{Rb}$	64.90 $\pm$ 0.09 [Oi01]	64.761 $\pm$ 0.031 [Ba01]			64.776 $\pm$ 0.043	1.5

transition in the manner already described, obtaining an average result with uncertainty in each case,  $\bar{x}_j \pm \delta\bar{x}_j$ , where the subscript  $j$  now designates a particular transition. For transitions unconnected by difference measurements, these uncertainties were scaled if necessary and then the values were quoted as final results. For those transitions involved in one or more difference measurements we combined their average  $Q_{EC}$  values,  $\bar{x}_j \pm \delta\bar{x}_j$ , with the difference measurements,  $d_k \pm \delta d_k$ , in a single fitting procedure. If  $M_1$  is the number of transitions that are connected by difference measurements and  $M_2$  is the number of those difference measurements, then we have a total of  $M_1 + M_2$  input data values from which we need to extract a final set of  $M_1$  average  $Q_{EC}$  values,  $\bar{x}_j \pm \delta\bar{x}_j$ . We accomplish this by minimizing  $\chi^2$ , where

$$\chi^2 = \sum_{j=1}^{M_1} \left( \frac{\bar{x}_j - \bar{x}_j}{\delta\bar{x}_j} \right)^2 + \sum_{k=1}^{M_2} \left( \frac{d_k - \bar{d}_k}{\delta d_k} \right)^2 \quad (7)$$

and

$$\bar{d}_k = \bar{x}_{j_1} - \bar{x}_{j_2},$$

with  $j_1$  and  $j_2$  designating the two transitions whose  $Q_{EC}$ -value difference is determined in a particular  $d_k$  measurement. For each of these individual  $Q_{EC}$  values, we obtained its scale

factor from Eq. (6), where the  $\chi^2$  used in that equation is now given by

$$\chi^2 = \sum_i \left( \frac{x_i - \bar{x}_j}{\delta x_i} \right)^2 + \sum_l \left( \frac{d_l - \bar{d}_l}{\delta d_l} \right)^2, \quad (8)$$

where  $j$  is the particular transition of interest. The sum in  $i$  extends over all individual  $Q_{EC}$ -value measurements of transition  $j$ , and the sum in  $l$  extends over all doublet measurements that include transition  $j$  as one component. The resultant value of  $S$  was applied to the uncertainty,  $\delta\bar{x}_j$ , with the same conventions as were described previously.

## B. Data tables

The  $Q_{EC}$ -value data appear in Tables I and II. For the best-known nine superallowed decays—those of  $^{10}\text{C}$ ,  $^{14}\text{O}$ ,  $^{26}\text{Al}^m$ ,  $^{34}\text{Cl}$ ,  $^{38}\text{K}^m$ ,  $^{42}\text{Sc}$ ,  $^{46}\text{V}$ ,  $^{50}\text{Mn}$ , and  $^{54}\text{Co}$ —the daughter nuclei are stable, and the most precise determinations of their  $Q_{EC}$  values have come from direct measurements of that property via, for example, ( $p,n$ ) or ( $^3\text{He},t$ ) reactions. Such measurements are identified in column 3 of Table I by “ $Q_{EC}(sa)$ ” and each individual result is itemized with its

TABLE IV. Branching ratios,  $R$ , for superallowed  $\beta$  transitions. (See Table VIII for the correlation between the alphabetical reference code used in this table and the actual reference numbers.)

Parent/daughter nuclei	Daughter state $E_x$ (MeV)	Measured branching ratio, $R$ (%)		Average value		
		1	2	$R$ (%)	Scale	
$\langle T_z = -1 \rangle$						
$^{10}\text{C}$	$^{10}\text{B}$	2.16 <b>1.74</b>	$0_{-0}^{+0.0008}$ [Go72] $1.468 \pm 0.014$ [Ro72] $1.465 \pm 0.009$ [Kr91] $1.4665 \pm 0.0038$ [Fu99]	$1.473 \pm 0.007$ [Na91] $1.4625 \pm 0.0025$ [Sa95]	$0_{-0}^{+0.0008}$ <b><math>1.4646 \pm 0.0019</math></b>	<b>1.0</b>
$^{14}\text{O}$	$^{14}\text{N}$	gs 3.95 <b>2.31</b>	$0.60 \pm 0.10$ [Sh55] $0.61 \pm 0.01$ [Si66] $0.062 \pm 0.007$ [Ka69] $0.053 \pm 0.002$ [He81]	$0.65 \pm 0.05$ [Fr63] $0.058 \pm 0.004$ [Wi80]	$0.611 \pm 0.010$ $0.0545 \pm 0.0019$ <b><math>99.334 \pm 0.010</math></b>	1.0 1.1
$^{18}\text{Ne}$	$^{18}\text{F}$	<b>1.04</b>	$9 \pm 3$ [Fr63]	$7.70 \pm 0.21^a$ [Ha75]	<b><math>7.70 \pm 0.21</math></b>	<b>1.0</b>
$^{22}\text{Mg}$	$^{22}\text{Na}$	<b>0.66</b>	$54.0 \pm 1.1$ [Ha75]	$53.15 \pm 0.12$ [Ha03]	<b><math>53.16 \pm 0.12</math></b>	<b>1.0</b>
$^{26}\text{Si}$	$^{26}\text{Al}$	1.06 <b>0.23</b>	$21.8 \pm 0.8$ [Ha75]		$21.8 \pm 0.8$ <b><math>75.09 \pm 0.92^a</math></b>	
$^{30}\text{S}$	$^{30}\text{P}$	gs <b>0.68</b>	$20 \pm 1$ [Fr63]		$20 \pm 1$ <b><math>77.4 \pm 1.0^a</math></b>	
$^{34}\text{Ar}$	$^{34}\text{Cl}$	0.67 gs	$2.49 \pm 0.10$ [Ha74a]		$2.49 \pm 0.10$ <b><math>94.45 \pm 0.25^a</math></b>	
$^{42}\text{Ti}$	$^{42}\text{Sc}$	0.61 gs	$56 \pm 14$ [Al69]		$56 \pm 14$ <b><math>43 \pm 14^a</math></b>	
$\langle T_z = 0 \rangle$						
$^{26}\text{Al}^m$	$^{26}\text{Mg}$	gs	$>99.997$ [Ki91]		$100.000_{-0.003}^{+0}$	
$^{34}\text{Cl}$	$^{34}\text{S}$	gs	$>99.988$ [Dr75]		$100.000_{-0.012}^{+0}$	
$^{38}\text{K}^m$	$^{38}\text{Ar}$	3.38 gs	$<0.0019$ [Ha94] $>99.998$		$0_{-0}^{+0.002}$ <b><math>100.000_{-0.002}^{+0}</math></b>	
$^{42}\text{Sc}$	$^{42}\text{Ca}$	1.84 gs	$0.0063 \pm 0.0026$ [In77] $0.0103 \pm 0.0031$ [Sa80]	$0.0022 \pm 0.0017$ [De78] $0.0070 \pm 0.0012$ [Da85]	$0.0059 \pm 0.0014$ <b><math>99.9941 \pm 0.0014</math></b>	1.6
$^{46}\text{V}$	$^{46}\text{Ti}$	2.61 4.32 $\Sigma\text{GT}^b$ gs	$0.0039 \pm 0.0004$ [Ha94] $0.0113 \pm 0.0012$ [Ha94] $<0.004$		$0.0039 \pm 0.0004$ $0.0113 \pm 0.0012$ $0_{-0}^{+0.004}$ <b><math>99.9848_{-0.0042}^{+0.0013}</math></b>	
$^{50}\text{Mn}$	$^{50}\text{Cr}$	3.63 3.85 5.00 gs	$0.057 \pm 0.003$ [Ha94] $<0.0003$ [Ha94] $0.0007 \pm 0.0001$ [Ha94]		$0.057 \pm 0.003$ $0_{-0}^{+0.0003}$ $0.0007 \pm 0.0001$ <b><math>99.9423 \pm 0.0030</math></b>	
$^{54}\text{Co}$	$^{54}\text{Fe}$	2.56 $\Sigma\text{GT}^b$ gs	$0.0045 \pm 0.0006$ [Ha94] $<0.03$		$0.0045 \pm 0.0006$ $0_{-0}^{+0.03}$ <b><math>99.9955_{-0.0300}^{+0.0006}</math></b>	
$^{62}\text{Ga}$	$^{62}\text{Zn}$	$\Sigma\text{GT}^b$ gs	$0.15_{-0.05}^{+0.15}$ [Hy03],[Bl02]		$0.15_{-0.05}^{+0.15}$ <b><math>99.85_{-0.15}^{+0.05}</math></b>	
$^{74}\text{Rb}$	$^{74}\text{Kr}$	$\Sigma\text{GT}^b$ gs	$0.50 \pm 0.10$ [Pi03]		$0.50 \pm 0.10$ <b><math>99.50 \pm 0.10</math></b>	

<sup>a</sup>Result also incorporates data from Table V.

<sup>b</sup>Total Gamow-Teller transitions to levels not explicitly listed; values were derived with the help of calculations in [Ha02].

appropriate reference in the next three columns. The weighted average [see Eq. (5)] of all measurements for a particular decay appears in column 7, with the corresponding scale factor [see Eq. (6)] in column 8. A few of these cases, such as  $^{34}\text{Cl}$  and  $^{46}\text{V}$ , have no further complications. There are other cases, however, in which  $Q_{EC}$ -value differences have been measured in addition to the individual  $Q_{EC}$  values.

These measurements are presented in Table II. They have been dealt with in combination with the direct  $Q_{EC}$ -value measurements, as described in Sec. II A [see, in particular, Eq. (7)], with the final average  $Q_{EC}$  value appearing in column 7 of Table I and the average difference in column 4 of Table II. Both are flagged with footnotes to indicate the interconnection.

TABLE V. Relative intensities of  $\beta$ -delayed  $\gamma$ -rays in the superallowed  $\beta$ -decay daughters. These data are used to determine some of the branching ratios presented in Table IV. (See Table VIII for the correlation between the alphabetical reference code used in this table and the actual reference numbers.)

Parent/daughter nuclei	Daughter ratios <sup>a</sup>	Measured $\gamma$ -ray ratio		Average value	
		1	2	Ratio	Scale
<sup>18</sup> Ne	<sup>18</sup> F	$\gamma_{660}/\gamma_{1042}$	0.021 ± 0.003 [Ha75] 0.0172 ± 0.0005 [Ad83]	0.0169 ± 0.0004 [He82]	0.0171 ± 0.0003 1.0
<sup>26</sup> Si	<sup>26</sup> Al	$\gamma_{1622}/\gamma_{829}$	0.149 ± 0.016 [Mo71] 0.1245 ± 0.0023 [Wi80]	0.134 ± 0.005 [Ha75]	0.1265 ± 0.0036 1.7
		$\gamma_{1655}/\gamma_{829}$	0.00145 ± 0.00032 [Wi80]		0.0015 ± 0.0003
		$\gamma_{1843}/\gamma_{829}$	0.013 ± 0.003 [Mo71]	0.016 ± 0.003 [Ha75]	
		$\gamma_{2512}/\gamma_{829}$	0.01179 ± 0.00027 [Wi80]		0.0118 ± 0.0003 1.0
		$\gamma_{total}/\gamma_{829}$	0.00282 ± 0.00010 [Wi80]		0.0028 ± 0.0001 0.1426 ± 0.0036
<sup>30</sup> S	<sup>30</sup> P	$\gamma_{709}/\gamma_{677}$	0.006 ± 0.003 [Mo71]	0.0037 ± 0.0009 [Wi80]	0.0039 ± 0.0009 1.0
		$\gamma_{2341}/\gamma_{677}$	0.033 ± 0.002 [Mo71]	0.0290 ± 0.0006 [Wi80]	0.0293 ± 0.0011 1.9
		$\gamma_{3019}/\gamma_{677}$	0.00013 ± 0.00006 [Wi80]		0.0001 ± 0.0001
		$\gamma_{total}/\gamma_{677}$			0.0334 ± 0.0014
<sup>34</sup> Ar	<sup>34</sup> S	$\gamma_{461}/\gamma_{666}$	0.28 ± 0.16 [Mo71]	0.365 ± 0.036 [Ha74a]	0.361 ± 0.035 1.0
		$\gamma_{2580}/\gamma_{666}$	0.38 ± 0.09 [Mo71]	0.345 ± 0.010 [Ha74a]	0.345 ± 0.010 1.0
		$\gamma_{3129}/\gamma_{666}$	0.67 ± 0.08 [Mo71]	0.521 ± 0.012 [Ha74a]	0.524 ± 0.022 1.8
		$\gamma_{total}/\gamma_{666}$			1.231 ± 0.043
<sup>42</sup> Ti	<sup>42</sup> Sc	$\gamma_{2223}/\gamma_{611}$	0.012 ± 0.004 [Ga69]		0.012 ± 0.004
		$\gamma_{total}/\gamma_{611}$	2 × 0.012 [En90]		0.024 ± 0.008

<sup>a</sup> $\gamma$ -Ray intensities are denoted by  $\gamma_E$ , where  $E$  is the  $\gamma$ -Ray energy in kiloelectron volts.

There are two cases, <sup>26</sup>Al<sup>m</sup> and <sup>38</sup>K<sup>m</sup>, in which the superallowed decay originates from an isomeric state. For both, there are  $Q_{EC}$ -value measurements that correspond to the ground state as well as to the isomer. Obviously, these

two sets of measurements are simply related to one another by the excitation energy of the isomeric state in the parent. In Table I, the set of measurements for the ground-state  $Q_{EC}$  value and for the excitation energy of the isomeric state appear

TABLE VI. References for which the original decay-energy results have been updated to incorporate the most recent calibration standards. (See Table VIII for the correlation between the alphabetical reference code used in this table and the actual reference numbers.)

References (parent nucleus) <sup>a</sup>	Update procedure
Bo64( <sup>18</sup> Ne), Ba84( <sup>10</sup> C), Br94( <sup>26</sup> Al <sup>m</sup> ) Ba98( <sup>10</sup> C), Ha98( <sup>38</sup> K <sup>m</sup> ), To03( <sup>14</sup> O)	We have converted all original ( $p, n$ ) threshold measurements to $Q$ values using the most recent mass excesses [Au03].
Ry73a( <sup>34</sup> Cl), Ho74( <sup>54</sup> Co), Sq76( <sup>46</sup> V) Ba77c( <sup>34</sup> Cl), Wh77( <sup>14</sup> O)	These ( $p, n$ ) threshold measurements have been adjusted to reflect recent calibration $\alpha$ -energies [Ry91] before being converted to $Q$ values.
Pr67( <sup>18</sup> Ne)	Before conversion to a $Q$ value, this ( $p, n$ ) threshold was adjusted to reflect a new value for the <sup>7</sup> Li( $p, n$ ) threshold [Wh85], which was used as calibration.
Ja78( <sup>38</sup> K <sup>m</sup> )	This ( $p, n$ ) threshold was measured relative to those for <sup>10</sup> C and <sup>14</sup> O; we have adjusted it based on average $Q$ values obtained for those decays in this work.
Bu79( <sup>38</sup> K <sup>m</sup> )	Before conversion to a $Q$ value, this ( $p, n$ ) threshold was adjusted to reflect the modern value for the <sup>35</sup> Cl( $p, n$ ) threshold [Au03], which was used as calibration.
Bu61( <sup>14</sup> O), Ba62( <sup>14</sup> O)	These <sup>12</sup> C( <sup>3</sup> He, $n$ ) threshold measurements have been adjusted for updated calibration reactions based on current mass excesses [Au03].
Ha74d( <sup>34</sup> Cl, <sup>50</sup> Mn, <sup>54</sup> Co)	These ( <sup>3</sup> He, $t$ ) reaction $Q$ values were calibrated by the <sup>27</sup> Al( <sup>3</sup> He, $t$ ) reaction to excited states in <sup>27</sup> Si; they have been revised according to modern mass excesses [Au03] and excited-state energies [En98].
Ki89( <sup>42</sup> Sc)	This <sup>41</sup> Ca( $p, \gamma$ ) reaction $Q$ value was measured relative to that for <sup>40</sup> Ca( $p, \gamma$ ); we have slightly revised the result based on modern mass excesses [Au03].
Ha74c( <sup>22</sup> Mg, <sup>26</sup> Si, <sup>30</sup> S, <sup>34</sup> Ar), Se74( <sup>38</sup> Ca)	These ( $p, t$ ) reaction $Q$ values have been adjusted to reflect the current $Q$ value for the <sup>16</sup> O( $p, t$ ) reaction [Au03], against which they were calibrated.

<sup>a</sup>These references all appear in Table I under the appropriate parent nucleus.

TABLE VII. References from which some or all results have been rejected. (See Table VIII for the correlation between the alphabetical reference code used in this table and the actual reference numbers.)

References (parent nucleus)	Reason for rejection
(Decay energies)	
Pa72( <sup>30</sup> S, <sup>38</sup> Ca)	No calibration is given for the measured ( <i>p,t</i> ) reaction <i>Q</i> values; update is clearly required but none is possible.
No74( <sup>22</sup> Mg)	Calibration reaction <i>Q</i> values have changed but calibration process is too complex to update.
Ro74( <sup>10</sup> C)	P. H. Barker (coauthor) later considered that inadequate attention had been paid to target surface purity [Ba84].
Ba77b( <sup>10</sup> C)	P. H. Barker (coauthor) later stated [Ba84] that the ( <i>p,t</i> ) reaction <i>Q</i> value could not be updated to incorporate modern calibration standards.
Wh81 and Ba98( <sup>14</sup> O)	The result in [Wh81] was updated in [Ba98] but then eventually withdrawn by P. H. Barker (coauthor) in [To03].
(Half-lives)	
Ja60( <sup>26</sup> Al <sup>m</sup> ), He61( <sup>14</sup> O), Ba62( <sup>14</sup> O), Ea62( <sup>10</sup> C), Ba63( <sup>10</sup> C), Fr63( <sup>14</sup> O, <sup>26</sup> Si), Fr65b( <sup>42</sup> Sc, <sup>46</sup> V, <sup>50</sup> Mn), Si72( <sup>14</sup> O)	Quoted uncertainties are too small, and results likely biased, in light of statistical difficulties more recently understood (see [Fr69a]). In particular, the “maximum-likelihood” analysis was not used.
Ha72a( <sup>26</sup> Al <sup>m</sup> , <sup>34</sup> Cl, <sup>38</sup> K <sup>m</sup> , <sup>42</sup> Sc)	All four quoted half-lives are systematically higher than more recent and accurate measurements.
Ro74( <sup>10</sup> C)	P. H. Barker (coauthor) later considered that pile-up had been inadequately accounted for [Ba90].
Ch84( <sup>38</sup> K <sup>m</sup> )	“Maximum-likelihood” analysis was not used.
(Branching ratios)	
Fr63( <sup>26</sup> S)	Numerous impurities present; result is obviously wrong.

TABLE VIII. Reference key relating alphabetical reference codes used in Tables I–III to the actual reference numbers.

Table code	Reference number	Table code	Reference number	Table code	Reference number	Table code	Reference number	Table code	Reference number	Table code	Reference number
Ad83	[8]	Ba98	[33]	Dr75	[58]	He61	[83]	Mc67	[108]	Se74	[133]
Aj88	[9]	Ba00	[34]	Ea62	[59]	He81	[84]	Mi67	[109]	Se04	[134]
Aj91	[10]	Ba01	[35]	En90	[60]	He82	[85]	Mo71	[110]	Sh55	[135]
Al69	[11]	Ba04	[36]	En98	[61]	He02	[86]	Mu04	[111]	Si66	[136]
Al70	[12]	Ba05	[37]	Fr63	[62]	Ho64	[87]	Na91	[112]	Si72	[137]
Al72	[13]	Be68	[38]	Fr65b	[63]	Ho74	[88]	Ni69	[113]	Sq75	[138]
Al75	[14]	Be78	[39]	Fr69a	[64]	Hu82	[89]	No74	[114]	Sq76	[139]
Al77	[15]	Be85	[40]	Fr75	[65]	Hy03	[90]	Oi01	[115]	Ti95	[140]
Al78	[16]	Bi03	[41]	Fu99	[66]	Ia03	[91]	Pa72	[116]	To03	[141]
Al82	[17]	Bi02	[42]	Ga69	[67]	In77	[92]	Pi03	[117]	Vo77	[142]
An70	[18]	Bi04a	[43]	Ga01	[68]	Is80	[93]	Pr67	[118]	Wa83	[143]
As70	[19]	Bi04b	[44]	Go72	[69]	Ja60	[94]	Pr90	[119]	Wa92	[144]
Au03	[20]	Bo64	[45]	Gi72	[70]	Ja78	[95]	Ra83	[120]	We68	[145]
Az74	[21]	Br94	[46]	Ha67	[71]	Ka68	[96]	Re85	[121]	Wh77	[146]
Az75	[22]	Bu61	[47]	Ha68	[72]	Ka69	[97]	Ro70	[122]	Wh81	[147]
Ba62	[23]	Bu79	[48]	Ha72a	[73]	Ke04	[98]	Ro72	[123]	Wh85	[148]
Ba63	[24]	Bu88	[49]	Ha74a	[74]	Ki89	[99]	Ro74	[124]	Wi76	[149]
Ba67	[25]	Ca05	[50]	Ha74b	[75]	Ki91	[100]	Ro75	[125]	Wi78	[150]
Ba77a	[26]	Ch84	[51]	Ha74c	[76]	Ko83	[101]	Ry73a	[126]	Wi80	[151]
Ba77b	[27]	Cl73	[52]	Ha74d	[77]	Ko87	[102]	Ry91	[127]	Zi72	[152]
Ba77c	[28]	Da79	[53]	Ha75	[78]	Ko97a	[103]	Sa80	[128]	Zi87	[153]
Ba84	[29]	Da80	[54]	Ha94	[79]	Ko97b	[104]	Sa95	[129]		
Ba88	[30]	Da85	[55]	Ha98	[80]	Kr91	[105]	Sa04	[130]		
Ba89	[31]	De69	[56]	Ha02	[81]	Li94	[106]	Sc05	[131]		
Ba90	[32]	De78	[57]	Ha03	[82]	Ma94	[107]	Se73	[132]		



in separate rows, each with its identifying property given in column 3 and its weighted average appearing in column 7. In the row below, the average value given in column 7 for the superallowed transition is the weighted average not only of the direct superallowed  $Q_{EC}$ -value measurements in that row but also of the result derived from the two preceding rows. Note that in all cases the  $Q_{EC}$  value for the superallowed transition appears in bold.

For those 11 superallowed decays that lead to radioactive daughter nuclei, there are very few direct measurements of the  $Q_{EC}$  value for the superallowed transition. In general, that  $Q_{EC}$  value must be deduced from the measured mass excesses of the parent and daughter nuclei, together with the excitation energy of the analog  $0^+$  state in the daughter. Each of these properties is identified in column 3 of Table I, with the individual measurements of that property, their weighted average and a scale factor appearing in columns to the right. The average  $Q_{EC}$  value listed for the corresponding superallowed transition is obtained from these separate averages. If a direct measurement of the superallowed  $Q_{EC}$  value exists, then it is also included in the final average.

Especially in these latter 11 cases, it might be imagined that it would have been sufficient for us to use the 2003 mass tables [20] to derive the  $Q_{EC}$  values of interest. There are, however, significant differences in our approach. We have included all pertinent measurements for each property as described in Sec. II A; typically, only a subset of the available data is included as input to the mass tables. Furthermore, we have examined each reference in detail and either accepted the result, updated it to modern calibration standards or rejected it for cause. The updating procedures are outlined, reference by reference, in Table VI and the rejected results are similarly documented in Table VII. With a comparatively small data set, we could afford to pay the kind of individual attention that is impossible when one is considering all nuclear masses.

The half-life data appear in Table III in similar format to Table I. For obvious reasons, half-life measurements do not lend themselves to being updated. Consequently, a number of mostly pre-1970 measurements have been rejected because they were not analyzed with the “maximum-likelihood” method. The importance of using this technique for precision measurements was not recognized until that time [64] and, without access to the primary data, there is no way a new analysis can be applied retroactively. All rejected half-life measurements are also documented in Table VII.

Finally, the branching-ratio measurements are presented in Table IV. The decays of the  $T_z = 0$  parents are the most straightforward because, in all these cases, the superallowed branch accounts for >99.5% of the total decay strength. Thus, even imprecise measurements of the weak nonsuperallowed branches can be subtracted from 100% to yield the superallowed branching ratio with good precision. For the higher- $Z$  parents of this type, particularly  $^{62}\text{Ga}$  and heavier, it has been shown theoretically [81] and experimentally [90,117] that numerous very weak Gamow-Teller transitions occur, which, in total, can carry significant decay strength. Where such unobserved transitions are expected to exist, we have used a combination of experiment and theory to account for

the unobserved strength, with uncertainties being adjusted accordingly.

The branching ratios for decays from  $T_z = -1$  parents are much more challenging to determine, because the superallowed branch is usually one of several strong branches—with the notable exception of  $^{14}\text{O}$ —and, in two of the measured cases, it actually has a branching ratio of less than 10%. The decays of  $^{18}\text{F}$ ,  $^{26}\text{Si}$ ,  $^{30}\text{S}$ ,  $^{34}\text{Ar}$ , and  $^{42}\text{Ti}$  required special treatment in our presentation. In each of these five cases, the absolute branching ratio for a single  $\beta$ -transition has been measured. The branching ratios for other  $\beta$  transitions must then be determined from the relative intensities of  $\beta$ -delayed  $\gamma$  rays in the daughter. The relevant  $\gamma$ -ray intensity measurements appear in Table V, with their averages then being used to determine the superallowed branching-ratio averages shown in bold type in Table IV. These cases are also flagged with a table footnote.

### III. THE $\mathcal{F}t$ VALUES

Having surveyed the primary experimental data, we now turn to producing a set of  $ft$  values for the 20 superallowed transitions being considered. The statistical rate function,  $f$ , for each transition depends primarily on the charge of the daughter nucleus,  $Z$ , and on the  $Q_{EC}$  value to the fifth power. Consequently the uncertainty in the value of  $f$  because of the experimental uncertainty in  $Q_{EC}$  is given by  $(\Delta f/f) \approx 5(\Delta Q_{EC}/Q_{EC})$ . Our goal in computing  $f$  therefore is to ensure that the computation itself yields percentage errors much less than those due to the uncertainty in the  $Q_{EC}$  value, which can be <0.02% in the best cases. To this end we have written a new code, the details of which are given in the Appendix. Our final  $f$  values and their uncertainties are recorded in column two of Table IX.

The partial half-life,  $t$ , for each transition is obtained from its total half-life,  $t_{1/2}$ , and branching ratio,  $R$ , according to the following formula:

$$t = \frac{t_{1/2}}{R} (1 + P_{EC}), \quad (9)$$

where  $P_{EC}$  is the calculated electron-capture fraction. The evaluation of  $P_{EC}$  is discussed by Bambynek *et al.* [155] and is based on the equation

$$P_{EC} = \frac{1}{2} \pi \left[ \sum_x \beta_x^2 (W_{EC} - |W_x|)^2 B_x \right] / f. \quad (10)$$

The sum extends over all atomic subshells from which an electron can be captured. The factor  $\beta_x$  is the Coulomb amplitude of the appropriate bound-state electron radial wave function;  $W_{EC}$  is the  $Q_{EC}$  value expressed in electron rest-mass units;  $W_x$  is the  $x$ -subshell binding energy also in electron rest-mass units; and  $B_x$  takes account of the effects of electron exchange and overlap. We have computed  $P_{EC}$  for the cases of interest here using our  $Q_{EC}$  values from Table I and the values of  $\beta_x^2 B_x$  and  $W_x$  from, respectively, Tables 1 and 2 of Appendix F in Ref. [156]. The  $P_{EC}$  results are shown (as percentages) in column 3 of Table IX. Based on experimental tests of such

TABLE IX. Derived results for superallowed Fermi  $\beta$  decays.

Parent nucleus	$f$	$P_{EC}$ (%)	Partial half-life $t$ (ms)	$ft$ (s)	$\delta'_R$ (%)	$\delta_C - \delta_{NS}$ (%)	$\mathcal{F}t$ (s)
$\langle T_z = -1 \rangle$							
$^{10}\text{C}$	$2.3009 \pm 0.0012$	0.297	$1321000 \pm 1900$	$3039.5 \pm 4.7$	$1.652 \pm 0.004$	$0.540 \pm 0.039$	$3073.0 \pm 4.9$
$^{14}\text{O}$	$42.772 \pm 0.024$	0.088	$71151 \pm 16$	$3043.3 \pm 1.9$	$1.520 \pm 0.008$	$0.570 \pm 0.056$	$3071.9 \pm 2.6$
$^{18}\text{Ne}$	$134.48 \pm 0.15$	0.081	$21730 \pm 590$	$2922 \pm 80$	$1.484 \pm 0.012$	$0.910 \pm 0.047$	$2938 \pm 80$
$^{22}\text{Mg}$	$418.44 \pm 0.18$	0.069	$7295 \pm 17$	$3052.4 \pm 7.2$	$1.446 \pm 0.017$	$0.505 \pm 0.024$	$3080.9 \pm 7.4$
$^{26}\text{Si}$	$1023.3 \pm 3.7$	0.064	$2978 \pm 40$	$3047 \pm 42$	$1.420 \pm 0.023$	$0.600 \pm 0.024$	$3072 \pm 42$
$^{30}\text{S}$	$1967.1 \pm 7.8$	0.066	$1524 \pm 21$	$2998 \pm 44$	$1.405 \pm 0.029$	$1.125 \pm 0.039$	$3006 \pm 44$
$^{34}\text{Ar}$	$3414.2 \pm 1.5$	0.069	$896.0 \pm 3.5$	$3059 \pm 12$	$1.394 \pm 0.035$	$0.825 \pm 0.044$	$3076 \pm 12$
$^{38}\text{Ca}$	$5338 \pm 22$	0.075			$1.397 \pm 0.042$	$0.910 \pm 0.053$	
$^{42}\text{Ti}$	$7043 \pm 30$	0.088	$470 \pm 160$	$3300 \pm 1100$	$1.412 \pm 0.050$	$1.015 \pm 0.110$	$3300 \pm 1100$
$\langle T_z = 0 \rangle$							
$^{26}\text{Al}^m$	$478.20 \pm 0.11$	0.082	$6350.2 \pm 1.9$	$3036.7 \pm 1.2$	$1.458 \pm 0.020$	$0.261 \pm 0.024$	$3072.9 \pm 1.5$
$^{34}\text{Cl}$	$1996.39 \pm 0.41$	0.080	$1527.99^{+0.44}_{-0.47}$	$3050.5 \pm 1.1$	$1.425 \pm 0.032$	$0.720 \pm 0.039$	$3071.7 \pm 1.9$
$^{38}\text{K}^m$	$3298.10 \pm 0.33$	0.085	$925.11 \pm 0.27$	$3051.1 \pm 1.0$	$1.423 \pm 0.039$	$0.720 \pm 0.047$	$3072.2 \pm 2.1$
$^{42}\text{Sc}$	$4470.03 \pm 1.46$	0.099	$681.43 \pm 0.26$	$3046.0 \pm 1.5$	$1.437 \pm 0.047$	$0.460 \pm 0.047$	$3075.6 \pm 2.5$
$^{46}\text{V}$	$7200.0 \pm 5.0$	0.101	$422.99 \pm 0.11$	$3045.5 \pm 2.2$	$1.429 \pm 0.054$	$0.465 \pm 0.033$	$3074.7 \pm 3.0$
$^{50}\text{Mn}$	$10731.2 \pm 1.8$	0.107	$283.71 \pm 0.13$	$3044.5 \pm 1.5$	$1.429 \pm 0.062$	$0.547 \pm 0.037$	$3071.1 \pm 2.7$
$^{54}\text{Co}$	$15749.3 \pm 3.0$	0.111	$193.495^{+0.063}_{-0.086}$	$3047.4^{+1.2}_{-1.5}$	$1.428 \pm 0.071$	$0.639 \pm 0.043$	$3071.2 \pm 2.9$
$^{62}\text{Ga}$	$26250 \pm 400$	0.137	$116.509^{+0.070}_{-0.179}$	$3058 \pm 47$	$1.445 \pm 0.087$	$1.42 \pm 0.16$	$3058 \pm 47$
$^{66}\text{As}$	$31610 \pm 890$	0.156			$1.457 \pm 0.095$	$1.45 \pm 0.16$	
$^{70}\text{Br}$	$38600 \pm 3600$	0.175			$1.47 \pm 0.11$	$1.41 \pm 0.20$	
$^{74}\text{Rb}$	$47280 \pm 110$	0.194	$65.227 \pm 0.078$	$3084.3 \pm 8.0$	$1.49 \pm 0.12$	$1.50 \pm 0.41$	$3083 \pm 15$
Average (best 12), $\overline{\mathcal{F}t}$							$3072.7 \pm 0.8$
$\chi^2/\nu$							0.42

$P_{EC}$  calculations [155], we expect these results to be accurate to a few parts in 100; thus they do not contribute perceptibly to the overall uncertainties. Partial half-lives derived from Eq. (9) and corresponding  $ft$  values appear in columns 4 and 5.

To obtain  $\mathcal{F}t$  values according to Eq. (4) we must now deal with the small correction terms. The term  $\delta'_R$  has been calculated from standard QED and is currently evaluated to order  $Z\alpha^2$  and estimated in order  $Z^2\alpha^3$  [157,158]; its values, listed in column six of Table IX, are around 1.4% and can be considered to be very reliable. The structure-dependent terms  $\delta_{NS}$  and  $\delta_C$  have also been calculated in the past but at various times over three decades and with a variety of different models. Their uncertainties are larger. This topic has been reviewed recently by Towner and Hardy [159], who presented new calculations of these corrections in which consistent model spaces and approximations have been used for both correction terms. The results of these new calculations are recorded in column 7 of Table IX. Finally, the resulting  $\mathcal{F}t$  values are listed in column 8.

#### A. CVC test

We are now ready to test the CVC assertion that the  $\mathcal{F}t$  values should be constant for all nuclear superallowed transitions of this type. The data in Table IX clearly satisfy the test; the weighted average of the 12 most precise results (with “statistical” uncertainty only) is

$$\overline{\mathcal{F}t} = 3072.7 \pm 0.8s, \quad (11)$$

with a corresponding chi-square-per-degree-of-freedom of  $\chi^2/\nu = 0.42$ . In Fig. 1 we plot the same 12 values, all of whose statistical accuracy is better than 0.5%. It is evident from both the figure and the table that the data form a consistent set, thus verifying the expectation of the CVC hypothesis at the level of  $3 \times 10^{-4}$ , which is the fractional uncertainty quoted in Eq. (11). This is a 30% improvement over the results from our last survey in 1990 [7] and is principally because of improvements in the experimental data themselves.

#### B. $\mathcal{F}t$ -value error budgets

In Figs. 2 and 3 we show the contributing factors to the individual  $\mathcal{F}t$ -value uncertainties. For the most precise data,  $^{14}\text{O}$  to  $^{54}\text{Co}$ , which appear in Fig. 2, the theoretical uncertainties are greater than, or comparable to, the experimental ones. The nuclear-structure-dependent correction,  $\delta_C - \delta_{NS}$ , contributes an almost constant uncertainty of four parts in  $10^4$  across these nuclei, whereas the nucleus-dependent radiative correction,  $\delta'_R$ , has an uncertainty that grows as  $Z^2$ . This is because the contribution to  $\delta'_R$  from order  $Z^2\alpha^3$  has been estimated only from its leading logarithm [157] and the magnitude of this estimate has been taken as the uncertainty in  $\delta'_R$ . In fact, for  $^{50}\text{Mn}$  and  $^{54}\text{Co}$  this becomes the leading uncertainty, indicating that a closer look at the order  $Z^2\alpha^3$  contribution would now be worthwhile. For the eight precise data, the experimental




FIG. 1.  $\mathcal{F}t$  values plotted as a function of the charge on the daughter nucleus,  $Z$ . The shaded horizontal band gives 1 standard deviation around the average  $\overline{\mathcal{F}t}$  value, Eq. (11).

branching ratios are  $>99\%$  and have very small associated uncertainties with the exception of  $^{54}\text{Co}$ , which has a  $3 \times 10^4$  fractional uncertainty attributed to its branching ratio. This is because  $^{54}\text{Co}$  is predicted to have several weak Gamow-Teller branches that have not yet been observed. We have used an estimate of the strength of the missing branches, taken from a shell-model calculation [81], to assign an uncertainty to the superallowed branching ratio. Missing weak branches become a larger issue for the heavier-mass nuclei with  $A \geq 62$ , where they contribute significantly to the branching-ratio uncertainty.

For the less precisely known decay of  $^{10}\text{C}$ , and for the 12 decays depicted in Fig. 3, the predominant uncertainties are all experimental in origin with the single exception of  $^{74}\text{Rb}$ , for

which the nuclear-structure calculation is quite difficult [159], resulting in a larger uncertainty on  $\delta_C - \delta_{NS}$ . Many of the experimental  $Q$  values, half-lives, and branching ratios have yet to be measured for the cases shown in Fig. 3, but recent advances in experimental techniques are likely to change this situation dramatically within the next few years.

**C. Accounting for systematic uncertainties**

So far, we have dealt with the internuclear behavior of  $\mathcal{F}t$  values, examining their constancy as a test of CVC. With that test passed at high precision, we are now in a position to use the average  $\mathcal{F}t$ -value obtained from these concordant nuclear data to go beyond nuclei, obtaining first




FIG. 2. Summary histogram of the fractional uncertainties attributable to each experimental and theoretical input factor that contributes to the final  $\mathcal{F}t$  values for the nine most precise superallowed decay data.




FIG. 3. Summary histogram of the fractional uncertainties attributable to each experimental and theoretical input factor that contributes to the final  $\mathcal{F}t$  values for the 12 other superallowed decay data. Where the error is shown as exceeding 60 parts in  $10^4$ , no useful experimental measurement has been made.

the vector coupling constant [see Eq. (2)] and then the  $V_{ud}$  matrix element. Before doing so, however, we must address one more possible source of uncertainty. Though the average  $\mathcal{F}t$  value given in Eq. (11) includes a full assessment of the uncertainties attributable to experiment and to the particular calculations used to obtain the correction terms, it does not incorporate any provision for a common systematic error that could arise from the type of calculation chosen to model the nuclear-structure effects. In this section we look more critically at the nuclear-structure-dependent corrections and in particular at the isospin-symmetry-breaking correction,<sup>1</sup>  $\delta_C$ .

There have been a number of previous calculations of  $\delta_C$  in addition to those of ours [159], including Hartree-Fock calculations of Ormand and Brown [160], RPA calculations of Sagawa, van Giai and Suzuki [161], R-matrix calculations of Barker [162], and Woods-Saxon calculations of Wilkinson [163], to name some of the more recent publications. Of these, we retain only the Ormand-Brown (OB) calculations because they, like ours (TH), are constrained to reproduce other isospin properties of the nuclei involved: They reproduce the measured coefficients of the relevant isobaric multiplet mass

<sup>1</sup>The reason we do not consider further the nuclear-structure-dependent radiative correction,  $\delta_{NS}$ , is that it is very small for the series of transitions that have  $T_z = 0$  parent states [159]. Of the nine precisely known transitions we are concentrating on, seven are of this type.

equation, the known proton and neutron separation energies, and the measured  $ft$  values of weak nonanalog  $0^+ \rightarrow 0^+$  transitions [79], where they are known. The other calculations are not constrained by experiment in any way and thus offer no independent means to assess their efficacy.

Unfortunately, calculations of  $\delta_C$  by OB are not available for all the cases listed in Table IX, so we must concentrate on the nine most precise data:  $^{10}\text{C}$ ,  $^{14}\text{O}$ ,  $^{26}\text{Al}^m$ ,  $^{34}\text{Cl}$ ,  $^{38}\text{K}^m$ ,  $^{42}\text{Sc}$ ,  $^{46}\text{V}$ ,  $^{50}\text{Mn}$ , and  $^{54}\text{Co}$ . When the TH values of  $\delta_C$  are used, the average  $\mathcal{F}t$  value for these nine cases alone is  $\overline{\mathcal{F}t} = 3072.6 \pm 0.8 \text{ s}$  with  $\chi^2/\nu = 0.35$ . When OB values are used for  $\delta_C$  instead, the weighted average is  $\overline{\mathcal{F}t} = 3074.5 \pm 0.8 \text{ s}$  with  $\chi^2/\nu = 0.92$ . Although the chi-square with the OB values is worse by nearly a factor of 3, we do not argue that this is sufficient reason to reject the OB calculation. Rather, we observe that the OB values of  $\delta_C$  are systematically smaller and hence the  $\mathcal{F}t$  values are systematically larger than ours. Evidently there is a systematic difference between our Woods-Saxon and OB's Hartree-Fock calculations of  $\delta_C$  and that difference should be accounted for in the final result. Thus, we adopt the average of these two results for our recommended  $\overline{\mathcal{F}t}$  and assign a systematic uncertainty equal to half the spread between them:

$$\begin{aligned} \overline{\mathcal{F}t} &= 3073.5 \pm 0.8_{\text{stat}} \pm 0.9_{\text{sys}}\text{s} \\ &= 3073.5 \pm 1.2\text{s}, \end{aligned} \tag{12}$$

where the two errors have been combined in quadrature.

#### IV. IMPACT ON WEAK-INTERACTION PHYSICS

##### A. The value of $V_{ud}$

With a mutually consistent set of  $\mathcal{F}t$  values, we can now use their average value in Eq. (12) to determine the vector coupling constant,  $G_V$ , from Eq. (2). The value of  $G_V$  itself is of little interest, but it can be related to the weak-interaction constant for the purely leptonic muon decay,  $G_F$ , to yield the much more interesting up-down matrix element of the CKM quark-mixing matrix<sup>2</sup>:  $G_V = G_F V_{ud}$ . The relation we use is the following one:

$$V_{ud}^2 = \frac{K}{2G_F^2(1 + \Delta_R^V)\overline{\mathcal{F}t}}, \tag{13}$$

where  $\Delta_R^V$  is the nucleus-independent radiative correction. The currently accepted value for this correction is derived from the expression [164,165]:

$$\Delta_R^V = \frac{\alpha}{2\pi} [4 \ln(m_Z/m_p) + \ln(m_p/m_A) + 2C_{\text{Born}}] + \dots, \tag{14}$$

where the ellipses represent further small terms of the order of 0.1%. Here  $m_Z$  is the Z-boson mass,  $m_p$  the proton mass,  $m_A$  the mass parameter in the dipole form of the axial-vector

<sup>2</sup>More completely we could write  $G_V = G_F V_{ud} g_V(q^2 \rightarrow 0)$ , where  $g_V$  is the vector form factor given in Eq. (A18), or as  $G_V = G_F V_{ud} C_V$ , where  $C_V$  is the vector coupling constant in the Jackson, Treiman, and Wyld [166] Hamiltonian in Eq. (22), with  $g_V(q^2 \rightarrow 0) = C_V = 1$ .

form factor, and  $C_{\text{Bom}}$  the universal order- $\alpha$  axial-vector contribution. The various terms have the following values:

$$\Delta_R^V = 2.12 - 0.03 + 0.20 + 0.10\%, \quad (15)$$

with the first term, the leading logarithm, being essentially unambiguous in value. The final value recommended by Sirlin [164] is

$$\Delta_R^V = (2.40 \pm 0.08)\%. \quad (16)$$

The uncertainty is almost entirely because of the value selected for the axial-vector form factor mass, which Sirlin argues should lie in the range  $(m_{a_1}/2) \leq m_A \leq 2m_{a_1}$ , where  $m_{a_1}$  is the physical  $a_1$  meson mass.

Using the Particle Data Group (PDG) [154] value for the weak interaction coupling constant from muon decay of  $G_F/(\hbar c)^3 = (1.16639 \pm 0.00001) \times 10^{-5} \text{ GeV}^{-2}$ , we obtain from Eq. (13) the result

$$|V_{ud}|^2 = 0.9482 \pm 0.0008. \quad (17)$$

Note that the total uncertainty here—0.00083, if the next significant figure is included—is almost entirely because of the uncertainties contributed by the theoretical corrections. By far the largest contribution, 0.00074, arises from the uncertainty in  $\Delta_R^V$ ; 0.00031 comes from the nuclear-structure-dependent corrections  $\delta_C - \delta_{NS}$  (principally from the systematic difference between the OB and TH calculations discussed in Sec. II C) and 0.00012 is attributable to  $\delta'_R$ . Only 0.00016 can be considered to be experimental in origin.

The corresponding value of  $V_{ud}$  is

$$|V_{ud}| = 0.9738 \pm 0.0004, \quad (18)$$

a result that differs by two units in the last quoted digit from our previously recommended result [1]. This shift, well within the quoted 1 standard deviation, is because of the improvements in the experimental data and to our recomputing of the statistical rate function (see the Appendix), in which a number of different parameter choices were made for the charge-density distribution, the oscillator length parameter for nuclear radial functions, and for the screening correction. Coincidentally, the value of  $V_{ud}$  quoted in Eq. (18) is identical to the currently recommended PDG value [154], although our uncertainty is one digit smaller.

### B. Unitarity of the CKM matrix

The CKM matrix yields the transformation equations for a change of basis from quark weak-interaction eigenstates to quark-mass eigenstates. As such, the CKM matrix must be unitary so the bases remain orthonormal. With the CKM matrix elements determined from experimental data, one important test they should satisfy is that they yield a unitary matrix. Currently, the sum of the squares of the top-row elements, which should equal 1, constitutes the most demanding available test. With our experimental value for  $|V_{ud}|^2$  given in Eq. (17) and the PDG's recommended values [154] of  $|V_{us}| = 0.2200 \pm 0.0026$  and  $|V_{ub}| = 0.00367 \pm 0.00047$ , this unitarity test yields

$$|V_{ud}|^2 + |V_{us}|^2 + |V_{ub}|^2 = 0.9966 \pm 0.0014. \quad (19)$$

The test fails by 2.4 standard deviations. The majority of the total uncertainty originates from the uncertainty in  $|V_{us}|^2$ , *viz.* 0.0011, with  $|V_{ud}|^2$  contributing 0.0008. The latter, as we have already demonstrated, is *not* predominantly experimental in origin but is dominated by the uncertainty in the nucleus-independent radiative correction,  $\Delta_R^V$ .

A recent measurement of the  $K^+ \rightarrow \pi^0 e^+ \nu_e (K_{e3}^+)$  branching ratio from the Brookhaven E865 experiment [2] obtains  $V_{us} = 0.2272 \pm 0.0030$ . Although this result is included in the PDG average value, it is considerably higher than the older experimental results from  $K_{e3}^+$  and  $K_{\mu 3}^0$  decays, with which it is inconsistent. An even more recent measurement [3] supports the higher value but has not yet been included in the PDG average; it comes from the Fermilab E832 experiment, which determines the six largest  $K_L$  branching fractions, including  $K_{e3}^0$  and  $K_{\mu 3}^0$  and yields  $V_{us} = 0.2252 \pm 0.0022$ . At least one more pertinent experiment is still in progress and should help clarify the situation. If, for the moment, we adopt the weighted average of the E865 and E832 experiments,  $V_{us} = 0.2259 \pm 0.0018$ , rather than the PDG average, then the result in Eq. (19) is modified to

$$|V_{ud}|^2 + |V_{us}|^2 + |V_{ub}|^2 = 0.9993 \pm 0.0011 \quad (20)$$

and unitarity is fully satisfied.

Another, at present, less demanding test is to examine the first column of the CKM matrix. The PDG value for  $V_{cd}$  is  $0.224 \pm 0.012$ , but little is known about  $V_{td}$  other than it is expected to lie in the range  $0.0048 \leq V_{td} \leq 0.014$ . In this range it has negligible impact on the unitarity sum. With our value of  $|V_{ud}|^2$ , this unitarity sum is

$$|V_{ud}|^2 + |V_{cd}|^2 + |V_{td}|^2 = 0.9985 \pm 0.0054. \quad (21)$$

Here the error is given entirely by the uncertainty in the value of  $V_{cd}$  and unitarity is evidently satisfied at this level of accuracy.

### C. Fundamental scalar interaction

For the past 40 years, the weak interaction has been described by an equal mix of vector and axial-vector interactions that maximizes parity violation. The theory is known colloquially as the  $V$ - $A$  theory. Despite the ever-increasing precision possible in weak-interaction experiments, no defect has been found in the  $V$ - $A$  theory. Prior to the establishment of the  $V$ - $A$  theory, other forms of fundamental couplings, notably scalar and tensor interactions, were popular. Today there is still interest in searching for scalar and tensor interactions, not because we expect them to contribute importantly, but rather because we wish to establish limits to their possible contribution.

A general form of the weak-interaction Hamiltonian was written down by Jackson, Treiman, and Wyld [166]. In examining superallowed Fermi transitions, we are only interested in scalar and vector couplings, for which that Hamiltonian becomes the following:

$$H_{S+V} = (\bar{\psi}_p \psi_n) (C_S \bar{\phi}_e \phi_{\bar{\nu}_e} + C'_S \bar{\phi}_e \gamma_5 \phi_{\bar{\nu}_e}) + (\bar{\psi}_p \gamma_\mu \psi_n) (C_V \bar{\phi}_e \gamma_\mu \phi_{\bar{\nu}_e} + C'_V \bar{\phi}_e \gamma_\mu \gamma_5 \phi_{\bar{\nu}_e}). \quad (22)$$

If we assume that the Hamiltonian is invariant under time reversal, then all the coupling constants must be real. Those coupling constants carrying a prime represent parity nonconserving interactions. If we further assume that parity violation is maximal, then  $C'_i = C_i$ . In this limit, the scalar and vector terms can be written as

$$H_{S+V} = (\bar{\psi}_p C_S \psi_n) [\bar{\phi}_e (1 + \gamma_5) \phi_{\bar{\nu}_e}] + (\bar{\psi}_p C_V \gamma_\mu \psi_n) [\bar{\phi}_e \gamma_\mu (1 + \gamma_5) \phi_{\bar{\nu}_e}]. \quad (23)$$

A nonrelativistic reduction of the hadron matrix element for the scalar and the time part of the vector interaction shows that they both reduce simply to constants in leading order. However, under charge conjugation the matrix element  $(\bar{\psi}_p C_S \psi_n)$  changes sign relative to  $(\bar{\psi}_p C_V \gamma_4 \psi_n)$ . Thus we write  $\pm C_S$  in the ensuing formulae with the upper sign being for electron emission and the lower sign for positron emission. The lepton matrix elements are different in the two terms in Eq. (23) so the contribution to the shape-correction function from the scalar interaction will involve a different combination of electron and neutrino radial functions than that from the vector interaction. The final formula for  $C(Z, W)$  is

$$C(Z, W) = \sum_{k_e, k_\nu, K} \lambda_{k_e} \left\{ [M_K(k_e, k_\nu) + \bar{m}_K(k_e, k_\nu)]^2 + [m_K(k_e, k_\nu) + \bar{M}_K(k_e, k_\nu)]^2 - \frac{2\mu_{k_e} \gamma_{k_e}}{k_e W} [M_K(k_e, k_\nu) + \bar{m}_K(k_e, k_\nu)] \times [m_K(k_e, k_\nu) + \bar{M}_K(k_e, k_\nu)] \right\}, \quad (24)$$

where  $M_K(k_e, k_\nu)$  and  $m_K(k_e, k_\nu)$  are the reduced matrix elements given in Eq. (A13), which incorporate the radial functions,  $F(r)$  and  $f(r)$ , defined in Eq. (A14). The reduced matrix elements  $\bar{M}_K(k_e, k_\nu)$  and  $\bar{m}_K(k_e, k_\nu)$  are the same as  $M_K(k_e, k_\nu)$  and  $m_K(k_e, k_\nu)$  except that the radial functions,  $F(r)$  and  $f(r)$ , are replaced by  $\bar{F}(r)$  and  $\bar{f}(r)$ , where

$$\begin{aligned} \bar{F}(r) &= H(r) \{-G_{--} j_{k_\nu-1}(p_\nu r) + G_{-+} j_{k_\nu}(p_\nu r)\} \\ &\quad + D(r) \{G_{+-} j_{k_\nu-1}(p_\nu r) - G_{++} j_{k_\nu}(p_\nu r)\} \\ \bar{f}(r) &= h(r) \{-G_{--} j_{k_\nu-1}(p_\nu r) + G_{-+} j_{k_\nu}(p_\nu r)\} \\ &\quad + d(r) \{G_{+-} j_{k_\nu-1}(p_\nu r) - G_{++} j_{k_\nu}(p_\nu r)\}. \end{aligned} \quad (25)$$

The functions  $H, D, h,$  and  $d$  are linear combinations of the electron functions,  $f_\kappa(r)$  and  $g_\kappa(r)$ , as given in Eq. (A15); and the angular momentum factors  $G_{\pm\pm}$  are defined in Eq. (A16).

### 1. Order of magnitude estimates

For a pure Fermi transition, the multipolarity of the transition operators is  $K = 0$ . Keeping only the lowest lepton partial waves,  $k_e = 1$  and  $k_\nu = 1$ , we expand the lepton radial functions in a power series in  $r$ . The order of magnitude of the lepton wave functions at small  $r$  are

$$\begin{aligned} f_1(r) &= 1 - \dots \\ g_{-1}(r) &= 1 - \dots \\ f_{-1}(r) &= \text{small} \end{aligned}$$

$$\begin{aligned} g_1(r) &= \text{small} \\ j_0(p_\nu r) &= 1 - \dots \\ j_1(p_\nu r) &= \text{small}. \end{aligned} \quad (26)$$

We retain only  $f_1(r), g_{-1}(r)$  and  $j_0(p_\nu r)$ , setting their values to unity, and drop the other small terms. The angular momentum factors for  $K = L = s = 0$  are  $G_{++} = G_{--} = 1$  and  $G_{+-} = G_{-+} = 0$ . Then the amplitudes become

$$\begin{aligned} M_0(1, 1) &= C_V M_F + \dots \\ m_0(1, 1) &= \text{small}, \\ \bar{M}_0(1, 1) &= \mp C_S M_F + \dots \\ \bar{m}_0(1, 1) &= \text{small}. \end{aligned} \quad (27)$$

where  $M_F$  is the Fermi matrix element. The shape-correction function is then

$$\begin{aligned} C(Z, W) &= |M_F|^2 \left( C_V^2 + C_S^2 \pm \frac{2\mu_1 \gamma_1}{W} C_S C_V \right) \\ &\simeq |M_F|^2 C_V^2 (1 + b_F \gamma_1 / W + \dots), \end{aligned} \quad (28)$$

where it is assumed that  $C_S \ll C_V$ . The term in  $b_F \gamma_1 / W$  is called the Fierz interference term, with  $b_F = \pm 2\mu_1 C_S / C_V$ . This is the well-known expression given by Jackson, Treiman, and Wyld [166]. Here  $\mu_1$  is one of the  $\beta$ -decay Coulomb functions, Eq. (A9), and is of order unity and  $\gamma_1 = (1 - (\alpha Z)^2)^{1/2}$ .

### 2. Determining a limit on $C_S/C_V$

With the results of our data survey, we can now search for any evidence of a  $1/W$  term in the shape-correction function and hence set a limit on  $C_S$ . The test is based on the corrected  $\mathcal{F}t$  values being a constant for all superallowed transitions between isospin  $T = 1$  analog states. For optimum sensitivity, we do not use Eq. (28) for  $C(Z, W)$  in the evaluation of the statistical rate function,  $f$ , because of the extreme nature of some of the approximations made in deriving that equation. Instead we use the exact numerically computed expression. Because this calculated value of  $f$  depends on the value of  $C_S$  we simply treat  $C_S$  as an adjustable parameter and seek a value that minimizes  $\chi^2$  in a least-squares fit to the expression  $\mathcal{F}t = \text{constant}$ . The result is

$$C_S/C_V = -(0.00005 \pm 0.00130). \quad (29)$$

The sign of  $C_S/C_V$  is determined from the fit, because the calculated  $f$  depends on the interference between vector and scalar interactions. The interpretation of the sign is a little more delicate. We define  $C_S$  to be the strength of the scalar interaction in electron-emission  $\beta$  decay, and this is the value quoted in Eq. (29). Because all the superallowed data involve positron emitters there is a sign change mentioned earlier due to charge conjugation that operationally is included in the

computations. The corresponding Fierz interference constant,  $b_F$ , is just  $-2$  times this quantity<sup>3</sup>: i.e.  $b_F = 0.0001 \pm 0.0026$ . Had we not assumed that parity was violated maximally then the outcome would be

$$\frac{C_S C_V + C'_S C'_V}{|C_V|^2 + |C'_V|^2 + |C_S|^2 + |C'_S|^2} = -(0.00005 \pm 0.00130). \quad (30)$$

This result shows a reduction by a factor of 30 in the central value compared to our previously published result [1], with the standard deviation being essentially unchanged. This is by far the most stringent limit on  $C_S/C_V$  ever obtained from nuclear  $\beta$  decay.

#### D. Induced scalar interaction

If we consider only the vector part of the weak interaction, for composite spin-1/2 nucleons the most general form of that interaction is written [167] as

$$H_V = \bar{\psi}_p (g_V \gamma_\mu - f_M \sigma_{\mu\nu} q_\nu + i f_S q_\mu) \psi_n \bar{\phi}_e \gamma_\mu (1 + \gamma_5) \phi_{\bar{\nu}_e}, \quad (31)$$

with  $q_\mu$  being the four-momentum transfer,  $q_\mu = (p_p - p_n)_\mu$ . The values of the coupling constants  $g_V$  (vector),  $f_M$  (weak magnetic), and  $f_S$  (induced scalar) are prescribed if the CVC hypothesis—that the weak vector current is just an isospin rotation of the electromagnetic vector current—is correct. In particular, because CVC implies that the vector current is divergenceless, we anticipate that  $f_S = 0$ . An independent argument [168], that there be no second-class currents in the hadronic weak interaction, also requires  $f_S$  to vanish. Our goal in this section is to use the data from superallowed  $\beta$  decay to set limits on the possible value of the induced scalar coupling constant,  $f_S$ . This will provide a test of the CVC hypothesis and simultaneously set limits on the presence of second-class currents in the hadronic vector weak interaction.

#### 1. Relation between $f_S$ and $C_S$

Considering, then, just the induced scalar term in the vector part of the weak interaction,

$$H_V(S) = \bar{\psi}_p (i f_S q_\mu) \psi_n \bar{\phi}_e \gamma_\mu (1 + \gamma_5) \phi_{\bar{\nu}_e}, \quad (32)$$

we see that this term can be reorganized to match closely the Hamiltonian from the fundamental scalar interaction shown in Eq. (22). The momentum transfer,  $q_\mu = (p_p - p_n)_\mu = -(p_e + p_{\bar{\nu}_e})_\mu$ , can be moved into the lepton matrix element where, in combination with  $\gamma_\mu$ , it can be replaced with the free-particle Dirac equation:  $\gamma_\mu(p_e)_\mu \phi_e = i m_e \phi_e$ ,  $\gamma_\mu(p_{\bar{\nu}_e})_\mu \phi_{\bar{\nu}_e} =$

$i m_{\bar{\nu}_e} \phi_{\bar{\nu}_e}$ , with  $m_e$  and  $m_{\bar{\nu}_e}$  being the electron and neutrino masses, respectively. On setting the neutrino mass to zero, we find that  $H_V(S)$  becomes

$$H_V(S) = \bar{\psi}_p m_e f_S \psi_n \bar{\phi}_e (1 + \gamma_5) \phi_{\bar{\nu}_e}. \quad (33)$$

This expression is equivalent to the fundamental scalar interaction in Eq. (22) with  $C_S$  simply replaced by  $m_e f_S$ . Thus, its effect on the shape-correction function can be described by the same replacement in Eq. (28). An equivalent result was obtained by Holstein [169].

#### 2. Determining a limit on $f_S$

We have now established the mathematical equivalence of the effects that  $f_S$  and  $C_S$  have on the shape-correction function,  $C(Z, W)$ . As a result, we can use Eq. (29) to conclude that

$$m_e f_S / g_V = -(0.00005 \pm 0.00130). \quad (34)$$

The sign of  $f_S/g_V$  follows the same convention as that described after Eq. (29). This result is a vindication for the CVC hypothesis, which predicts  $g_V = 1$  and  $f_S = 0$ . Our result confirms this prediction at the level of 13 parts in  $10^4$ . As already mentioned, this result can also be interpreted as setting a limit on vector second-class currents in the semileptonic weak interaction, which therefore have not been observed here at the same level of precision.

#### E. Right-hand currents

Let us no longer consider parity violation to be maximal. The general form of the weak interaction Hamiltonian [166] for just the vector couplings of relevance for superallowed  $\beta$  decay is

$$H_V = (\bar{\psi}_p \gamma_\mu \psi_n) (C_V \bar{\phi}_e \gamma_\mu \phi_{\bar{\nu}_e} + C'_V \bar{\phi}_e \gamma_\mu \gamma_5 \phi_{\bar{\nu}_e}). \quad (35)$$

With  $C'_V \neq C_V$  we cannot associate the coupling constants with the hadron matrix elements as we did in Eq. (23). Instead, the lepton and neutrino radial functions remain combined with  $C_V$  or  $C'_V$ . The final formula for the shape-correction function then becomes

$$C(Z, W) = \sum_{k_e k_\nu K} \lambda_{k_e} \left\{ \frac{1}{2} (M_K^2(k_e, k_\nu) + m_K^2(k_e, k_\nu) + N_K^2(k_e, k_\nu) + n_K^2(k_e, k_\nu)) - \frac{2\mu_{k_e} \gamma_{k_e}}{k_e W} \frac{1}{2} (M_K(k_e, k_\nu) m_K(k_e, k_\nu) + N_K(k_e, k_\nu) n_K(k_e, k_\nu)) \right\}, \quad (36)$$

where  $M_K(k_e, k_\nu)$ ,  $m_K(k_e, k_\nu)$ ,  $N_K(k_e, k_\nu)$ , and  $n_K(k_e, k_\nu)$  are reduced matrix elements as defined in Eq. (A13), with their respective radial functions being  $F(r)$ ,  $f(r)$ ,  $G(r)$ , and  $g(r)$ . These radial functions are expressed as follows:

$$F(r) = H(r) \{ C_V G_{--} j_{k_\nu-1}(p_\nu r) - C'_V G_{-+} j_{k_\nu}(p_\nu r) \} + D(r) \{ C'_V G_{-+} j_{k_\nu-1}(p_\nu r) - C_V G_{++} j_{k_\nu}(p_\nu r) \}$$

<sup>3</sup>In our previous work described in Ref. [5] and adopted in our subsequent publications, we explicitly included a minus sign in the formulae in recognition that all the superallowed Fermi transitions involved positron emitters. Thus the shape-correction function  $C(Z, W)$  was modified to  $C(Z, W)(1 - \gamma_1 b_F / W)$  and a fit of  $\mathcal{F}t(1 - \gamma_1 b_F / (W))$  to a constant yielded a value of  $b_F$  that was negative. Currently in Eq. (28) we have defined  $b_F$  such that  $C(Z, W)$  is modified to  $C(Z, W)(1 + \gamma_1 b_F / W)$  and hence we are now quoting  $b_F$  with a positive sign.

$$\begin{aligned}
f(r) &= h(r)\{C_V G_{--} j_{k_v-1}(p_\nu r) - C'_V G_{-+} j_{k_v}(p_\nu r)\} \\
&\quad + d(r)\{C'_V G_{-+} j_{k_v-1}(p_\nu r) - C_V G_{++} j_{k_v}(p_\nu r)\} \\
G(r) &= H(r)\{-C'_V G_{--} j_{k_v-1}(p_\nu r) + C_V G_{-+} j_{k_v}(p_\nu r)\} \\
&\quad + D(r)\{-C_V G_{-+} j_{k_v-1}(p_\nu r) + C'_V G_{++} j_{k_v}(p_\nu r)\} \\
g(r) &= h(r)\{-C'_V G_{--} j_{k_v-1}(p_\nu r) + C_V G_{-+} j_{k_v}(p_\nu r)\} \\
&\quad + d(r)\{-C_V G_{-+} j_{k_v-1}(p_\nu r) + C'_V G_{++} j_{k_v}(p_\nu r)\},
\end{aligned} \tag{37}$$

where the functions  $H$ ,  $D$ ,  $h$ , and  $d$  are linear combinations of the electron functions,  $f_\kappa(r)$  and  $g_\kappa(r)$ , as given in Eq. (A15). The angular momentum factors  $G_{\pm\pm}$  are defined in Eq. (A16).

### 1. Order-of-magnitude estimates

Consider a pure Fermi transition for which the multipolarity is  $K = 0$  and only the lowest lepton partial waves,  $k_e = 1$  and  $k_\nu = 1$ , are kept. Then, as in Sec. IV C 1, the amplitudes become

$$\begin{aligned}
M_0(1, 1) &= C_V M_F + \dots \\
m_0(1, 1) &= \text{small} \\
N_0(1, 1) &= -C'_V M_F + \dots \\
n_0(1, 1) &= \text{small}.
\end{aligned} \tag{38}$$

The shape-correction function is then

$$C(Z, W) = |M_F|^2 \frac{1}{2} (C_V^2 + C_V'^2). \tag{39}$$

We see that the dominant impact of the right-hand current is simply to scale the statistical rate function by  $\frac{1}{2}(1 + C_V'^2/C_V^2)$ . This has no impact on the CVC test that demonstrates that  $\mathcal{F}t = \text{constant}$ , but it does shift the value of the vector coupling constant and thus the deduced value of  $V_{ud}^2$ . However,  $V_{ud}^2$  is obtained from the ratio of  $\beta$  decay to muon-decay rates, so before we can make any definitive statement on the effect of a right-hand current on  $V_{ud}^2$ , we must first examine the impact of that current on muon decay. We will show next that the correction due to a right-hand current is second order in small quantities in muon decay but first order in vector  $\beta$  decay. To this end we examine a more general Hamiltonian presented by Herczeg [170].

### 2. The effect on $V_{ud}^2$

In the  $SU(2)_L \times U(1)$  standard model, the semileptonic weak interaction Hamiltonian can be written schematically as

$$H_{SM} = \frac{G_F}{\sqrt{2}} V_{ud} (V - A)(V - A), \tag{40}$$

where the first factor of  $V - A$  represents the lepton currents,  $V = \bar{\phi}_e \gamma_\mu \phi_{\nu_e}$  and  $-A = \bar{\phi}_e \gamma_\mu \gamma_5 \phi_{\nu_e}$ , whereas the second  $V - A$  represents the hadron currents,  $V = \bar{\psi}_p \gamma_\mu \psi_n$  and  $-A = \bar{\psi}_p \gamma_\mu \gamma_5 \psi_n$ . The weak interaction coupling constant is  $G_F/\sqrt{2} = g^2/8M_W^2$ , where  $g$  is the basic coupling constant of the Weinberg-Salam standard model and  $M_W$  is the mass of the exchanged  $W$  boson.

Herczeg [170,171] considers an extension that is the most general form for nonderivative local four-fermion couplings:

$$\begin{aligned}
H &= a_{LL}(V - A)(V - A) + a_{LR}(V - A)(V + A) \\
&\quad + a_{RL}(V + A)(V - A) + a_{RR}(V + A)(V + A),
\end{aligned} \tag{41}$$

where, again, the first factor represents the lepton currents and the second the hadron currents. The lepton fields are now written as  $V = \bar{\phi}_e \gamma_\mu \phi_{\nu_e}^L$  or  $\bar{\phi}_e \gamma_\mu \phi_{\nu_e}^R$  depending on whether the chirality of the neutrino is left-handed for  $V - A$  coupling or right-handed for  $V + A$  coupling. The neutrino states are in general linear combinations of mass eigenstates,

$$\phi_{\nu_e}^L = \sum_i U_{ei} \phi_i^L \quad \phi_{\nu_e}^R = \sum_i V_{ei} \phi_i^R, \tag{42}$$

where  $U_{ei}$  and  $V_{ei}$  are first-row elements of the neutrino mixing matrix. The observed  $\beta$  decay probability is the sum of the probabilities of decays into the energetically allowed neutrino mass eigenstates. We follow Herczeg [170,171] in assuming that the neutrinos produced in  $\beta$  decay are light enough that the effects of their masses can be neglected. In particular, the terms that arise from the interference between amplitudes involving neutrinos of different chirality are dropped. Then the effect of neutrino mass mixing can be taken into account by our multiplying the coupling constants  $a_{LL}$  and  $a_{LR}$  by  $\sqrt{u_e}$ , and  $a_{RL}$  and  $a_{RR}$  by  $\sqrt{v_e}$ , where

$$u_e = \sum_i' |U_{ei}|^2 \quad v_e = \sum_i' |V_{ei}|^2. \tag{43}$$

The prime on the summation indicates that the sum extends only over the neutrinos that are light enough to be produced in  $\beta$  decay. Note that if all the neutrinos are light for both left-handed and right-handed chiralities, then  $u_e = v_e = 1$  as a consequence of the unitarity of the neutrino mixing matrix.

Herczeg's Hamiltonian, Eq. (41), can be rewritten as follows:

$$\begin{aligned}
H &= (a_{LL} + a_{LR} + a_{RL} + a_{RR})VV \\
&\quad + (-a_{LL} - a_{LR} + a_{RL} + a_{RR})AV \\
&\quad + (-a_{LL} + a_{LR} - a_{RL} + a_{RR})VA \\
&\quad + (a_{LL} - a_{LR} - a_{RL} + a_{RR})AA.
\end{aligned} \tag{44}$$

We can compare this with the Jackson, Treiman, and Wyld (JTW) Hamiltonian [166], which in the current notation becomes

$$\begin{aligned}
H_{JTW} &= (C_V V - C'_V A)V + (-C_A A + C'_A V)A \\
&= C_V VV - C'_V AV + C'_A VA - C_A AA.
\end{aligned} \tag{45}$$

Thus we identify the correspondences<sup>4</sup> as

$$\begin{aligned}
C_V &= a_{LL} + a_{LR} + a_{RL} + a_{RR} \\
C'_V &= a_{LL} + a_{LR} - a_{RL} - a_{RR} \\
C_A &= -a_{LL} + a_{LR} + a_{RL} - a_{RR} \\
C'_A &= -a_{LL} + a_{LR} - a_{RL} + a_{RR}.
\end{aligned} \tag{46}$$

<sup>4</sup>Herczeg [170,171] employs a metric that leads to a different sign on the  $\gamma_5$  matrix, so his correspondences yield a different overall sign from ours for  $C'_V$  and  $C_A$ .



For Fermi  $\beta$  decay, only the vector part of the weak hadron current contributes, so the decay rate,  $\Gamma_\beta$ , as shown earlier in Eq. (39), is given by the following proportionality:

$$\begin{aligned}\Gamma_\beta &\propto \frac{1}{2}(|C_V|^2 + |C'_V|^2) \\ &= |a_{LL} + a_{LR}|^2 + |a_{RL} + a_{RR}|^2 \\ &= |a_{LL}|^2(|1 + \bar{a}_{LR}|^2 + |\bar{a}_{RL} + \bar{a}_{RR}|^2) \\ &\simeq |a_{LL}|^2(1 + 2Re\bar{a}_{LR} + \dots),\end{aligned}\quad (47)$$

where  $\bar{a}_{ij} = a_{ij}/a_{LL}$ .

To continue our determination of  $V_{ud}$  we need to consider the purely leptonic muon decay. Herczeg [170] writes the effective Hamiltonian for muon decay in analogy to Eq. (41) as

$$\begin{aligned}H &= c_{LL}(V - A)(V - A) + c_{LR}(V - A)(V + A) \\ &\quad + c_{RL}(V + A)(V - A) + c_{RR}(V + A)(V + A).\end{aligned}\quad (48)$$

The coupling constants in Eqs. (48) and (41) are related by the CKM matrix elements by

$$\begin{aligned}a_{LL} &= c_{LL}V_{ud}^L \\ a_{LR} &= c_{LR}e^{i\alpha}V_{ud}^R \\ a_{RL} &= c_{RL}V_{ud}^L \\ a_{RR} &= c_{RR}e^{i\alpha}V_{ud}^R.\end{aligned}\quad (49)$$

Here  $V_{ud}^L$  is the  $ud$ -matrix element of the CKM matrix for left-handed chirality quarks, and  $V_{ud}^R$  is for right-handed chirality quarks. The phase  $\alpha$  is a CP-violating phase in the right-handed CKM matrix. The decay rate,  $\Gamma_\mu$ , for muon decay is given by

$$\begin{aligned}\Gamma_\mu &\propto |c_{LL}|^2 + |c_{LR}|^2 + |c_{RL}|^2 + |c_{RR}|^2 \\ &= |c_{LL}|^2(1 + |\bar{c}_{LR}|^2 + |\bar{c}_{RL}|^2 + |\bar{c}_{RR}|^2),\end{aligned}\quad (50)$$

where  $\bar{c}_{ij} = c_{ij}/c_{LL}$ .

Combining Eqs. (47) and (50), we obtain an expression that connects the ratio of  $\beta$ -decay to muon-decay rates with the value of  $|V_{ud}^L|^2$ , viz.:

$$\frac{\Gamma_\beta}{\Gamma_\mu} = |V_{ud}^L|^2 \frac{|1 + \bar{a}_{LR}|^2 + |\bar{a}_{RL} + \bar{a}_{RR}|^2}{1 + |\bar{c}_{LR}|^2 + |\bar{c}_{RL}|^2 + |\bar{c}_{RR}|^2}.\quad (51)$$

In the standard model, only  $a_{LL}$  and  $c_{LL}$  are nonzero; in any case, the quantities  $\bar{a}_{ij}$  and  $\bar{c}_{ij}$  with  $ij = LR, RL$ , or  $RR$  can certainly be considered small. The correction to the muon decay rate from right-handed interactions is therefore seen to be second order in small quantities, whereas the correction to Fermi beta decay rate is first order. Keeping only first-order small quantities, Eq. (51) reduces to<sup>5</sup>

$$\frac{\Gamma_\beta}{\Gamma_\mu} = |V_{ud}^L|^2(1 + 2Re\bar{a}_{LR}).\quad (52)$$

If the neutrino masses are such that  $u_e \neq 1$  and  $v_e \neq 1$  then this equation is modified to

$$\frac{\Gamma_\beta}{\Gamma_\mu} = |V_{ud}^L|^2 \frac{(u_e)_\beta}{[(u_e)_\mu(u_\mu)_\mu]^{1/2}}(1 + 2Re\bar{a}_{LR}),\quad (53)$$

where  $(u_e)_\beta$  in the numerator is given by the  $u_e$  in Eq. (43), with the sum extended over neutrinos light enough to be produced in  $\beta$  decay, whereas in the denominator  $(u_e)_\mu$  is given by the same expression but with the sum extended over neutrinos light enough to be produced in muon decay. Note that the  $Q$  value for muon decay is 105 MeV, a factor of 10 times larger than the  $Q$  value for any Fermi  $\beta$  decay we are considering. Also  $u_\mu$  in Eq. (53) is defined as  $\sum_i |U_{\mu i}|^2$ , where  $U_{\mu i}$  are second-row elements of the neutrino mixing matrix. In what follows, we will assume  $(u_e)_\beta = (u_e)_\mu = (u_\mu)_\mu = 1$ .

Before proceeding to numeric limits, we show how the current formulae relate to the simpler and more restrictive manifest left-right symmetric model [172]. In this model the departure from maximal parity violation is entirely because of the presence of a second  $W$  boson whose mass is much heavier than the usual  $W$  boson. If left-hand couplings are mediated by the boson  $W_L$ , and right-hand couplings by  $W_R$ , then  $W_L$  and  $W_R$  will be linear combinations of the mass eigenstates  $W_1$  and  $W_2$ : viz.

$$\begin{aligned}W_L &= W_1 \cos \zeta + W_2 \sin \zeta \\ W_R &= e^{i\omega}(-W_1 \sin \zeta + W_2 \cos \zeta)\end{aligned}\quad (54)$$

and  $\omega$  is a CP violating phase. If it is further assumed that, apart from the different masses of the  $W_1$  and  $W_2$  bosons, the coupling constants and CKM matrix elements are identical for left-hand and right-hand couplings, then there are only two parameters in this model. These parameters are  $\delta = (m_1/m_2)^2$  and  $\zeta$ , where  $m_1$  and  $m_2$  are the masses of  $W_1$  and  $W_2$  respectively. Both parameters are small and, of course, are zero in the standard model. The parameters of Herczeg's Hamiltonian, Eq. (41), can be expressed in terms of  $\delta$  and  $\zeta$  [171] as follows:

$$\begin{aligned}a_{LL} &= \frac{g^2}{8m_1^2}V_{ud}^L & \bar{a}_{RR} &= \delta \\ \bar{a}_{LR} &= \bar{a}_{RL} = -e^{i\omega}\zeta & &\rightarrow -\zeta\end{aligned}\quad (55)$$

for negligible CP-violating effects. In this limit, the expression for the ratio of Fermi  $\beta$  to muon decay rates, Eq. (52), reduces to the following:

$$\frac{\Gamma_\beta}{\Gamma_\mu} = |V_{ud}^L|^2(1 - 2\zeta).\quad (56)$$

This is the expression we used in our earlier work [1] to set limits on the extent of right-hand currents.

### 3. Numeric limit

Let us now insert the experimental values from our survey data for the  $\beta$  decay and muon-decay rates to determine an experimental value for  $|V_{ud}^L|^2$ , which we will write as  $|V_{ud}^L|_{\text{expt}}^2$ . This is the value we recorded earlier in Eq. (17). If analogous interactions exist in all three generations, then Eq. (52) can be

<sup>5</sup>Herczeg [171] also considers the possibility that the relation between purely leptonic and semileptonic Hamiltonians, Eq. (49), is not sufficiently general. He writes  $a_{LL} = (a_{LL})_{SM} + a'_{LL}$ , with  $(a_{LL})_{SM} = c_{LL}V_{ud}^L$ . Then Eq. (52) becomes  $\Gamma_\beta/\Gamma_\mu = |V_{ud}^L|^2(1 + 2Re(\bar{a}'_{LL} + \bar{a}_{LR}))$ , where  $\bar{a}'_{LL} = a'_{LL}/(a_{LL})_{SM}$ . We pursue this no further, but it is obvious the formulae above can accommodate this extension with a simple replacement of  $\bar{a}_{LR}$  with  $\bar{a}'_{LL} + \bar{a}_{LR}$ .

written as follows:

$$\begin{aligned} \sum_i |V_{ui}|_{\text{expt}}^2 &= \sum_i |V_{ui}^L|^2 (1 + 2\text{Re}\bar{a}_{LR}) \\ &= (1 + 2\text{Re}\bar{a}_{LR}), \end{aligned} \quad (57)$$

where in the second line we have inserted the condition for unitarity of the CKM matrix. Adopting the PDG's recommended values [154] for  $|V_{us}^L|$  and  $|V_{ub}^L|$  [see text preceding Eq. (19)], we obtain the following result from Eq. (57):

$$\begin{aligned} (0.9966 \pm 0.0014) &= (1 + 2\text{Re}\bar{a}_{LR}) \\ \text{Re}\bar{a}_{LR} &= -0.0017 \pm 0.0007. \end{aligned} \quad (58)$$

Within the context of the manifest left-right symmetric model [see Eq. (56)], this result corresponds to  $\zeta = 0.0017 \pm 0.0007$ , a similar value to the one we reported previously [1]. The result of a nonzero  $a_{LR}$  or  $\zeta$  simply reflects the fact that the experimental values of the first-row CKM matrix elements do not satisfy the unitarity requirement.

If, instead, we adopt the average of the E865 and the E832 values [2,3] for  $V_{us}$  rather than the PDG average, then the result in Eq. (59) is modified to the following:

$$\begin{aligned} (0.9993 \pm 0.0012) &= (1 + 2\text{Re}\bar{a}_{LR}) \\ \text{Re}\bar{a}_{LR} &= -0.0004 \pm 0.0006. \end{aligned} \quad (59)$$

This result is consistent with no right-hand currents and unitarity being satisfied in the experimental CKM matrix elements.

## V. CONCLUSIONS

Previous surveys of superallowed Fermi  $\beta$  decay have at times noted disagreement [4,6] among the derived  $\mathcal{F}t$  values and at other times agreement [5,7]. When disagreement was evident, subsequent attention paid to the problem led to both theoretical and experimental advances. As presented here, in Sec. III, the status now is of excellent agreement among all  $\mathcal{F}t$  values—to better than 3 parts in  $10^4$  over a wide range of nuclei from  $A = 10$  to  $A = 74$ . Such agreement confirms the expectations of CVC, allows very restrictive limits to be set on the possible presence of scalar currents, and makes it possible to go forward with confidence to the next steps—the determination of  $V_{ud}$  and the unitarity test of the CKM matrix.

The outstanding challenge at this time is that the value obtained for  $V_{ud}$ , when combined with the current PDG-recommended values of  $V_{us}$  and  $V_{ub}$ , leads to a unitarity test that fails by more than 2 standard deviations. There are no evident defects in the calculated radiative and isospin-symmetry-breaking corrections that could remove this problem and, indeed, a shift in any one of these corrections large enough to restore unitarity would be almost impossible to justify [1]. Moreover, the derived value of  $V_{ud}$  from nuclear decays has been remarkably stable for three decades despite a vast increase in the quantity of high-quality data and many theoretical refinements in the calculations of the correction terms.

So if any progress is to be made in firmly establishing (or eliminating) the discrepancy with unitarity, both theory

and experiment must be brought to bear afresh on the principal sources of uncertainty. Although we focus here on improving the nuclear contribution to the unitarity test, additional experiments are also required for neutron, pion, and kaon decays. The first two provide independent, though so far much less precise, values for  $V_{ud}$ ; the third establishes the value of  $V_{us}$ , which may ultimately turn out to be solely responsible for restoring the CKM matrix to unitarity. Whatever the outcome for unitarity, however, the results of all these studies will provide crucial information, either in characterizing new physics beyond the standard model or in setting a tight limit on its possible existence.

We have taken pains throughout this work to pay careful attention to all uncertainties, theoretical and experimental. In Sec. II A we detailed the various contributions to the uncertainty in  $|V_{ud}|^2$ . Of these, by far the largest is because of the nucleus-independent radiative correction,  $\Delta_R^V$ . Its uncertainty arises primarily from a box diagram involving the exchange of one  $W$  boson and one photon between the hadron and the lepton. To make the loop integration tractable, it is divided by a scale parameter into high- and low-energy portions. The high-energy contribution can be computed reliably [173] but the low-energy one, as calculated originally by Sirlin [174], depends on the choice of scale parameter. Sirlin chose [164,165] a reasonable range for this parameter, which has been retained by subsequent authors [175,176]. It is this choice of range that drives the overall uncertainty on  $\Delta_R^V$ . Recent work [177] with effective field theories based on chiral perturbation theory has been unable to improve the situation: although this approach replaces the low-energy contributions to the loop diagrams by well-defined low-energy constants, the values of these constants are not known *a priori*. How to obtain a more refined, first-principles computation of the low-energy contribution remains an open theoretical problem [173] but one of considerable importance and urgency. Not only is this uncertainty the principal limitation on the precision with which  $V_{ud}$  can be determined from nuclear superallowed  $\beta$  decay, but it will have a similar limiting effect on its determination from neutron and pion decays as well.

The next largest contributor to the error budget on  $|V_{ud}|^2$  is the isospin-symmetry-breaking correction,  $\delta_C$ . Although uncertainties have been individually determined for the most recent calculations [159] of  $\delta_C - \delta_{NS}$  (see Table IX), the dominant source of  $|V_{ud}|^2$  uncertainty attributable to  $\delta_C$  arises from the small systematic difference between the results from different theoretical techniques used to calculate  $\delta_C$ . Our approach [159], using Woods-Saxon functions, yields larger  $\delta_C$  values than the Ormand-Brown one [160], using Hartree-Fock functions. Here we have taken the democratic approach, considering that these two sets of calculations are equally likely to be correct and letting the difference between their results determine a systematic uncertainty that we apply to the final result [see Eq. (12)].

If reducing the uncertainty on  $\Delta_R^V$  must rank as the first priority for future theoretical work, then improving our confidence in  $\delta_C$  can be taken as the top priority challenge for experiment. Although there is no way to check the correctness of the absolute values of  $\delta_C$  from experiment, it is possible to check the nucleus-to-nucleus variations in the calculated




FIG. 4. Experimental  $ft$  values plotted as a function of the charge on the daughter nucleus,  $Z$ . The bands represent the theoretical quantity  $\overline{\mathcal{F}t}/[(1 + \delta'_R)(1 - \delta_C + \delta_{NS})]$ . The two groups distinguish those  $\beta$  emitters whose parent nuclei have isospin  $T_z = -1$  (darker shading) from those with  $T_z = 0$  (lighter shading).

values. The method, which is illustrated in Fig. 4, is based on the validity of the CVC hypothesis that the corrected  $\mathcal{F}t$  values for the superallowed  $0^+ \rightarrow 0^+$  decays should be constant. In the figure we compare the uncorrected measured  $ft$  values (points and error bars) with the theoretical quantity  $\overline{\mathcal{F}t}/[(1 + \delta'_R)(1 - \delta_C + \delta_{NS})]$  shown as a band, the width of which represents its estimated error. With the average  $\mathcal{F}t$  value,  $\overline{\mathcal{F}t}$ , taken from Table IX, this comparison specifically tests the collective ability of all three calculated correction terms to reproduce the variations in  $ft$  from one transition to another. However, because  $\delta'_R$  is almost independent of  $Z$  when  $Z > 10$ , this test really probes directly the effectiveness of the calculated values of  $\delta_C - \delta_{NS}$ .

It can be seen that there is remarkable agreement between theory and experiment. In assessing the significance of this agreement, it is important to recognize that the calculations of  $\delta_C$  and  $\delta_{NS}$  for  $Z \leq 26$  are based on well-established shell-model wave functions and were further tuned to reproduce measured binding energies, charge radii and coefficients of the isobaric multiplet mass equation [159]. The origins of the calculated correction terms for all cases are completely independent of the superallowed decay data. Thus, the agreement in the figure between the measured superallowed data points and the theoretical band is already a powerful validation of the calculated corrections used in determining that band. The validation becomes even more convincing when we consider that it would require a pathological fault indeed in the theory to allow the observed nucleus-to-nucleus variations in  $\delta_C$  to be reproduced in such detail while failing to obtain the *absolute* values to comparable precision. Pleasing as the agreement in Fig. 4 is, though, new experiments can still improve the test, making it even more demanding, and can ultimately reduce the uncertainty in  $\delta_C$  further.

These new experiments can follow different paths. One is to improve the precision on the nine superallowed transitions whose  $ft$  values are already known to within 0.15% or better. On the one hand, these are the easiest cases to study, all having stable daughters and all, except  $^{10}\text{C}$ , decaying predominantly (>99%) through a single superallowed transition. On the other hand, they have been the subject of intense scrutiny for at least the past four decades and, given the number of careful measurements already published, the prospects for really significant improvements in these cases, at least in the near future, do not seem bright. Nevertheless, a glance at Fig. 2 shows that some modest improvements are certainly possible. If we accept as a goal that experiment should be a factor of two more precise than theory, then we see that the  $Q_{EC}$  values for  $^{10}\text{C}$ ,  $^{14}\text{O}$ ,  $^{26}\text{Al}^m$ ,  $^{42}\text{Sc}$ , and  $^{46}\text{V}$ , the half-lives of  $^{10}\text{C}$ ,  $^{26}\text{Al}^m$ ,  $^{42}\text{Sc}$ , and  $^{50}\text{Mn}$ , and the branching ratio for  $^{10}\text{C}$  can all bear improvement.

A second path is to expand the number of precisely measured superallowed emitters to include cases for which the calculated nuclear-structure-dependent corrections are larger, or show larger variations from nuclide to nuclide, than the values applied to the nine currently best known transitions. We argue that if the calculations reproduce the experimentally observed variations where they are large, then that must surely verify their reliability for the original nine transitions whose corrections are considerably smaller. The recent results for  $^{22}\text{Mg}$ ,  $^{34}\text{Ar}$ , and  $^{74}\text{Rb}$  are the first cases of this type to reach sufficient precision that they can contribute to the test (see Fig. 4) but more are sure to follow. We have included in our survey all cases that we believe are potential candidates within the next few years.

Without doubt, these new cases present serious experimental challenges. In general, the parent nuclei are more exotic

and thus more difficult to produce in pure and statistically useful quantities. They also exhibit more complex branching patterns, which for the cases with  $A \geq 62$  include Gamow-Teller transitions that may be unobservable individually but collectively can play a nonnegligible role [81]. These heavier nuclei also have very short half-lives, which currently limit the precision with which their  $Q_{EC}$  values can be measured. Even so, all these obstacles are obviously now being overcome and we may reasonably hope that before long there will not only be more cases with precisely measured parameters, but there will be more than one measurement of each parameter, an essential prerequisite for reliable results at the level of precision needed to constrain the correction terms.

Although it would not impact significantly on the unitarity question, there is an additional reason to improve the precision with which the  $ft$  values are known, particularly for the cases with  $A \leq 26$ . A scalar current, if it exists, would manifest itself as a  $1/W$  dependence in the shape-correction function used in the  $f$ -value calculations [see Eq. (28)]. Because the superallowed transition energies increase with increasing  $A$ , this effect would be strongest for the lightest nuclei,  $^{10}\text{C}$  and  $^{14}\text{O}$ . Improved precision in the  $\mathcal{F}t$  values for those two nuclei would act directly to reduce the limits set on a possible scalar current.

In conclusion, we can assert that world data for superallowed  $0^+ \rightarrow 0^+$   $\beta$  decays strongly support the CVC expectation of an unrenormalized vector coupling constant and also set a new limit on the possible existence of scalar currents. The nuclear-structure-dependent corrections used in the analyses of these data have so far stood up very favorably to experimental tests, and the value currently obtained with them for  $V_{ud}$  is deemed to be very robust, even though it is an important component of the top-row test of CKM unitarity that, until recently, evidently failed by more than 2 standard deviations. We have indicated the improvements required from both theory and experiment to increase the precision in future so as to produce a more definitive result for the unitarity test.

### ACKNOWLEDGMENTS

The work of J.C.H. was supported by the U.S. Department of Energy under Grant DE-FG03-93ER40773 and by the Robert A. Welch Foundation. I.S.T. thanks the Cyclotron Institute of Texas A&M University for its hospitality during several 2-month visits.

### APPENDIX: STATISTICAL RATE FUNCTION

The statistical rate function is an integral over phase space,

$$f = \int_1^{W_0} pW(W_0 - W)^2 F(Z, W) S(Z, W) dW, \quad (\text{A1})$$

where  $W$  is the electron total energy in electron rest-mass units,  $W_0$  is the maximum value of  $W$ ,  $p = (W^2 - 1)^{1/2}$  is the electron momentum,  $Z$  is the charge number of the daughter nucleus (positive for electron emission, negative for positron emission),  $F(Z, W)$  is the Fermi function, and  $S(Z, W)$  is the

shape-correction function. If the shape-correction function is put to unity, the integral becomes the customarily defined one for  $\beta$  decay, which we will denote as  $f_{\text{stat}}$  as follows:

$$f_{\text{stat}} = \int_1^{W_0} pW(W_0 - W)^2 F(Z, W) dW. \quad (\text{A2})$$

The exact evaluation of  $f$  differs from  $f_{\text{stat}}$  by 0.2% at  $A = 10$  up to 5.7% at  $A = 74$ . Thus, to maintain 0.1% accuracy for  $f$  over that range, we must determine the shape-correction function itself with 2% accuracy. Obtaining this accuracy requires consideration of the following issues:

- The electron wave functions can no longer be simply those of the lowest partial wave ( $j = 1/2$ ) generated by a point nuclear charge and evaluated at radius  $R$ , the nuclear surface, but must be the exact functions for some chosen nuclear charge-density distribution.
- The lepton wave functions exhibit some  $r^2$  dependence over the nuclear volume, leading to what are called second-forbidden corrections. Furthermore, a more accurate treatment of the weak interaction leads to relativistic and induced-current corrections. All these effects must be incorporated because they impact on the nuclear matrix elements and inject a mild nuclear-structure dependence into the evaluation of  $f$ .
- The atomic electrons cannot be ignored but must be accommodated approximately in a screening correction.

In what follows we describe the ingredients of a code we have written that incorporates these effects. It is based on the formalism of Behrens and Bühring [167]. Note that they define

$$F(Z, W)S(Z, W) = F_0 L_0 C(Z, W), \quad (\text{A3})$$

where  $F_0 = 2F(Z, W)/(1 + \gamma_1)$ . The purpose of their redefining the shape correction factor in this way was to remove the historic requirement to evaluate the electron wave functions at the nuclear surface. The product  $F_0 L_0$  is given entirely in terms of the amplitudes of the electron wave function at the origin [see Eq. (A9) below], and  $C(Z, W)$  is the shape-correction function defined with respect to this choice.

### Electron radial wave functions

The wave function for the electron emitted in  $\beta$  decay is given by the solution to the Dirac equation with an external electromagnetic field present but restricted to the special case where the vector potential vanishes identically and the scalar potential is static and spherically symmetric. We solve in spherical coordinates and introduce a partial wave expansion such that the basis states are written:

$$\psi_\kappa^\mu = \begin{pmatrix} \text{sign}(\kappa) f_\kappa(r) \chi_{-\kappa}^\mu \\ g_\kappa(r) \chi_\kappa^\mu \end{pmatrix}, \quad (\text{A4})$$

where  $f_\kappa$  and  $g_\kappa$  are radial functions and  $\chi_\kappa^\mu$  are the usual spin-angular-momentum wave functions describing the coupling of the orbital angular momentum  $l$  and spin  $\frac{1}{2}$  to give a total

angular momentum  $j$  with  $z$ -component  $\mu$ : i.e.

$$\chi_\kappa^\mu = i^l \sum_{m_l m_s} \left\langle l m_l \frac{1}{2} m_s \middle| j \mu \right\rangle Y_{l m_l}(\hat{\mathbf{r}}) \chi_{m_s}. \quad (\text{A5})$$

The eigenvalue  $\kappa$  is

$$\kappa = -j(j+1) + l(l+1) - \frac{1}{4} \quad (\text{A6})$$

and has values

$$\begin{aligned} \kappa &= -(l+1) = -(j + \frac{1}{2}) \quad \text{if } j = l + \frac{1}{2} \\ \kappa &= l = (j + \frac{1}{2}) \quad \text{if } j = l - \frac{1}{2}. \end{aligned} \quad (\text{A7})$$

The radial functions are solutions to the following pair of coupled equations:

$$\begin{aligned} \frac{d g_\kappa}{d r} + \frac{(\kappa+1)}{r} g_\kappa - [W+1-V(r)] f_\kappa &= 0 \\ \frac{d f_\kappa}{d r} - \frac{(\kappa-1)}{r} f_\kappa + (W-1-V(r)) g_\kappa &= 0. \end{aligned} \quad (\text{A8})$$

Here  $V(r)$  is a spherically symmetric static potential that represents the interaction of the electron with the charge distribution of the nucleus.

Our task is to solve the pair of coupled radial equations, Eq. (A8), in three regions,  $0 \leq r \leq R_1$ ,  $R_1 \leq r \leq R_2$ ,  $R_2 \leq r \leq \infty$ , matching the solutions at each region boundary. The first region is the one over which the nuclear charge density is nonzero. Here, we establish a power-series solution, regular at the origin, as the starting solution and integrate numerically to  $R_1$ . In the second region, we have a pure Coulomb potential, for which an analytic solution can be found in terms of confluent hypergeometric functions of complex argument. The asymptotic solution in the third region is expressed in terms of the desired outgoing waves and a phase shift. The unknowns in the calculation are the phase shift,  $\Delta_\kappa$ , and the normalization of the interior solution,  $\alpha_\kappa$ : they are determined from the matching conditions.

In the derivation of expressions for  $\beta$ -decay observables, certain combinations of amplitudes and phase shifts characterizing the electron wave functions appear. These combinations are called the  $\beta$ -decay Coulomb functions and are generally organized so that they are of order unity, with corrections of order  $(\alpha Z)^2$ . Because we are interested here only in the shape-correction function, the only such functions we need are  $F_0 L_0$ ,  $\lambda_k$  and  $\mu_k$ , where  $k = |\kappa|$ , which are defined as follows:

$$\begin{aligned} F_0 L_0 &= \frac{\alpha_{-1}^2 + \alpha_{+1}^2}{2p^2} \\ \lambda_k &= \frac{\alpha_{-k}^2 + \alpha_{+k}^2}{\alpha_{-1}^2 + \alpha_{+1}^2} \\ \mu_k &= \frac{\alpha_{-k}^2 - \alpha_{+k}^2}{\alpha_{-k}^2 + \alpha_{+k}^2} \frac{kW}{\gamma_k}, \end{aligned} \quad (\text{A9})$$

where  $\gamma_k = \{k^2 - (\alpha Z)^2\}^{1/2}$ . Although these functions have actually been tabulated by Behrens and Jänecke [178], we have not used the tables but, in the interests of precision, have explicitly computed the functions from the calculated values of  $\alpha_\kappa$ .

### Shape-correction function

Behrens and Bühring [167] give the following expression for the shape-correction function:

$$\begin{aligned} C(Z, W) &= \sum_{k_e k_\nu K} \lambda_{k_e} \left\{ M_K^2(k_e, k_\nu) + m_K^2(k_e, k_\nu) \right. \\ &\quad \left. - \frac{2\mu_{k_e} \gamma_{k_e}}{k_e W} M_K(k_e, k_\nu) m_K(k_e, k_\nu) \right\}, \end{aligned} \quad (\text{A10})$$

where the sums over  $k_e$  and  $k_\nu$  are partial-wave expansions of the electron and neutrino wave functions with  $k_e = j_e + \frac{1}{2}$  and  $k_\nu = j_\nu + \frac{1}{2}$ ,  $j_e$  and  $j_\nu$  being the electron and neutrino total angular momenta. The integer  $K$  represents the multipolarity of the transition operators and is limited to the range  $|j_e - j_\nu| \leq K \leq j_e + j_\nu$ . The functions  $M_K(k_e, k_\nu)$  and  $m_K(k_e, k_\nu)$  are given in terms of form factors [167] and we evaluate these form factors in the ‘‘impulse approximation.’’ In this approximation, the nucleus is treated as a collection of noninteracting nucleons so it is necessary to consider only the weak interaction as acting on a single nucleon. All the many-body aspects of the nucleus can thus be handled in the standard shell-model way. Let  $O$  be a one-body operator, which can be written

$$O = \sum_{\alpha\beta} \langle \alpha | O | \beta \rangle a_\alpha^\dagger a_\beta, \quad (\text{A11})$$

where  $a_\alpha^\dagger$  is the creation operator for a nucleon in quantum state  $\alpha$ , and  $a_\beta$  is an annihilation operator destroying a nucleon in state  $\beta$ . The matrix element of  $O$  for many-body states becomes

$$\langle f | O | i \rangle = \sum_{\alpha\beta} \langle \alpha | O | \beta \rangle \langle f | a_\alpha^\dagger a_\beta | i \rangle; \quad (\text{A12})$$

that is, the matrix element is a linear combination of single-particle matrix elements,  $\langle \alpha | O | \beta \rangle$ . The coefficients in the expansion,  $\langle f | a_\alpha^\dagger a_\beta | i \rangle$ , are called one-body density matrix elements (OBDME). We leave it, then, as the job of the shell model to provide the OBDMEs and deal here only with the single-particle matrix elements. The functions  $M_K(k_e, k_\nu)$  and  $m_K(k_e, k_\nu)$  are now given in terms of reduced nuclear matrix elements for a single-particle transition  $j_\beta \rightarrow j_\alpha$  by

$$\begin{aligned} M_K(k_e, k_\nu) &= \frac{\sqrt{4\pi}}{\hat{K} \hat{J}_i} \sum_{L_S} (-)^{K-L} \langle j_\alpha \| F(r) \hat{T}_{K L_S} \| j_\beta \rangle \\ m_K(k_e, k_\nu) &= \frac{\sqrt{4\pi}}{\hat{K} \hat{J}_i} \sum_{L_S} (-)^{K-L} \langle j_\alpha \| f(r) \hat{T}_{K L_S} \| j_\beta \rangle, \end{aligned} \quad (\text{A13})$$

where  $\hat{j}$  is a short-hand notation for  $(2j+1)^{1/2}$  and  $J_i$  is the spin of the decaying nucleus. The radial functions are

$$\begin{aligned} F(r) &= H(r) \{ G_{--} j_{k_\nu-1}(p_\nu r) - G_{-+} j_{k_\nu}(p_\nu r) \} \\ &\quad + D(r) \{ G_{+-} j_{k_\nu-1}(p_\nu r) - G_{++} j_{k_\nu}(p_\nu r) \} \\ f(r) &= h(r) \{ G_{--} j_{k_\nu-1}(p_\nu r) - G_{-+} j_{k_\nu}(p_\nu r) \} \\ &\quad + d(r) \{ G_{+-} j_{k_\nu-1}(p_\nu r) - G_{++} j_{k_\nu}(p_\nu r) \}, \end{aligned} \quad (\text{A14})$$

with

$$\begin{aligned} H(r) &= \frac{1}{2} [f_{k_e}(r) + g_{-k_e}(r)] \\ D(r) &= \frac{1}{2} [g_{k_e}(r) - f_{-k_e}(r)] \\ h(r) &= \frac{1}{2} [f_{k_e}(r) - g_{-k_e}(r)] \\ d(r) &= \frac{1}{2} [g_{k_e}(r) + f_{-k_e}(r)]. \end{aligned} \quad (\text{A15})$$

Here  $f_\kappa(r)$  and  $g_\kappa(r)$  are radial electron functions,<sup>6</sup> whereas the spherical Bessel functions represent radial neutrino wave functions. The functions  $G_{++}$ ,  $G_{+-}$ ,  $G_{-+}$ , and  $G_{--}$  are short-hand notations for  $G_{KLS}(k_e, k_\nu)$ ,  $G_{KLS}(k_e, -k_\nu)$ ,  $G_{KLS}(-k_e, k_\nu)$ , and  $G_{KLS}(-k_e, -k_\nu)$  respectively, where the functions  $G_{KLS}(k_e, \kappa_\nu)$  contain all the angular momentum factors for the leptons and are given by the equation

$$G_{KLS}(k_e, \kappa_\nu) = i^{l_\nu+L+l_e} (-)^{j_e-j_\nu} \hat{K} \hat{J}_e \hat{J}_\nu \hat{l}_e \hat{l}_\nu (l_e 0 l_\nu 0 | L 0) \begin{Bmatrix} K & s & L \\ j_e & \frac{1}{2} & l_e \\ j_\nu & \frac{1}{2} & l_\nu \end{Bmatrix}. \quad (\text{A16})$$

Last, the operators  $\hat{T}_{KLS}$  depend on the angle and spin coordinates and are defined as

$$\begin{aligned} \hat{T}_{KLS}^M(\hat{\mathbf{r}}) &= (V_0 + A_0) i^L Y_{LM}(\hat{\mathbf{r}}) \delta_{K,L} \quad \text{if } s = 0 \\ &= (\mathbf{V} + \mathbf{A}) \cdot i^L \mathbf{Y}_{KLM}(\hat{\mathbf{r}}) \quad \text{if } s = 1, \end{aligned} \quad (\text{A17})$$

where  $V_0$  and  $A_0$  are the time parts of the vector and axial-vector hadronic currents and  $\mathbf{V}$  and  $\mathbf{A}$  are the space parts. Further  $\mathbf{Y}_{KLM}(\hat{\mathbf{r}})$  is a vector spherical harmonic [179], which in Eq. (A17) forms a scalar product with vectors  $\mathbf{V}$  and  $\mathbf{A}$ .

### Hadronic matrix element

For nucleons, the vector and axial-vector interactions are written

$$\begin{aligned} V_\mu &= g_V \gamma_\mu - f_M \sigma_{\mu\nu} q_\nu + i f_S q_\mu \\ A_\mu &= g_A \gamma_\mu \gamma_5 - f_T \sigma_{\mu\nu} q_\nu \gamma_5 + i f_P q_\mu \gamma_5. \end{aligned} \quad (\text{A18})$$

Were we discussing the weak interaction between pointlike spin-1/2 fermions, then we would set  $g_V = g_A = 1$  and  $f_M = f_S = f_T = f_P = 0$ . However, in considering nucleons, we recognize that they are not pointlike and furthermore they are influenced by the presence of the strong interaction. Thus Eq. (A18) presents the most general form of a vector and axial-vector interaction that is consistent with Lorentz invariance and excludes momentum operators beyond the first power. Here  $q_\mu = (p_f - p_i)_\mu$  is the momentum transfer. The coefficients, in principle, could be functions of  $q^2$  but, because of the low four-momentum transfer in  $\beta$  decay, this  $q^2$  dependence can be neglected and the coefficients are referred to as coupling constants with individual titles:  $g_V$  being vector;  $g_A$ , axial-vector;  $f_M$ , weak magnetic;

<sup>6</sup>The internal normalizations at the origin are here set to unity. Recall that these normalizations,  $\alpha_\kappa$ , have been separated out into the  $\beta$ -decay Coulomb functions,  $F_0 L_0$  [see Eq. (A9)].

$f_S$ , induced scalar;  $f_T$ , induced tensor; and  $f_P$ , induced pseudoscalar.

Our aim is to reduce the matrix element  $i\bar{u}_f(V_\mu + A_\mu)u_i$ , where  $\bar{u}_f$  and  $u_i$  are Dirac spinors characterizing nucleons of momentum  $\mathbf{p}_f$  and  $\mathbf{p}_i$ , to the nonrelativistic form involving Pauli two-component spinors,

$$\chi_{m_f}^\dagger (V_0 + \mathbf{V} + A_0 + \mathbf{A}) \chi_{m_i}, \quad (\text{A19})$$

by simply multiplying out the Dirac matrices involved and keeping terms to first order in  $|\mathbf{p}|/M_N$  and dropping terms in  $|\mathbf{p}^2|/M_N^2$  and higher order. This multiplication yields

$$V_0 = g_V + f_S [W_0 - V(r)] \quad (\text{A20})$$

$$\mathbf{V} = \frac{g_V}{2M_N} [\mathbf{p} + i\boldsymbol{\sigma} \times \mathbf{q}] + f_M i\boldsymbol{\sigma} \times \mathbf{q} - f_S \mathbf{q} \quad (\text{A21})$$

$$A_0 = -\frac{g_A}{2M_N} \boldsymbol{\sigma} \cdot \mathbf{p} - f_T \boldsymbol{\sigma} \cdot \mathbf{q} \quad (\text{A22})$$

$$\mathbf{A} = -g_A \boldsymbol{\sigma} + f_T [W_0 - V(r)] \boldsymbol{\sigma}, \quad (\text{A23})$$

where  $\mathbf{p} = \mathbf{p}_f + \mathbf{p}_i$  and  $\mathbf{q} = \mathbf{p}_f - \mathbf{p}_i$ . Note that the large terms of order unity occur in  $V_0$  and  $\mathbf{A}$ . Each of the four coupling constants denoted by an  $f$  are small and of order  $1/M_N$  and, as a consequence, terms in  $f/M_N$  have been dropped. Eqs. (A20) to (A23) are the quantities needed in the operators  $\hat{T}_{KLS}^M(\hat{\mathbf{r}})$  in Eq. (A17).

### Reduced matrix elements

All the  $\beta$ -decay observables [167] can be expressed in terms of the functions  $M_K(k_e, k_\nu)$  and  $m_K(k_e, k_\nu)$  defined in Eq. (A13). Here we are only interested in the shape-correction function, Eq. (A10), which is a particularly simple combination of these quantities. Before proceeding to evaluate  $M_K(k_e, k_\nu)$  and  $m_K(k_e, k_\nu)$ , we note that the expressions for both differ only in the presence of  $F(r)$  in one case and  $f(r)$  in the other. For simplicity in what follows, we will only explicitly display formulae incorporating  $f(r)$ ; obviously an equivalent set can be written with  $F(r)$ .

The operators  $\hat{T}_{KLS}^M(\hat{\mathbf{r}})$  are shown in Eq. (A17), but it is tidier if we incorporate the phase,  $(-)^{K-L}$  [see Eq. (A13)] into the operator. All the operators (some after rearrangement) can then be expressed as a product of spherical tensors, one in orbital space and one in spin space. So, generically the operators take the following form:

$$T_{KLS}^M(\mathbf{r}) = f(r) (-)^{K-L} T_{KM}(\Lambda_L \otimes \Sigma_S), \quad (\text{A24})$$

where  $\Lambda_L$  is a spherical tensor in orbital space of multipolarity  $L$  and  $\Sigma_S$  a spherical tensor in spin space.<sup>7</sup> We have introduced notation for a composite spherical tensor obtained from the combination of two other spherical tensors: i.e.

$$T_{KM}(\Lambda_L \otimes \Sigma_S) = \sum_{M_L M_S} \langle L M_L S M_S | K M \rangle \Lambda_{L M_L} \Sigma_{S M_S}. \quad (\text{A25})$$

<sup>7</sup>Note, the uppercase  $S$  in Eq. (A24) referring to the multipolarity of the spin operator is not the same as the lowercase  $s$  in Eq. (A17).

The single-particle wave functions from the shell model can also be expressed as products of orbital and spin space functions:

$$|jm\rangle = \sum_{m_l m_s} \left\langle l m_l \frac{1}{2} m_s \middle| j m \right\rangle R_{nl}(r) i^l Y_{lm_l}(\hat{\mathbf{r}}) \chi_{m_s}, \quad (\text{A26})$$

where  $n$  is the principal quantum number designating the number of nodes in the radial function. Notice the presence of  $i^l$  with the spherical harmonics.<sup>8</sup>

The first step in evaluating the reduced matrix element is to factorize it into orbital and spin reduced matrix elements:

$$\begin{aligned} & \left\langle \left( l_\alpha \frac{1}{2} \right) j_\alpha \middle| T_{KLS} \middle| \left( l_\beta \frac{1}{2} \right) j_\beta \right\rangle \\ &= (-)^{K-L} A_{(LS)K} \langle l_\alpha \| f(r) \Lambda_L \| l_\beta \rangle \langle \frac{1}{2} \| \Sigma_S \| \frac{1}{2} \rangle, \end{aligned} \quad (\text{A27})$$

where

$$A_{(LS)K} = \hat{j}_\alpha \hat{K} \hat{j}_\beta \begin{Bmatrix} l_\alpha & \frac{1}{2} & j_\alpha \\ l_\beta & \frac{1}{2} & j_\beta \\ L & S & K \end{Bmatrix}. \quad (\text{A28})$$

Our conventions on reduced matrix elements are those of Edmonds [179]. Next, we need to define two spin matrix elements, denoted  $S_0$  and  $S_1$  and given by

$$\begin{aligned} S_0 &\equiv \langle \frac{1}{2} \| 1 \| \frac{1}{2} \rangle = \sqrt{2} \delta_{S,0} \\ S_1 &\equiv \langle \frac{1}{2} \| \boldsymbol{\sigma} \| \frac{1}{2} \rangle = \sqrt{6} \delta_{S,1}; \end{aligned} \quad (\text{A29})$$

and two orbital matrix elements denoted  $L_L$  and  $L_{(J_1)L}(\mathbf{Q})$ . The first is

$$\begin{aligned} L_L &\equiv \langle l_\alpha \| f(r) i^L Y_L(\hat{\mathbf{r}}) \| l_\beta \rangle \\ &= i^{l_\alpha + L + l_\beta} \frac{\hat{l}_\alpha \hat{L} \hat{l}_\beta}{\sqrt{4\pi}} \begin{pmatrix} l_\alpha & L & l_\beta \\ 0 & 0 & 0 \end{pmatrix} \langle R_\alpha \| f(r) | R_\beta \rangle, \end{aligned} \quad (\text{A30})$$

where the last factor is the radial integral:

$$\langle R_\alpha \| f(r) | R_\beta \rangle \equiv \int_0^\infty R_\alpha(r) f(r) R_\beta(r) r^2 dr. \quad (\text{A31})$$

The second involves derivative operators and requires a little more care. The matrix element is

$$L_{(J_1)L}(\mathbf{Q}) \equiv \langle l_\alpha \| f(r) i^J T_L(Y_J \otimes \mathbf{Q}) \| l_\beta \rangle, \quad (\text{A32})$$

where  $\mathbf{Q}$  is either  $\mathbf{p} = \mathbf{p}_f + \mathbf{p}_i$  or  $\mathbf{q} = \mathbf{p}_f - \mathbf{p}_i$ . Thus we write  $\mathbf{Q}$  as  $\mathbf{p}_f \pm \mathbf{p}_i$  with the upper sign appropriate for  $\mathbf{p}$  and the lower sign for  $\mathbf{q}$ . Now  $\mathbf{Q}$  is also  $-i(\nabla_f \pm \nabla_i)$ , where the gradient operator acts on either the initial or final nuclear wave function but not on the integrand,  $f(r)$ . Thus the interpretation is as follows:  $\langle \phi_f | f(\mathbf{Q}) \phi_i \rangle = -i \{ \pm \langle \phi_f | f | \nabla \phi_i \rangle - \langle \nabla \phi_f | f | \phi_i \rangle \}$ . The result for  $L_{(J_1)L}(\mathbf{Q})$  is

$$\begin{aligned} L_{(J_1)L}(\mathbf{Q}) &= i^{l_\alpha + l_\beta + J + 1} (-)^{J-L} \frac{\hat{j}}{\sqrt{4\pi}} \\ &\times \left\{ \pm U(l_\beta 1 l_\alpha J; l_\beta + 1 L) \begin{pmatrix} l_\alpha & J & l_\beta + 1 \\ 0 & 0 & 0 \end{pmatrix} \hat{l}_\alpha (l_\beta + 1)^{1/2} \right. \\ &\times \left\langle R_\alpha \| f \left| \left( \frac{d}{dr} - \frac{l_\beta}{r} \right) R_\beta \right. \right\rangle \mp U(l_\beta 1 l_\alpha J; l_\beta - 1 L) \\ &\times \begin{pmatrix} l_\alpha & J & l_\beta - 1 \\ 0 & 0 & 0 \end{pmatrix} \hat{l}_\alpha (l_\beta)^{1/2} \left\langle R_\alpha \| f \left| \left( \frac{d}{dr} + \frac{l_\beta + 1}{r} \right) R_\beta \right. \right\rangle \\ &- (-)^{J+1-L} U(l_\beta J l_\alpha 1; l_\alpha + 1 L) \begin{pmatrix} l_\alpha + 1 & J & l_\beta \\ 0 & 0 & 0 \end{pmatrix} \hat{l}_\beta (l_\alpha + 1)^{1/2} \\ &\times \left\langle \left( \frac{d}{dr} - \frac{l_\alpha}{r} \right) R_\alpha \middle| f \middle| R_\beta \right\rangle + (-)^{J+1-L} U(l_\beta J l_\alpha 1; l_\alpha - 1 L) \\ &\times \begin{pmatrix} l_\alpha - 1 & J & l_\beta \\ 0 & 0 & 0 \end{pmatrix} \hat{l}_\beta (l_\alpha)^{1/2} \left\langle \left( \frac{d}{dr} + \frac{l_\alpha + 1}{r} \right) R_\alpha \middle| f \middle| R_\beta \right\rangle \right\}, \end{aligned} \quad (\text{A33})$$

where the upper sign is used for  $\mathbf{Q} = \mathbf{p}$  and the lower sign for  $\mathbf{Q} = \mathbf{q}$ . The  $U$  coefficient is a recoupling of three angular momenta and is related to a  $6j$ -symbol by

$$U(abcd; ef) = (-)^{a+b+c+d} \hat{e} \hat{f} \begin{Bmatrix} a & b & e \\ d & c & f \end{Bmatrix}. \quad (\text{A34})$$

Finally, we are ready to write down the specific reduced matrix elements for all the different hadronic components of the weak interaction, Eqs. (A20) to (A23) as follows:

- *Spacelike Axial Current*

$$\begin{aligned} & \left\langle \left( l_\alpha \frac{1}{2} \right) j_\alpha \middle\| f(r) \mathbf{A} \cdot i^L \mathbf{Y}_{KL} \middle\| \left( l_\beta \frac{1}{2} \right) j_\beta \right\rangle \\ &= (-)^{K-L} \left[ -g_A + f_T \left( W_0 + \frac{6}{5} \frac{\alpha Z}{R} \right) \right] A_{(L_1)K} L_L S_1. \end{aligned} \quad (\text{A35})$$

- *Timelike Vector Current*

$$\begin{aligned} & \left\langle \left( l_\alpha \frac{1}{2} \right) j_\alpha \middle\| f(r) V_0 i^K Y_K \middle\| \left( l_\beta \frac{1}{2} \right) j_\beta \right\rangle \\ &= \left[ g_V + f_S \left( W_0 + \frac{6}{5} \frac{\alpha Z}{R} \right) \right] A_{(K_0)K} L_K S_0. \end{aligned} \quad (\text{A36})$$

In Eqs. (A35) and (A36), for simplicity we have replaced the function  $V(r)$  [see Eqs. (A20) and (A23)] with the potential due to a uniform charge distribution for small  $r$ , with  $r^2$  replaced by its expectation value. In our computations we actually included the function  $V(r)$  in the integrand of the appropriate radial integral.

<sup>8</sup>If one-body density matrix elements are imported from a shell-model calculation into this  $\beta$ -decay environment, then it is important that these OBDME be computed with similar  $i^l$  phases included in the definition of single-particle wave functions.

• *Spacelike Vector Current*

$$\begin{aligned} & \left\langle \left( l_\alpha \frac{1}{2} \right) j_\alpha \left\| f(r) \mathbf{V} \cdot i^L \mathbf{Y}_{KL} \left\| \left( l_\beta \frac{1}{2} \right) j_\beta \right\rangle \right. \\ &= (-)^{K-L} \left\{ \frac{g_V}{2M_N} A_{(K0)K} L_{(L1)K}(\mathbf{p}) S_0 \right. \\ & \quad - f_S A_{(K0)K} L_{(L1)K}(\mathbf{q}) S_0 - \sqrt{2} \sum_J U(11KL; 1J) \\ & \quad \left. \times \left( \frac{g_V}{2M_N} + f_M \right) A_{(J1)K} L_{(L1)J}(\mathbf{q}) S_1 \right\}. \end{aligned} \quad (\text{A37})$$

• *Timelike Axial Current*

$$\begin{aligned} & \left\langle \left( l_\alpha \frac{1}{2} \right) j_\alpha \left\| f(r) A_0 i^K Y_K \left\| \left( l_\beta \frac{1}{2} \right) j_\beta \right\rangle \right. \\ &= \sum_J (-)^{J-K} \frac{\hat{J}}{\hat{K}} \left\{ -\frac{g_A}{2M_N} A_{(J1)K} L_{(K1)J}(\mathbf{p}) S_1 \right. \\ & \quad \left. - f_T A_{(J1)K} L_{(K1)J}(\mathbf{q}) S_1 \right\}. \end{aligned} \quad (\text{A38})$$

**Numerical results**

The key ingredient for the computation of exact electron wave functions in  $\beta$  decay is the charge-density distribution of the daughter nucleus. There are various parameterizations available in the literature, of which the following are the most common:

- *Two-parameter Fermi distribution (2pF)*. This charge density distribution,

$$\rho(r) = \frac{\rho_0}{1 + \exp\{(r - c)/a\}}, \quad (\text{A39})$$

has two parameters,  $c$  and  $a$ , other than its normalization.

- *Three-parameter Fermi distribution (3pF)*. This is an extension of the two-parameter model, which introduces a dimensionless “wine-bottle” parameter,  $w$ , that impacts on the small- $r$  behavior of the density distribution. The functional form is as follows:

$$\rho(r) = \frac{\rho_0(1 + wr^2/c^2)}{1 + \exp\{(r - c)/a\}}. \quad (\text{A40})$$

- *Three-parameter Gaussian distribution (3pG)*. This is an alternative three-parameter model with a Gaussian rather than a Fermi distribution:

$$\rho(r) = \frac{\rho_0(1 + wr^2/c^2)}{1 + \exp\{(r^2 - c^2)/a^2\}}. \quad (\text{A41})$$

- *Harmonic-oscillator distribution (HO)*. In light  $p$ -shell nuclei, where only  $s$  and  $p$  orbitals are occupied, a density distribution can be constructed from the harmonic oscillator radial wavefunctions. Its form is as follows:

$$\rho(r) = \rho_0(1 + \alpha r^2/b^2) \exp(-r^2/b^2), \quad (\text{A42})$$

where  $b$  is the harmonic oscillator length parameter and  $\alpha$  is related to the number of  $p$ -shell protons,  $\alpha = (Z - 2)/3$ . However, in practice both  $b$  and  $\alpha$  are treated as free parameters and adjusted to fit the elastic-electron scattering data.

TABLE X. Charge-density distributions from elastic electron-scattering data [180]. The radius parameter, in some cases, has been adjusted to reproduce  $\langle r^2 \rangle^{1/2}$ . Parameters  $c$  and  $a$  are in femtometers units; parameter  $w$  is dimensionless.

Daughter nucleus	$\langle r^2 \rangle^{1/2}$ fm	Model <sup>a</sup>	$c$	$a$	$w$	$\Delta f^b$ %
$\langle T_z = -1 \rangle$						
<sup>10</sup> B	2.45(10)	HO	1.709 <sup>c</sup>	0.837 <sup>d</sup>		0.001
<sup>14</sup> N	2.52(2)	3pF	2.572	0.5052	-0.180	0.000
<sup>18</sup> F	2.90(3)	2pF	2.574	0.567		0.001
<sup>22</sup> Na	2.95(5)	2pF	2.750	0.549		0.001
<sup>26</sup> Al	3.03(2)	2pF	2.791	0.569		0.001
<sup>30</sup> P	3.18(3)	3pF	3.350	0.582	-0.173	0.002
<sup>34</sup> Cl	3.39(2)	3pF	3.479	0.599	-0.100	0.001
<sup>38</sup> K	3.41(4)	3pF	3.738	0.585	-0.201	0.004
<sup>42</sup> Sc	3.50(5)	3pF	3.794	0.586	-0.161	0.004
$\langle T_z = 0 \rangle$						
<sup>26</sup> Mg	3.06(5)	2pF	3.049	0.523		0.002
<sup>34</sup> S	3.29(1)	3pG	2.810	2.191	0.160	0.001
<sup>38</sup> Ar	3.36(5)	2pF	3.590	0.507		0.004
<sup>42</sup> Ca	3.48(3)	3pF	3.765	0.586	-0.161	0.002
<sup>46</sup> Ti	3.61(3)	2pF	3.711	0.588		0.003
<sup>50</sup> Cr	3.66(4)	2pF	3.868	0.566		0.004
<sup>54</sup> Fe	3.69(2)	3pG	3.541	2.270	0.403	0.003
<sup>62</sup> Zn	3.90(2)	3pG	3.570	2.465	0.342	0.005
<sup>66</sup> Ge	4.04(4)	2pF	4.398	0.585		0.011
<sup>70</sup> Se	4.07(5)	2pF	4.442	0.585		0.011
<sup>74</sup> Kr	4.10(5)	2pF	4.489	0.585		0.013

<sup>a</sup>See Eqs. (A39) to (A42).

<sup>b</sup>Percentage uncertainty in  $f$  because the uncertainty in  $\langle r^2 \rangle^{1/2}$ .

<sup>c</sup>This is parameter  $b$  of Eq. (A42) in femtometers.

<sup>d</sup>This is the dimensionless parameter  $\alpha$  of Eq. (A42).

These model distributions typically contain two or three parameters. Where possible, the parameters are determined from experimental data on elastic electron scattering, because the measured electron-scattering form factors are just the Fourier transforms of the charge-density distributions. A compilation of charge-density distributions determined from electron scattering is given by De Vries *et al.* [180]. We have assessed these data and selected for each daughter nucleus what we believe to be the “best” value of the rms radius,  $\langle r^2 \rangle^{1/2}$ , and its probable error. In cases where data are not available on the isotope of interest, we have examined the nearest isotope that is available and applied a modest isotope shift to its value of  $\langle r^2 \rangle^{1/2}$ . Our final selected values are listed in Table X. We also list the percentage uncertainty in the exact value of  $f$  solely because of the uncertainty in  $\langle r^2 \rangle^{1/2}$ . Clearly, the uncertainty in the charge-density distribution is not a factor in the determination of  $f$  to 0.1% accuracy.

Before the final evaluation of the statistical rate function,  $f$ , there are two further corrections to consider: screening and recoil. To accommodate these corrections and to remove—as



TABLE XI. Comparison of statistical rate functions  $f_{\text{stat}}$  and  $f_{\text{approx}}$  and the exact value  $f$ .

Parent	$Q_{EC}$ (keV)	$W_0$	$f_{\text{stat}}$	$f$	$\Delta f_{\text{stat}}^a$ %
$\langle T_z = -1 \rangle$					
$^{10}\text{C}$	1907.9	2.7336	2.29484	2.30089	-0.26
$^{14}\text{O}$	2831.2	4.5405	42.6380	42.7719	-0.31
$^{18}\text{Ne}$	3402.0	5.6575	133.867	134.484	-0.46
$^{22}\text{Mg}$	4124.6	7.0716	415.827	418.441	-0.62
$^{26}\text{Si}$	4836.9	8.4656	1014.75	1023.28	-0.83
$^{30}\text{S}$	5459.5	9.6840	1945.49	1967.05	-1.10
$^{34}\text{Ar}$	6062.8	10.8647	3366.22	3414.21	-1.41
$^{38}\text{Ca}$	6614.2	11.9437	5247.54	5338.46	-1.70
$^{42}\text{Ti}$	7000.9	12.7004	6904.47	7042.83	-1.96
$\langle T_z = 0 \rangle$					
$^{26m}\text{Al}$	4232.6	7.2829	474.718	478.203	-0.73
$^{34}\text{Cl}$	5491.8	9.7472	1971.95	1996.39	-1.22
$^{38m}\text{K}$	6044.4	10.8286	3248.45	3298.10	-1.51
$^{42}\text{Sc}$	6425.6	11.5747	4391.33	4470.03	-1.76
$^{46}\text{V}$	7050.7	12.7979	7044.04	7199.96	-2.17
$^{50}\text{Mn}$	7632.4	13.9363	10455.9	10731.2	-2.57
$^{54}\text{Co}$	8242.6	15.1304	15280.6	15749.3	-2.98
$^{62}\text{Ga}$	9171.0	16.9472	25187.9	26247.6	-4.04
$^{66}\text{As}$	9550.0	17.6889	30146.4	31613.7	-4.64
$^{70}\text{Br}$	9970.0	18.5108	36605.4	38602.2	-5.17
$^{74}\text{Rb}$	10416.8	19.3852	44612.8	47285.0	-5.65

$$^a \Delta f_{\text{stat}} = 100 * (f_{\text{stat}} - f)/f.$$

is customary—the leading matrix element from the definition of  $f$ , we rewrite  $f$  as follows:

$$f = \xi R(W_0) \int_1^{W_0} p W (W_0 - W)^2 F_0 L_0 C(Z, W) Q(Z, W) dW. \quad (\text{A43})$$

Comparison with Eqs. (A1) and (A3) reveals three new factors,  $Q(Z, W)$ ,  $R(W_0)$ , and  $\xi$ . We deal with them in that order.

The calculation of the Fermi function presented so far makes no allowance for the screening of the atomic electrons. Rose [181] was the first to find a simple analytic prescription to obtain the Fermi function for a screened field from the Fermi function for the corresponding unscreened field. That prescription is to incorporate a correction factor into the integrand for  $f$  as follows:

$$Q(Z, W) = \frac{\tilde{p} \tilde{W} F(Z, \tilde{W})}{p W F(Z, W)}, \quad (\text{A44})$$

where  $\tilde{W} = W - V_0$ ,  $\tilde{p} = (\tilde{W}^2 - 1)^{1/2}$  and  $V_0 = N(\tilde{Z}) \alpha^2 \tilde{Z}^{4/3}$ , with  $\tilde{Z}$  being the electronic charge of the parent

atom and  $N(\tilde{Z})$  being a weak function of  $\tilde{Z}$ , which varies from  $N = 1.42$  at  $\tilde{Z} = 8$  to  $N = 1.56$  at  $\tilde{Z} = 29$  (see Matese and Johnson [182]). Because the factor  $Q(Z, W)$  yields a correction to  $f$  of order  $\sim 0.2\%$ , we need only Rose's screening correction to be accurate to within 50% of its central value to assure us an accuracy in  $f$  of 0.1%. Matese and Johnson [182] have tested the Rose formula by comparing it with numerical solutions of the Dirac equation for a self-consistent Hartree-Fock-Slater potential. They conclude that  $Q(Z, W)$  has an accuracy of four significant figures or better for all energies except the very lowest in positron emitters. Because we integrate  $Q(Z, W)$  over the whole  $\beta$  spectrum, which actually deemphasizes the lowest positron energies, we conclude that the Rose formula has far more than sufficient accuracy for our purpose.

The second new factor in Eq. (A43) is  $R(W_0)$ , which is the correction for recoil: it recognizes that the daughter nucleus is not at rest but has a small amount of recoiling kinetic energy. As a result, the leptons' maximum energy is actually slightly less than  $W_0$ . The recoil correction [167] is given by

$$R(W_0) = 1 - \frac{3W_0}{2M_A}, \quad (\text{A45})$$

where  $M_A$  is the average of the initial and final nuclear masses. For use in eq. (A45),  $M_A$  must, like  $W_0$ , be expressed in electron rest-mass units. The resulting correction is very small, being of the order 0.02% for the superallowed  $\beta$  decays from  $A = 10$  to  $A = 74$ .

Last, for allowed transitions it is customary to remove the leading matrix element from the definition of  $f$ . Thus, we have introduced  $\xi$  in Eq. (A43), where  $\xi = 1/|M_F|^2$  for superallowed Fermi transitions,  $M_F$  being the Fermi matrix element. For pure Gamow-Teller transitions  $\xi = 1/|M_{GT}|^2$ , with  $M_{GT}$  being the Gamow-Teller matrix element.

In Table XI we list both the values of  $f_{\text{stat}}$ , Eq. (A2), and the exact values of  $f$  for cases of interest in superallowed  $\beta$  decay. The relevant  $Q_{EC}$  value is listed as well. For the exact calculations, we imported one-body density matrix elements, OBDME, from a shell-model code. For each case we performed several shell-model calculations for various sets of effective interactions and model spaces. We used, in fact, the same wave functions that we used [159] to compute the nuclear-structure corrections  $\delta_C$  and  $\delta_{NS}$ . Thus our  $f$  calculations can be considered to be entirely consistent with the calculation of the nuclear-structure-dependent corrections. The  $f$  calculation, however, is not very sensitive to the shell-model inputs. In light nuclei, different shell-model OBDME gave changes in  $f$  at the 0.01% level, increasing to around 0.1% in  $A = 74$ , our heaviest-mass case. Where we have more than one shell-model calculation for a given nucleus, we have averaged the  $f$  values for the entry in Table XI.

- [1] I. S. Towner and J. C. Hardy, J. Phys. G: Nucl. Part. Phys. **29**, 197 (2003).  
 [2] A. Sher *et al.*, Phys. Rev. Lett. **91**, 261802 (2003).  
 [3] T. Alexopoulos *et al.*, Phys. Rev. Lett. **93**, 181802 (2004).

- [4] I. S. Towner and J. C. Hardy, Nucl. Phys. **A205**, 33 (1973).  
 [5] J. C. Hardy and I. S. Towner, Nucl. Phys. **A254**, 221 (1975).  
 [6] V. T. Koslowsky, E. Hagberg, J. C. Hardy, H. Schmeing, R. E. Azuma, and I. S. Towner, in Proceedings of the

- 7th International Conference on Atomic Masses and Fundamental Constants, Darmstadt-Seeheim, edited by O. Klepper (T. H. Darmstadt, 1984) p. 572.
- [7] J. C. Hardy, I. S. Towner, V. T. Koslowsky, E. Hagberg, and H. Schmeing, *Nucl. Phys.* **A509**, 429 (1990).
- [8] E. G. Adelberger, M. M. Hindi, C. D. Hoyle, H. E. Swanson, R. D. Von Lintig, and W. C. Haxton, *Phys. Rev. C* **27**, 2833 (1983); this reference replaces the result reported in E. G. Adelberger, C. D. Hoyle, H. E. Swanson, and R. D. Von Lintig, *Phys. Rev. Lett.* **46**, 695 (1981).
- [9] F. Ajzenberg-Selove, *Nucl. Phys.* **A490**, 1 (1988).
- [10] F. Ajzenberg-Selove, *Nucl. Phys.* **A523**, 1 (1991).
- [11] A. M. Aldridge, K. W. Kemper, and H. S. Plendl, *Phys. Lett.* **B30**, 165 (1969).
- [12] D. E. Alburger and D. H. Wilkinson, *Phys. Lett.* **B32**, 190 (1970).
- [13] D. E. Alburger, *Phys. Rev. C* **5**, 274 (1972).
- [14] D. E. Alburger and F. P. Calaprice, *Phys. Rev. C* **12**, 1690 (1975).
- [15] D. E. Alburger and D. H. Wilkinson, *Phys. Rev. C* **15**, 2174 (1977); this reference replaces the  $^{46}\text{V}$  half-life from Ref. [150].
- [16] D. E. Alburger, *Phys. Rev. C* **18**, 1875 (1978).
- [17] P. F. A. Alkemade, C. Alderliesten, P. De Wit, and C. Van der Leun, *Nucl. Instr. Methods* **197**, 383 (1982).
- [18] A. Anttila, M. Bister, and E. Arminen, *Z. Phys.* **234**, 455 (1970).
- [19] E. Aslanides, F. Jundt, and A. Gallmann, *Nucl. Phys.* **A152**, 251 (1970).
- [20] G. Audi, A. H. Wapstra, and C. Thibault, *Nucl. Phys.* **A729**, 337 (2003).
- [21] G. Azuelos, J. E. Crawford, and J. E. Kitching, *Phys. Rev. C* **9**, 1213 (1974).
- [22] G. Azuelos and J. E. Kitching, *Phys. Rev. C* **12**, 563 (1975).
- [23] R. K. Barden, C. A. Barnes, W. A. Fowler, and P. G. Seeger, *Phys. Rev.* **127**, 583 (1962).
- [24] F. J. Bartis, *Phys. Rev.* **132**, 1763 (1963).
- [25] P. H. Barker, N. Drysdale, and W. R. Phillips, *Proc. Phys. Soc.* **91**, 587 (1967).
- [26] P. H. Barker, C. J. Scofield, R. J. Petty, J. M. Freeman, S. D. Hoath, W. E. Burcham, and G. T. A. Squier, *Nucl. Phys.* **A275**, 37 (1977); the same result also appears in G. T. A. Squier, W. E. Burcham, S. D. Hoath, J. M. Freeman, P. H. Barker, and R. J. Petty, *Phys. Lett.* **B65**, 122 (1976).
- [27] P. H. Barker and J. A. Nolen, in *Proceedings of the International Conference on Nuclear Structure, Tokyo, Japan, 1977*, edited by the Organizing Committee (International Academic Printing Co. Ltd., Tokyo, 1977).
- [28] P. H. Barker, R. E. White, H. Naylor, and N. S. Wyatt, *Nucl. Phys.* **A279**, 199 (1977).
- [29] P. H. Barker and R. E. White, *Phys. Rev. C* **29**, 1530 (1984).
- [30] P. H. Barker and S. M. Ferguson, *Phys. Rev. C* **38**, 1936 (1988).
- [31] S. C. Baker, M. J. Brown, and P. H. Barker, *Phys. Rev. C* **40**, 940 (1989).
- [32] P. H. Barker and G. D. Leonard, *Phys. Rev. C* **41**, 246 (1990).
- [33] P. H. Barker and P. A. Amundsen, *Phys. Rev. C* **58**, 2571 (1998); this reference updates the  $^{10}\text{C}$   $Q_{EC}$  value from [62]; its value for the  $^{14}\text{O}$   $Q_{EC}$  value was later withdrawn in Ref. [168].
- [34] P. H. Barker and M. S. Wu, *Phys. Rev. C* **62**, 054302 (2000).
- [35] G. C. Ball, S. Bishop, J. A. Behr, G. C. Boisvert, P. Bricault, J. Cerny, J. M. D'Auria, M. Dombsky, J. C. Hardy, V. Iacob, J. R. Leslie, T. Lindner, J. A. Macdonald, H.-B. Mak, D. M. Moltz, J. Powell, G. Savard, and I. S. Towner, *Phys. Rev. Lett.* **86**, 1454 (2001).
- [36] P. H. Barker, I. C. Barnett, G. J. Baxter, and A. P. Byrne, *Phys. Rev. C* **70**, 024302 (2004).
- [37] G. C. Ball *et al.* (to be published).
- [38] E. Beck and H. Daniel, *Z. Phys.* **216**, 229 (1968).
- [39] J. A. Becker, R. A. Chalmers, B. A. Watson, and D. H. Wilkinson, *Nucl. Instr. Methods* **155**, 211 (1978).
- [40] F. J. Bergmeister, K. P. Lieb, K. Pampus, and M. Uhrmacher, *Z. Phys. A* **320**, 693 (1985).
- [41] S. Bishop *et al.*, *Phys. Rev. Lett.* **90**, 162501 (2003).
- [42] B. Blank, *Eur. Phys. J. A* **15**, 121 (2002).
- [43] B. Blank, G. Savard, J. Doring, A. Blazhev, G. Canchel, M. Chartier, D. Henderson, Z. Janas, R. Kirchner, I. Mukha, E. Roeckl, K. Schmidt, and J. Zylicz, *Phys. Rev. C* **69**, 015502 (2004).
- [44] K. Blaum, G. Audi, D. Beck, G. Bollen, C. Guenaut, P. Delahaye, F. Herfurth, A. Kellerbauer, H.-J. Kluge, D. Lunney, D. Rodriguez, S. Schwarz, L. Schweikhard, C. Weber, and C. Yazidjian, *Nucl. Phys.* **A746**, 305c (2004).
- [45] R. O. Bondelid and J. W. Butler, *Nucl. Phys.* **53**, 618 (1964).
- [46] S. A. Brindhaban and P. H. Barker, *Phys. Rev. C* **49**, 2401 (1994); reference replaces earlier conference proceedings from the same laboratory.
- [47] J. W. Butler and R. O. Bondelid, *Phys. Rev.* **121**, 1770 (1961).
- [48] W. E. Burcham and G. T. A. Squier, *Nucl. Instr. Methods* **164**, 533 (1979).
- [49] R. H. Burch, C. A. Gagliardi, and R. E. Tribble, *Phys. Rev. C* **38**, 1365 (1988).
- [50] G. Canchel, B. Blank, M. Chartier, F. Delalee, P. Dendooven, C. Dossat, J. Giovinozzo, J. Huikari, A. S. Lalleman, M. J. Lopez Jimenez, V. Madec, J. L. Pedroza, H. Penttila, and J. C. Thomas, *Eur. Phys. J. A* **23**, 409 (2005).
- [51] N. M. Chaudri, *Fizika* **16**, 297 (1984).
- [52] G. J. Clark, J. M. Freeman, D. C. Robinson, J. S. Ryder, W. E. Burcham, and G. T. A. Squier, *Nucl. Phys.* **A215**, 429 (1973); this reference replaces the half-life value in G. J. Clark, J. M. Freeman, D. C. Robinson, J. S. Ryder, W. E. Burcham, and G. T. A. Squier, *Phys. Lett.* **B35**, 503 (1971).
- [53] C. N. Davids, C. A. Gagliardi, M. J. Murphy, and E. B. Norman, *Phys. Rev. C* **19**, 1463 (1979).
- [54] C. N. Davids, in *Atomic Masses and Fundamental Constants 6*, edited by J. A. Nolen and W. Benenson (Plenum, New York, 1980), p. 419.
- [55] W. W. Daehnick and R. D. Rosa, *Phys. Rev. C* **31**, 1499 (1985).
- [56] P. De Wit and C. Van der Leun, *Phys. Lett.* **B30**, 639 (1969).
- [57] R. M. DelVecchio and W. W. Daehnick, *Phys. Rev. C* **17**, 1809 (1978).
- [58] M. A. van Driel, H. Klijnman, G. A. P. Engelbertink, H. H. Eggehuysen, and J. A. J. Hermans, *Nucl. Phys.* **A240**, 98 (1975).
- [59] L. G. Earwaker, J. G. Jenkin, and E. W. Titterton, *Nature* **195**, 271 (1962).
- [60] P. M. Endt, *Nucl. Phys.* **A521**, 1 (1990).
- [61] P. M. Endt, *Nucl. Phys.* **A633**, 1 (1998).
- [62] G. Frick, A. Gallmann, D. E. Alburger, D. H. Wilkinson, and J. P. Coffin, *Phys. Rev.* **132**, 2169 (1963).
- [63] J. M. Freeman, G. Murray, and W. E. Burcham, *Phys. Lett.* **17**, 317 (1965).

- [64] J. M. Freeman, J. G. Jenkin, G. Murray, D. C. Robinson, and W. E. Burcham, Nucl. Phys. **A132**, 593 (1969); this reference replaces the half-life values in J. M. Freeman, J. H. Montague, G. Murray, R. E. White, and W. E. Burcham, Nucl. Phys. **69**, 433 (1965) and J. M. Freeman, J. G. Jenkin, G. Murray, and W. E. Burcham, Phys. Rev. Lett. **16**, 959 (1966).
- [65] J. M. Freeman, R. J. Petty, S. D. Hoath, G. T. A. Squier, and W. E. Burcham, Phys. Lett. **B53**, 439 (1975).
- [66] B. K. Fujikawa, S. J. Asztalos, R. M. Clark, M. A. Deleplanque-Stephens, P. Fallon, S. J. Freeman, J. P. Greene, I.-Y. Lee, L. J. Lising, A. O. Macchiavelli, R. W. MacLeod, J. C. Reich, M. A. Rowe, S.-Q. Shang, F. S. Stephens, and E. G. Wasserman, Phys. Lett. **B449**, 6 (1999).
- [67] A. Gallmann, E. Aslanides, F. Jundt, and E. Jacobs, Phys. Rev. **186**, 1160 (1969).
- [68] M. Gaelens, J. Andrzejewski, J. Camps, P. Decroock, M. Huyse, K. Kruglov, W. F. Mueller, A. Piechaczek, N. Severijns, J. Szerypo, G. Vancraeynest, P. Van Duppen, and J. Wauters, Eur. Phys. J. A **11**, 413 (2001).
- [69] D. R. Goosman and D. E. Alburger, Phys. Rev. C **5**, 1893 (1972); the branching-ratio upper limit set in this reference is considered to replace the much higher value reported by D. R. Brown, S. M. Ferguson, and D. H. Wilkinson, Nucl. Phys. **A135**, 159 (1969).
- [70] H. J. Gils, D. Flothmann, R. Loehken, and W. Wiesner, Nucl. Instr. Methods **105**, 179 (1972).
- [71] G. I. Harris and A. K. Hyder, Phys. Rev. **157**, 958 (1967).
- [72] M. Hagen, K. H. Maier, and R. Michaelsen, Phys. Lett. **B26**, 432 (1968).
- [73] J. C. Hardy and D. E. Alburger, Phys. Lett. **B42**, 341 (1972).
- [74] J. C. Hardy, H. Schmeing, J. S. Geiger, and R. L. Graham, Nucl. Phys. **A223**, 157 (1974); this reference replaces results in J. C. Hardy, H. Schmeing, J. S. Geiger, R. L. Graham, and I. S. Towner, Phys. Rev. Lett. **29**, 1027 (1972).
- [75] J. C. Hardy, H. R. Andrews, J. S. Geiger, R. L. Graham, J. A. Macdonald, and H. Schmeing, Phys. Rev. Lett. **33**, 1647 (1974).
- [76] J. C. Hardy, H. Schmeing, W. Benenson, G. M. Crawley, E. Kashy, and H. Nann, Phys. Rev. C **9**, 252 (1974).
- [77] J. C. Hardy, G. C. Ball, J. S. Geiger, R. L. Graham, J. A. Macdonald, and H. Schmeing, Phys. Rev. Lett. **33**, 320 (1974); the value for the  $^{46}\text{V}$   $Q_{EC}$  value from this reference was later withdrawn by J. C. Hardy and I. S. Towner, in *Atomic Masses and Fundamental Constants 5*, edited by J. H. Sanders and A. H. Wapstra (Plenum, New York, 1976), p. 66.
- [78] J. C. Hardy, H. Schmeing, J. S. Geiger, and R. L. Graham, Nucl. Phys. **A246**, 61 (1975); this reference replaces results in J. C. Hardy, H. Schmeing, J. S. Geiger, R. L. Graham, and I. S. Towner, Phys. Rev. Lett. **29**, 1027 (1972).
- [79] E. Hagberg, V. T. Koslowsky, J. C. Hardy, I. S. Towner, J. G. Hykawy, G. Savard, and T. Shinozuka, Phys. Rev. Lett. **73**, 396 (1994); uncertainties on the Gamow-Teller decays observed from  $^{46}\text{V}$  and  $^{50}\text{Mn}$  did not appear in this reference but have been derived from the original data and added here.
- [80] P. D. Harty, N. S. Bowden, P. H. Barker, and P. A. Amundsen, Phys. Rev. C **58**, 821 (1998).
- [81] J. C. Hardy and I. S. Towner, Phys. Rev. Lett. **88**, 252501 (2002).
- [82] J. C. Hardy, V. E. Iacob, M. Sanchez-Vega, R. G. Neilson, A. Azhari, C. A. Gagliardi, V. E. Mayes, X. Tang, L. Trache, and R. E. Tribble, Phys. Rev. Lett. **91**, 082501 (2003).
- [83] D. L. Hendrie and J. B. Gerhart, Phys. Rev. **121**, 846 (1961).
- [84] A. M. Hernandez and W. W. Daehnick, Phys. Rev. C **24**, 2235 (1981).
- [85] A. M. Hernandez and W. W. Daehnick, Phys. Rev. C **25**, 2957 (1982).
- [86] F. Herfurth *et al.*, Eur. Phys. J. A **15**, 17 (2002).
- [87] I. Hofmann, Acta Phys. Aust. **18**, 309 (1964).
- [88] S. D. Hoath, R. J. Petty, J. M. Freeman, G. T. A. Squier, and W. E. Burcham, Phys. Lett. **B51**, 345 (1974).
- [89] P. Hungerford and H. H. Schmidt, Nucl. Instr. Methods **192**, 609 (1982).
- [90] B. C. Hyman, V. E. Iacob, A. Azhari, C. A. Gagliardi, J. C. Hardy, V. E. Mayes, R. G. Neilson, M. Sanchez-Vega, X. Tang, L. Trache, and R. E. Tribble, Phys. Rev. C **68**, 015501 (2003).
- [91] V. E. Iacob, E. Mayes, J. C. Hardy, R. G. Neilson, M. Sanchez-Vega, A. Azhari, C. A. Gagliardi, L. Trache, and R. E. Tribble, Bul. Am. Phys. Soc. **48**, 34 (2003).
- [92] P. D. Ingalls, J. C. Overley, and H. S. Wilson, Nucl. Phys. **A293**, 117 (1977).
- [93] M. A. Islam, T. J. Kennett, S. A. Kerr, and W. V. Prestwich, Can. J. Phys. **58**, 168 (1980).
- [94] J. Jaenecke, Z. Naturf. **15a**, 593 (1960).
- [95] A. N. James, F. J. Sharpey-Schafer, A. M. Al-Naser, A. H. Behbehani, C. J. Lister, P. J. Nolan, P. H. Barker, and W. E. Burcham, J. Phys. G **4**, 579 (1978).
- [96] R. W. Kavanagh, A. Gallmann, E. Aslanides, F. Jundt, and E. Jacobs, Phys. Rev. **175**, 1426 (1968).
- [97] R. W. Kavanagh, Nucl. Phys. **A129**, 172 (1969).
- [98] A. Kellerbauer, G. Audi, D. Beck, K. Blaum, G. Bollen, B. A. Brown, P. Delahaye, C. Guenaut, F. Herfurth, H.-J. Kluge, D. Lunney, S. Schwarz, L. Schweikhard, and C. Yazidjian, Phys. Rev. Lett. **93**, 072502 (2004); the mass excesses quoted in this reference for  $^{74}\text{Rb}$  and  $^{74}\text{Kr}$  include the data reported earlier in [86] and so replace the values quoted there.
- [99] S. W. Kikstra, C. van der Leun, S. Raman, E. T. Jurney, and I. S. Towner, Nucl. Phys. **A496**, 429 (1989).
- [100] S. W. Kikstra, Z. Guo, C. Van der Leun, P. M. Endt, S. Raman, Walkiewicz, J. W. Starner, E. T. Jurney, and I. S. Towner, Nucl. Phys. **A529**, 39 (1991).
- [101] V. T. Koslowsky, E. Hagberg, J. C. Hardy, R. E. Azuma, E. T. H. Clifford, H. C. Evans, H. Schmeing, U. J. Schrewe, and K. S. Sharma, Nucl. Phys. **A405**, 29 (1983).
- [102] V. T. Koslowsky, J. C. Hardy, E. Hagberg, R. E. Azuma, G. C. Ball, E. T. H. Clifford, W. G. Davies, H. Schmeing, U. J. Schrewe, and K. S. Sharma, Nucl. Phys. **A472**, 419 (1987); the  $^{14}\text{O}$ - $^{26}\text{Al}^m$   $Q_{EC}$  value-difference result reported in this reference replaces an earlier value given in V. T. Koslowsky, J. C. Hardy, R. E. Azuma, G. C. Ball, E. T. H. Clifford, W. G. Davies, E. Hagberg, H. Schmeing, U. J. Schrewe, and K. S. Sharma, Phys. Lett. **B119**, 57 (1982).
- [103] V. T. Koslowsky, E. Hagberg, J. C. Hardy, G. Savard, H. Schmeing, K. S. Sharma, and X. J. Sun, Nucl. Instr. Methods in Phys. Res. A **401**, 289 (1997).
- [104] V. T. Koslowsky, E. Hagberg, J. C. Hardy, H. Schmeing, and I. S. Towner, Nucl. Phys. **A624**, 293 (1997).
- [105] M. A. Kroupa, S. J. Freeman, P. H. Barker, and S. M. Ferguson, Nucl. Instr. Methods in Phys. Res. A **310**, 649 (1991).
- [106] S. Lin, S. A. Brindhaban, and P. H. Barker, Phys. Rev. C **49**, 3098 (1994).

- [107] P. V. Magnus, E. G. Adelberger, and A. Garcia, *Phys. Rev. C* **49**, R1755 (1994).
- [108] W. R. McMurray, P. Van der Merwe, and I. J. Van Heerden, *Nucl. Phys.* **A92**, 401 (1967).
- [109] R. G. Miller and R. W. Kavanagh, *Nucl. Phys.* **A94**, 261 (1967).
- [110] C. E. Moss, C. Detraz, and C. S. Zaidins, *Nucl. Phys.* **A174**, 408 (1971).
- [111] M. Mukherjee, A. Kellerbauer, D. Beck, K. Blaum, G. Bollen, F. Carrel, P. Delahaye, J. Dilling, S. George, C. Guenaut, F. Herfurth, A. Herlert, H.-J. Kluge, U. Koster, D. Lunney, S. Schwarz, L. Schweikhard, and C. Yazidjian, *Phys. Rev. Lett.* **93**, 150801 (2004).
- [112] Y. Nagai, K. Kunihiro, T. Toriyama, S. Harada, Y. Torii, A. Yoshida, T. Nomura, J. Tanaka, and T. Shinozuka, *Phys. Rev. C* **43**, 9(R) (1991).
- [113] F. M. Nichols, N. Lawley, I. G. Main, M. F. Thomas, and P. J. Twin, *Nucl. Phys.* **A124**, 97 (1969).
- [114] J. A. Nolen, G. Hamilton, E. Kashy, and I. D. Proctor, *Nucl. Instr. Methods* **115**, 189 (1974).
- [115] M. Oinonen *et al.*, *Phys. Lett.* **B511**, 145 (2001).
- [116] R. A. Padock, *Phys. Rev. C* **5**, 485 (1972).
- [117] A. Piechaczek, E. F. Zganjar, G. C. Ball, P. Bricault, J. M. D'Auria, J. C. Hardy, D. F. Hodgson, V. Iacob, P. Klages, W. D. Kulp, J. R. Leslie, M. Lipoglavsek, J. A. Macdonald, H.-B. Mak, D. M. Moltz, G. Savard, J. von Schwarzenberg, C. E. Svensson, I. S. Towner, and J. L. Wood, *Phys. Rev. C* **67**, 051305(R) (2003); the branching-ratio results from this measurement are considered to replace the contradictory upper limit set in an earlier less-precise measurement [143].
- [118] F. W. Prosser, G. U. Din, and D. D. Tolbert, *Phys. Rev.* **157**, 779 (1967).
- [119] W. V. Prestwich and T. J. Kennett, *Can. J. Phys.* **68**, 261 (1990); (erratum) **68**, 1352 (1990).
- [120] S. Raman, E. T. Jurney, D. A. Outlaw, and I. S. Towner, *Phys. Rev. C* **27**, 1188 (1983).
- [121] J. P. L. Reinecke, F. B. Waanders, P. Oberholtzer, P. J. C. Janse van Rensburg, J. A. Cilliers, J. J. A. Smit, M. A. Meyer, and P. M. Endt, *Nucl. Phys.* **A435**, 333 (1985).
- [122] M. L. Roush, L. A. West, and J. B. Marion, *Nucl. Phys.* **A147**, 235 (1970).
- [123] D. C. Robinson, J. M. Freeman, and T. T. Thwaites, *Nucl. Phys.* **A181**, 645 (1972); this reference replaces the  $^{10}\text{C}$  branching-ratio from J. M. Freeman, J. G. Jenkin, and G. Murray, *Nucl. Phys.* **A124**, 393 (1969).
- [124] D. C. Robinson and P. H. Barker, *Nucl. Phys.* **A225**, 109 (1974).
- [125] C. Rolfs, W. S. Rodney, S. Durrance, and H. Winkler, *Nucl. Phys.* **A240**, 221 (1975).
- [126] J. S. Ryder, G. J. Clark, J. E. Draper, J. M. Freeman, W. E. Burcham, and G. T. A. Squier, *Phys. Lett.* **B43**, 30 (1973).
- [127] A. Rytz, *At. Data Nucl. Data Tables* **47**, 205 (1991).
- [128] A. M. Sandorfi, C. J. Lister, D. E. Alburger, and E. K. Warburton, *Phys. Rev. C* **22**, 2213 (1980).
- [129] G. Savard, A. Galindo-Uribarri, E. Hagberg, J. C. Hardy, V. T. Koslowsky, D. C. Radford, and I. S. Towner, *Phys. Rev. Lett.* **74**, 1521 (1995).
- [130] G. Savard, J. A. Clark, F. Buchinger, J. E. Crawford, S. Gulick, J. C. Hardy, A. A. Hecht, V. E. Iacob, J. K. P. Lee, A. F. Levand, B. F. Lundgren, N. D. Scielzo, K. S. Sharma, I. Tanihata, I. S. Towner, W. Trimble, J. C. Wang, Y. Wang, and Z. Zhou, *Phys. Rev. C* **70**, 042501(R) (2004).
- [131] R. J. Scott, R. P. Rassool, M. N. Thompson, and D. V. Webb, *Nucl. Instr. Methods A* **539**, 191 (2005).
- [132] J. C. Sens, A. Pape, and R. Armbruster, *Nucl. Phys.* **A199**, 241 (1973).
- [133] K. K. Seth, A. Saha, W. Benenson, W. A. Lanford, H. Nann, and B. H. Wildenthal, *Phys. Rev. Lett.* **33**, 233 (1974).
- [134] D. Seweryniak, P. J. Woods, M. P. Carpenter, T. Davinson, R. V. F. Janssens, D. J. Jenkins, T. Lauritsen, C. J. Lister, C. Ruiz, J. Shergur, S. Sinha, and A. Woehr, *Phys. Rev. Lett.* **94**, 032501 (2005).
- [135] R. Sherr, J. B. Gerhart, H. Horie, and W. F. Hornyak, *Phys. Rev.* **100**, 945 (1955).
- [136] G. S. Sidhu and J. B. Gerhart, *Phys. Rev.* **148**, 1024 (1963).
- [137] J. Singh, *Ind. J. Pure Appl. Phys.* **10**, 289 (1972).
- [138] G. T. A. Squier, W. E. Burcham, J. M. Freeman, R. J. Petty, S. D. Hoath, and J. S. Ryder, *Nucl. Phys.* **A242**, 62 (1975).
- [139] G. T. A. Squier, W. E. Burcham, S. D. Hoath, J. M. Freeman, P. H. Barker, and R. J. Petty, *Phys. Lett.* **B65**, 122 (1976).
- [140] D. R. Tilley, H. R. Weller, C. M. Cheves, and R. M. Chasteler, *Nucl. Phys.* **A595**, 1 (1995).
- [141] N. R. Tolich, P. H. Barker, P. D. Harty, and P. A. Amundsen, *Phys. Rev. C* **67**, 035503 (2003).
- [142] H. Vonach, P. Glaessel, E. Huenges, P. Maier-Komor, H. Roesler, H. J. Scheerer, H. Paul, and D. Semrad, *Nucl. Phys.* **A278**, 189 (1977).
- [143] F. B. Waanders, J. L. Reinecke, H. N. Jacobs, J. J. A. Smit, M. A. Meyer, and P. M. Endt, *Nucl. Phys.* **A411**, 81 (1983).
- [144] T. A. Walkiewicz, S. Raman, E. T. Jurney, J. W. Starmer, and J. E. Lynn, *Phys. Rev. C* **45**, 1597 (1992).
- [145] H. Wenninger, J. Stiewe, and H. Leutz, *Nucl. Phys.* **A109**, 561 (1968).
- [146] R. E. White and H. Naylor, *Nucl. Phys.* **A276**, 333 (1977).
- [147] R. E. White, H. Naylor, P. H. Barker, D. M. J. Lovelock, and R. M. Smythe, *Phys. Lett.* **B105**, 116 (1981).
- [148] R. E. White, P. H. Barker, and D. J. Lovelock, *Metrologia* **21**, 193 (1985).
- [149] D. H. Wilkinson and D. E. Alburger, *Phys. Rev. C* **13**, 2517 (1976).
- [150] D. H. Wilkinson, A. Gallmann, and D. E. Alburger, *Phys. Rev. C* **18**, 401 (1978).
- [151] H. S. Wilson, R. W. Kavanagh, and F. M. Mann, *Phys. Rev. C* **22**, 1696 (1980).
- [152] J. Zioni, A. A. Jaffe, E. Friedman, N. Haik, R. Schectman, and D. Nir, *Nucl. Phys.* **A181**, 465 (1972).
- [153] F. Zijderhand, R. C. Markus, and C. van der Leun, *Nucl. Phys.* **A466**, 280 (1987).
- [154] S. Eidelman *et al.*, *Phys. Lett.* **B592**, 1 (2004).
- [155] W. Bambynek, H. Behrens, M. H. Chen, B. Crasemann, M. L. Fitzpatrick, K. D. Ledingham, H. Genz, M. Mutterer, and R. L. Intemann, *Rev. Mod. Phys.* **49**, 77 (1977).
- [156] R. B. Firestone, *Table of Isotopes*, 8th ed. (Wiley, New York, 1996).
- [157] A. Sirlin, *Phys. Rev. D* **35**, 3423 (1987); A. Sirlin and R. Zucchini, *Phys. Rev. Lett.* **57**, 1994 (1986).
- [158] W. Jaus and G. Rasche, *Phys. Rev. D* **35**, 3420 (1987).
- [159] I. S. Towner and J. C. Hardy, *Phys. Rev. C* **66**, 035501 (2002).
- [160] W. E. Ormand and B. A. Brown, *Phys. Rev. C* **52**, 2455 (1995); W. E. Ormand and B. A. Brown, *Phys. Rev. Lett.* **62**, 866 (1989); W. E. Ormand and B. A. Brown, *Nucl. Phys.* **A440**, 274 (1985).

- [161] H. Sagawa, N. Van Giai, and T. Suzuki, Phys. Rev. C **53**, 2163 (1996).
- [162] F. C. Barker, Nucl. Phys. **A537**, 134 (1992).
- [163] D. H. Wilkinson, Nucl. Instr. Methods in Phys. Res. A **488**, 654 (2002); **526**, 386 (2004).
- [164] A. Sirlin, in *Precision Tests of the Standard Electroweak Model*, edited by P. Langacker (World-Scientific, Singapore, 1994).
- [165] W. J. Marciano and A. Sirlin, Phys. Rev. Lett. **56**, 22 (1986).
- [166] J. D. Jackson, S. B. Treiman, and H.W. Wyld, Jr., Phys. Rev. **106**, 517 (1957).
- [167] H. Behrens and W. Bühring, *Electron Radial Wave Functions and Nuclear Beta-Decay* (Clarendon Press, Oxford, 1982).
- [168] S. Weinberg, Phys. Rev. **112**, 1375 (1958).
- [169] B. R. Holstein, Phys. Rev. C **29**, 623 (1984); W. E. Ormand, B. A. Brown, and B. R. Holstein, Phys. Rev. C **40**, 2914 (1989).
- [170] P. Herczeg, Phys. Rev. D **34**, 3449 (1986).
- [171] P. Herczeg, Prog. Part. Nucl. Phys. **46**, 413 (2001).
- [172] M. A. B. Bég, R. V. Budny, R. Mohapatra, and A. Sirlin, Phys. Rev. Lett. **38**, 1252 (1977).
- [173] J. Erler and M. J. Ramsey-Musolf, Prog. Part. Nucl. Phys. **54**, 351 (2005).
- [174] A. Sirlin, Rev. Mod. Phys. **50**, 573 (1978).
- [175] I. S. Towner, Nucl. Phys. **A540**, 478 (1992).
- [176] I. S. Towner, Phys. Lett. **B333**, 13 (1994).
- [177] S. Ando, H. W. Fearing, V. Gudkov, K. Kubodera, F. Myhrer, S. Nakamura, and T. Sato, Phys. Lett. **B595**, 250 (2004).
- [178] H. Behrens and J. Jänecke, *Numerical Tables for Beta Decay and Electron Capture, Landolt-Börnstein, New Series I/4* (Springer-Verlag, Berlin, 1969).
- [179] A. R. Edmonds, *Angular Momentum in Quantum Mechanics* (Princeton University Press, 1964).
- [180] H. De Vries, C. W. De Jager, and C. De Vries, At. Data Nucl. Data Tables **36**, 495 (1987).
- [181] M. E. Rose, Phys. Rev. **49**, 727 (1936).
- [182] J. J. Matese and W. R. Johnson, Phys. Rev. **150**, 846 (1966).