

Generalized Courant-Snyder Theory for Charged-Particle Dynamics in General Focusing Lattices

Hong Qin,^{1,2} Ronald C. Davidson,¹ Moses Chung,³ and Joshua W. Burby¹

¹Plasma Physics Laboratory, Princeton University, Princeton, New Jersey 08543, USA

²Department of Modern Physics, University of Science and Technology of China, Hefei, Anhui 230026, China

³Accelerator Physics Center, Fermi National Accelerator Laboratory, Batavia, Illinois 60510, USA

(Received 10 June 2013; published 5 September 2013)

The Courant-Snyder (CS) theory for one degree of freedom is generalized to the case of coupled transverse dynamics in general linear focusing lattices with quadrupole, skew-quadrupole, dipole, and solenoidal components, as well as torsion of the fiducial orbit and variation of beam energy. The envelope function is generalized into an envelope matrix, and the phase advance is generalized into a 4D symplectic rotation. The envelope equation, the transfer matrix, and the CS invariant of the original CS theory all have their counterparts, with remarkably similar expressions, in the generalized theory.

DOI: [10.1103/PhysRevLett.111.104801](https://doi.org/10.1103/PhysRevLett.111.104801)

PACS numbers: 29.27.-a, 52.20.Dq

For a charged particle in an uncoupled quadrupole focusing lattice, the classical Courant-Snyder (CS) theory [1] gives a complete description of its dynamics. Because the dynamics in the x direction and y direction are decoupled, the equations of motion in both directions are given by

$$\ddot{q} + \kappa_q(t)q = 0, \quad (1)$$

where q represents either of the transverse coordinates x or y , and $\kappa_q(t)$ is the time-dependent linear focusing strength. For a quadrupole lattice, $\kappa_x(t) = -\kappa_y(t)$. The CS theory states that the solution of Eq. (1) can be expressed as a time-dependent symplectic map applied to the initial phase-space coordinates, i.e., $(q, \dot{q})^T = M_1(t)(q_0, \dot{q}_0)^T$,

$$M_1(t) = \begin{pmatrix} \sqrt{\frac{\beta}{\beta_0}}[\cos\phi + \alpha_0\sin\phi] & \sqrt{\beta\beta_0}\sin\phi \\ -\frac{1+\alpha\alpha_0}{\sqrt{\beta\beta_0}}\sin\phi + \frac{\alpha_0-\alpha}{\sqrt{\beta\beta_0}}\cos\phi & \sqrt{\frac{\beta_0}{\beta}}[\cos\phi - \alpha\sin\phi] \end{pmatrix}, \quad (2)$$

where the superscript “ T ” denotes the transpose operation, and $\alpha(t)$, $\beta(t)$, and $\phi(t)$ are time-dependent functions defined in terms of the envelope function $w(t)$ by

$$\beta(t) \equiv w^2(t), \quad \alpha(t) \equiv -w\dot{w}, \quad \phi(t) \equiv \int_0^t \frac{dt}{\beta(t)}. \quad (3)$$

The envelope function $w(t)$ is determined by the nonlinear envelope equation

$$\ddot{w} + \kappa_q(t)w = w^{-3}. \quad (4)$$

In Eq. (2) $q_0 = q(t=0)$, $\dot{q}_0 = \dot{q}(t=0)$, $\beta_0 = \beta(t=0)$, and $\alpha_0 = \alpha(t=0)$ are initial conditions. The transfer matrix $M_1(t)$ can be decomposed as [2]

$$M_1(t) = \begin{pmatrix} w & 0 \\ \dot{w} & \frac{1}{w} \end{pmatrix} \begin{pmatrix} \cos\phi & \sin\phi \\ -\sin\phi & \cos\phi \end{pmatrix} \begin{pmatrix} w_0^{-1} & 0 \\ -\dot{w}_0 & w_0 \end{pmatrix}, \quad (5)$$

from which it is clear that the physical meaning of ϕ is the phase advance, and $\beta^{-1} = w^{-2}$ is the rate of phase advance. The Courant-Snyder invariant [1,3] is given by

$$I_{\text{CS}} = \frac{q^2}{w^2} + (w\dot{q} - \dot{w}q)^2 = (q, \dot{q}) \begin{pmatrix} \gamma & \alpha \\ \alpha & \beta \end{pmatrix} \begin{pmatrix} q \\ \dot{q} \end{pmatrix}, \quad (6)$$

$$\gamma(t) \equiv w^{-2} + \dot{w}^2. \quad (7)$$

The functions $\alpha(t)$, $\beta(t)$, and $\gamma(t)$ defined by Eqs. (3) and (7) are known as the Twiss parameters. These main results of the CS theory constitute a parametrization of the symplectic transfer map $M_1(t)$ for a standard uncoupled quadrupole lattice, and provide the foundation for accelerator and beam physics. Mathematically, there are different schemes to parametrize a time-dependent symplectic map. Among these, the CS parametrization is unique because it describes the physics of charged particle dynamics. The main components of the CS theory, i.e., the phase advance, the envelope equation, the transfer matrix, and the CS invariant, are of considerable importance to beam physics. The envelope function describes the transverse dimensions in configuration space while the CS invariant defines the emittance in phase space. This theoretical framework also makes it possible to investigate collective effects associated with high-intensity beams. For instance, it is used to construct the Kapchinskij-Vladimirskij distribution [4–6].

However, the CS theory can only be applied to the ideal case of uncoupled quadrupole focusing lattices. In realistic accelerators, there also exist bending magnets, torsion of the design orbit (fiducial orbit), and skew-quadrupole components introduced intentionally or by misalignment. In certain applications, such as the NDCX-II experiment [7], solenoidal magnets are also used. When these additional linear components are included, the transverse dynamics are coupled, and the dynamics of a charged particle relative to the fiducial orbit are governed by a general time-dependent Hamiltonian [8]

$$H = \frac{1}{2} z^T A z, \quad A = \begin{pmatrix} \kappa(t) & R(t) \\ R(t)^T & m^{-1}(t) \end{pmatrix}. \quad (8)$$

Here, $z = (x, y, p_x, p_y)^T$ are the transverse phase-space coordinates, and $\kappa(t)$, $R(t)$, and $m^{-1}(t)$ are time-dependent 2×2 matrices. The matrices $\kappa(t)$ and $m^{-1}(t)$ are also symmetric. The skew-quadrupole and dipole components are included in the off-diagonal terms of $\kappa(t)$, and the solenoidal component and the torsion of the fiducial orbit are included in $R(t)$. There are several different methods to include the effect of torsion. A summary is given by Hoffstaetter [9]. The variation of beam energy along the fiducial orbit is reflected in the mass matrix $m^{-1}(t)$. For complete generality, we further allow $m^{-1}(t)$ to be any real symmetric matrix. The transfer matrix $M(t)$ corresponding to H is a time-dependent 4×4 symplectic matrix, which has ten time-dependent parameters and admits many different schemes for parametrization. Teng and Edwards [10–12] first derived a set of parametrization schemes for $M(t)$, some of which have been adopted in lattice design and particle tracking codes, such as the MAD code [13]. A class of different parametrizations has also been developed by Dattoli *et al.* [14]. However, these parametrization schemes are rather mathematical and fail to connect with important physical parameters of the beam, in contrast to the original CS theory. For example, they cannot be used effectively to describe the instability properties of the coupled dynamics, and are not directly related to beam envelopes and emittance. In order to describe beam envelopes for coupled dynamics, Ripken [15,16] developed a method without using these parametrization schemes. In short, the elegant and much-needed connection with the physics of beam dynamics in the original CS theory for one degree of freedom is lost in these parametrization schemes.

Recently we have generalized the CS theory for uncoupled dynamics to treat coupled dynamics induced by a skew-quadrupole lattice [17–20]. However, this generalization is valid only for cases without torsion and solenoidal components, i.e., cases with $R = 0$ in Eq. (8). For cases with torsion and/or solenoidal components, $R \neq 0$ and we had to perform a phase-space rotation first to transform away the nonvanishing R , and then apply the generalized CS theory. While this procedure worked technically, it does not fit into the framework of the CS theory. This is because the first phase-space rotation is not consistent with the CS theory, and the envelope matrix and phase advance in the rotated frame lost their original meaning and physical importance. In this Letter, we complete the development of the generalized CS theory by allowing the focusing lattice to assume the most general form in Eq. (8), including bending magnets, torsion of the design orbit, and solenoidal magnets, in addition to quadrupole and skew-quadrupole components. In this generalized theory, the physics components of the original CS theory, i.e., the phase advance, the envelope equation, the transfer

matrix, and the CS invariant are all generalized to the 2D coupled case with identical structure.

We will use a time-dependent symplectic transformation technique [17,20,21]. A generalized envelope equation in 2×2 matrix form for the most general form of coupled dynamics is developed [see Eq. (16)]. In the original CS theory, the 1D envelope equation (4) plays a central role. It is also an important equation in other branches of physics, and has been discovered or rediscovered many times [22–26]. In quantum physics, it is also known as the Ermakov-Milne-Pinney equation [22–24], which has been utilized to study nonadiabatic Berry phases [27] in 1D time-dependent quantum systems [28,29]. A brief account of the history of the 1D envelope equation can be found in Ref. [3]. The generalization of the 1D envelope equation to higher dimensions for the most general Hamiltonian is also expected to have many possible applications in areas other than beam physics. In addition, the 1D CS invariant given by Eq. (6), also known as the Lewis invariant [25,26] in quantum physics, is generalized to higher dimension in Eq. (22).

The 1D phase advance concept is generalized to a time-dependent matrix P , which belongs to the symplectic rotation group $Sp(4) \cap SO(4) = U(2)$. Here, $Sp(4)$, $SO(4)$, and $U(2)$ denote the groups of 4×4 symplectic matrices, 4×4 rotation matrices, and 2×2 unitary matrices, respectively.

The generalized decomposition for the symplectic map $M(t)$ is given by Eq. (21), which has exactly the same structure as the original 1D CS theory given by Eq. (5). In addition to its aesthetic appearance, the generalized CS theory provides an effective tool to describe the beam physics determined by the most general Hamiltonian. The 2×2 envelope matrix w defines the transverse dimension of the beam, and the generalized CS invariant defines the emittance. We will also show that the stability properties of a charged particle are completely determined by envelope matrix and the generalized phase advance $P(t)$, which is a time-dependent curve in $Sp(4) \cap SO(4) = U(2)$.

The most critical part of the generalized theory is the decomposition of the time-dependent symplectic coordinate transformation G given by Eq. (20). The factorization of G as a product PS is closely related to the Iwasawa decomposition for a semisimple Lie group [30], which has played an important role in phase-space optics [31] and phase-space quantum mechanics [32]. However, the unique feature of the theory developed here is that the factorization is provided from the viewpoint of dynamics and self-consistently constructed from the generalized envelope equation. It is a pleasant surprise to find the deep connection between the original CS theory for charged particle dynamics [1] and the Iwasawa decomposition for Lie groups [30], two theoretical formalisms developed concurrently. This realization also demonstrates that beam dynamics, phase-space optics, and quantum dynamics have a similar dynamical structure at the fundamental level.

For the present application to beam transverse dynamics, there are two degrees of freedom. But the theory developed is valid for any degree of freedom. For a system with n degrees of freedom, the time-dependent matrix $A(t)$ specifying the Hamiltonian in Eq. (8) will be $2n \times 2n$, the envelope matrix will be $n \times n$, and the phase advance will belong to $Sp(2n) \cap SO(2n) = U(n)$.

We now present the detailed derivation of the generalized CS theory. We start by quickly reviewing the time-dependent symplectic transformation technique developed by Leach [21]. For a Hamiltonian given by $H_1 = (1/2)z_1^T A_1(t) z_1$, we seek a time-dependent linear symplectic coordinate transformation $z_2 = S(t)z_1$ such that in the z_2 coordinates, the dynamics are governed by $H_2 = (1/2)z_2^T A_2(t) z_2$. Here H_1 is specified by a $2n \times 2n$ symmetric matrix $A_1(t)$, and H_2 is a targeted Hamiltonian specified by a desired $2n \times 2n$ symmetric matrix $A_2(t)$. The required symplectic matrix $S(t)$ must satisfy the matrix equation [21]

$$\dot{S} = JA_2S - SJA_1, \quad (9)$$

where $J = \begin{pmatrix} 0 & I \\ -I & 0 \end{pmatrix}$ is the $2n \times 2n$ unit symplectic matrix. The derivation of Eq. (9) can be found in Refs. [17,18,21], and its geometric interpretation is given in Ref. [20].

We now apply this technique to the Hamiltonian given by Eq. (8). We proceed in two steps. First, we seek a coordinate transformation $\bar{z} = S_1 z$ such that, in the \bar{z} coordinates, the Hamiltonian assumes the form

$$\bar{H} = \frac{1}{2} \bar{z}^T \bar{A} \bar{z}, \quad \bar{A} = \begin{pmatrix} \mu(t) & 0 \\ 0 & \mu(t) \end{pmatrix}, \quad (10)$$

where $\mu(t)$ is a 2×2 matrix to be determined. In terms of 2×2 blocks, $S = \begin{pmatrix} S_1 & S_2 \\ S_3 & S_4 \end{pmatrix}$, and Eq. (9) splits into four matrix equations. Including $\mu(t)$, we have five 2×2 matrices to be determined. Based on the analogy with Eq. (5), we define $S_2 \equiv 0$. We define $w \equiv S_4$ because it will be clear later that S_4 is the envelope matrix. Equation (9) becomes

$$\dot{S}_1 = \mu S_3 - S_1 R^T, \quad (11)$$

$$S_1 = \mu w m, \quad (12)$$

$$\dot{S}_3 = -\mu S_1 - S_3 R^T + w \kappa, \quad (13)$$

$$S_3 = -\dot{w} m + w R m. \quad (14)$$

Because Eqs. (11)–(14) describe a curve in $Sp(4)$, they are consistent with the symplectic condition $S_1 w^T = I$, which implies

$$\mu = (w m w^T)^{-1}. \quad (15)$$

Then, from Eqs. (13) and (14) we immediately obtain the following matrix differential equation for the envelope matrix w ,

$$\frac{d}{dt} \left(\frac{dw}{dt} m - w R m \right) + \frac{dw}{dt} m R^T + w (\kappa - R m R^T) - (w^T w m w^T)^{-1} = 0. \quad (16)$$

This is the generalized envelope equation. It generalizes the 1D envelope equation (4) (or Ermakov-Milne-Pinney equation [22–24]), as well as the previous matrix envelope equation for cases with only quadrupole and skew-quadrupole magnets ($R = 0$) [17–20]. The envelope equation developed here [Eq. (16)] applies to the most general Hamiltonian with arbitrary R , and it includes Eq. (59) of Ref. [17] as a special case. By comparison, Eq. (16) is much more general than Eq. (59) of Ref. [17]. For its application in quantum mechanics, both of the envelope equations can be viewed as a generalized Ermakov-Milne-Pinney equation. However, Eq. (16) obviously is more general and has more applications. For example, it is applicable to quantum systems in a time-dependent magnetic field. For n degrees of freedom, the envelope matrix w will be $n \times n$, and the generalized envelope equation has the same format as Eq. (16). Once w is solved for from the envelope equation, we can determine S_1 from Eq. (12) and S_3 from Eq. (14). In terms of the envelope matrix w , the first symplectic transformation S and its inverse are given by

$$S = \begin{pmatrix} w^{-T} & 0 \\ (wR - \dot{w})m & w \end{pmatrix}, \quad (17)$$

$$S^{-1} = \begin{pmatrix} w^T & 0 \\ (w^{-1}\dot{w} - R)m w^T & w^{-1} \end{pmatrix}.$$

The second step is to use another coordinate transformation $\bar{\bar{z}} = P(t)\bar{z}$ to transform \bar{H} into a vanishing Hamiltonian $\bar{\bar{H}} \equiv 0$ for all time, thereby rendering the dynamics trivial in the new coordinates. In this case, the determining equation for the transformation $P(t)$ is

$$\dot{P} = -PJ\bar{A} = P \begin{pmatrix} 0 & -\mu \\ \mu & 0 \end{pmatrix}. \quad (18)$$

It is straightforward to verify that $-J\bar{A} \in sp(4) \cap so(4)$, where $sp(4)$ and $so(4)$ are the Lie algebras of $Sp(4)$ and $SO(4)$, respectively. Thus $P(t)$ is a curve in the group of 4D symplectic rotations, i.e., $P(t) \in Sp(4) \cap SO(4) = U(2)$, provided the initial condition of $P(t)$ is chosen such that $P(0) \in Sp(4) \cap SO(4) = U(2)$. We call $P(t)$ the generalized phase advance, an appropriate title in light of the fact that $P(t)$ is a symplectic rotation. The Lie algebra element (infinitesimal generator) $-J\bar{A} = \begin{pmatrix} 0 & -\mu \\ \mu & 0 \end{pmatrix}$ is therefore the phase advance rate, and it is determined by the envelope matrix through Eq. (15). Since $Sp(4) \cap SO(4) = U(2)$, P and its inverse must have the form of

$$P = \begin{pmatrix} P_1 & P_2 \\ -P_2 & P_1 \end{pmatrix}, \quad P^{-1} = P^T = \begin{pmatrix} P_1^T & -P_2^T \\ P_2^T & P_1^T \end{pmatrix}. \quad (19)$$

Combining the two symplectic coordinate transformations, we obtain the transformation

$$\bar{z} = G(t)z = P(t)S(t)z. \quad (20)$$

As mentioned above, the factorization of G as a product of a symplectic rotation (unitary) matrix P and a lower-triangular matrix S of the specified form is closely related to the Iwasawa decomposition for a semisimple Lie group [30]. Given that we have derived the factorization from purely dynamical considerations, the theory developed here can be viewed as a dynamical interpretation of the factorization for the symplectic group.

Because $\bar{H} \equiv 0$, $\bar{z} = \text{const}$. Therefore, the symplectic matrix specifying the map between z_0 and $z = M(t)z_0$ is

$$M(t) = S^{-1}P^{-1}P_0S_0 \\ = \begin{pmatrix} w^T & 0 \\ (w^{-1}\dot{w} - R)mw^T & w^{-1} \end{pmatrix} P^T \begin{pmatrix} w^{-T} & 0 \\ (wR - \dot{w})m & w \end{pmatrix}_0, \quad (21)$$

where subscript “0” denotes initial conditions at $t = 0$, and P_0 is taken to be I without loss of generality. This expression for $M(t)$ generalizes the decomposition of the symplectic map for the original 1D CS theory given by Eq. (5). Specifically, the 2D rotation matrix, which is also symplectic, in Eq. (5) is generalized into a 4D symplectic rotation P^T . The phase advance P is generated by the infinitesimal generator determined by the envelope matrix through $\mu = (wmw^T)^{-1}$, just as in the original 1D CS theory where the infinitesimal generator of the phase advance is w^{-2} for a scalar envelope w . The first and the third matrices in Eq. (5) obviously have the same construction as their counterparts in Eq. (21). It is important to understand the physical meaning of the decomposition in Eqs. (21) and (5). The first matrix from the right is a matching transformation at $t = 0$ of the initial conditions to an equivalent, homogeneously focusing system, where the external forces act constantly on the beam. The second matrix from the right is a transformation along the time axis in this homogeneously focusing system, with the phase advance playing the role of a timelike evolution parameter. And the third matrix from the right is a back transformation to the original coordinate system at $t > 0$.

Given the lattice function $A(t)$, we need to solve the envelope equation (16) and the phase advance equation (18) to construct $M(t)$ according to Eq. (21). The initial conditions for w and \dot{w} can be arbitrary except that they have to satisfy the symplectic condition $wS_3^T = S_3w^T$, which is a single constraint since w is a 2×2 matrix. The envelope equation (16) admits a 1D gauge freedom as well. For any constant matrix $c \in O(2)$, we can show by straightforward calculation that if w is a solution of Eq. (16), then cw is also a solution. However, $P(t)$ and $M(t)$ are independent of this gauge. The number of initial

conditions for Eq. (16) is eight, and the dimension of $P(t)$ as the $U(2)$ group is four. Subtracting the one symplectic constraint on the initial conditions and the one gauge freedom, the total number of independent parameters for $M(t)$ is ten, as expected.

Because $\bar{z} = \text{const}$,

$$I_\xi = z^T S^T P^T \xi P S z \quad (22)$$

is a constant of motion for any constant 4×4 positive definite matrix ξ . This expression generalizes the CS invariant [1] (or Lewis invariant [25,26]) for one degree of freedom in Eq. (6). In particular, we can define I_ξ , with ξ being the unit matrix, to be the special invariant corresponding to the original CS invariant. In this case, the phase advance P in Eq. (22) drops out, and

$$I_{\text{CS}} \equiv z^T S^T S z = z^T \begin{pmatrix} \gamma & \alpha \\ \alpha^T & \beta \end{pmatrix} z,$$

where α , β , and γ are the generalized Twiss parameters defined by

$$\alpha \equiv w^T S_3, \quad \beta \equiv w^T w, \quad \gamma \equiv S_3^T S_3 + w^{-1} w^{-T}.$$

It is straightforward to verify that as time-dependent 2×2 matrices, α , β , and γ satisfy

$$\beta\gamma = I + \alpha^2, \quad (23)$$

which is a familiar relationship in the original CS theory between the scalar Twiss parameters defined by Eqs. (3) and (7). Use has been made of the symplectic condition $wS_3^T = S_3w^T$ in obtaining Eq. (23).

The envelope matrix w and the invariant I_ξ define the beam dimensions and emittance for both low intensity beams and high intensity beams with strong space-charge potential. Investigation of these properties will be reported on in future publications. Here we demonstrate how to use the generalized theory to study the instability properties of the transverse dynamics governed by the Hamiltonian $H(t)$ specified in Eq. (8) for a general periodic lattice. After solving for a matched envelope matrix, the one-turn map $M_c(t)$ is given by

$$M_c(t) = S^{-1}(t)P_c^T(t)S(t), \quad (24)$$

where $P_c^T(t)$ is the one-turn phase advance transposed. Equation (24) states that $M_c(t)$ is similar to $P_c^T(t)$, and thus we reach the conclusion that the stability properties of the dynamics are determined by the one-turn phase advance $P_c^T(t)$. This significantly simplifies the stability analysis and showcases the physical relevance of the generalized CS parametrization scheme developed here. Since $P_c^T(t)$ is a symplectic rotation, we obtain the following stability criterion: a sufficient condition for the general

transverse dynamics to be stable is that the envelope equation has a matched solution.

This research was supported by the U.S. Department of Energy (Contracts No. DE-AC02-09CH11466 and No. DE-AC02-07CH11359).

-
- [1] E. Courant and H. Snyder, *Ann. Phys. (N.Y.)* **3**, 1 (1958).
- [2] S. Y. Lee, *Accelerator Physics* (World Scientific, Singapore, 1999), p. 47.
- [3] H. Qin and R. C. Davidson, *Phys. Rev. ST Accel. Beams* **9**, 054001 (2006).
- [4] I. Kapchinskij and V. Vladimirskij, in *Proceedings of the International Conference on High Energy Accelerators and Instrumentation* (CERN Scientific Information Service, Geneva, 1959), p. 274.
- [5] H. Qin, M. Chung, and R. C. Davidson, *Phys. Rev. Lett.* **103**, 224802 (2009).
- [6] H. Qin and R. C. Davidson, *Phys. Rev. Lett.* **110**, 064803 (2013).
- [7] A. Friedman, J. J. Barnard, R. H. Cohen, D. P. Grote, S. M. Lund, W. M. Sharp, A. Faltens, E. Henestroza, J.-Y. Jung, J. W. Kwan *et al.*, *Phys. Plasmas* **17**, 056704 (2010).
- [8] L. Michelotti, *Intermediate Classical Dynamics with Applications to Beam Physics* (John Wiley & Sons, New York, 1995), Chap. 3, p. 166.
- [9] G. H. Hoffstaetter, in *Proceedings of 1995 Particle Accelerator Conference* (IEEE, Piscataway, NJ, 1995).
- [10] L. C. Teng, National Accelerator Laboratory Report No. FN-229 (1971).
- [11] D. A. Edwards and L. C. Teng, *IEEE Trans. Nucl. Sci.* **20**, 885 (1973).
- [12] L. C. Teng, in *Proceedings of the 2003 Particle Accelerator Conference* (IEEE, Piscataway, NJ, 2003), p. 2895.
- [13] H. Grote, F. Iselin, E. Keil, and J. Niederer, in *Proceedings of 1989 Particle Accelerator Conference* (IEEE, Piscataway, NJ, 1989), p. 1292.
- [14] G. Dattoli, C. Mari, M. Richetta, and A. Torre, *Nuovo Cimento Soc. Ital. Fis.* **107B**, 269 (1992).
- [15] G. Ripken, DESY Tech. Report No. R1-70/04 (1970).
- [16] H. Wiedemann, *Particle Accelerator Physics* (Springer-Verlag, Berlin, 2007), pp. 614–620.
- [17] H. Qin and R. C. Davidson, *Phys. Rev. ST Accel. Beams* **12**, 064001 (2009).
- [18] H. Qin and R. C. Davidson, *Phys. Plasmas* **16**, 050705 (2009).
- [19] M. Chung, H. Qin, and R. C. Davidson, *Phys. Plasmas* **17**, 084502 (2010).
- [20] H. Qin and R. C. Davidson, *Phys. Plasmas* **18**, 056708 (2011).
- [21] P. Leach, *J. Math. Phys. (N.Y.)* **18**, 1608 (1977).
- [22] V. Ermakov, *Univ. Izv. Kiev* **20**, 1 (1880).
- [23] W. E. Milne, *Phys. Rev.* **35**, 863 (1930).
- [24] E. Pinney, *Proc. Am. Math. Soc.* **1**, 681 (1950).
- [25] H. R. Lewis, *J. Math. Phys. (N.Y.)* **9**, 1976 (1968).
- [26] H. R. Lewis and W. B. Riesenfeld, *J. Math. Phys. (N.Y.)* **10**, 1458 (1969).
- [27] M. V. Berry, *J. Phys. A* **18**, 15 (1985).
- [28] D. A. Morales, *J. Phys. A* **21**, L889 (1988).
- [29] D. B. Monteoliva, H. J. Korsch, and J. A. Nunez, *J. Phys. A* **27**, 6897 (1994).
- [30] K. Iwasawa, *Ann. Math.* **50**, 507 (1949).
- [31] K. B. Wolf, *Geometric Optics on Phase Space* (Springer, Berlin, 2004), pp. 173–177.
- [32] M. de Gosson, *Symplectic Geometry and Quantum Mechanics* (Birkhäuser Verlag, Basel, 2006), pp. 42–44.