



Monitoring microvascular perfusion variations with laser speckle contrast imaging using a view-based temporal template method

Submitted by Anne Humeau-Heurtier on Fri, 12/09/2016 - 21:08

Titre	Monitoring microvascular perfusion variations with laser speckle contrast imaging using a view-based temporal template method
Type de publication	Article de revue
Auteur	Ansari, Mohammad Zaheer [1], Kang, Eun-Jeung [2], Manole, Mioara D [3], Dreier, Jens P [4], Humeau-Heurtier, Anne [5]
Pays	Etats-Unis
Editeur	Elsevier
Type	Article scientifique dans une revue à comité de lecture
Année	2017
Langue	Anglais
Date	Mai 2017
Pagination	49-59
Volume	111
Titre de la revue	Microvascular Research
ISSN	0026-2862
Mots-clés	Biomedical signal processing [6], Cerebral blood flow [7], Microcirculation [8], Motion history image [9]
Résumé en anglais	<p>Purpose Laser speckle contrast imaging (LSCI) continues to gain an increased interest in clinical and research studies to monitor microvascular perfusion. Due to its high spatial and temporal resolutions, LSCI may lead to a large amount of data. The analysis of such data, as well as the determination of the regions where the perfusion varies, can become a lengthy and tedious task. We propose here to analyze if a view-based temporal template method, the motion history image (MHI) algorithm, may be of use in detecting the perfusion variations locations.</p> <p>Methods LSCI data recorded during three different kinds of perfusion variations are considered: (i) cerebral blood flow during spreading depolarization (SD) in a mouse; (ii) cerebral blood flow during SD in a rat; (iii) cerebral blood flow during cardiac arrest in a rat. Each of these recordings was processed with MHI.</p> <p>Results We show that, for the three pathophysiological situations, MHI identifies the area in which perfusion evolves with time. The results are more easily obtained compared with a visual inspection of all of the frames constituting the recordings. MHI also has the advantage of relying on a rather simple algorithm.</p> <p>Conclusions MHI can be tested in clinical and research studies to aid the user in perfusion analyses.</p>

URL de la notice <http://okina.univ-angers.fr/publications/ua15279> [10]
DOI [10.1016/j.mvr.2016.12.004](https://doi.org/10.1016/j.mvr.2016.12.004) [11]
Lien vers le document <http://www.sciencedirect.com/science/article/pii/S0026286216301765> [12]

Liens

- [1] <http://okina.univ-angers.fr/publications?f%5Bauthor%5D=24850>
- [2] <http://okina.univ-angers.fr/publications?f%5Bauthor%5D=25618>
- [3] <http://okina.univ-angers.fr/publications?f%5Bauthor%5D=25619>
- [4] <http://okina.univ-angers.fr/publications?f%5Bauthor%5D=24852>
- [5] <http://okina.univ-angers.fr/a.hum/publications>
- [6] <http://okina.univ-angers.fr/publications?f%5Bkeyword%5D=22108>
- [7] <http://okina.univ-angers.fr/publications?f%5Bkeyword%5D=22107>
- [8] <http://okina.univ-angers.fr/publications?f%5Bkeyword%5D=1229>
- [9] <http://okina.univ-angers.fr/publications?f%5Bkeyword%5D=21187>
- [10] <http://okina.univ-angers.fr/publications/ua15279>
- [11] <http://dx.doi.org/10.1016/j.mvr.2016.12.004>
- [12] <http://www.sciencedirect.com/science/article/pii/S0026286216301765>

Publié sur *Okina* (<http://okina.univ-angers.fr>)