

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23

Biocontrol of Ochratoxigenic moulds (*Aspergillus ochraceus* and *Penicillium nordicum*) by *Debaryomyces hansenii* and *Saccharomycopsis fibuligera* during speck production.

Lucilla Iacumin, Marisa Manzano, Debbie Andyanto and Giuseppe Comi*

Department Department of Agricultural, Food, Animal and Environmental Sciences

Via Sondrio 2/a, 33100 Udine, Italy.

*Corresponding author:

Giuseppe Comi

Department Department of Agricultural, Food, Animal and Environmental Sciences

Via Sondrio 2/A

33100 Udine, Italy

phone: +39 0432 558129; mobile: +39 338 9918561; fax: +39 0432 558130;

Abstract

24 Speck is a meat product obtained from the deboned leg of pork that is salted, smoked and
25 seasoned for four to six months. During speck seasoning, *Eurotium rubrum* and
26 *Penicillium solitum* grow on the surface and collaborate with other moulds and tissue
27 enzymes to produce the typical aroma. Both of these strains usually predominate over other
28 moulds. However, moulds producing ochratoxins, such as *Aspergillus ochraceus* and
29 *Penicillium nordicum*, can also co-grow on speck and produce ochratoxin A (OTA).
30 Consequently, speck could represent a potential health risk for consumers. Because *A.*
31 *ochraceus* and *P. nordicum* could represent a problem for artisanal speck production, the
32 aim of this study was to inhibit these mould strains using *Debaryomyces hansenii* and
33 *Saccharomycopsis fibuligera*. Six *D. hansenii* and six *S. fibuligera* strains were tested *in*
34 *vitro* to inhibit *A. ochraceus* and *P. nordicum*. The *D. hansenii* DIAL 1 and *S. fibuligera*
35 DIAL 3 strains demonstrated the highest inhibitory activity and were selected for *in vivo*
36 tests. The strains were co-inoculated on fresh meat cuts for speck production with both of
37 the OTA-producing moulds prior to drying and seasoning. At the end of seasoning (six
38 months), OTA was not detected in the speck treated with both yeast strains. Because the
39 yeasts did not adversely affect the speck odour or flavour, the strains are proposed as
40 starters for the inhibition of ochratoxigenic moulds.

41 42 43 44 45 46 47 48 49 **1. Introduction**

50 Speck is a typical meat product in the European alpine area, particularly North Tyrol
51 (Austria) and South Tyrol (Alto Adige, Italy), regions that utilize different recipes but the
52 same technology. A flow chart showing the traditional speck production process is
53 presented in Table 1. Fresh meat cuts (FMC) obtained from deboned legs are carefully
54 brined with a mixture of salt and spices, such as pepper, juniper, laurel and rosemary, and
55 the percentages of these ingredients depend on the facility. The meat cuts are then cold
56 smoked (20 °C), dehydrated and seasoned for four to six months at 10-15 °C with a
57 relative humidity (R.H.) between 60 and 90%. During this period, moulds grow on the
58 meat surface and give it a distinctive flavour. The colonization of speck during drying and
59 seasoning occurs spontaneously from the mycobiota of the production environment or by
60 contamination with spices (Peintner et al., 2000). The indigenous mycobiota are valued for
61 their beneficial seasoning effects (Peintner et al., 2000). Moulds prevent excessive drying
62 of the surface, improve the texture, and limit excessive hardness of the flesh due to their
63 proteolytic activity (Sunesen and Stahnke, 2003). They also exerts an antioxidant effect,
64 contribute to colour production (Spotti et al., 2008) and improve the aroma and flavour of
65 speck (Rojas et al., 1991, Nunez et al., 1996). The surface moulds of meat products have
66 been studied in great detail (Dragoni et al., 1980; Huerta et al., 1987; Spotti et al., 1989;
67 Rojas et al., 1991; Nunez et al., 1996; Battilani et al., 2007; Iacumin et al., 2011). *Eurotium*
68 spp. appears to be the most prevalent during all stages of meat production (Peintner et al.,
69 2000; Comi et al., 2004; Battilani et al., 2010; Comi and Iacumin, 2013). In North and
70 South Tyrol, *Eurotium rubrum* and *Penicillium solitum* are the dominant species in all
71 parts of speck (crust, meat and fat). Other Pennicillia, including *P. nalgiovense*, *P.*
72 *verrucosum*, *P. canescens*, *P. brevicompactum*, *P. chrysogenum*, *P. glabrum*, *P. commune*
73 and *P. waksmanii*, are less frequently observed (Peintner et al., 2000). The development of
74 mould can also be related to negative effects, such as the production of mycotoxin (Comi
75 and Iacumin, 2013). Several recent investigations have highlighted the presence of

76 ochratoxin A (OTA) in different meat products, and several fungi, including *Penicillium*
77 *verrucosum*, *P. nordicum*, and *Aspergillus ochraceus*, can produce OTA following growth
78 on the surface of meat products during seasoning and storage (Matrella et al. 2006; Pietri et
79 al. 2006; Iacumin et al., 2011). OTA is undesirable because it is classified by IARC
80 (International Agency for Research of Cancer) into "Group B" as a molecule with possible
81 carcinogenic activity in humans (IARC, 1993). For this reason, limits have been introduced
82 to foods. In Italy, the maximum concentration of OTA allowed in meat and meat products
83 is 1 µg/kg (Ministry of Health Circular no. 10-93 09/06/1999). Different artisanal
84 production lots of speck were recently contaminated with *A. ochraceus* and *P. nordicum*.
85 Both species grow during drying and seasoning and produce up to 1 µg/kg OTA. Many
86 technologies must be used to eliminate the growth of OTA-producing moulds on meat
87 products. The most interesting methods developed to date are represented by bioprotective
88 cultures. Exploring the antagonistic activity of autochthonous yeast strains against moulds
89 could be of great interest. Several researchers (Andrade et al., 2014; Virgili et al., 2012)
90 have experimented with the ability of native yeasts to protect meat products against mould
91 growth. *D. hansenii* is the main yeast present in meat products from the early stage of
92 seasoning to the end, when it becomes the predominant strain (Comi and Cantoni, 1980;
93 Nunez et al., 1996; Asefa et al., 2010; Simoncini et al., 2007). Consequently, *D. hansenii* is
94 largely used as a starter culture to improve the quality and sensorial characteristics of
95 European sausages (Hammes and Knauf, 1994). Conversely, *S. fibuligera* is rarely present
96 in meat products because it is typical of starchy substrates and vegetables (Chi et al.,
97 2009). However, it is used because it grows in the presence of high salt and sugar
98 concentrations similarly to *D. hansenii*. Because yeasts are capable of growing and
99 colonizing meat products, they could represent the main strains for food biocontrol against
100 moulds and pathogenic microorganisms (Virgili et al., 2012; Andrade et al., 2014;
101 Simoncini et al., 2014). The efficacy of the protection provided by yeasts against moulds

102 on meat products must be seriously evaluated because moulds often grow on meat product
103 surfaces despite the presence of yeasts (Virgili et al., 2012). Little information regarding
104 the antagonistic activity of yeasts on dry cured ham has been reported (Virgili et al., 2012),
105 and different researchers have recently investigated and demonstrated the inhibitory effect
106 of yeasts *in vitro* against OTA-producing moulds (Andrade et al., 2014; Simoncini et al.,
107 2014). These researchers have demonstrated that it is possible to reduce *P. nordicum*
108 growth and activity using different yeast strains. The inhibitory mechanism is not well
109 understood, but it has been suggested that yeasts restrict nutrient availability and sites for
110 colonization and consequently predominate and limit the growth of OTA-producing
111 moulds (Bjönberg and Schnürer, 1993; Spotti et al., 2009). Considering the many
112 opportunities that yeasts can have in mould inhibition, this study aimed to evaluate the
113 abilities of *Debaryomyces hansenii* and *Saccharomycopsis fibuligera* to control
114 ochratoxigenic mould growth (*Aspergillus ochraceus* and *Penicillium nordicum*) during
115 speck production.

116

117 **2. Materials and methods**

118 Six strains of *Saccharomycopsis fibuligera* (DIAL 1, DIAL 2, DIAL 3, DIAL 4, DIAL 5,
119 and DIAL 6) and six strains of *Debaryomyces hansenii* (DIAL 1, DIAL 2, DIAL 3, DIAL
120 4, DIAL 5, and DIAL 6) were selected from a total of 90 isolates from meat products and
121 stored in the Collection of the Department of Food Science of the University of Udine
122 (DIAL, Udine, Italy). The twelve yeast strains were selected for potential competition
123 against *Penicillium nordicum* and *Aspergillus ochraceus*. The strains were maintained in
124 malt extract modified agar (MEMA, Oxoid, Italy) supplemented with dextrose (1%) and
125 peptone (1%).

126

127 *2.1. Preparation of yeast inocula*

128 The yeast strains were grown on MEMA at 25 °C for 48 h. Suspensions were then
129 prepared by adding a loop full of cells to peptonised water (0.7% NaCl and 0.1% peptone
130 in 1.000 mL of water). The density of the yeast cultures was determined
131 spectrophotometrically by measuring their optical density at 600 nm. Serial dilutions were
132 prepared to obtain the concentration used for the experiments.

133

134 2.2. Preparation of the *P. nordicum* and *A. ochraceus* inocula

135 One *P. nordicum* and one *A. ochraceus* strain were used in this study. These strains were
136 isolated from sausages in a previous study and identified at the molecular level. Both
137 strains produce OTA and belong to the Food Science Department of the University of
138 Udine. Both strains were grown in Czapek yeast extract agar plates (Oxoid, Italy) and
139 incubated at 25 °C for seven days. At the end of the incubation period, the conidia were
140 removed from the culture surface according to the method reported by Virgili et al. (2012).
141 The conidial suspension was adjusted to 10^2 , 10^4 and 10^6 conidia/mL using a
142 haemocytometer.

143

144 2.3. Inhibition of *P. nordicum* and *A. ochraceus* by yeast strains

145 The assay was performed using the method described by Virgili et al. (2012) and Bleve et
146 al. (2006) and modified as follows: the experiments were performed on MEMA adjusted to
147 pH 6.0. A top agar was prepared by mixing 10 mL of MEMA broth with 0.7% agar and 1
148 mL of a yeast suspension (*D. hansenii* or *S. fibuligera*) containing 10^6 CFU/mL to obtain a
149 thick, continuous layer on the plate surface. This suspension was distributed into Petri
150 plates containing 15 mL of MEMA. Three 10 μ L portions of *P. nordicum* or *A. ochraceus*
151 corresponding to 10^6 conidia/mL were then spotted onto each plate and incubated at 20 °C.
152 Top agar containing different yeast concentrations (10^2 , 10^4 , or 10^6 CFU/mL) was also
153 prepared to investigate differences among different yeast inocula.

154 Three replicate experiments were performed for each yeast. Plates with culture medium
155 (MEMA) that were not inoculated with yeast were included as controls.

156 Fungal growth was expressed as the average measurement (mm) of two orthogonal
157 diameters per colony after 14 days of incubation at 20 °C. The inhibitory activity was
158 calculated using the equation reported by Lima et al. (1999):

$$\% \text{ Inhibitory activity} = \frac{(\text{Fungal growth in control plate} - \text{Fungal growth in treated plate}) \times 100}{\text{Fungal growth in control plate}}$$

162 2.4. Inhibitory activity at different concentrations of the selected yeasts

163 The above-mentioned method (Virgili et al., 2012) was used the following modifications,
164 and the concentration of yeast in top agar was changed. Briefly, top agars containing
165 different concentrations of the yeasts (10^2 , 10^4 , or 10^6 CFU/mL) were used to examine
166 differences among yeast inocula concentrations. *D. hansenii* (strain DIAL 1) and *S.*
167 *fibuligera* (DIAL 3), which demonstrated the best performance in the above-mentioned
168 test, were used. Ten millilitres of inoculated top agar were distributed onto 15 mL of
169 MEMA, and three 10- μ L suspensions containing 10^6 conidia/mL of the moulds were then
170 spotted separately onto each plate and incubated at 20 °C for 14 days. Three replicate
171 experiments were performed for each yeast concentration. MEMA plates that were not
172 inoculated with yeast were included as controls.

173

174 2.5. Inhibitory activity at different concentrations of moulds (*P. nordicum* and *A.* 175 *ochraceus*)

176 The above-mentioned method (Virgili et al., 2012) was used with the following
177 modifications. The concentration of yeasts in top agar was 10^4 CFU/mL, and *D. hansenii*
178 (strain DIAL 1) and *S. fibuligera* (DIAL 3), which demonstrated the best performance in
179 the above-described test, were used. Ten millilitres of inoculated top agar were distributed

180 onto 15 mL of MEMA, and three 10- μ L suspensions containing different concentrations of
181 conidia (10^2 , 10^4 , or 10^6 conidia/mL) of both moulds were then separately spotted onto
182 each plate and incubated at 20 °C for 14 days. Three replicate experiments were performed
183 for each mould concentration. MEMA plates that were not inoculated with yeast were
184 included as controls.

185

186 *2.6. Inhibitory activity of yeasts on OTA production in the speck model system* (Battilani et
187 al., 2010, modified)

188 Five specks with different water activity (A_w) values were collected from a South
189 Tyrolean facility. From each speck, samples (50 mm in diameter and 5 mm in height) were
190 excised using a hollow metal sampler with a cylindrical cutting edge. The A_w values of the
191 samples measured with an AquaLab CX-2 instrument (Steroglass, Pullman, WA, USA)
192 were 0.960 ± 0.005 , 0.940 ± 0.005 , 0.920 ± 0.005 , 0.900 ± 0.005 , and 0.880 ± 0.005 . After
193 dipping in absolute ethanol for 2 min, the samples were removed and flamed to sterilize
194 them prior to inoculation (Rojas et al., 1991). The samples were placed in Petri plates,
195 inoculated and placed in sealed boxes equipped with beakers containing NaCl solutions
196 (Multon and Bizot, 1980) with the same A_w as the five speck samples. The salt solutions
197 were prepared with distilled water (w/w) with 6.57% NaCl (A_w 0.96), 9.38% NaCl (A_w
198 0.94), 11.90% NaCl (A_w 0.92), 14.18% NaCl (A_w 0.90) and 16.28% NaCl (A_w 0.88).
199 Each A_w value was confirmed using AquaLab CX-2.

200 The samples were incubated in the dark at 14 °C for 30 days. Six different inocula or co-
201 inocula were tested: 1) Control A, 10^4 conidia/cm 2 *Aspergillus ochraceus*; 2) Control B,
202 10^4 conidia/cm 2 *Penicillium nordicum*; 3) 10^6 CFU/g *D. hansenii* DIAL 1 vs. 10^4
203 conidia/cm 2 *Aspergillus ochraceus*; 4) 10^6 CFU/g *D. hansenii* DIAL 1 vs. 10^4 conidia/cm 2
204 *Penicillium nordicum*; 5) 10^6 CFU/g *S. fibuligera* DIAL 3 vs. 10^4 conidia/cm 2 *Aspergillus*
205 *ochraceus*; and 6) 10^6 CFU/g *S. fibuligera* DIAL 3 vs. 10^4 conidia/cm 2 *P. nordicum*. Each

206 condition was evaluated in triplicate. At the end of the incubation period, the 90 samples
207 were collected and analysed for OTA according to the method reported by Matrella et al.
208 (2006).

209

210 *2.7. Inhibitory activity of yeasts on OTA production during speck seasoning*

211 One hundred and twenty fresh meat cuts (FMCs) were collected. Each FMC (6.5 kg) was
212 trimmed and squared according to the traditional procedure and was then salted and
213 smoked. After smoking, the FMCs were inoculated with a mix of yeast and mould
214 suspensions or with a mould suspension alone according to the following experimental
215 design: 20 FMCs were inoculated with a suspension of *P. nordicum* at a final concentration
216 10^4 conidia/cm²; 20 FMCs were inoculated with a suspension of *A. ochraceus* at a final
217 concentration 10^4 conidia/cm²; 20 FMCs were inoculated with a mix of *A. ochraceus* and
218 *D. hansenii* (final concentration 10^4 conidia/ 10^6 CFU/cm²); 20 FMCs were inoculated with
219 a mix of *A. ochraceus* and *S. fibuligera* (10^4 conidia/ 10^6 CFU/cm²); 20 FMCs were
220 inoculated with a mix of *P. nordicum* and *D. hansenii* (10^4 conidia/ 10^6 CFU/cm²); and 20
221 FMCs were inoculated with a mix of *P. nordicum* and *S. fibuligera* (10^4 conidia/ 10^6
222 CFU/cm²). The *S. fibuligera* DIAL 3 and *D. hansenii* DIAL 1 strains were used. All of the
223 FMCs were seasoned for six months according to the traditional procedure (Table 1). At
224 the end of the seasoning period, 100-cm² samples were taken from a depth of 0.5 cm below
225 the speck surface and analysed for OTA. Briefly, meat collected from a depth of 0.5 cm
226 below the slime was homogenized using a Stomacher instrument (Lab Blender 400, PBI,
227 Italy), and 10 g of the homogenate was then used for the analysis. OTA was extracted and
228 evaluated according to the method described by Matrella et al. (2006). Before sampling for
229 OTA analysis, the surface of each speck was observed with a Stereoscope (320X; WILD
230 M 420, Heerbrugg, CH) to determine the presence of mould growth (i.e., hyphae).

231

232 2.8. *Sensorial analysis*

233 To evaluate the influence of the yeast culture starter on the organoleptic characteristics of
234 the product, a sensory analysis was performed using the triangle test methodology (ISO
235 4120:2004). FMCs were divided into three lots. After smoking, lot A (10 cuts) was
236 inoculated with a suspension of *S. fibuligera* DIAL 3 (10^6 CFU/cm²), and Lot B (10 cuts)
237 was inoculated with *D. hansenii* DIAL 1 (10^6 CFU/cm²). Lot C (10 cuts) served as the
238 control and was not inoculated with yeast cultures. All cuts were seasoned according to the
239 traditional procedure (Table 1). At the end of seasoning (6 months), all of the lots were
240 subjected to the triangle test, which was used to compare Lot A to Lot B, Lot A to Lot C,
241 and Lot B to Lot C. Twenty non-professional assessors were presented with three products,
242 two of which were identical. The assessors were asked to state which product they believed
243 was a unique sample. The assessors who indicated the existence of two distinct samples
244 were asked to identify the best sample.

245

246 2.9. *Statistical analysis*

247 The values of the various parameters were compared through one-way analysis of
248 variance. The averages were compared with Tukey's honest significance test using the
249 Statistical Graphics software package (Rockville, MD, USA).

250

251 **3. Results**

252 A biocontrol test was performed using six *Saccharomycopsis fibuligera* and six
253 *Debaryomyces hansenii* strains isolated from meat products. Four out of the six *D.*
254 *hansenii* and three out of the six *S. fibuligera* strains were detected in dry cured ham. The
255 other strains were isolated from sausages. All of the strains were tested for their ability to
256 grow *in vitro* and *in vivo* in the presence of low Aw (0.88) and 6% NaCl and on speck and
257 dry cured ham (data not shown).

258 Different *in vitro* inhibitory activities against *P. nordicum* and *A. ochraceus* were observed
259 among the tested strains ($p < 0.05$). *D. hansenii* DIAL 1 and *S. fibuligera* DIAL 3 showed
260 the highest inhibitory activities against *A. ochraceus* (76.0% and 86.1%, respectively) and
261 *P. nordicum* (78.8% and 85.2%, respectively; Tables 2-3). The diameters of *A. ochraceus*
262 colonies were reduced from 32.5 mm (control) to 4.5 mm by *S. fibuligera* and to 7.8 mm
263 by *D. hansenii*. The diameters of *P. nordicum* were reduced from 20.3 mm (control) to 3.0
264 and 4.3 mm by *S. fibuligera* and *D. hansenii*, respectively.

265 For this reason, both strains were selected for the assessment of their *in vitro* or *in vivo*
266 inhibitory effects. Different concentrations of both yeast strains were tested separately to
267 identify the best strain. As expected, the inhibitory activity was dependent on the yeast
268 concentration. Indeed, an *in vitro* concentration of 10^6 CFU/mL of agar showed the best
269 inhibitory effect for both of the tested yeasts ($p < 0.05$; Table 4). The antagonistic activities
270 of the yeasts (10^4 CFU/mL) were evaluated separately against different concentrations of
271 both moulds (10^2 , 10^4 , and 10^6 conidia/mL). The percentage of inhibition varied according
272 to the mould concentration (Table 5). The lowest percentage of inhibition was observed
273 after inoculation with the highest mould concentration ($p < 0.05$). The activity of *S.*
274 *fibuligera* was higher than the activity of *D. hansenii*. A higher yeast concentration (i.e.,
275 10^6 CFU/mL) completely inhibited mould at concentrations of 10^2 and 10^4 conidia/mL
276 (data not shown), and the same results, which are shown in Table 4, were obtained with
277 similar yeast and mould concentrations ($10^6/10^6$ CFU/mL). The measured diameters
278 demonstrated that the inhibitory activity of *S. fibuligera* was higher than that of *D. hansenii*
279 ($p < 0.05$), even though the media used appeared to be more productive for *D. hansenii*
280 than for *S. fibuligera*.

281 A reduction in fungal growth and consequently a decrease in OTA production were
282 observed in speck model systems (Photograph 1). Table 6 displays the effects of the yeast
283 strains on the OTA production of the tested moulds in pieces of speck with different A_w

284 levels. As shown, a low OTA concentration was detected in the pieces of speck inoculated
285 with both yeast and mould strains. The OTA concentration in the speck co-inoculated with
286 yeast and mould strains was less than 1 µg/kg (Table 6), which is the limit allowed in meat
287 and meat products by the Italian Ministry of Health (Circular no. 10-93 09/06/1999). In
288 contrast, OTA concentrations in speck inoculated only with mould reached 1 µg/kg. As
289 expected, pieces of speck with the higher A_w values (i.e., 0.96 and 0.94) exhibited the
290 highest OTA concentrations (Table 6).

291 The inhibitory activity of both yeasts against OTA-producing moulds during speck
292 production was tested *in vivo*. Both yeasts were co-inoculated separately in FMCs with the
293 *A. ochraceus* and *P. nordicum* strains. The strains inhibited both of the moulds *in vivo*. An
294 inoculum with the tested strains was prepared after the smoking phase. Less than 0.1 µg/kg
295 OTA was detected in speck inoculated with yeast and mould (Table 7). Conversely, OTA
296 concentrations of 65.5 µg/kg and 68.1 µg/kg were detected in speck inoculated with *A.*
297 *ochraceus* and *P. nordicum*, respectively. No *hyphae* or OTA-producing moulds were
298 observed on the surface of speck using a stereoscope, as demonstrated in Photograph 2.2.
299 The data demonstrated that both of the tested yeast strains were able to reduce or
300 completely inhibit the growth of *A. ochraceus* and *P. nordicum* both *in vitro* and *in vivo*.
301 Consequently, the yeasts can inhibit OTA production.

302 Based on the promising results obtained, the use of starter cultures of yeast to control
303 OTA-producing moulds might be recommended. However, it was necessary to evaluate the
304 influence of the starter yeasts on the acceptability of speck prior to implementing this
305 suggestion. Speck produced with and without the yeast starter were seasoned using the
306 traditional method (Table 1). At the end of seasoning, the speck samples were subjected to
307 a triangular test by a panel of 20 non-professional assessors. The entire panel confirmed
308 the acceptability of the speck supplemented with the starter cultures and found no
309 difference among Lots A, B (with bio-protective cultures) and C (without protective

310 cultures). The tasted speck slices were compact and homogeneous. The lean part was ruby
311 red in colour, and the colour of the fat was white, which is typical of the product. The
312 consistency was compact but not elastic, and the bouquet was delicate and distinctive. The
313 taste was delicate, and there was no perception of spices or flavours.

314

315 **4. Discussion**

316 The potential biopreservative activities of six different *D. hansenii* and six *S. fibuligera*
317 strains isolated from meat products against ochratoxigenic moulds were investigated in the
318 present study. These species were selected because they are widespread on meat and meat
319 products (Cocolin et al., 2006; Andrade et al., 2009) and are considered safe by the food
320 industry (EFSA, 2012).

321 Speck is a typical meat product of North Tyrol (Austria) and South Tyrol (Alto Adige,
322 Italy). In Italy, speck is also produced in the Veneto and Friuli Venezia Giulia regions.
323 During the seasoning process, a large population of microorganisms (mainly moulds and
324 yeasts) develops on the outer layers of the speck (Peintner et al., 2000). Some of these
325 moulds could produce OTA, such as *P. nordicum* and *A. ochraceus*. Under some
326 environmental conditions, their growth and ability to synthesize OTA could represent a
327 real risk for human health. OTA has been found in different meats and meat products
328 contaminated by ochratoxigenic moulds (Rojas et al., 1991; Battilani et al., 2010;
329 Dall'Asta et al., 2010; Iacumin et al., 2011). However, the presence of OTA is not always
330 correlated with OTA-producing moulds in meat products (Rodríguez et al., 2012).
331 Different methods can be used to inhibit OTA-producing moulds (Iacumin et al., 2011;
332 Comi et al., 2013). The most commonly used method consists of spreading starter cultures
333 on the meat product surface in the late drying and seasoning stages (Comi et al., 2013). In
334 sausage production, mould and yeast starters are mainly used because they limit the growth
335 of pathogenic microorganisms and because they improve the sensorial quality of the

336 product. Our study focused on the use of two yeast strains, and their activities were
337 evaluated either *in vitro* or *in vivo*. Under *in vitro* conditions, *D. hansenii* and *S. fibuligera*
338 significantly inhibited the growth of the tested ochratoxigenic moulds, and differences
339 were observed among the *D. hansenii* and *S. fibuligera* strains. *D. hansenii* DIAL 1 and *S.*
340 *fibuligera* DIAL 3 showed the highest efficiencies for the inhibition of *P. nordicum* and *A.*
341 *ochraceus* on solid media.

342 The inhibitory effect of *S. fibuligera* DIAL 3 against both of the mould strains was higher
343 and significantly different ($p < 0.05$) compared with the effect of *D. hansenii* DIAL 1. The
344 inhibitory activity of *D. hansenii* strains against *P. nordicum* was also found to be reduced
345 by Virgili et al. (2012), which is in agreement with the results of our investigation. These
346 researchers found that two strains, which were selected for their ability to grow in dry
347 cured ham-like substrates (*Candida zeylanoides* and *Hyphopichia burtonii*), were more
348 effective against *P. nordicum* than against *D. hansenii* and *Candida famata* (the
349 anamorphic form of *D. hansenii*). Indeed, both of these strains inhibited *P. nordicum*
350 growth and OTA production (Virgili et al., 2012). Differences in inhibitory activity against
351 *P. nordicum* were also found among *D. hansenii* strains among Andrade et al. (2014).
352 These researchers demonstrated that the *in vitro* efficiency depends on the strains, the
353 medium and the A_w value and that the lysed yeast cells might provide nutrients for *P.*
354 *nordicum* growth. These researchers concluded that the inhibition observed appears to be
355 due to compounds produced by yeast strains with activity against the tested moulds.
356 Several researchers (Marquina et al., 2001; Masoud et al., 2005) have demonstrated that *D.*
357 *hansenii*, *Saccharomyces cerevisiae* and *Pichia anomala* produce killer proteins against
358 sensitive strains of yeasts and moulds. Again, the observed mould inhibition could be due
359 to the competition activity of nutrients, as previously suggested (Björnberg and Schnürer,
360 1993; Zhao et al., 2008); however, researchers have speculated the existence of a minor
361 effect on the production of extracellular volatile compounds and killer proteins.

362 Due to the large variability in OTA-producing moulds on speck and meat products (Virgili
363 et al., 2012; Asefa et al., 2010), yeasts inoculated with a lower concentration were
364 associated with a lower level of mould inhibition. Additionally, the *in vitro* inhibitory
365 activity of *S. fibuligera* against OTA-producing moulds was higher than the inhibitory
366 activity of *D. hansenii*. The use of different yeast concentrations *in vitro* resulted in
367 different percentages of mould inhibition, demonstrating that mould inhibition depends on
368 the competition for the nutrients but not on antimycotic production. Consequently, the
369 highest yeast concentration (10^6 CFU/g) induced the largest inhibitory action against both
370 of the OTA-producing moulds. The experiment confirmed the potential biocontrol effect of
371 both of the yeast strains used. Our data agree with the results reported by Virgili et al.
372 (2012). In fact, the inhibitory activity of yeast was affected by the mould concentration.
373 Virgili et al. (2012) hypothesized that the concentration of OTA-producing moulds in meat
374 products is a key factor for the effectiveness of yeast starters. Consequently, the meat
375 production industry should apply the most severe controls during the seasoning process to
376 limit the contamination and growth of OTA-producing moulds. Additionally, the raw
377 material, temperatures, and R.H. of the drying and seasoning rooms should be controlled.
378 In the speck model system, the data demonstrated the possibility of the effective biocontrol
379 of the OTA-producing moulds by the yeast. The results did not appear to depend on the
380 A_w , the yeasts or the mould strains. The reduced OTA concentration in the speck model
381 system with different A_w values compared with the control samples was quite obvious.
382 The antagonistic activities of *D. hansenii* and *S. fibuligera* were not dependent on the A_w
383 and were similar to data reported by Simoncini et al. (2014), who demonstrated that the
384 variability in the antagonistic effect was affected by the strains of the different species used
385 (*D. hansenii* and *C. zeylanoides*). A positive effect of both of the yeasts tested was
386 observed on the inhibition of mould growth, OTA production and accumulation. However,
387 no association was found between the presence of mycotoxins and the biomass of the

388 OTA-producing moulds. As demonstrated by Xu et al. (2007), the production and
389 concentration of some mycotoxins in food are not necessarily proportional to the biomass
390 of the OTA-producing mould. The biocontrol activity of *D. hansenii* has been
391 demonstrated by various researchers (Hernández-Montiel et al., 2010; Virgili et al., 2012;
392 Andrade et al., 2014). There are no data on the biocontrol activity of *S. fibuligera* against
393 OTA-producing moulds, even though different yeast species isolated from cheese, milk,
394 grape must, wine, fruits and vegetables have been demonstrated to inhibit many
395 mycotoxigenic moulds and to reduce the presence of mycotoxins (Bleve et al., 2006; Zhao
396 et al., 2008; Virgili et al., 2012; Simoncini et al., 2014). Thus, this study provides the first
397 demonstration of the use of this yeast strain for the biocontrol of OTA-producing moulds.
398 In addition, this study used *D. hansenii* DIAL 1 and *S. fibuligera* DIAL 3 as starter cultures
399 for speck production to eliminate the growth of OTA-producing moulds and to eliminate or
400 reduce the presence of OTA. OTA was found at a high level ($>1 \mu\text{g}/\text{kg}$) only in the speck
401 produced without the addition of yeasts. Inoculation of the FMCs with either *A. ochraceus*
402 or *P. nordicum* resulted in the production of OTA at a level that was higher than the limit
403 proposed by the Italian Ministry of Health ($1 \mu\text{g}/\text{kg}$). In the absence of yeasts, both of the
404 inoculated moulds grew on the FMCs from the late phase of drying to the end of the
405 seasoning phase. OTA production was detected in the meat. *P. nordicum* produced a white-
406 and-green cottoned slime (Photograph 2.1), whereas *A. ochraceus* produced a yellow
407 cottoned slime. Both cottoned slimes entirely covered the meat portion of the FMCs, but
408 no growth was observed on the skin of the FMCs. The co-inoculation of yeasts and moulds
409 did not permit mould growth on the FMCs. Therefore, the dominance of the inoculated
410 yeasts was the main parameter involved in the elimination of the growth of OTA-
411 producing moulds and consequently the elimination of OTA presence and production. In
412 the co-inoculated FMCs at the end of the seasoning phase, the amount of OTA was less
413 than the limit of detection of the method ($< 0.1 \mu\text{g}/\text{kg}$). The speck obtained with the starter

414 yeasts were safe and in accordance with the OTA limit proposed by the Italian Ministry of
415 Health (Circolare Ministero Sanità No. 30 10-09/06/1999).

416 Several researchers (Virgili et al., 2012; Andrade et al., 2014; Simoncini et al., 2014) have
417 demonstrated that inoculated and native yeasts are able to dominate *P. nordicum* and OTA-
418 producing moulds. Specifically, the antagonistic effects of some strains, such as *D.*
419 *hansenii*, *C. zeylanoides* and *H. burtonii*, are independent of the Aw, temperature and R.H.

420 However, these researchers recognized the existence of an antagonistic variability at the
421 level of the tested strains. Therefore, it was necessary to select the strain with the highest
422 antagonistic effect among the isolated species. This finding was confirmed in our study.

423 The significant absence of OTA appeared to be related to the inhibition of the growth of
424 both mould strains by the co-inoculated yeasts. No other hypotheses might be formulated.

425 In fact, no mould hyphae were observed on the surfaces of the co-inoculated specks under
426 a stereomicroscope. For this reason, the possibility that the absence of OTA in the co-
427 cultured samples might be due to mycotoxin degradation by the yeasts into less toxic
428 compounds was excluded (Andrade et al., 2014; Simoncini et al., 2014). Additionally, the
429 possibility that either of the yeasts could have adsorbed OTA on their cell wall was
430 excluded. Various researchers (Gil-Serna et al., 2011; Shetty et al., 2007) recently reported
431 that *S. cerevisiae* and *D. hansenii* could influence the regulation of mycotoxin
432 biosynthesis. The mechanisms involved in the reduction of OTA in the presence of yeasts
433 need to be clarified (Andrade et al., 2014). However, the results obtained by the
434 stereoscope observations made in this work demonstrate that the absence of OTA must be
435 due to the dominating effect of the co-inoculated yeasts over the OTA-producing moulds
436 (Photograph 2.2).

437 Despite the antagonistic effect observed in the speck model system at all of the Aw levels
438 tested, it is reasonable to propose that both yeasts should be inoculated after the smoking
439 phase and before the drying and seasoning steps, when the Aw is 0.95 ± 0.01 and the

440 moulds have not started to grow. The suggested concentration of the inoculum should be
441 10^6 CFU/cm² to absolutely ensure that the inoculum can predominate over the moulds.
442 Different researchers (Rodriguez et al., 1994; Andrade et al., 2014) have suggested that
443 yeast starter should be inoculated at the beginning of the production process for dry cured
444 ham (specifically at the end of the post-salting stage), when the *A_w* of the product is still
445 high (0.94) and can support OTA-producing mould growth. In our study, both of the
446 inoculated strains grew rapidly on the FMCs at this *A_w* value and blocked the growth of
447 OTA-producing moulds. Therefore, the biopreservative effect of the tested *D. hansenii* and
448 *S. fibuligera* strains was obvious. Consequently, both strains may be proposed as
449 antagonistic agents to prevent the presence of OTA-producing moulds and the
450 bioaccumulation of OTA during speck production.

451 The complete prevention of OTA in speck can be obtained by the application of yeast co-
452 inocula and of an adequate hygienic system based on Hazard Analysis and Critical Control
453 Points (HACCP). The HACCP system can reduce the level of OTA-producing mould
454 contamination on FMCs and speck and favour the antagonistic effect of both yeasts.

455 The sensorial acceptability of the speck inoculated with the starter cultures was confirmed
456 by a triangular test using a panel composed of 20 non-professional assessors. These
457 assessors did not find any difference between Lots A and B (with bioprotective cultures)
458 and Lot C (uninoculated control).

459 In conclusion, *D. hansenii* and *S. fibuligera* are potential biopreservative agents for
460 elimination of the growth of ochratoxigenic moulds in speck, a typical meat product of
461 North Italy and Austria. The use of selected *D. hansenii* and *S. fibuligera* starter cultures,
462 the control of raw meat and the technological (temperature and R.H.) and hygienic
463 parameters are fundamental for the reduction of health hazards due to the development of
464 OTA-producing moulds in dry-cured meat products, such as speck. Consequently, the

465 inoculation of *D. hansenii* or *S. fibuligera* strains after the smoking stage and during the
466 drying and seasoning phases could improve the safety and quality of speck.

467

468 **References**

469

470 Andrade, M.J., Rodriguez, A., Casado, E.M., Bermúdez, E., Cordoba, J.J., 2009.

471 Differentiation of yeasts growing on dry-cured meat Iberian ham by mitochondrial
472 DNA restriction analysis, RAPD-PCR and their volatile compounds production. *Food*
473 *Microbiology* 26, 578-586.

474 Andrade, M.J., Thorsen, L., Rodriguez, A., Cordoba, J.J., Jespersen, L., 2014. Inhibition of
475 ochratoxigenic moulds by *Debaryomyces hansenii* strains for biopreservation of dry-
476 cured meat products. *International Journal of Food Microbiology* 170, 70-77.

477 Asefa, D.T., Kure, C.F., Gjerde, R.O., Omer, M.K., Langrud, S., Nesbakken, T., Skarar, I.,
478 2010. Fungal growth pattern, sources and factors of mould contamination in a dry-cured
479 meat production facility. *International Journal of Food Microbiology* 140, 131-135.

480 Battilani, P., Formenti, S., Toscani, T., Virgili, R., 2010. Influence of abiotic parameters on
481 ochratoxin A production by a *P. nordicum* strain in dry-cured meat model systems. *Food*
482 *Control* 21, 1379-1744.

483 Battilani, P., Pietri, V. A., Giorni, P., Formenti, S., Bertuzzi, T., Toscani, T., Virgili, R., &
484 Kozakiewicz, Z., 2007. *Penicillium* populations in dry-cured ham manufacturing plants.
485 *Journal of Food Protection* 70, 975-980.

486 Björnberg, A., Schnürer, J., 1993. Inhibition of the growth of the grain-storage molds in
487 vitro by the yeast *Pichia anomala* (Hansen) Kurtzman. *Canadian Journal of*
488 *Microbiology* 39, 623–628.

489 Bleve, G., Grieco, F., Cozzi, G., Logrieco, A., Visconti, A., 2006. Isolation of epiphytic
490 yeasts with potential for biocontrol of *Aspergillus carbonarius* and *A. niger* on grape.
491 International Journal of Food Microbiology 108, 204-209.

492 Chi, Z., Chi, Z., Liu, G., Wang, F., Ju, L., Zhang, T. 2009. *Saccharomyces fibuligera*
493 and its applications in biotechnology. Biotechnology Advances 27 (4), 423-431.

494 Cocolin, L., Urso, R., Rantsiou, K., Cantoni, C., Comi, G., 2006. Dynamics and
495 characteriza- tion of yeasts during natural fermentation of Italian sausages. FEMS Yeast
496 Research 6, 692–701.

497 Comi, G., Cantoni, C., 1980. I lieviti di insaccati crudi stagionati. Industrie Alimentari, 19,
498 Novembre, 857-860.

499 Comi, G., Orlic, S., Redzepovic, S., Urso, R., Iacumin, L., 2004. Moulds isolated from
500 Istrian dried ham at the pre-seasoning and seasoning level. International Journal of Food
501 Microbiology 96, 29–34. [L]
[SEP]

502 Comi, G., Iacumin, L., 2013. Ecology of moulds during the pre-seasoning and seasoning of
503 San Daniele dry cured ham. Food Research International 54, 1113-1119.

504 Comi, G., Chiesa, L., Panseri, S., Orlic, S., Iacumin, L., 2013. Evaluation of different
505 methods to prevent *Penicillium nordicum* growth and Ochratoxin A (OTA) presence on
506 country-style sausages. World Mycotoxin Journal 6 (4), 411-418.

507 Commission of the European Communities 2006. Commission Regulation (EC)
508 No1881/2006 of 19 December 2006 setting maximum levels for certain contaminants in
509 foodstuffs. Official Journal of the European Union L364, 5–24. [L]
[SEP]

510 Dall'Asta, C., Galaverna, G., Bertuzzi, T., Moseriti, A., Pietri, A., Dossena, A., Marchelli,
511 R., 2010. Occurrence of ochratoxin A in raw ham muscle, salami and dry-cured ham
512 from pigs fed with contaminated diet. Food Chemistry 120, 978–983. [L]
[SEP]

513 Dragoni, I., Marino, C., & Cantoni, C., 1980). Muffe in prodotti carnei salati e stagionati
514 (bresaole e prosciutti crudi). Industrie Alimentari 19, 405-407.

515 EFSA Panel on Biological Hazards (BIOHAZ) 2012. Scientific opinion on the
516 maintenance of the list of QPS biological agents intentionally added to food and feed
517 (2012 up- date). EFSA Journal 10 (12), 3020.

518 Gil-Serna, J., Vázquez, C., Sardiñas, N., González-Jaén, M. T., Patiño, B., 2011. Revision
519 of ochratoxin A production capacity by the main species of *Aspergillus* section
520 *Circumdati*. *Aspergillus steynii* revealed as the main risk of OTA contamination. *Food*
521 *Control* 22, 343–345.

522 Hammes, W.P., Knauf H. J., 1994. Starters in the processing of meat products. *Meat*
523 *Science* 36 (1-2), 155-68.

524 Hernández-Montiel, L.G., Ochoa, J.L., Troyo-Diéguéz, E., Larralde-Corona, C.P., 2010.
525 Biocontrol of postharvest blue mold (*Penicillium italicum* Wehmer) on Mexican lime
526 by marine and citrus *Debaryomyces hansenii* isolates. *Postharvest Biology and*
527 *Technology* 56, 181–187.

528 Huerta, T., Sanchis, V., Hernandez, J., Hernandez, E., 1987. Mycoflora of dry-salted
529 Spanish ham. *Microbiologia Alimentare and Nutrition* 5, 247–252.

530 Iacumin, L., Milesi, S., Pirani, S., Comi, G., Chiesa, L. M. 2011. Ochratoxigenic moulds
531 and Ochratoxin A in Sausages from Different Areas in Northern Italy: Occurrence,
532 Elimination or Prevention with ozonated air. *Journal of Food Safety* 31, 538-545.

533 I.A.R.C., 1993. Ochratoxin A. In *Some naturally occurring substances: food items and*
534 *constituents, heterocyclic aromatic amines and mycotoxins*. IARC Monograph on the
535 *evolution of carcinogenic risks to humans* (vol. 56, pp. 489-521) Geneva: International
536 Agency for Research on Cancer.

537 Lima, G., Arru, S., DE Curtis, F., Arras, G., 1999. Influence of antagonistic, host fruit and
538 pathogen on the biological control of postharvest fungal diseases by yeasts. *Journal of*
539 *Industrial Microbiology and Biotechnology* 23, 223-229.

540 Marquina, D., Barroso, J., Santos, A., Peinado, J.M., 2001. Production and characteristics
541 of *Debaryomyces hansenii* killer toxin. Microbiological Research 156, 387–391.

542 Masoud, W., Poll, L., Jakobsen, M., 2005. Influence of volatile compounds produced by
543 yeasts predominant during processing of *Coffea arabica* in East Africa on growth and
544 ochratoxin A (OTA) production by *Aspergillus ochraceus*. Yeast 22, 1133-1142.

545 Matrella, R., Monaci, L., Milillo, M.A., Palmisano, F., Tantillo, M.G. (2006). Ochratoxin
546 A determination in paired kidneys and muscle samples from swines slaughtered in
547 southern Italy. Food Control 17, 114-117.

548 Multon, J. L., Bizot, H., 1980. Water adsorption in cereal foods and water activity
549 measurement. In Ingletti, G. E., Munck, L. (Eds.), Cereals for food and beverages (pp.
550 97-119). New York: Academic Press.

551 Núñez F., Rodríguez M.M., Bermudez M.E, Córdoba J.J., Asensio M.A., 1996.
552 Composition and toxigenic potential of the mould population on dry-cured Iberian ham.
553 International Journal of Food Microbiology 32, 185-187.

554 Peintner, U., Geiger, J., Pöder, R., 2000. The mycobiota of Speck, a traditional Tyrolean
555 smoked and cured ham. Journal of Food Protection 63(10), 1399-1403.

556 Pietri, A., Bertuzzi, T., Gualla, A., Piva, G., 2006. Occurrence of Ochratoxin A in raw ham
557 muscles and in pork products from Northern Italy. Italian Journal of Food Science
558 18(1), 99-106.

559 Rodríguez, M., Núñez, F., Córdoba, J.J., Sanabria, C., Bermúdez, E., Asensio, M.A., 1994.
560 Characterization of *Staphylococcus* spp. and *Micrococcus* spp. isolated from Iberian
561 ham throughout the seasoning process. International Journal Food Microbiology 24,
562 329-335.

563 Rodríguez, A., Rodríguez, M., Martín, A., Delgado, J., Córdoba, J.J., 2012. Presence of
564 ochratoxin A on the surface of dry-cured Iberian ham after initial fungal growth in the
565 drying stage. Meat Science 92, 728–734.

566 Rojas F. I., Jodral M., Gosalvez F., Pozo R. 1991. Mycoflora and toxigenic *Aspergillus*
567 *flavus* in Spanish dry-cured ham. International Journal of Food Microbiology 13, 249-
568 256.

569 Schmidt-Heydt, M., Richter, W., Michulec, M., Buttinger, G., Geisen, R., 2008.
570 Comprehensive molecular system to study the presence, growth and ochratoxin A bio-
571 synthesis of *Penicillium verrucosum* in wheat. Food Additives and Contaminants, Part
572 A, Chemistry Analysis, Control, Exposure and Risk Assessment, 25 (2), 989–996.

573 Shetty, P. H., Hald, B., Jespersen, L., 2007. Surface binding of aflatoxin B1 by
574 *Saccharomyces cerevisiae* strains with potential decontaminating abilities in indigenous
575 fermented foods. International Journal of Food Microbiology 113 (1), 41-46.

576 Simoncini, N., Rotelli, D., Virgili, R., & Quintavalla, S., 2007. Dynamics and
577 characterization of yeasts during seasoning of typical Italian dry-cured ham. Food
578 Microbiology 24 (6), 577-584.

579 Simoncini, N., Virgili, R., Spadola, G., Battilani, P., 2014. Autochthonous yeasts as
580 potential biocontrol agents in dry-cured meat products. Food Control 46, 160-167.

581 Spotti, E., Mutti, P., Campanini, M., 1989. Presenza di muffe sui prosciutti durante la
582 prestagionatura e la stagionatura: contaminazione degli ambienti e sviluppo sulla
583 porzione muscolare. Industria Conserve 64, 110-113.

584 Spotti E., Berni E., Cacchioli C. 2008. Characteristics and application of moulds. Meat
585 Biotechnology, Toldrà (ed.) 181-195.

586 Spotti, E., Berni, E., Cacchioli, C., Simoncini, N., Quintavalla, S., 2009. Growth and
587 antagonistic activity of *Hyphopichia burtonii* against other fungal species frequently
588 found on meat products during maturation. Industria delle Conserve 84, 11-22.

589 Sunesen, L.O., Stahnke, L. H., 2003. Mould starter cultures for dry sausages-selection,
590 application and effects. Meat Science 65, 935-948.

591 Virgili, R., Simoncini, N., Toscani, T., Leggieri, M.C., Formenti, S., Battilani, P., 2012.
592 Biocontrol of *Penicillium nordicum* growth and Ochratoxin A production by Native
593 Yeasts of Dry Cured Ham. *Toxins* 4 (2), 68-82.

594 Xu, X.M, Monger, W., Ritieni, A., Nicholson, P., 2007. Effect of temperature and duration
595 of wetness during initial periods on disease development, fungal biomass and mycotoxin
596 concentrations on wheat inoculated with single, or combinations of *Fusarium* species.
597 *Plant Pathology* 56, 943-956.

598 Zhao, Y., Tu, K., Shao, X.F., Jing, W., Su, Z.P., 2008. Effects of the yeast *Pichia*
599 *guilliermondii* against *Rhizopus nigricans* on tomato fruit. *Postharvest Biology*
600 *Technology* 49, 113–120.

601

602

603 **Conflict of interest**

604 None of the authors of this paper has a financial or personal relationship with other people
605 or organisations that could inappropriately influence or bias the content of the paper.

606

607

608

609

610

611

612

613

614

615

616

617

618

619

620
621
622
623
624
625

Table 1: Flowchart of speck production

Production phases	Time	Temperature	R.H. %
Raw meat -	24 h	1-7 °C	50-60
Trimming			
Brining	8 day	1-5 °C	75-88
Smoking	170 h	15-22 °C	70-80
Drying	1 month	10-20 °C	50-90
Seasoning	5 months	15-20 °C	50-90

Legend: R.H.: Relative Humidity.

626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644

645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695

Table 2: Inhibitory activity of *Saccharomycopsis fibuligera* vs OTA producing moulds co-inoculated in agar plate.

Starter yeast strain	% Inibitory activity on	
	<i>A. ochraceus</i>	<i>P. nordicum</i>
<i>S. fibuligera</i> DIAL 1	70.6 ± 0.7a	71.0 ± 0.3a
<i>S. fibuligera</i> DIAL 2	75.5 ± 0.7b	75.4 ± 0.2b
<i>S. fibuligera</i> DIAL 3	86.1 ± 0.3c	85.2 ± 0.7c
<i>S. fibuligera</i> DIAL 4	69.1 ± 0.6d	68.5 ± 0.7d
<i>S. fibuligera</i> DIAL 5	80.2 ± 0.1e	78.5 ± 0.2e
<i>S. fibuligera</i> DIAL 6	76.5 ± 0.3b	75.2 ± 0.2b

Legend: Data represent the means ± standard deviations of the total samples; Mean with the same letters within the same column (following the values) are not significantly differently (P< 0.05).

696
697
698
699
700
701
702
703

Table 3: Percentage of inhibitory activity of *Debaryomyces hansenii* vs OTA producing moulds co-inoculated in agar plates.

Starter yeast strain	% Inibitory activity on	
	<i>A. ochraceus</i>	<i>P. nordicum</i>
<i>D. hansenii</i> DIAL 1	76.0 ± 0.7a	78.8 ± 0.5a
<i>D. hansenii</i> DIAL 2	70.2 ± 0.3b	73.0 ± 0.7b
<i>D. hansenii</i> DIAL 3	70.3 ± 0.5b	72.2 ± 0.5b
<i>D. hansenii</i> DIAL 4	69.3 ± 0.7b	72.2 ± 0.3b
<i>D. hansenii</i> DIAL 5	68.0 ± 0.3c	72.0 ± 0.4b
<i>D. hansenii</i> DIAL 6	72.3 ± 0.5d	73.0 ± 0.5b

Legend: Data represent the means ± standard deviations of the total samples; Mean with the same letters within the same column (following the values) are not significantly differently ($p < 0.05$).

722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746

704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721

747
748
749

Table 4: Percentage of inhibitory effect of different concentration of yeasts vs OTA producing moulds co-inoculated in agar plates

Starter yeast strain	% Inibitory activity on	
	<i>A. ochraceus</i>	<i>P. nordicum</i>
<i>D. hansenii</i> DIAL 1		
10 ² CFU/mL	38.4 ± 0.7a	31.0 ± 0.2a
10 ⁴ CFU/mL	53.3 ± 0.5b	50.4 ± 0.3b
10 ⁶ CFU/mL	76.0 ± 0.7c	78.8 ± 0.5c
<i>S. fibuligera</i> DIAL 3		
10 ² CFU/mL	26.4 ± 0.2a	26.3 ± 0.5a
10 ⁴ CFU/mL	54.2 ± 0.3b	51.1 ± 0.5b
10 ⁶ CFU/mL	86.1 ± 0.3c	85.2 ± 0.7c

Legend: Data represent the means ± standard deviations of the total samples; Mean with the same letters within the same column (following the values) are not significantly differently ($p < 0.05$). *A. ochraceus* and *P. nordicum* concentration: 10⁶ conidia/mL.

750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774

775
776
777
778
779
780
781
782
783

Table 5: Percentage of inhibitory effect of yeasts vs different concentration of OTA producing moulds co-inoculated in agar plates

Starter yeast strain	% Inibitory activity on	
	<i>D. hansenii</i>	<i>S. fibuligera</i>
<i>P. nordicum</i>		
10 ² CFU/mL	76.2 ± 0.3a	79.3 ± 0.1a
10 ⁴ CFU/mL	55.4 ± 0.4b	59.0 ± 0.2b
10 ⁶ CFU/mL	38.5 ± 0.3c	42.5 ± 0.9c
<i>A. ochraceus</i>		
10 ² CFU/mL	78.5 ± 0.5a	85.0 ± 0.6a
10 ⁴ CFU/mL	60.3 ± 0.4b	65.6 ± 0.2b
10 ⁶ CFU/mL	42.0 ± 0.3c	48.0 ± 0.2c

Legend: Data represent the means ± standard deviations of the total samples; Mean with the same letters within the same column (following the values) are not significantly differently ($p < 0.05$). *D. hansenii* and *S. fibuligera* concentration: 10⁴ CFU/mL.

784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802

803
804
805
806
807
808
809
810
811

Table 6: Mean production of Ochratoxin A by *P. nordicum* and *A. ochraceus* in control and selected yeast co-inoculated in speck model system with different Aw

Ochratoxin A (µg/kg) in Speck						
Aw	<i>P. nordicum</i> Control	<i>P. nordicum</i> / <i>S. fibuligera</i>	<i>P. nordicum</i> / <i>D. hansenii</i>	<i>A.</i> <i>ochraceus</i> Control	<i>A.</i> <i>ochraceus</i> / <i>S. fibuligera</i>	<i>A.</i> <i>ochraceus</i> / <i>D. hansenii</i>
0.96	7.7	0.4	0.3	8.9	0.7	0.7
0.94	5.6	0.3	0.2	6.4	< 0.1	< 0.1
0.92	3.8	0.2	0.2	3.9	0.7	0.3
0.90	2.9	0.7	0.3	2.5	0.4	0.3
0.88	1.9	0.2	0.3	2.0	0.5	0.3

Legend: OTA Mean µg/kg.; LOD < 0.1 µg/kg; Samples were represented by meat pieces of 50 mm diameter and 5 mm in height.

812
813

Table 7: Mean production of Ochratoxin A by *Penicillium nordicum* and *Aspergillus ochraceus* in control and selected yeast co-inoculated after smoking stage and valued during the seasoning of speck.

814

Toxigenic moulds	Ochratoxin A (µg/kg) in Speck			
	<i>Speck inoculated with S. fibuligera</i>	<i>Speck inoculated with D. hansenii</i>	Control <i>A. ochraceus</i>	Control <i>P. nordicum</i>
<i>A. ochraceus</i>	< 0.1	< 0.1	65.5 ± 1.5	-
<i>P. nordicum</i>	< 0.1	< 0.1	-	68.1 ± 1.9

Data: OTA mean ± standard deviations of 20 replicates: µg/kg.; LOD < 0.1 µg/kg; Samples taken from a depth of 0.5 cm below the surface; Inoculum after smoking.

815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843

844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860

Photograph 1: Competition yeasts versus OTA producing moulds





861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879

Legend: 1-*D. hansenii* vs *P. nordicum*; 2-*E. fibuligera* vs *P. nordicum*
3-*D. hansenii* vs *A. ochraceus*; 4-*E. fibuligera* vs *A. ochraceus*

Photograph 2: Mould growth on speck without (1) and with yeast added (2)



Legend: 1 - *P. nordicum*; 2 - *D. hansenii* vs *P. nordicum*

880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896

898
899
900
901
902

Photograph 1: Competition yeasts versus OTA producing moulds





903
904 Legend: 1-*D. hansenii* vs *P. nordicum*; 2-*E. fibuligera* vs *P. nordicum*
905 3-*D. hansenii* vs *A. ochraceus*; 4-*E. fibuligera* vs *A. ochraceus*
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922

923
924
925

Photograph 2: Mould growth on speck without (1) and with yeast added (2)



926
927
928
929
930
931
932
933

Legend: 1 - *P. nordicum*; 2 - *D. hansenii* vs *P. nordicum*