



UNIVERSITÀ
DEGLI STUDI
DI PADOVA

Università degli Studi di Padova

Padua Research Archive - Institutional Repository

Derived equivalences induced by nonclassical tilting objects

Original Citation:

Availability:

This version is available at: 11577/3216497 since: 2017-01-30T15:29:25Z

Publisher:

Published version:

DOI: 10.1090/proc/13368

Terms of use:

Open Access

This article is made available under terms and conditions applicable to Open Access Guidelines, as described at <http://www.unipd.it/download/file/fid/55401> (Italian only)

(Article begins on next page)

DERIVED EQUIVALENCES INDUCED BY NONCLASSICAL TILTING OBJECTS

LUISA FIOROT, FRANCESCO MATTIELLO, AND MANUEL SAORÍN

ABSTRACT. Suppose that \mathcal{A} is an abelian category whose derived category $\mathcal{D}(\mathcal{A})$ has *Hom* sets and arbitrary (small) coproducts, let T be a (not necessarily classical) $(n-)$ tilting object of \mathcal{A} and let \mathcal{H} be the heart of the associated t -structure on $\mathcal{D}(\mathcal{A})$. We show that there is a triangulated equivalence of unbounded derived categories $\mathcal{D}(\mathcal{H}) \xrightarrow{\cong} \mathcal{D}(\mathcal{A})$ which is compatible with the inclusion functor $\mathcal{H} \hookrightarrow \mathcal{D}(\mathcal{A})$. The result admits a straightforward dualization to cotilting objects in abelian categories whose derived category has *Hom* sets and arbitrary products.

INTRODUCTION

Tilting theory first appeared in the eighties of last century in the context of Representation Theory of finite dimensional algebras (see [BB80], [HR82], [Bo81]) as a generalization of Morita Theory. Instead of the classical equivalence of module categories given by the theorem of Morita, for a tilting A -module T , which was always finite dimensional in these seminal papers, the so-called Brenner-Butler theorem gave a counter-equivalence between the torsion pairs $(\text{Ker}(\text{Ext}_A^1(T, ?)), \text{Ker}(\text{Hom}(T, ?)))$ and $(\text{Ker}(? \otimes_B T), \text{Ker}(\text{Tor}_1^B(? , T)))$ of $\text{mod} - A$ and $\text{mod} - B$, respectively, where $B = \text{End}(T_A)$ is the endomorphism algebra and mod denotes the category of finite dimensional modules. This theorem was extended in [CF90] by considering A to be an arbitrary ring, T to be a finitely presented A -module and the module categories over A and B to be those of all modules, not just the finitely presented ones. By definition, in all these papers tilting modules had projective dimension less or equal than one. A generalization of the concept to modules of finite projective dimension greater than one was given in [M86], where the new tilting modules had a finite projective resolution with finitely generated terms (we will call these modules strongly finitely presented). The substitute of Brenner-Butler theorem in this context was a series of equivalences of full subcategories $\text{Ker}(\text{Ext}_A^k(T, ?)) \xleftarrow{\cong} \text{Ker}(\text{Tor}_k^B(? , T))$, for all integers $k \in \mathbb{Z}$.

Up to here, even working over arbitrary rings, tilting modules were assumed to be strongly finitely presented. The historical development brought a new notion

2010 *Mathematics Subject Classification.* Primary 18E30; Secondary, 18E10, 18G55.

Mattiello is supported by Assegno di ricerca “Tilting theory in triangulated categories” del Dipartimento di Matematica dell’Università degli Studi di Padova and by Progetto di Eccellenza della fondazione Cariparo.

Saorín is supported by research projects from the Spanish Ministerio de Economía y Competitividad (MTM2013-46837-P) and from the Fundación ‘Séneca’ of Murcia (19880/GERM/15), with a part of FEDER funds.

The authors thank their institutions for their help. They also thank Jorge Vitória and Alexandra Zvonareva for pointing out some omissions in a first draft of the paper.

of tilting module, where finite projective dimension was still required but not the strongly finitely presented condition. This was first done in [CT95], for modules of projective dimension less or equal than one, and in [AC01] for modules of arbitrary finite projective dimension. Since then the most commonly used terminology reserves the term *classical (n-)tilting module* for the strongly finitely presented modules of finite projective dimension (less or equal than n), while the term *(n-)tilting module* applies to all modules, not just the strongly finitely presented ones.

Even from the beginning, tilting theory expanded from studying tilting modules to study tilting objects in more general abelian categories, and this extension has played a fundamental role in several branches of Mathematics, as Algebraic Geometry and Representation Theory of Groups and Algebras (see [AHK07]).

Soon after their introduction, it became clear that tilting modules and tilting objects had to do with derived categories. In [CPS86] and [H87] (see also [H88]), the authors proved that any classical tilting module gives an equivalence of bounded derived categories $\mathcal{D}^b(A) \xrightarrow{\cong} \mathcal{D}^b(B)$. The general Morita theory for derived categories, developed immediately after by Rickard and Keller (see [Ri89] and [Ke94]), showed that this equivalence actually extended to one between the corresponding unbounded derived categories. It was then the turn for nonclassical tilting modules to be related to derived categories. This was first done implicitly in [HRS96], within the study of quasi-tilted algebras. The authors proved that each torsion pair $\mathbf{t} = (\mathcal{T}, \mathcal{F})$ in an abelian category \mathcal{A} defined a t-structure in $\mathcal{D}^b(\mathcal{A})$ (and also in $\mathcal{D}(\mathcal{A})$), in the sense of [BBD82], whose heart \mathcal{H} admitted $(\mathcal{X}, \mathcal{Y}) := (\mathcal{F}[1], \mathcal{T})$ as a torsion pair (counterequivalent to \mathbf{t}). Then they showed that if either \mathcal{T} was a cogenerating or \mathcal{F} was a generating class, then the inclusion $\mathcal{H} \hookrightarrow \mathcal{D}^b(\mathcal{A})$ extended to a triangulated equivalence $\mathcal{D}^b(\mathcal{H}) \xrightarrow{\cong} \mathcal{D}^b(\mathcal{A})$. This result can be then applied when $\mathcal{A} = \text{Mod} - A$ and the torsion pair is either $(\text{Ker}(\text{Ext}_A^1(T, ?)), \text{Ker}(\text{Hom}(T, ?)))$, the one associated to any 1-tilting module, or $(\text{Ker}(\text{Hom}_A(? , Q)), \text{Ker}(\text{Ext}_A^1(? , Q)))$, the one associated to a 1-cotilting module (defined dually). Furthermore, the module T is a projective generator of \mathcal{H} , which is a progenerator when T is classical 1-tilting (see [BR07, III.4]). Therefore in this latter case \mathcal{H} is equivalent to $\text{Mod} - B$ and, as a consequence, the equivalence of [CPS86] or [H87] for classical 1-tilting modules is just a particular case of the mentioned result of [HRS96]. For another example of the relationship between nonclassical tilting modules and derived equivalences, we refer to [BMT11].

When T is an arbitrary n -tilting module, we no longer have a torsion pair in $\text{Mod} - A$, although the pair $\tau_T = (T^{\perp > 0}, T^{\perp < 0})$ in $\mathcal{D}(A)$ (or $\mathcal{D}^b(A)$) is still a t-structure (see, for example [NSZ15]). The mentioned result of [HRS96] naturally leads to two questions. The first one would ask whether the equivalence $\mathcal{D}^b(\mathcal{H}) \cong \mathcal{D}^b(A)$ for 1-tilting modules also holds at the unbounded level. The second one would ask whether this equivalence (at the bounded or unbounded level) also holds when we replace 1-tilting (resp. 1-cotilting) by n -tilting (n -cotilting), for n arbitrary, and \mathcal{H} is the heart of τ_T . The first question was answered in the affirmative by Xiao Wu Chen [Ch10], for any 1-tilting object in an arbitrary abelian category. As for the second question, an affirmative answer in the case of a classical n -tilting object in a Grothendieck category, both in the bounded and unbounded ed situations, follows from [FMT14]. When T is a nonclassical tilting object, the second question has an affirmative answer at the bounded level, in essentially any AB3 abelian category, due to a stronger recent result of Psaroudakis and Vitória (see

[PV15][Corollary 5.2]). The main goal of this short note is to prove that the second question has an affirmative answer also at the unbounded level for any (n -)tilting object in essentially any AB3(=cocomplete) abelian category \mathcal{A} . More concretely, the following result and its dual will be direct consequences of our main theorem (see Theorem 1.7):

Proposition 2.3. *Let \mathcal{A} be an abelian category such that its derived category $\mathcal{D}(\mathcal{A})$ has Hom sets and arbitrary (small) coproducts, let T be a tilting object in \mathcal{A} and let \mathcal{H} be the heart of the associated t -structure $\tau_T = (T^{\perp > 0}, T^{\perp < 0})$ in $\mathcal{D}(\mathcal{A})$. Then there is a triangulated equivalence $\mathcal{D}(\mathcal{H}) \xrightarrow{\cong} \mathcal{D}(\mathcal{A})$ which is compatible with the inclusion functor $\mathcal{H} \hookrightarrow \mathcal{D}(\mathcal{A})$.*

In this statement and in the rest of the paper, the compatibility with the inclusion $\mathcal{H} \hookrightarrow \mathcal{D}(\mathcal{A})$ means that the restriction of the equivalence to \mathcal{H} is naturally isomorphic to the inclusion functor.

The organization of the paper goes as follows. In Section 1, we recall some basic fact about t -structures in triangulated categories and we prove the Tilting Theorem 1.7. Here the setup is very general. We consider a t -structure $\tau = (\mathcal{U}, \mathcal{U}^{\perp}[1])$ in $\mathcal{D}(\mathcal{A})$ such that $\mathcal{D}^{\leq -n}(\mathcal{A}) \subseteq \mathcal{U} \subseteq \mathcal{D}^{\leq 0}(\mathcal{A})$ and consider the class $\mathcal{Y} := \mathcal{A} \cap \mathcal{H}_\tau$, where \mathcal{H}_τ is the heart of τ . Theorem 1.7 states that if \mathcal{Y} is cogenerating in \mathcal{A} , then there is a triangulated equivalence $\mathcal{D}(\mathcal{H}_\tau) \xrightarrow{\cong} \mathcal{D}(\mathcal{A})$ which is compatible with the inclusion functor $\mathcal{H}_\tau \hookrightarrow \mathcal{D}(\mathcal{A})$. In Section 2 we apply the Tilting Theorem to nonclassical tilting objects. Firstly, we give the definition of (nonclassical) tilting object in any abelian category as in the statement and we give characterizations suitable for our purposes. Both, definition and characterizations, are taken from [NSZ15]. Then we show that the above Proposition 2.3 and its dual (see Proposition 2.4, which includes a recent result of Stovicek about derived equivalences induced by n -cotilting modules) are direct consequences of the Tilting Theorem. Finally, as a byproduct, we prove that a (n -)tilting object as in the situation of Proposition 2.3 is self-small if and only if it is classical tilting (Corollary 2.5). Except for the case $n = 1$ (see [CT95][Proposition 1.3]), this fact seems to be unknown even for tilting modules.

1. TILTING THEOREM

Let \mathcal{D} be an additive category. For any full subcategory \mathcal{D}' of \mathcal{D} we denote by ${}^{\perp}\mathcal{D}'$ the following full subcategory of \mathcal{D} :

$${}^{\perp}\mathcal{D}' := \{X \in \mathcal{D} \mid \text{Hom}_{\mathcal{D}}(X, Y) = 0, \text{ for all } Y \in \mathcal{D}'\},$$

and by \mathcal{D}'^{\perp} the following full subcategory of \mathcal{D} :

$$\mathcal{D}'^{\perp} := \{X \in \mathcal{D} \mid \text{Hom}_{\mathcal{D}}(Y, X) = 0, \text{ for all } Y \in \mathcal{D}'\}.$$

Let \mathcal{D} be a triangulated category and denote its suspension functor by $[1]$. If \mathcal{U}, \mathcal{V} are full subcategories of \mathcal{D} , then we denote by $\mathcal{U} \star \mathcal{V}$ the category of extensions of \mathcal{V} by \mathcal{U} , that is, the full subcategory of \mathcal{D} consisting of objects X which may be included in a distinguished triangle $U \rightarrow X \rightarrow V \xrightarrow{+}$ in \mathcal{D} , with $U \in \mathcal{U}$ and $V \in \mathcal{V}$.

Recall that a t -structure in \mathcal{D} is a pair $\tau = (\mathcal{U}, \mathcal{V})$ of full additive subcategories of \mathcal{D} which satisfy the following properties:

- (i) $\text{Hom}_{\mathcal{D}}(U, V[-1]) = 0$, for all $U \in \mathcal{U}$ and $V \in \mathcal{V}$.
- (ii) $\mathcal{U}[1] \subseteq \mathcal{U}$.

(iii) $\mathcal{D} = \mathcal{U} \star \mathcal{V}[-1]$, that is, for each object X of \mathcal{D} , there is a distinguished triangle $U \rightarrow X \rightarrow V \xrightarrow{\pm}$ in \mathcal{D} , where $U \in \mathcal{U}$ and $V \in \mathcal{V}[-1]$.

In such case one has $\mathcal{V} = \mathcal{U}^\perp[1]$ and $\mathcal{U} = {}^\perp(\mathcal{V}[-1]) = {}^\perp(\mathcal{U}^\perp)$. Thus, we may write a t -structure as $\tau = (\mathcal{U}, \mathcal{U}^\perp[1])$. The objects U and V in the above triangle are uniquely determined by X , up to isomorphism, and define functors $\tau_{\mathcal{U}}: \mathcal{D} \rightarrow \mathcal{U}$ and $\tau^{\mathcal{U}^\perp}: \mathcal{D} \rightarrow \mathcal{U}^\perp$ which are right and left adjoints to the respective inclusion functors. The full subcategory $\mathcal{H}_\tau = \mathcal{U} \cap \mathcal{V} = \mathcal{U} \cap \mathcal{U}^\perp[1]$ is called the *heart* of the t -structure and it is an abelian category, where the short exact sequences “are” the triangles in \mathcal{D} with their three terms in \mathcal{H}_τ . The assignments $X \rightsquigarrow (\tau_{\mathcal{U}} \circ \tau^{\mathcal{U}^\perp[1]})(X)$ and $X \rightsquigarrow (\tau^{\mathcal{U}^\perp[1]} \circ \tau_{\mathcal{U}})(X)$ define naturally isomorphic functors $\mathcal{D} \rightarrow \mathcal{H}$ which are cohomological (see [BBD82]). In the sequel, we will use the notation $\tau^{[a,b]} = \mathcal{U}[-b] \cap \mathcal{U}^\perp[1-a]$, where $a \leq b$ are integers. Given an abelian category \mathcal{A} , its derived category $\mathcal{D}(\mathcal{A})$ is a triangulated category which admits a t -structure $\delta = (\mathcal{D}(\mathcal{A})^{\leq 0}, \mathcal{D}(\mathcal{A})^{\geq 0})$, called the *natural t -structure*, where $\mathcal{D}(\mathcal{A})^{\leq 0}$ (resp. $\mathcal{D}(\mathcal{A})^{\geq 0}$) is the full subcategory of complexes without cohomology in positive (resp. negative) degrees. The heart of δ is (naturally equivalent to) \mathcal{A} .

For the rest of this section, we make the following assumptions and set the following notation:

Setup 1. \mathcal{A} is any abelian category such that $\mathcal{D}(\mathcal{A})$ has *Hom* sets. We denote by $\delta = (\mathcal{D}(\mathcal{A})^{\leq 0}, \mathcal{D}(\mathcal{A})^{>0}[1])$ the natural t -structure on $\mathcal{D}(\mathcal{A})$ and by $\tau = (\mathcal{U}, \mathcal{U}^\perp[1])$ another t -structure on $\mathcal{D}(\mathcal{A})$, whose heart is \mathcal{H}_τ . We shall assume that

$$\mathcal{D}(\mathcal{A})^{\leq -n} \subseteq \mathcal{U} \subseteq \mathcal{D}(\mathcal{A})^{\leq 0} \quad (\text{equivalently, } \mathcal{D}(\mathcal{A})^{>0} \subseteq \mathcal{U}^\perp \subseteq \mathcal{D}(\mathcal{A})^{> -n}).$$

Consider the class $\mathcal{Y} = \mathcal{A} \cap \mathcal{H}_\tau$ in the sequel. We denote by $\mathcal{K}_{ac}^{\mathcal{A}}(\mathcal{Y})$ (resp. $\mathcal{K}_{ac}^{\mathcal{H}_\tau}(\mathcal{Y})$) the additive full subcategory of the homotopy category $\mathcal{K}(\mathcal{Y})$ of cochain complexes on \mathcal{Y} whose objects are the complexes of $\mathcal{K}(\mathcal{Y})$ which are acyclic in \mathcal{A} (resp. in \mathcal{H}_τ).

Lemma 1.1. *Let s be a natural number and let Y^\bullet be the complex*

$$\dots \longrightarrow 0 \longrightarrow Y^{-s} \xrightarrow{d^{-s}} Y^{-s+1} \longrightarrow \dots \longrightarrow Y^{-1} \xrightarrow{d^{-1}} Y^0 \longrightarrow 0 \longrightarrow \dots$$

where Y_i is an object of \mathcal{Y} , for every $i = -s, \dots, -1, 0$. Then $Y^\bullet \in \delta^{[-s,0]} \cap \tau^{[-s,0]}$.

Proof. The complex Y^\bullet belongs to the following full subcategory of $\mathcal{D}(\mathcal{A})$:

$$(\mathcal{D}(\mathcal{A})^{>0}[1] \cap \mathcal{U})[0] \star (\mathcal{D}(\mathcal{A})^{>0}[1] \cap \mathcal{U})[1] \star \dots \star (\mathcal{D}(\mathcal{A})^{>0}[1] \cap \mathcal{U})[s-1] \star (\mathcal{D}(\mathcal{A})^{>0}[1] \cap \mathcal{U})[s],$$

The latter subcategory is contained in $\mathcal{D}(\mathcal{A})^{>0}[1+s] \cap \mathcal{U}$. By the assumption made in Setup 1, $\mathcal{U} \subseteq \mathcal{D}(\mathcal{A})^{\leq 0}$ and $\mathcal{D}(\mathcal{A})^{>0}[1+s] \subseteq \mathcal{U}^\perp[1+s]$. From this we conclude that Y^\bullet belongs to

$$(\mathcal{D}(\mathcal{A})^{\leq 0} \cap \mathcal{D}(\mathcal{A})^{>0}[1+s]) \cap (\mathcal{U} \cap \mathcal{U}^\perp[1+s]) = \delta^{[-s,0]} \cap \tau^{[-s,0]}.$$

□

Lemma 1.2. *Let $Y^{-n} \xrightarrow{f_{-n}} Y^{-n+1} \rightarrow \dots \rightarrow Y^{-1} \xrightarrow{f_{-1}} Y^0$ be a sequence of morphisms in \mathcal{Y} such that $f_{-k} \circ f_{-k-1} = 0$, for every $k = 1, \dots, n-1$, where n is the natural number such that $\mathcal{D}(\mathcal{A})^{\leq -n} \subseteq \mathcal{U} \subseteq \mathcal{D}(\mathcal{A})^{\leq 0}$. The following assertions hold.*

- (1) *If the sequence is exact in \mathcal{H}_τ , then $\text{Ker}_{\mathcal{H}_\tau} f_{-n} \in \mathcal{Y}$.*
- (2) *If the sequence is exact in \mathcal{A} , then $\text{Coker}_{\mathcal{A}} f_{-1} \in \mathcal{Y}$.*

Proof. Let Y^\bullet be the complex in $\mathcal{D}(\mathcal{A})$ obtained from the sequence $Y^{-n} \xrightarrow{f_{-n}^-} Y^{-n+1} \rightarrow \dots \rightarrow Y^{-1} \xrightarrow{f_{-1}^-} Y^0$ by completing with zeros on both sides. By Lemma 1.1, $Y^\bullet \in \delta^{[-n,0]} \cap \tau^{[-n,0]}$.

1) For every $k = 0, 1, \dots, n$ there is a distinguished triangle in $\mathcal{D}(\mathcal{A})$

$$\mathrm{Ker}_{\mathcal{H}_\tau}(f_{-k}) \longrightarrow Y^{-k} \longrightarrow \mathrm{Ker}_{\mathcal{H}_\tau}(f_{-k+1}) \xrightarrow{+}$$

(with the convention that $f_0: Y^0 \rightarrow L := \mathrm{Coker}_{\mathcal{H}_\tau}(f_{-1}) = \mathrm{Ker}_{\mathcal{H}_\tau}(f_1)$ is the cokernel morphism of f_{-1}). By an iterated use of the octahedral axiom, we obtain the following distinguished triangle in $\mathcal{D}(\mathcal{A})$:

$$\mathrm{Ker}_{\mathcal{H}_\tau}(f_{-n})[n] \longrightarrow Y^\bullet \longrightarrow L \xrightarrow{+}$$

thus by shifting:

$$\mathrm{Ker}_{\mathcal{H}_\tau}(f_{-n}) \longrightarrow Y^\bullet[-n] \longrightarrow L[-n] \xrightarrow{+}$$

Now, $Y^\bullet[-n] \in \mathcal{D}(\mathcal{A})^{>0}[1]$ and $L[-n] \in \mathcal{H}_\tau[-n] \subseteq \mathcal{D}(\mathcal{A})^{>0}[1]$. Thus $\mathrm{Ker}_{\mathcal{H}_\tau}(f_{-n}) \in \mathcal{D}(\mathcal{A})^{>0}[1]$. Therefore $\mathrm{Ker}_{\mathcal{H}_\tau}(f_{-n}) \in \mathcal{D}(\mathcal{A})^{\leq 0} \cap \mathcal{D}(\mathcal{A})^{>0}[1] = \mathcal{A}$ since $\mathcal{H}_\tau \subseteq \mathcal{D}(\mathcal{A})^{\leq 0}$. It follows that $\mathrm{Ker}_{\mathcal{H}_\tau}(f_{-n}) \in \mathcal{A} \cap \mathcal{H}_\tau = \mathcal{Y}$.

2) This is the symmetric argument of the previous one, using the fact that $Y^\bullet \in \tau^{[-n,0]}$. \square

Proposition 1.3. *Let Y^\bullet be a complex of objects in \mathcal{Y} . The following are equivalent:*

- (1) Y^\bullet is acyclic in \mathcal{A} .
- (2) Y^\bullet is acyclic in \mathcal{H}_τ .

Proof. 1) \implies 2) The complex Y^\bullet is obtained by gluing the following short exact sequences in \mathcal{A} :

$$(1.1) \quad 0 \longrightarrow \mathrm{Coker}_{\mathcal{A}}(d_{Y^\bullet}^{k-2}) \longrightarrow Y^k \longrightarrow \mathrm{Coker}_{\mathcal{A}}(d_{Y^\bullet}^{k-1}) \longrightarrow 0, \quad k \in \mathbb{Z}.$$

Now the sequences $Y^{k-n} \xrightarrow{d_{Y^\bullet}^{k-n}} Y^{k-n+1} \rightarrow \dots \rightarrow Y^k$, $k \in \mathbb{Z}$, satisfy the hypothesis of Lemma 1.2, 2), hence $\mathrm{Coker}_{\mathcal{A}}(d_{Y^\bullet}^{k-1}) \in \mathcal{Y}$, for all $k \in \mathbb{Z}$. Then we have distinguished triangles in $\mathcal{D}(\mathcal{A})$

$$\mathrm{Coker}_{\mathcal{A}}(d_{Y^\bullet}^{k-2}) \longrightarrow Y^k \longrightarrow \mathrm{Coker}_{\mathcal{A}}(d_{Y^\bullet}^{k-1}) \xrightarrow{+}, \quad k \in \mathbb{Z}.$$

with all the terms in \mathcal{H}_τ . From this we can conclude that the sequences (1.1) are short exact in \mathcal{H}_τ and hence that Y^\bullet is acyclic in \mathcal{H}_τ .

2) \implies 1) This is just a symmetric argument using Lemma 1.2, 1). \square

In order to construct resolutions of complexes, we will need the following two auxiliary results and their duals, which hold true since the opposite category of an abelian category is again abelian.

Lemma 1.4. [KS05, Lemma 13.2.1] *Let \mathcal{A} be an abelian category, let \mathcal{I} be an additive full subcategory of \mathcal{A} and let $X \in \mathcal{C}^{\geq a}(\mathcal{A})$ for some $a \in \mathbb{Z}$. Assume that \mathcal{I} is cogenerating in \mathcal{A} . Then there exist $Y \in \mathcal{C}^{\geq a}(\mathcal{I})$ and a quasi-isomorphism $X \rightarrow Y$.*

Proposition 1.5. [KS05, Proposition 13.2.6] *Let \mathcal{A} be an abelian category and let \mathcal{I} be an additive full subcategory of \mathcal{A} . Assume that:*

- (i) \mathcal{I} is cogenerating,
- (ii) there exists a non-negative integer d such that, for any exact sequence $Y^d \rightarrow \dots \rightarrow Y^1 \rightarrow Y \rightarrow 0$, with $Y^j \in \mathcal{I}$ for $1 \leq j \leq d$, we have $Y \in \mathcal{I}$.

Then for any complex $X \in C(\mathcal{A})$, there exist $Y \in C(\mathcal{I})$ and a quasi-isomorphism $X \rightarrow Y$. In particular, there is an equivalence of triangulated categories

$$K(\mathcal{I})/\mathcal{K}_{ac}^{\mathcal{A}}(\mathcal{I}) \xrightarrow{\cong} \mathcal{D}(\mathcal{A}),$$

where $\mathcal{K}_{ac}^{\mathcal{A}}(\mathcal{I})$ is the class of complexes on \mathcal{I} which are acyclic in \mathcal{A} .

Lemma 1.6. *If the class $\mathcal{Y} = \mathcal{A} \cap \mathcal{H}_\tau$ is cogenerating in \mathcal{A} , then \mathcal{Y} is generating in \mathcal{H}_τ .*

Proof. Let us suppose that \mathcal{Y} cogenerates \mathcal{A} and let us prove that \mathcal{Y} generates \mathcal{H}_τ . Let $B \in \mathcal{H}_\tau$. First we show that we can represent the object B as a complex

$$Y^\bullet := \dots \rightarrow 0 \rightarrow Y^{-n} \xrightarrow{d^{-n}} Y^{-n+1} \xrightarrow{d^{-n+1}} \dots \xrightarrow{d^{-2}} Y^{-1} \xrightarrow{f} A \rightarrow 0 \rightarrow \dots$$

whose terms Y_i lie in \mathcal{Y} for any $-n \leq i \leq -1$, $A \in \mathcal{A}$ and it is placed in degree zero. Indeed, since $\mathcal{H}_\tau \subseteq \delta^{[-n,0]}$, Lemma 1.4 gives that B is quasi-isomorphic to a complex $\dots \rightarrow 0 \rightarrow Y^{-n} \rightarrow \dots \rightarrow Y^0 \rightarrow Y^1 \rightarrow \dots$, with Y^i in \mathcal{Y} for all $i \geq -n$. Then using the classical truncation at 0 by taking $A = \text{Ker}_{\mathcal{A}}(Y^0 \rightarrow Y^1)$, gives a complex $\dots \rightarrow 0 \rightarrow Y^{-n} \rightarrow \dots \rightarrow Y^{-1} \rightarrow A \rightarrow 0 \rightarrow \dots$, which is clearly quasi-isomorphic to Y^\bullet , and hence isomorphic to B in $\mathcal{D}(\mathcal{A})$.

Now let us consider the following distinguished triangle in $\mathcal{D}(\mathcal{A})$ (see on the left in the vertical sense the distinguished triangle):

$$\begin{array}{ccccccccccc}
C & & 0 & \longrightarrow & Y^{-n} & \longrightarrow & Y^{-n+1} & \longrightarrow & \dots & \longrightarrow & Y^{-2} & \longrightarrow & Y^{-1} \\
\downarrow & & \downarrow & & \downarrow & & \downarrow & & \downarrow & & \downarrow & & \downarrow f \\
A[0] & & 0 & \longrightarrow & 0 & \longrightarrow & 0 & \longrightarrow & 0 & \longrightarrow & 0 & \longrightarrow & A \\
\downarrow & & \downarrow & & \downarrow & & \downarrow & & \downarrow & & \downarrow & & \downarrow \\
B & & Y^{-n} & \longrightarrow & Y^{-n+1} & \longrightarrow & \dots & \longrightarrow & \dots & \longrightarrow & Y^{-1} & \longrightarrow & A.
\end{array}$$

By Lemma 1.1, we have $C \in \delta^{[-n,0]} \cap \tau^{[-n,0]}$. Since $B \in \mathcal{H}_\tau$ we deduce that $A \in \mathcal{A} \cap \mathcal{U} = \mathcal{Y}$. Now $A, B \in \mathcal{H}_\tau$ hence $C \in \mathcal{U}^\perp[1] \cap \tau^{[-n,0]} = \mathcal{H}_\tau$. We can conclude that \mathcal{Y} generates \mathcal{H}_τ since for any $B \in \mathcal{H}_\tau$ we have found $A \in \mathcal{Y}$ and the previous distinguished triangle proves that $0 \rightarrow C \rightarrow A \rightarrow B \rightarrow 0$ is an exact sequence in \mathcal{H}_τ . \square

Theorem 1.7 (Tilting Theorem). *Let \mathcal{A} be an abelian category such that $\mathcal{D}(\mathcal{A})$ has Hom sets and let $\tau = (\mathcal{U}, \mathcal{U}^\perp[1])$ be a t -structure in $\mathcal{D}(\mathcal{A})$ such that $\mathcal{D}^{\leq -n}(\mathcal{A}) \subseteq \mathcal{U} \subseteq \mathcal{D}^{\leq 0}(\mathcal{A})$, for some $n \in \mathbb{N}$, and let \mathcal{H}_τ be its heart. If the class $\mathcal{Y} = \mathcal{A} \cap \mathcal{H}_\tau$ cogenerates \mathcal{A} , then there is a triangulated equivalence*

$$\mathcal{D}(\mathcal{H}_\tau) \xrightarrow{\cong} \mathcal{D}(\mathcal{A})$$

which is compatible with the inclusion functor $\mathcal{H}_\tau \hookrightarrow \mathcal{D}(\mathcal{A})$.

Proof. By assumption, the full subcategory \mathcal{Y} is cogenerating in \mathcal{A} and by Lemma 1.2,1) \mathcal{Y} satisfies assumption (ii) of Proposition 1.5. Therefore we can apply the latter result, which gives that the canonical composition of functors $\mathcal{K}(\mathcal{Y}) \hookrightarrow \mathcal{K}(\mathcal{A}) \hookrightarrow \mathcal{D}(\mathcal{A})$ induces a triangulated equivalence $\mathcal{K}(\mathcal{Y})/\mathcal{K}_{ac}^{\mathcal{A}}(\mathcal{Y}) \xrightarrow{\cong} \mathcal{D}(\mathcal{A})$.

On the other hand, by Lemma 1.6 the full subcategory \mathcal{Y} is generating in \mathcal{H}_τ and by Lemma 1.2, 2) \mathcal{Y} satisfies the assumptions of the dual of Proposition 1.5. Therefore by the dual of Proposition 1.5, the canonical composition of functors $\mathcal{K}(\mathcal{Y}) \hookrightarrow \mathcal{K}(\mathcal{H}_\tau) \hookrightarrow \mathcal{D}(\mathcal{H}_\tau)$ induces a triangulated equivalence $\mathcal{K}(\mathcal{Y})/\mathcal{K}_{ac}^{\mathcal{H}_\tau}(\mathcal{Y}) \xrightarrow{\cong} \mathcal{D}(\mathcal{H}_\tau)$. By Proposition 1.3, $\mathcal{K}_{ac}^{\mathcal{A}}(\mathcal{Y}) = \mathcal{K}_{ac}^{\mathcal{H}_\tau}(\mathcal{Y}) =: \mathcal{K}_{ac}(\mathcal{Y})$. Hence we obtain the following diagram

$$\begin{array}{ccc} & \mathcal{K}(\mathcal{Y})/\mathcal{K}_{ac}(\mathcal{Y}) & \\ \cong \swarrow & & \searrow \cong \\ \mathcal{D}(\mathcal{H}_\tau) & \xrightarrow{\cong} & \mathcal{D}(\mathcal{A}) \end{array}$$

Finally, consider any object X^\bullet of \mathcal{H}_τ and denote by F the just defined equivalence $\mathcal{D}(\mathcal{H}_\tau) \xrightarrow{\cong} \mathcal{D}(\mathcal{A})$. By the dual of Proposition 1.5, we have a quasi-isomorphism $Y^\bullet \rightarrow X^\bullet$ in $\mathcal{K}(\mathcal{H}_\tau)$, whence an isomorphism in $\mathcal{D}(\mathcal{H}_\tau)$, where $Y^\bullet \in \mathcal{K}(\mathcal{Y})$. By the above commutative diagram, we have an isomorphism $F(X^\bullet) \cong F(Y^\bullet) \cong Y^\bullet$ in $\mathcal{D}(\mathcal{A})$, which shows the compatibility of F with the inclusion functor $\mathcal{H}_\tau \hookrightarrow \mathcal{D}(\mathcal{A})$. \square

Remark 1.8. Lemma 1.6 can be dualized in the following way: let \mathcal{A} be an abelian category such that $\mathcal{D}(\mathcal{A})$ has Hom sets and let $\tau = (\mathcal{U}, \mathcal{U}^\perp[1])$ be a t -structure in $\mathcal{D}(\mathcal{A})$ such that $\mathcal{D}^{\leq 0}(\mathcal{A}) \subseteq \mathcal{U} \subseteq \mathcal{D}^{\leq n}(\mathcal{A})$, for some $n \in \mathbb{N}$, and let \mathcal{H}_τ be its heart. If the class $\mathcal{Y} = \mathcal{A} \cap \mathcal{H}_\tau$ is generating in \mathcal{A} , then \mathcal{Y} is cogenerating in \mathcal{H}_τ . The proof of this statement follows by using the dual arguments of those presented in the proof of Lemma 1.6. Under these assumptions, Theorem 1.7 yields a triangle equivalence $\mathcal{D}(\mathcal{H}_\tau) \xrightarrow{\cong} \mathcal{D}(\mathcal{A})$ which is compatible with the inclusion $\mathcal{H}_\tau \hookrightarrow \mathcal{D}(\mathcal{A})$.

2. APPLICATIONS TO NONCLASSICAL TILTING OBJECTS

In this section we apply the Tilting Theorem 1.7 to the case of nonclassical tilting objects.

The following definition appears in [NSZ15].

Definition 2.1. [NSZ15] Let \mathcal{A} be an abelian category whose derived category has Hom sets and arbitrary (small) coproducts. Let n be a natural number. An object T of \mathcal{A} is called n -tilting when the following conditions hold:

- T0 For each set I , the object $T^{(I)}$ is the coproduct of I copies of T both in \mathcal{A} and $\mathcal{D}(\mathcal{A})$;
- T1 $\text{Ext}_{\mathcal{A}}^k(T, T^{(I)}) = 0$, for all integers $k > 0$ and all sets I ;
- T2 The projective dimension of T is $\leq n$, that is, $\text{Ext}_{\mathcal{A}}^k(T, N) = 0$ for all integers $k > n$ and all objects N of \mathcal{A} ;
- T3 There is a generating class \mathcal{G} of \mathcal{A} such that, for each $G \in \mathcal{G}$, there is an exact sequence $0 \rightarrow G \rightarrow T^0 \rightarrow T^1 \rightarrow \dots \rightarrow T^n \rightarrow 0$, where all the T^k are in $\text{Add}(T)$.

T is called a *tilting* object when it is n -tilting, for some natural number n . A *classical* (n -)tilting object is a (n -)tilting object which is compact as an object of $\mathcal{D}(\mathcal{A})$.

As proved in [NSZ15], if \mathcal{A} be an abelian category whose derived category has Hom sets and arbitrary (small) coproducts, as in Definition 2.1, then \mathcal{A} is automatically cocomplete. When \mathcal{A} is an abelian category whose derived category has

Hom sets and arbitrary (small) products, the dual concepts of *n-cotilting*, *cotilting* and *classical (n-)cotilting* object appear naturally.

Recall from [PV15] that a *tilting* object in $\mathcal{D}(\mathcal{A})$ is a complex T^\bullet such that the pair $(T^{\bullet\perp>0}, T^{\bullet\perp<0})$ is a *t-structure* in $\mathcal{D}(\mathcal{A})$ and $T^\bullet \in T^{\bullet\perp>0}$. Here $T^{\bullet\perp>0}$ (resp. $T^{\bullet\perp<0}$) is the full subcategory of $\mathcal{D}(\mathcal{A})$ consisting of those Y^\bullet such that $\mathrm{Hom}_{\mathcal{D}(\mathcal{A})}(T^\bullet, Y^\bullet[k]) = 0$, for all $k > 0$ (resp. $k < 0$).

In the next result, when $m \in \mathbb{N}$, the subcategory of objects Y of \mathcal{A} which admit an exact sequence $T^{-m} \rightarrow \dots \rightarrow T^{-1} \rightarrow T^0 \rightarrow Y \rightarrow 0$, with all the T^{-k} in $\mathrm{Add}(T)$, will be denoted by $\mathrm{Pres}^m(T)$.

Theorem 2.2. [NSZ15] *Let \mathcal{A} be an abelian category such that its derived category $\mathcal{D}(\mathcal{A})$ has *Hom* sets and arbitrary (small) coproducts, and let T be an object of \mathcal{A} such that the coproduct $T^{(I)}$ is the same in \mathcal{A} and in $\mathcal{D}(\mathcal{A})$. The following are equivalent:*

- (1) T is a tilting object in \mathcal{A} .
- (2) T has finite projective dimension and is a tilting object of $\mathcal{D}(\mathcal{A})$.
- (3) If $\mathcal{Y} = \bigcap_{k>0} \mathrm{Ker}(\mathrm{Ext}_{\mathcal{A}}^k(T, ?))$, then:
 - (a) $\mathcal{Y} = \mathrm{Pres}^m(T)$, for some integer $m \in \mathbb{N}$;
 - (b) \mathcal{Y} is a cogenerating class of \mathcal{A} .

Moreover, when one of the previous assertion holds, the pair $\tau_T = (T^{\perp>0}, T^{\perp<0})$ is a *t-structure* in $\mathcal{D}(\mathcal{A})$ and $\mathcal{D}(\mathcal{A})^{\leq -n} \subseteq T^{\perp>0} \subseteq \mathcal{D}(\mathcal{A})^{\leq 0}$.

The main application of the Tilting Theorem 1.7 is the following proposition, which generalizes the derived equivalence induced by a classical tilting object (at the level of unbounded derived categories) to the non-classical case.

Proposition 2.3. *Let \mathcal{A} be an abelian category whose derived category has *Hom* sets and arbitrary (small) coproducts, let T be a tilting object in \mathcal{A} and let \mathcal{H}_T be the heart of the associated *t-structure* $\tau_T = (T^{\perp>0}, T^{\perp<0})$ in $\mathcal{D}(\mathcal{A})$. Then there is a triangulated equivalence $\mathcal{D}(\mathcal{H}_T) \xrightarrow{\cong} \mathcal{D}(\mathcal{A})$ which is compatible with the inclusion functor $\mathcal{H}_T \hookrightarrow \mathcal{D}(\mathcal{A})$.*

Proof. The proof follows directly from Theorems [NSZ15] and Theorem 1.7, after we realize that $\mathcal{A} \cap \mathcal{H}_T = \mathcal{A} \cap T^{\perp>0} = \bigcap_{k>0} \mathrm{Ker}(\mathrm{Ext}_{\mathcal{A}}^k(T, ?))$. \square

It is worth stating explicitly the dual of the last proposition, which is known when \mathcal{A} is a module category (see [Sto14][Theorems 4.5 and 5.21]).

Proposition 2.4. *Let \mathcal{A} be an abelian category such that $\mathcal{D}(\mathcal{A})$ has *Hom* sets and arbitrary (small) products, let Q be a cotilting object in \mathcal{A} and let \mathcal{H}_Q be the heart of the associated *t-structure* $({}^{\perp<0}Q, {}^{\perp>0}Q)$ in $\mathcal{D}(\mathcal{A})$. Then there is a triangulated equivalence $\mathcal{D}(\mathcal{H}_Q) \xrightarrow{\cong} \mathcal{D}(\mathcal{A})$ which is compatible with the inclusion $\mathcal{H}_Q \hookrightarrow \mathcal{D}(\mathcal{A})$.*

Recall that an object A of \mathcal{A} is *self-small* when the canonical morphism $\mathrm{Hom}_{\mathcal{A}}(A, A)^{(I)} \rightarrow \mathrm{Hom}_{\mathcal{A}}(A, A^{(I)})$ is an isomorphism, for all sets I . The following result seems to be unknown even when \mathcal{A} is a module category, and is an easy consequence of Theorem 1.7.

Corollary 2.5. *Let \mathcal{A} be an abelian category whose derived category has *Hom* sets and arbitrary (small) coproducts and let T be a tilting object of \mathcal{A} . The following assertions are equivalent:*

- (1) T is a self-small object of \mathcal{A} .

(2) T is a classical tilting object of \mathcal{A} .

Proof. 2) \implies 1) On the one hand, by definition of tilting object, the coproduct $T^{(I)}$ is the same in \mathcal{A} and in $\mathcal{D}(\mathcal{A})$. On the other hand, since T is classical by assumption, then it is a compact object of $\mathcal{D}(\mathcal{A})$. Hence, for any set I , we have a chain of isomorphisms

$$\begin{aligned} \mathrm{Hom}_{\mathcal{A}}(T, T)^{(I)} &\cong \mathrm{Hom}_{\mathcal{D}(\mathcal{A})}(T, T)^{(I)} \\ &\cong \mathrm{Hom}_{\mathcal{D}(\mathcal{A})}(T, T^{(I)}) \\ &\cong \mathrm{Hom}_{\mathcal{A}}(T, T^{(I)}), \end{aligned}$$

whose composition is the canonical morphism $\mathrm{Hom}_{\mathcal{A}}(T, T)^{(I)} \longrightarrow \mathrm{Hom}_{\mathcal{A}}(T, T^{(I)})$.

1) \implies 2) By [NSZ15], we know that the heart \mathcal{H}_T of the associated t-structure $\tau = (T^{\perp > 0}, T^{\perp < 0})$ is a module category, with $\tilde{H}(T) = T$ as a progenerator. In particular, T is a compact generator of $\mathcal{D}(\mathcal{H}_T)$. If now $F : \mathcal{D}(\mathcal{H}_T) \xrightarrow{\cong} \mathcal{D}(\mathcal{A})$ is a triangulated equivalence which is compatible with the inclusion functor $\mathcal{H}_T \hookrightarrow \mathcal{D}(\mathcal{A})$, then $F(T) \cong T$ is compact in $\mathcal{D}(\mathcal{A})$. \square

REFERENCES

- [AC01] ANGELERI-HÜGEL, L.; COELHO, F.U.: Infinitely generated tilting modules of finite projective dimension. *Forum Math.* **13** (2001), 239-250.
- [AHK07] ANGELERI-HÜGEL, L.; HAPPEL, D.; KRAUSE, H. (EDTS): Handbook of tilting theory. *London Math. Soc. Lect Note Ser.* **332**. Cambridge Univ. Press (2007).
- [BMT11] BAZZONI, S.; MANTESE, F.; TONOLO, A.: Derived equivalence induced by infinitely generated n -tilting modules. *Proceedings AMS* **139** (2011), 4225-4234.
- [BR07] BELIGIANNIS, A.; REITEN, I.: Homological and homotopical aspects of torsion theories. *Mem. Amer. Math. Soc.* **188** (2007), no. 883, viii+207.
- [Bo81] BONGARTZ, K.: Tilted Algebras. *Representations of Algebras*, Puebla 1980. Springer LNM **903** (1981), 26-38.
- [BB80] BRENNER, S.; BUTLER, M.: Generalizations of the Bernstein-Gelfand-Ponomarev reflection functors. *Representation Theory II*, Ottawa 1979. Springer LNM **832** (1980), 103-169.
- [BBD82] BEILINSON, A.; BERNSTEIN, J.; DELIGNE, P.: Faisceaux Pervers. *Analysis and Topology on Singular Spaces I*, Luminy 1981. *Astérisque* **100**, Soc. Math. France (1982).
- [Ch10] CHEN, X.W.: A short proof of HRS-tilting. *Proceedings AMS* **138**(2) (2010), 455-459.
- [CF90] COLBY, R.R.; FULLER, K.R.: Tilting, cotilting and serially tilting rings. *Comm. Algebra* **18**(5) (1990), 1585-1615.
- [CT95] COLPI, R.; TRILIFAJ, J.: Tilting modules and tilting torsion theories. *J. Algebra* **178** (1995), 614-634.
- [CPS86] CLINE, E.; PARSHALL, B.; SCOTT, L.: Derived categories and Morita theory. *J. Algebra* **104**(2) (1986), 397-409.
- [FMT14] FIOROT, L.; MATTIELLO, F.; TONOLO, A.: A classification theorem for t-structures. Preprint available at <http://arxiv.org/pdf/1412.8679.pdf>
- [H87] HAPPEL, D.: On the derived category of a finite dimensional algebra. *Coment. Math. Helvet.* **62**(1) (1987), 339-389.
- [H88] HAPPEL, D.: Triangulated categories in the representation theory of finite dimensional algebras. *London Math. Soc. Lect Note Ser.* **119** (1988).
- [HRS96] HAPPEL, D.; REITEN, I.; SMALO, S.O.: Tilting in abelian categories and quasi-tilted algebras. *Memoirs AMS* **575** (1996).
- [HR82] HAPPEL, D.; RINGEL, C.M.: Tilted algebras. *Transactions AMS* **274** (1982), 399-443
- [KS05] KASHIWARA, M.; SCHAPIRA, P.: *Categories and Sheaves*, Springer Berlin Heidelberg, 2005. *Grundlehren der mathematischen Wissenschaften*. Isbn: 9783540279495.
- [Ke94] KELLER, B.: Deriving DG categories. *Ann. Sci. Ec. Norm. Sup* **27** (1994), 63-102.
- [M86] MIYASHITA, Y.: Tilting modules of finite projective dimension. *Math. Zeitschr.* **193** (1986), 113-146.
- [NSZ15] NICOLÁS, P.; SAORÍN, M.; ZVONAREVA, A.: Silting objects in arbitrary triangulated categories. Preprint available at <http://arxiv.org/pdf/1512.04700v3.pdf>.

- [PV15] PSAROUDAKIS, C.; VITÓRIA, J.: Realisation functors in tilting theory. Preprint available at <http://arxiv.org/pdf/1511.02677v1.pdf>
- [Ri89] RICKARD, J.: Morita theory for derived categories. *J. London Math. Soc.* **39** (1989), 436-456.
- [Sto14] STOVICEK, J.: Derived equivalences induced by big cotilting modules. *Adv. Math.* **263** (2014), 45-87. doi:10.1016/j.aim.2014.06.007. [link to the journal] [arXiv:1308.1804]

DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DEGLI STUDI DI PADOVA, VIA TRIESTE 63, 35121 PADOVA (ITALY)

E-mail address: fiorot@math.unipd.it

DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DEGLI STUDI DI PADOVA, VIA TRIESTE 63, 35121 PADOVA (ITALY)

E-mail address: mattiell@math.unipd.it

DEPARTAMENTO DE MATEMÁTICAS, UNIVERSIDAD DE MURCIA, APTDO. 4021, 30100 ESPINARDO, MURCIA (SPAIN)

E-mail address: msaorinc@um.es