

Multi-omics approach reveals the secrets of metabolism of clear cell—renal cell carcinoma

Giuseppe Lucarelli¹, Matteo Ferro², Michele Battaglia¹

¹Department of Emergency and Organ Transplantation, Urology, Andrology and Kidney Transplantation Unit, University of Bari, Bari, Italy;

²Department of Urology, European Institute of Oncology, Milan, Italy

Correspondence to: Giuseppe Lucarelli, Department of Emergency and Organ Transplantation, Urology, Andrology and Kidney Transplantation Unit, Piazza G. Cesare 11, 70124 Bari, Italy. Email: giuseppe.lucarelli@inwind.it.

Submitted May 23, 2016. Accepted for publication May 24, 2016.

doi: 10.21037/tau.2016.06.12

View this article at: <http://dx.doi.org/10.21037/tau.2016.06.12>

Renal cell carcinoma (RCC) accounts for 2–3% of all adult malignant tumors, having the highest incidence in Western countries. It is estimated that in 2016, about 62,700 new cases will be diagnosed (3.7% of all new cancer cases) in the United States, and nearly 14,240 patients will die of this cancer (2.4% of all cancer deaths) (1). Patients with low stage, localized disease have an excellent prognosis after surgical treatment, but a significant percentage of subjects still present with a surgically unresectable tumor or will subsequently develop metastatic disease (2,3). Recent studies have provided novel insights into the molecular mechanisms involved in the RCC pathogenesis, leading to the identification of potential biomarkers for early diagnosis, risk assessment, and outcome prediction. Many molecular markers such as C-reactive protein, CA15-3, α Klotho and some metabolic enzymes have been investigated, although none of these molecules has improved the predictive accuracy of current prognostic systems and their use is not recommended in clinical practice (4–8). The recent introduction of large-scale methods to purify, identify, and characterize DNA, RNA, proteins, metabolites and other molecules, has led to an in-depth exploration of the molecular bases underlying the development of urological cancer, together with the identification of new biomarkers and potential therapeutic targets (9,10). In this scenario, analysis of cancer cell metabolism has shown that tumor cells exhibit an atypical reprogramming of the energy metabolism that serves to promote cell growth and proliferation (11). RCC is fundamentally a metabolic disease (12) and many studies have suggested that in some diseases characterized by an altered metabolism there is an increased risk to develop this tumor (13–15). Moreover, a bird's eye view of the

genetic alterations involved in the RCC pathogenesis shows that many genes play an important role in controlling cell metabolism (16).

The recent comprehensive molecular characterization of clear cell RCC by the cancer genome atlas (TCGA) research network has confirmed that oncogenic metabolism and epigenetic reprogramming are two main features of RCC (17). In particular, an inverse correlation has been shown between patients' survival and the activation of a metabolic shift characterized by a Warburg effect-like state (i.e., the shift to aerobic glycolysis with lactate production), an increased dependence on the pentose phosphate pathway (PPP), a reduced Krebs cycle activity, increased glutaminolysis and fatty acid production (17). A later study explored the specific role of glycolysis and PPP in sustaining RCC cell proliferation, the maintenance of NADPH levels, production of reactive oxygen species (ROS) and in reducing chemotherapy-induced cytotoxicity (7). Clear cell RCC was characterized by high levels of glucose and other sugars, in association with an increase in upstream glycolytic intermediates (glucose 6-phosphate and fructose 6-phosphate), a reduction in downstream intermediates (3-phosphoglycerate, 2-phosphoglycerate, and phosphoenolpyruvate), and an increased lactate production. Other important findings were the higher expression in neoplastic tissue of two cancer-specific isoenzymes, namely Pyruvate kinase isoform M2 (PKM2) and L-lactate dehydrogenase isoform 5 (LDH5). These metabolic alterations, in association with high levels of PPP enzymes (Glucose-6-phosphate dehydrogenase-G6PDH and Transketolase-TKT) and metabolic intermediates (sedoheptulose 7-phosphate, ribose 5-phosphate, and ribulose

5-phosphate/xylulose 5-phosphate), suggested a rerouting of the sugar metabolism toward the PPP, with the aim of promoting both anabolic reactions and redox homeostasis in RCC (7). Ribose-5-phosphate is a sugar used in the synthesis of nucleotides, so the increased flux of metabolites through the PPP provides an advantage for cell growth and survival. Since G6PDH is the rate-limiting enzyme of the PPP, its inhibition causes a significant decrease in cancer cell growth, confirming the importance of this pathway in RCC. Another important byproduct of PPP is NADPH, a reducing compound that is used for biosynthetic reactions and to control the redox state. Cancer cells pre-treatment with a G6PDH inhibitor induces a significant reduction in NADPH levels and an increased production of ROS in renal cancer cells, suggesting that G6PDH, and hence the PPP, have a fundamental role in maintaining redox homeostasis in RCC (7).

These findings were recently confirmed by Hakimi *et al.* that showed how clear cell RCC is characterized by a reprogramming of central carbon metabolism, one-carbon metabolism, and the antioxidant response (18). In particular, consistent with previous studies (7,17), these authors showed increased levels of metabolites in upper glycolysis, and a reduction of metabolic derivatives in lower glycolysis, downstream of fructose 6-phosphate. This different abundance in metabolites production between upper and lower glycolysis suggests that the metabolic flux through this pathway may be differentially partitioned. In fact, while the sugars produced in the upper part of glycolysis are diverted to the PPP, the triose phosphates generated in the lower part are rerouted towards the Krebs cycle or one-carbon metabolism. In accordance with these results, analysis of tricarboxylic acid (Krebs) cycle metabolites (reduced levels of malate and fumarate, and accumulation of succinate) suggests that in RCC the mitochondrial bioenergetics and oxidative phosphorylation processes are impaired. It has recently been demonstrated that NADH dehydrogenase (ubiquinone) 1 alpha subcomplex 4-like 2 (NDUFA4L2) is a HIF-1 target gene that encodes for a component of the electron transport chain (ETC) Complex I (19). In particular, Complex I catalyzes the first step of mitochondrial respiratory chain reactions, and it couples the Krebs cycle to oxidative phosphorylation. Tello *et al.* have shown that hypoxia-induced NDUFA4L2 attenuates mitochondrial oxygen consumption through inhibiting Complex I activity, and reduces intracellular ROS production under low-oxygen conditions (19). The most surprising fact is that NDUFA4L2 is one of the most highly

expressed genes in clear cell RCC, while its knockdown impairs cell proliferation, alters metabolic pathways and induces autophagy in renal cancer cells (20). Taken together, these findings outline a clear cell RCC metabolic signature characterized by an anaerobic switch that favors the upper part of glycolysis and the rerouting of the sugar metabolism toward the PPP, and attenuates the mitochondrial activity through the overexpression of NDUFA4L2.

Another important advance highlighted by the Hakimi study (18) was the development of a metabologram, a web-based application that integrates transcriptomic and metabolomics data (<http://sanderlab.org/kidneyMetabProject>), with the aim of providing a rapid evaluation of metabolic changes in RCC. This visual tool explores the metabolic pathways using both gene expression (derived from TCGA database) and metabolite abundance data (derived from MSK Institute). Using metabolograms, it is possible to simultaneously evaluate changes in gene expression and metabolite abundance between tumor and normal tissues, as well as between different tumor stages. In addition, this web platform makes it possible to explore the association between the abundance of particular metabolic intermediates and 24 clinical parameters such as gender, age, the BMI, pathological characteristics, the onset of recurrence or metastasis.

In conclusion, the field of cancer metabolomics is evolving very rapidly. Recent amazing discoveries have cast new light on the molecular regulation of cancer cells metabolism, and novel biochemical pathways are now under investigation. In the next years we may expect a progressive integration of data derived from different platforms and other omics approaches and a more extensive use of the systems biology approach to identify a panel of molecular factors as biomarkers for diagnosis, prognosis, and as potential targets for specific therapies.

Acknowledgements

This work was supported by a grant from “Regione Puglia – ARTI” (Progetto FiR 4HE9F93 to G Lucarelli).

Footnote

Provenance: This is a Guest Editorial commissioned by Editorial Board Member Dr. Xiongbing Zu, MD, PhD (Department of Urology, Xiangya Hospital, Central South University, Changsha, China).

Conflicts of Interest: The authors have no conflicts of interest to declare.

Comment on: Hakimi AA, Reznik E, Lee CH, *et al.* An integrated metabolic atlas of clear cell renal cell carcinoma. *Cancer Cell* 2016;29:104-16.

References

1. Siegel RL, Miller KD, Jemal A. Cancer statistics, 2016. *CA Cancer J Clin* 2016;66:7-30.
2. Battaglia M, Lucarelli G. The role of renal surgery in the era of targeted therapy: the urologist's perspective. *Urologia* 2015;82:137-8.
3. Fischer S, Gillessen S, Rothermundt C. Sequence of treatment in locally advanced and metastatic renal cell carcinoma. *Transl Androl Urol* 2015;4:310-25.
4. Zhou L, Cai X, Liu Q, *et al.* Prognostic role of C-reactive protein in urological cancers: a meta-analysis. *Sci Rep* 2015;5:12733.
5. Lucarelli G, Ditunno P, Bettocchi C, *et al.* Diagnostic and prognostic role of preoperative circulating CA 15-3, CA 125, and beta-2 microglobulin in renal cell carcinoma. *Dis Markers* 2014;2014:689795.
6. Gigante M, Lucarelli G, Divella C, *et al.* Soluble serum α Klotho is a potential predictive marker of disease progression in clear cell renal cell carcinoma. *Medicine (Baltimore)* 2015;94:e1917.
7. Lucarelli G, Galleggiante V, Rutigliano M, *et al.* Metabolomic profile of glycolysis and the pentose phosphate pathway identifies the central role of glucose-6-phosphate dehydrogenase in clear cell-renal cell carcinoma. *Oncotarget* 2015;6:13371-86.
8. Lucarelli G, Rutigliano M, Sanguedolce F, *et al.* Increased expression of the autocrine motility factor is associated with poor prognosis in patients with clear cell-renal cell carcinoma. *Medicine (Baltimore)* 2015;94:e2117.
9. von Rundstedt FC, Rajapakshe K, Ma J, *et al.* Integrative pathway analysis of metabolic signature in bladder cancer: a linkage to the cancer genome atlas project and prediction of survival. *J Urol* 2016;195:1911-9.
10. Lucarelli G, Rutigliano M, Galleggiante V, *et al.* Metabolomic profiling for the identification of novel diagnostic markers in prostate cancer. *Expert Rev Mol Diagn* 2015;15:1211-24.
11. Hanahan D, Weinberg RA. Hallmarks of cancer: the next generation. *Cell* 2011;144:646-74.
12. Linehan WM, Srinivasan R, Schmidt LS. The genetic basis of kidney cancer: a metabolic disease. *Nat Rev Urol* 2010;7:277-85.
13. Vavallo A, Simone S, Lucarelli G, *et al.* Pre-existing type 2 diabetes mellitus is an independent risk factor for mortality and progression in patients with renal cell carcinoma. *Medicine (Baltimore)* 2014;93:e183.
14. Breda A, Lucarelli G, Rodriguez-Faba O, *et al.* Clinical and pathological outcomes of renal cell carcinoma (RCC) in native kidneys of patients with end-stage renal disease: a long-term comparative retrospective study with RCC diagnosed in the general population. *World J Urol* 2015;33:1-7.
15. Chen L, Li H, Gu L, *et al.* The impact of diabetes mellitus on renal cell carcinoma prognosis: a meta-analysis of cohort studies. *Medicine (Baltimore)* 2015;94:e1055.
16. Von Roemeling CA, Marlow LA, Radisky DC, *et al.* Functional genomics identifies novel genes essential for clear cell renal cell carcinoma tumor cell proliferation and migration. *Oncotarget* 2014;5:5320-34.
17. Cancer Genome Atlas Research Network. Comprehensive molecular characterization of clear cell renal cell carcinoma. *Nature* 2013;499:43-9.
18. Hakimi AA, Reznik E, Lee CH, *et al.* An integrated metabolic atlas of clear cell renal cell carcinoma. *Cancer Cell* 2016;29:104-16.
19. Tello D, Balsa E, Acosta-Iborra B, *et al.* Induction of the mitochondrial NDUFA4L2 protein by HIF-1 α decreases oxygen consumption by inhibiting Complex I activity. *Cell Metab* 2011;14:768-79.
20. Minton DR, Fu L, Mongan NP, *et al.* Role of NADH dehydrogenase (ubiquinone) 1 alpha subcomplex 4-like 2 in clear cell renal cell carcinoma. *Clin Cancer Res* 2016;22:2791-801.

Cite this article as: Lucarelli G, Ferro M, Battaglia M. Multi-omics approach reveals the secrets of metabolism of clear cell—renal cell carcinoma. *Transl Androl Urol* 2016;5(5):801-803. doi: 10.21037/tau.2016.06.12