


This article was downloaded by:[University of Torino]
On: 24 October 2007
Access Details: [subscription number 776098302]
Publisher: Taylor & Francis
Informa Ltd Registered in England and Wales Registered Number: 1072954
Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



British Poultry Science

Publication details, including instructions for authors and subscription information:
<http://www.informaworld.com/smpp/title~content=t713408216>

Breast meat traits of Muscovy ducks fed on a microalga (*Cryptothecodinium cohnii*) meal supplemented diet

A. Schiavone ^a; R. Chiarini ^b; M. Marzoni ^b; A. Castillo ^b; S. Tassone ^c; I. Romboli ^b

^a Faculty of Veterinary Medicine, Dipartimento di Produzioni Animali, Epidemiologia ed Ecologia, Università degli Studi di Torino,

^b Faculty of Veterinary Medicine, Dipartimento di Produzioni Animali, Università degli Studi di Pisa,

^c Faculty of Agriculture, Dipartimento di Scienze Zootecniche, Università degli Studi di Torino, Italy

Online Publication Date: 01 October 2007

To cite this Article: Schiavone, A., Chiarini, R., Marzoni, M., Castillo, A., Tassone,

S. and Romboli, I. (2007) 'Breast meat traits of Muscovy ducks fed on a microalga (*Cryptothecodinium cohnii*) meal supplemented diet', *British Poultry Science*, 48:5, 573 - 579

To link to this article: DOI: 10.1080/00071660701615796

URL: <http://dx.doi.org/10.1080/00071660701615796>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article maybe used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

Breast meat traits of Muscovy ducks fed on a microalga (*Cryptocodinium cohnii*) meal supplemented diet

A. SCHIAVONE, R. CHIARINI¹, M. MARZONI¹, A. CASTILLO¹, S. TASSONE² AND I. ROMBOLI¹

Faculty of Veterinary Medicine, Dipartimento di Produzioni Animali, Epidemiologia ed Ecologia, Università degli Studi di Torino, ¹Faculty of Veterinary Medicine, Dipartimento di Produzioni Animali, Università degli Studi di Pisa, and ²Faculty of Agriculture, Dipartimento di Scienze Zootecniche, Università degli Studi di Torino, Italy

Abstract 1. A trial was conducted in order to increase the docosahexaenoic acid (DHA) content in the meat of Muscovy ducks (*Cairina moschata domestica* L.) fed on a diet supplemented with the microalga *Cryptocodinium cohnii*.

2. Two diets were provided to 48 male and 48 female ducks, belonging to an Italian rural strain during the last 3 weeks of life: a maize–soybean based diet as the control diet and the same diet supplemented with 5 g/kg microalga meal.

3. Dietary treatment did not induce differences in growth performances and slaughter traits. Similarly, chemical composition, colour, pH, oxidative stability and sensory characteristics of breast muscle were not influenced by the diet.

4. A significant increase of DHA content in breast meat of ducks fed on the *Cryptocodinium cohnii* enriched diet was observed.

INTRODUCTION

In human nutrition, the biological effect of n-3 long-chain polyunsaturated fatty acids (LC-PUFAs) has received great interest as they play an active role in the prevention and management of several pathologies such as coronary heart disease, hypertension, type 2 diabetes, renal disease, ulcerative colitis, chronic obstructive pulmonary disease and Crohn's disease (Simopoulos, 2000). The main molecules of the n-3 LC-PUFA family involved in the beneficial biological effects are eicosapentaenoic acid (EPA, C20:5 n-3), docosapentaenoic acid (DPA, C22:5 n-3) and docosahexaenoic acid (DHA, 22:6 n-3). The current Western diet is characterised by low fish consumption, representing the main source of n-3 LC-PUFAs, and a high consumption of products from terrestrial animal origin, characterised by a

large content of PUFAs of the n-6 series (Simopoulos, 2000). On the other hand, the nutritional value of these products rich in n-6 PUFAs, such as poultry meat, could be enhanced by increasing the n-3 LC-PUFA content. In fact, several studies aimed to enrich poultry products with n-3 LC-PUFAs indicated fish oil as the most effective source for this purpose (Hargis and Van Elswyk, 1993; Leskanich and Noble, 1997). Our previous experience showed an increase in the amount of LC-PUFAs in duck meat with dietary fish oil (Schiaivone *et al.*, 2004). The recommendation of the European Union (decision 00/766/EU), concerning the ban on the use of animal proteins in animal nutrition, and the consumers' preferences, stimulated the interest in making diets for birds using exclusively vegetable ingredients.

This investigation aimed to increase the DHA content of the breast muscle of

Correspondence to: A. Schiaivone, Dipartimento di Produzioni Animali, Epidemiologia ed Ecologia, Facoltà di Medicina Veterinaria, 10095, Grugliasco (TO), Italy. Tel: +39-011-6709208. Fax: +39-011-6709240. E-mail: achille.schiaivone@unito.it
Accepted for publication 15th May 2007.

Muscovy ducks. Therefore, ducks were fed on a diet supplemented with marine microalga meal at the end of the finisher period. The microalga *Cryptocodinium cohnii*, a non-photosynthetic marine dinoflagellate, rich in DHA (Jiang *et al.*, 1999) was included in the diet and its effect evaluated at the minimum slaughter age for Muscovy ducks. Besides, meat sensory evaluation was performed to exclude the presence of a disagreeable taste.

MATERIALS AND METHODS

Animals and diets

Muscovy ducklings (*Cairina moschata domestica* L.) of both sexes of an Italian rural strain, hatched at the Experimental Avian Station of the Department of Animal Production of Pisa (Italy), were used for this trial. The experimental protocol was approved by the Ethics Committee of the Faculty of Veterinary Medicine of Pisa, Italy. All animal housing conformed to European Union guidelines.

A total of 48 males and 48 females were distributed by sex in 6 pens, respectively. Density was 3.5 males/m² and 4 females/m². All ducks received the same diet before the beginning of the experimental trial: a starter diet (12.1 MJ ME/kg and 240 g of crude protein/kg) and a finisher diet (12.0 MJ ME/kg and 200 g of crude protein/kg). The trial began when male ducks and female ducks were 50 and 43 d old, respectively. During the 21-d experimental period the finisher diet (Control diet, C) was compared with the same diet enriched with 5 g/kg dried microalga *Cryptocodinium cohnii* (Microalga diet, MA) (Algamix-AG DHA[®]). The two diets were formulated to be isoenergetic and isonitrogenous (Table 1). Each diet was randomly assigned to three pens for each sex. Olive oil was used as fat source for both diets. During the trial, individual duck body weight and feed consumption per pen were recorded weekly to calculate the feed conversion ratio (FCR).

Carcass assessment and meat quality

At the minimum slaughter age, seven 71-d-old males and seven 64-d-old females per dietary treatment were killed by electrical stunning followed by neck-cutting, after a 12-h starvation period. Eviscerated and plucked carcasses were weighed after removal of the feet and abdominal fat to obtain ready to cook carcasses (RCC). Breast fillets and liver were excised from refrigerated RCC (6 h at 4°C) and weighed to evaluate slaughter traits.

Table 1. Ingredients, chemical and fatty acid composition of the diets

	Experimental period (the last 3 weeks of life)	
	C diet	MA diet
<i>Ingredients (g/kg)</i>		
Maize meal	598	602
Soybean meal	340	333
Dicalcium phosphate	20	20
Calcium carbonate	17	17
Sodium chloride	2	2
DL-Methionine	2	2
L-Lysine	1	1
Vitamin and mineral premix ¹	5	5
Olive oil	15	13
<i>Cryptocodinium cohnii</i> meal	–	5
<i>Chemical composition (g/kg)</i>		
Dry matter	878.1	878.6
Crude protein	200.1	202.6
Ether extract	34.4	37.3
Crude fibre	29.2	31.4
Ash	67.7	63.2
Metabolisable energy (MJ/kg)	12.09	12.07
<i>Fatty acid composition (g/100 g fatty acid)</i>		
C14:0	0.11	0.31
C16:0	13.89	13.92
C16:1n7	0.33	0.34
C18:0	3.06	2.81
C18:1n9ct	31.84	31.80
C18:2n6c	45.19	45.45
C18:3n3	1.99	1.72
C22:6n3	–	0.23
SFAs ²	17.96	17.98
UFAs ³	79.81	79.65
SFAs/UFAs	0.23	0.23
MUFAs ⁴	32.38	32.45
PUFAs ⁵	47.43	47.20
n6	45.48	45.22
n3	1.95	1.99
n6/n3	23.00	23.00

¹Provided per kg of diet: retinol 3 mg, cholecalciferol 45 mg, DL- α -tocopheryl acetate 30 mg, thiamine 1.5 mg, riboflavin 3 mg, pyridoxine 1.5 mg, cobalamin 0.015 mg, pantothenic acid 8.0 mg, niacin 25 mg, choline chloride 500 mg, Fe (FeSO₄·7H₂O) 30 mg, Cu (CuSO₄·5H₂O) 1.5 mg, Mn (MnSO₄·H₂O) 80 mg, Zn (ZnSO₄·7H₂O) 30.0 mg, I (KI) 1.4 mg.

²Saturated fatty acids.

³Unsaturated fatty acids.

⁴Monounsaturated fatty acids.

⁵Polyunsaturated fatty acids.

Just before the dissection, pH and colour measurements of *Pectoralis major* muscles were determined. pH was measured using a Hanna Instruments 8417 pH-meter supplied with a Hamilton Biotrode electrode. Meat colour (CIELAB system: *L**, *a**, *b**) was measured on breast muscle surface using a Minolta Chroma-Meter CR-300 colour analyser.

Pectoralis major muscle was vacuum-packaged and frozen (–20°C) for analytical and sensory determinations described below. AOAC methods (1990) were used to assess moisture, ash, protein

and ether extract and results were expressed as a percentage on a fresh matter basis. Fatty acid composition of both breast muscle and diets was ascertained by capillary gas chromatography after lipid extraction (Folch *et al.*, 1957) and esterification (Christie, 1982), using sodium methoxide as catalyst. Fatty acid methyl esters were analysed by a Perkin-Elmer AutoSystem gas chromatograph, equipped with a flame ionisation detector and a Supelco Omegawax 320 capillary column (30 m × 0.32 mm, 0.25 µm film). Each fatty acid peak was identified by pure methyl ester standards (Supelco and Restek Corporation, Bellefonte, PA, USA) and data were conveyed as relative values.

Susceptibility to lipid oxidation was estimated from thiobarbituric acid reactive substances (TBARS) according to the iron-induced TBARS procedure, described by Huang and Miller (1993): 3 g of minced breast were homogenised in 57 ml of a chilled 1.15% KCl solution; 30 ml of the homogenate were incubated at 37°C in a shaking water bath with 8.34 mg FeSO₄·7H₂O (final concentration 1 mM Fe³⁺) as oxidative agent. The iron-induced TBARS assay was performed at 0, 60 and 120 min of incubation and the absorbance was read at 532 nm. Liquid malonaldehyde bis(diethyl acetal) (MDA) (Aldrich Chemical Co. Ltd, Gillingham, UK) was used as the standard to determine the linear standard response and recovery. TBARS values were calculated by multiplying absorbance by a constant coefficient *K* (23.58), combining standard response, recovery (93.4%), molecular weight of the MDA and sample weight. TBARS values were expressed as mg MDA/kg fresh meat.

A sensory panel test was performed on *Pectoralis major* muscle, without skin, roasted in a hot air oven at 165°C until an internal temperature of 70°C, without salt or spices and skin. Cooked samples were immediately sliced into 8 pieces and randomly offered to 6 trained panellists. The trial consisted of

4 sessions and the traits assessed were: tenderness, juiciness, fibrousness and acceptability. A 5-point scale was used: 1 referring to very disagreeable, very tough, very dry, very fibrous and 5 to very agreeable, extremely tender, very juicy, without fibre (Cross *et al.*, 1986).

Statistical analysis

Measurements of *in vivo* performances, slaughter traits, chemical composition, fatty acid profile and sensory analysis of meat were split by sex and analysed with a linear model including the fixed effect of diet (SPSS, 1997). Significance of differences was evaluated by *t*-test.

RESULTS

Fatty acid composition of microalga meal was characterised by a 30.6% DHA content. The 5 g/kg inclusion of microalga meal in MA diet resulted in a 2.3 g/kg content of DHA of total fatty acids, while DHA in the C diet was not detected (Table 1). Growth performances, slaughter traits and breast muscle traits (colour, pH and chemical composition) were not influenced by dietary treatments (Tables 2, 3 and 4). Similarly, susceptibility to lipid oxidation was not affected by the diet (Table 4). Besides, the inclusion of microalga did not influence tenderness, juiciness, fibrousness and acceptability of meat (Table 4).

Fatty acid composition of breast meat (Table 5) was affected ($P < 0.05$) by the diet. In particular DHA content was significantly higher in birds fed on the MA diet (2.9-fold in males and 2.8-fold in females, for MA and C diet, respectively). The improvement of DHA content in breast muscles due to MA diet significantly influenced the total n-3 PUFA amount (1.9-fold in males and 1.8-fold in females, for MA and C diet, respectively) and the n-6/n-3 ratio (−1.8-fold in males and −1.9-fold in females, for MA and C diet, respectively).

Table 2. Effects of microalga meal incorporation in diets for 3 weeks on growth performance of Muscovy ducks (means, $n = 24$)

		Females				Males			
		C	MA	SEM	P	C	MA	SEM	P
Initial age	d	43	43			50	50		
Final age	d	64	64			71	71		
Initial LW ¹	g	1343	1369	21.05	NS	2314	2318	32.64	NS
Final LW	g	1893	1909	22.72	NS	3314	3339	41.55	NS
Total body weight gain	g	550	540	11.44	NS	1000	1021	24.34	NS
ADG ²	g	26.20	25.71	0.54	NS	47.64	48.62	1.16	NS
FCR ³		5.31	5.03	0.11	NS	4.36	4.00	0.11	NS

¹Live body weight.

²Average daily weight gain.

³Feed conversion ratio ($n = 3$ for females; $n = 4$ for males).

Table 3. Effects of microalga meal incorporation in diets for 3 weeks on slaughter traits of Muscovy ducks (means, n = 7)

		Females (64 d old)				Males (71 d old)			
		C	MA	SEM	P	C	MA	SEM	P
LW ¹	g	1895	1918	24.89	NS	3443	3454	23.12	NS
RCC ²	g	1171	1193	16.07	NS	2163	2181	20.72	NS
	%LW	61.78	62.23	0.46	NS	62.82	63.14	0.53	NS
BM ³	g	244.40	249.26	4.98	NS	431.20	432.17	14.67	NS
	%RCC	20.88	20.90	0.42	NS	21.27	21.69	0.32	NS
Liver	g	27.54	28.60	1.31	NS	67.76	70.53	3.93	NS
	%RCC	2.35	2.40	0.11	NS	3.18	3.24	0.19	NS

¹Live body weight.²Ready to cook carcass.³Breast muscles.**Table 4.** Effects of microalga meal incorporation in diets for 3 weeks on physico-chemical characteristics, oxidation susceptibility and sensory traits of the Pectoralis major muscle from Muscovy ducks (means, n = 7 or 5 for chemical composition)

	Females				Males				
	C	MA	SEM	P	C	MA	SEM	P	
pH ¹	5.64	5.67	0.04	NS	5.75	5.69	0.03	NS	
Chemical composition ¹									
Moisture (%)	74.61	75.20	0.22	NS	76.77	76.57	0.09	NS	
Protein (%)	21.15	20.90	0.11	NS	20.59	20.17	0.20	NS	
Fat (%)	1.06	1.00	0.09	NS	0.99	1.02	0.06	NS	
Ash (%)	1.35	1.36	0.02	NS	1.23	1.26	0.02	NS	
Iron-induced TBARS (mg MDA/kg meat) ¹									
0 min	0.54	0.41	0.10	NS	0.88	1.20	0.22	NS	
60 min	0.97	0.83	0.11	NS	1.89	2.46	0.37	NS	
120 min	1.62	1.41	0.15	NS	1.63	2.18	0.27	NS	
Colour ¹									
L*	44.95	45.14	0.55	NS	46.28	46.84	0.41	NS	
a*	17.71	17.98	0.54	NS	17.07	16.68	0.30	NS	
b*	5.46	5.54	0.29	NS	4.53	4.70	0.36	NS	
Sensory traits ²									
Tenderness	3.48	3.35	0.09	NS	3.35	3.21	0.13	NS	
Juiciness	3.39	3.08	0.14	NS	2.73	2.78	0.17	NS	
Fibrousness	3.32	3.04	0.13	NS	2.62	2.55	0.10	NS	
Acceptability	3.03	2.96	0.15	NS	2.48	2.83	0.10	NS	

¹Results are means of two replicate analyses.²Each value is the mean of the judgement of 4 panellists.

DISCUSSION

Our results of moisture, protein and lipid contents in breast meat, 74.6 to 76.8%, 20.2 to 21.2% and 0.99 to 1.06%, respectively, are similar to those reported by other authors (Koci *et al.*, 1982; Paquin, 1988; Baéza *et al.*, 1997, 2002).

Avian fatty acids are typically monounsaturated, due to an active hepatic delta-9 desaturase, and an oleic and palmitoleic acid predominance. Domestic granivores use dietary carbohydrate for *de novo* fatty acid synthesis. Birds usually, except for some species, lack the enzymatic capacity to introduce double bonds past the ninth carbon of the fatty acid. Thus, they cannot use stearic acid to synthesise linoleic acid or alpha-linolenic acid, which are essential fatty acids in birds (Klasing, 2000). Dietary provision

of lipids was shown to influence the composition of phospholipid membranes and adipose depots. In fact, susceptibility of avian tissues and yolk lipids to fatty acid manipulation has been reviewed by Leskanich and Noble (1997) and Hargis and Van Elswyk (1993). Thus, the enrichment of poultry products with n-3 fatty acids, especially those with a number of carbons up to 20, is the main goal of fatty acid manipulation. Maldijan *et al.* (1996) showed that phospholipid fatty acid composition of eggs from chicken and duck (*Anas platyrhynchos* L.) markedly differed. The phospholipid fraction of duck eggs has higher proportions of arachidonic acid and lower proportions of DHA than chicken eggs. This shows that duck liver is quite efficient at converting dietary 18:2 n-6 into 20:4 n-6, whereas in chicken the conversion of 18:3 n-3

Table 5. Effects of microalga meal incorporation in diets for 3 weeks on fatty acid composition (% of total fatty acids) of the *Pectoralis* major muscle from Muscovy ducks (means, n = 3, results are means of three replicates analyses)

	Female				Male			
	C	MA	SEM	P	C	MA	SEM	P
C14:0	0.35	0.38	0.02	NS	0.36	0.33	0.04	NS
C16:0	24.57	24.57	0.71	NS	24.00	22.36	0.55	NS
C16:1n7	1.27	1.17	0.16	NS	2.09	1.40	0.45	NS
C18:0	12.82	12.48	0.78	NS	12.29	13.17	0.76	NS
C18:1n9ct	23.75	23.11	1.48	NS	23.73	23.31	1.73	NS
C18:1n7	2.81	2.52	0.08	NS	2.64	2.44	0.16	NS
C18:2n6c	16.85	17.03	0.45	NS	15.08	16.38	1.12	NS
C18:3n3	0.36	0.36	0.05	NS	0.33	0.34	0.03	NS
C20:4n6	8.87	9.10	0.96	NS	9.90	9.85	0.42	NS
C20:5n3	0.06	0.11	0.02	NS	0.08	0.12	0.01	NS
C22:4n6	1.99	1.80	0.21	NS	2.14	1.77	0.13	NS
C22:5n6	1.24	1.04	0.15	NS	1.20	0.96	0.11	NS
C22:5n3	0.40	0.39	0.07	NS	0.47	0.44	0.04	NS
C22:6n3	0.70	1.97	0.11	<0.05	0.74	2.12	0.18	<0.05
SFAs ¹	38.08	37.82	0.26	NS	37.01	36.25	0.37	NS
UFAs ²	60.06	61.73	0.40	NS	60.28	61.54	0.75	NS
SFAs/UFAs	0.63	0.62	0.01	NS	0.61	0.59	0.01	NS
MUFAs ³	28.55	27.80	1.52	NS	29.13	27.80	2.05	NS
PUFAs ⁴	31.51	32.93	1.40	NS	31.16	33.74	1.80	NS
n6	29.98	30.10	1.25	NS	29.50	30.65	1.61	NS
n3	1.53	2.83	0.16	<0.05	1.66	3.09	0.21	<0.05
n6/n3	19.85	10.65	1.05	<0.05	17.84	9.98	0.67	<0.001

¹Saturated fatty acids.²Unsaturated fatty acids.³Monounsaturated fatty acids.⁴Polyunsaturated fatty acids.

into 22:6 n-3 has priority. Thus, variations in the fatty acid profiles of yolk lipids may be derived from dietary provision of PUFAs as well as genetic differences among avian species regarding liver metabolic activities. However, in broiler chicken the capability to transfer LC-PUFAs from liver to peripheral tissue seems to be less effective than the mechanism involved in the incorporation of these in the yolk lipids. In fact Lopez-Ferrer *et al.* (2001) showed that dietary linolenic acid (28% of total dietary fatty acid) stimulates hepatic metabolic pathways involved in the elongation and desaturation of C18:3 n-3 into 22:6 n-3 (5.67 to 7.16% of total fatty acid), resulting in an inefficient build-up in thigh muscles (0.25% of total fatty acid). Similarly, in turkey, the conversion of C18:3 n-3 to EPA and DHA is weak. According to Komprda *et al.* (2005) the administration of three different diets containing 1.9%, 36.5% and 1.1% alpha-linolenic acid resulted in 6.8%, 9.0% and 5.7% EPA + DHA in breast meat.

Dietary algae have been successfully used in the nutrition of laying hens (Herber and Van Elswyk, 1996; Nitsan *et al.*, 1999; Tallarico *et al.*, 2002), broiler chickens (Mooney *et al.*, 1998; González-Esquerra and Leeson, 2001; Sirri *et al.*, 2003a, b) to obtain enriched poultry products, characterised by a significant amount of DHA or EPA, which could contribute to the novel

concept of 'functional food' in human nutrition (Almeida *et al.*, 2006; Garg *et al.*, 2006). In biological tissues and fluids, susceptibility to lipid oxidation is related to fatty acid chain length, number of double bonds (Liu *et al.*, 1997) and amount of antioxidant molecules (Surai, 1999). Our data on iron-induced TBARS showed no differences between groups in spite of the increase in DHA amount in animals fed on the diet supplemented with the microalga meal. This might be explained by the fact that the content of naturally occurring carotenoids, such as beta-carotene and canthaxanthin in the *Cryptocodinium cohnii* (Barclay *et al.*, 1994), may provide LC-PUFA stabilisation due to their antioxidant properties (Surai, 2002). Other authors studied the effect of other dietary microalgae on lipid stability of avian meat. No differences were found by Mooney *et al.* (1998) in breast meat TBARS values among chickens fed on 0%, 2.8% and 5.5% of dried *Schizochytrium*. Sirri *et al.* (2003a, b) studied the effect of 0.0%, 0.5% and 1.0% dietary *Schizochytrium* in broiler chicken on lipid stability of both drumstick and breast meat. TBARS values of drumstick meat were not influenced by dietary *Schizochytrium*, oppositely breast meat TBARS values of the same birds receiving 1.0% *Schizochytrium* appeared to be significantly higher than those from birds fed on 0.0 and 0.5% alga meal.

The 5 g/kg inclusion of *Cryptocodinium cohnii* did not affect the sensory panel test, agreeing with the results shown by Sirri *et al.* (2003b) when *Schizochytrium* were administered to chickens. Likewise Mooney *et al.* (1998) reported no modification of flavour in fresh chicken meat, while cooked meat samples from the experimental group with the highest level of *Schizochytrium* (5.5% of diet) were judged 'undesirable'.

In conclusion the 5 g/kg inclusion of microalga *Cryptocodinium cohnii* in the diet, provided for the last 21 d of life, improved DHA content in the duck meat without negative influences on sensory traits of the breast meat.

Also, the dietary microalga did not negatively influence growth performances or slaughter traits. In addition, chemical composition, colour, pH and oxidative stability of breast muscle were not influenced by the microalga supplemented diet. Muscovy duck appeared to react positively to this dietary supplementation. Thus, the use of *Cryptocodinium cohnii* could be used as a dietary strategy to enhance n-3 PUFA in duck meat.

In conclusion this DHA enriched meat could contribute to human nutrition, providing an increase of about threefold in DHA content compared to meat from ducks fed on a diet without microalga meal supplementation.

ACKNOWLEDGEMENTS

We are grateful to Mrs Manuela Baglini for duck breeding and Mrs Chiara Bianchi for technical support. Acknowledgements are due to Bios Pan Srl (Concordia sulla Secchia, MO, Italy) for providing the Algamix-AG DHA[®].

REFERENCES

- ALMEIDA, M.D.V., PINHÃO, S., STEWART-KNOX, B., PARR, H.J. & GIBNEY, M.J. (2006) An overview of findings from a six-country European survey on consumer attitudes to the metabolic syndrome, genetics in nutrition, and potential agro-food technologies. *British Nutrition Foundation Nutrition Bulletin*, **31**: 239–246.
- AOAC (1990) Official Methods of Analysis, 14th edn (Arlington, V.A., Association of Official Analytical Chemists).
- BAÉZA, E., DE CARVILLE, H., SALICHON, M.R. & LECLERQ, B. (1997) Effect of selection, over three. *British Poultry Science*, **38**: 359–365.
- BAÉZA, E., DESSAY, C., WACRENIER, G., MARCHE, G. & LISTRAT, A. (2002) Effect of selection for improved body weight and composition on muscle and meat characteristics in Muscovy duck. *British Poultry Science*, **43**: 560–568.
- BARCLAY, W.R., MEAGER, K.M. & ABRIL, J.R. (1994) Heterotrophic production of long chain omega-3 fatty acid utilizing algae and algae-like microorganisms. *Journal of Applied Phycology*, **6**: 123–129.
- CHRISTIE, W.W. (1982) A simple procedure for rapid transmethylation of glycerolipids and cholesteryl esters. *Journal of Lipid Research*, **23**: 1072–1075.
- Cielab Colour System, (1976) Commission International de l'Eclairage (Paris, CIE Publication).
- CROSS, H.R., DURLAND, P.R. & SEIDEMAN, S.C. (1986) Sensory qualities of meat, in: BECHTEL, P.J. (Ed.) *Muscle as Food*, pp. 279–320 (Orlando, Harcourt Brace Jovanovich).
- FOLCH, J., LEES, M. & SLOANE-STANLEY, H. (1957) A simple method for the isolation and purification of total lipids from animal tissue. *Journal of Biological Chemistry*, **226**: 479–509.
- GARG, M.L., WOOD, L.G., SINGH, H. & MOUNGHAN, P.J. (2006) Mean of delivering recommended levels of long chain n-3 polyunsaturated fatty acids in human diets. *Journal of Food Science*, **71**: 66–71.
- GONZÁLEZ-ESQUERRA, R. & LEESON, S. (2001) Alternatives for enrichment of eggs and chicken meat with omega-3 fatty acids. *Journal of Animal Science*, **81**: 295–305.
- HARGIS, P.S. & VAN ELSWYK, M.E. (1993) Manipulation of fatty acid composition of poultry meat and eggs for the health conscious consumer. *World's Poultry Science Journal*, **49**: 257–264.
- HERBER, S.M. & VAN ELSWYK, M.E. (1996) Dietary marine algae promotes efficient deposition of n-3 fatty acids for the production of enriched shell eggs. *Poultry Science*, **75**: 1501–1507.
- HUANG, Y.X. & MILLER, E.L. (1993) Iron-induced TBARS as an indicator of oxidative stability of fresh chicken meat. *Proceedings of the 11th European Symposium on the Quality of Poultry Meat*, Tours, pp. 478–484.
- JIANG, Y., CHEN, F. & LIANG, S.-Z. (1999) Production potential of docosahexaenoic acid by the heterotrophic marine dinoflagellate. *Cryptocodinium cohnii*. *Process Biochemistry*, **34**: 633–637.
- KLASING, K.C. (2000) Lipids, in: KLASING, K.C. (Ed.) *Comparative Avian Nutrition*, pp. 171–173 (Cambridge, CAB International).
- KOCI, E., BAUMGARTNER, J., ILLES, V. & PALANSKA, O. (1982) Carcass and nutritive value of the small white broiler duck and its crosses. *Archiv für Geflügelkunde*, **46**: 157–163.
- KOMPRDA, T., ZELENKA, J., FAJMONOVA, E., FIALOVA, M. & KLADROBA, D. (2005) Arachidonic acid and long-chain n-3 polyunsaturated fatty acid contents in meat of selected poultry and fish species in relation to dietary fat sources. *Journal of Agricultural and Food Chemistry*, **53**: 6804–6812.
- LESKANICH, C.O. & NOBLE, R.C. (1997) Manipulation of n-3 polyunsaturated fatty acid composition of avian eggs and meat. *World's Poultry Science Journal*, **53**: 155–183.
- LIU, J., YEO, H.C., DONIGER, S.J. & AMES, B.N. (1997) Assay of aldehydes from lipid peroxidation: gas chromatography-mass spectrometry compared to thiobarbituric acid. *Analytical Biochemistry*, **245**: 161–166.
- LOPEZ-FERRER, S., BAUCCELLS, M.D., BARROETA, A.C., GALOBART, J. & GRASHORN, M.A. (2001) n-3 enrichment of chicken meat. Use of precursors of long-chain polyunsaturated fatty acids: linseed oil. *Poultry Science*, **80**: 753–761.
- MALDIJAN, A., CRISTOFORI, C., NOBLE, R.C. & SPEAKE, B.K. (1996) The fatty acid composition of brain phospholipids from chicken and duck embryos. *Comparative Biochemistry and Physiology B*, **115B**: 153–158.
- MOONEY, J.W., HIRSCHLER, E.M., KENNEDY, A.K., SAMS, A.R. & VAN ELSWYK, M.E. (1998) Lipid and flavour quality of stored breast meat from broilers fed marine algae. *Journal of the Science of Food and Agriculture*, **78**: 134–140.
- NITSAN, Z., MOKADY, S. & SUKENIK, A. (1999) Enrichment of poultry products with n-3 fatty acids by dietary supplementation with the alga *Nannochloropsis* and mantur oil. *Journal of Agricultural Food and Chemistry*, **47**: 5127–5132.
- PAQUIN, J. (1988) Valeur nutritionnelle des viandes de volailles. *L'aviculture française*, ROSSET, R. (Ed.) *L'aviculture française*,

- (Paris, Informations Techniques des Services Vétérinaires).
- SCHIAVONE, A., ROMBOLI, I., CHIARINI, R. & MARZONI, M. (2004) Influence of dietary lipid source and genotype on fatty acid composition of Muscovy duck meat. *Journal of Animal Physiology and Animal Nutrition*, **88**: 88–93.
- SIMOPOULOS, A.P. (2000) Human requirement for n-3 polyunsaturated fatty acid. *Poultry Science*, **79**: 961–970.
- SIRRI, F., MINELLI, G., IAFFALDANO, N., TALLARICO, N. & FRANCHINI, A. (2003a) Oxidative stability and quality traits of n-3 PUFA enriched chicken meat. *Italian Journal of Animal Science*, **2**: 450–452.
- SIRRI, F., MINELLI, G., MELUZZI, A. & FRANCHINI, A. (2003b) Quality traits and oxidative stability of n-3 PUFA enriched meat, in: BAEZA, E. & FERNANDEZ, X. (Eds) *Proceedings of the XVI European Symposium on the Quality of Poultry Meat*, Saint-Brieuc, pp. 258–264.
- SPSS, (1997) Guida dell'utente SPSS, base 7.5 per Windows® (Chicago).
- SURAI, P.F. (1999) Vitamin E in avian reproduction. *Poultry and Avian Biology Reviews*, **10**(1): 1–60.
- SURAI, P.F. (2002) *Carotenoids in Natural Antioxidants in Avian Nutrition and Reproduction*, pp. 129–200 (Nottingham, Nottingham University Press).
- TALLARICO, N., SIRRI, F., MELUZZI, A., PITTIA, P., PERPINELLO, G.P. & FRANCHINI, A. (2002) Effect of dietary vegetable lipids on functional and sensory properties of chicken eggs. *Italian Journal of Food Science*, **14**: 159–166.