



University  
of Glasgow

Swift, D.A., Sanderson, D., Nienow, P.W., Bingham, R.G., and  
Cochrane, I.C. (2010) *Anomalous luminescence of subglacial sediment at  
Haut glacier d'Arolla, Switzerland - a consequence of resetting at the  
glacier bed?* *Boreas*, 40 (3). pp. 446-458. ISSN 0300-9483

<http://eprints.gla.ac.uk/43183/>

Deposited on: 23 February 2012



**Anomalous luminescence of subglacial sediment at Haut Glacier d’Arolla, Switzerland – a consequence of resetting at the glacier bed?**

|                               |   |
|-------------------------------|---|
| Journal:                      | <i>Boreas</i>   |
| Manuscript ID:                | BOR-017-2010.R3   |
| Manuscript Type:              | Original Article  |
| Date Submitted by the Author: | 06-Oct-2010   |
| Complete List of Authors:     | Swift, Darrel; University of Sheffield, Geography<br>Sanderson, David; Scottish Universities Environmental Research Centre<br>Nienow, Peter; University of Edinburgh, Geography<br>Bingham, Robert; University of Aberdeen, Geography |
| Keywords:                     | subglacial sediment, sediment transport, sediment tracing, geomechanical resetting, optically stimulated luminescence, thermoluminescence   |
|                               |   |

1  
2  
3 1 **SWIFT, D.A., SANDERSON, D.C.W., NIENOW, P.W., BINGHAM, R.G. AND**  
4  
5 2 **COCHRANE, I.C.: Anomalous luminescence of subglacial sediment at Haut**  
6  
7 3 **Glacier d’Arolla, Switzerland – a consequence of resetting at the glacier bed?**  
8

9  
10 4 Luminescence has the potential to elucidate glacial geomorphic processes because primary  
11  
12 5 glacial sediment sources and transport pathways are associated with contrasting degrees of  
13  
14 6 exposure to light. Most notably, sediment entrained from extraglacial sources should be at  
15  
16 7 least partially reset, whereas sediment produced by glacial erosion of subglacial bedrock  
17  
18 8 should retain substantial luminescence commensurate with a geological irradiation history.  
19  
20 9 We set out to test the validity of this assumption at Haut Glacier d’Arolla, Switzerland  
21  
22 10 using sediment sampled extraglacially and from the glacier bed. Contrary to our  
23  
24 11 expectations, the subglacial samples exhibited natural signals that were substantially lower  
25  
26 12 than those of other sample groups, and further (albeit limited) analyses have indicated no  
27  
28 13 obvious differences in sample group luminescence characteristics or behaviour that could  
29  
30 14 account for this observation. For glaciological reasons, we can eliminate both the possibility  
31  
32 15 that the subglacial sediment has been extraglacially-reset or exposed *in situ* to heat or light.  
33  
34 16 We therefore advocate investigation of possible resetting processes related to subglacial  
35  
36 17 crushing and grinding, and speculate that such processes, if more generally present, may  
37  
38 18 enable the dating of subglacially-deposited tills using luminescence-based techniques.  
39  
40  
41  
42  
43

44 19 **Keywords:** Subglacial sediment, sediment transport, sediment tracing, geomechanical  
45  
46 20 resetting, optically stimulated luminescence, thermoluminescence.  
47  
48  
49

50 21 *Darrel A. Swift (D.A.Swift@sheffield.ac.uk), Department of Geography, University of*  
51  
52 22 *Sheffield, Winter Street, Sheffield, S10 2TN, UK*  
53  
54

55 23 *David C.W. Sanderson, Scottish Universities Environmental Research Centre, Rankine*  
56  
57 24 *Avenue, Scottish Enterprise Technology Park, East Kilbride G75 0QF, UK*  
58  
59  
60

BOR-017-2010.R2

*Luminescence of subglacial sediment*

1  
2  
3 25 *Peter W. Nienow, Department of Geography, School of GeoSciences, University of*  
4  
5 26 *Edinburgh, Drummond Street, Edinburgh, EH8 9XP, UK*  
6  
7

8  
9 27 *Robert G. Bingham, Geography & Environment, School of Geosciences, University of*  
10  
11 28 *Aberdeen, Elphinstone Road, Aberdeen, AB24 3UF, UK*  
12  
13

14 29 *Ian C. Cochrane, Department of Geographical and Earth Science, University of Glasgow,*  
15  
16 30 *Glasgow, G12 8QQ, UK*  
17  
18

19  
20 31 Luminescence properties of sedimentary deposits have the potential to further  
21  
22 32 understanding of complex geomorphic systems and processes by elucidating their sediment  
23  
24 33 sources and transport pathways. Firstly, luminescence behaviour could be exploited in  
25  
26 34 situations where quantifiable differences in sensitivity, fading or bleaching characteristics,  
27  
28 35 for example, are produced by mineralogically distinct sediment sources or transport  
29  
30 36 pathways characterised by contrasting bleaching-dosing histories. Secondly, residual dose  
31  
32 37 could be exploited where sediment sources or transport pathways are associated with  
33  
34 38 varying degrees of luminescence accumulation or resetting. The latter approach should be  
35  
36 39 particularly applicable to glaciated catchments, where exposure to daylight should result in  
37  
38 40 extraglacial sources being substantially bleached, whilst sediment eroded from bedrock  
39  
40 41 beneath many metres of glacier ice should carry substantial luminescence commensurate  
41  
42 42 with a purely geological irradiation history (cf. Fuchs & Owen 2008).  
43  
44  
45  
46  
47

48 43 Minerals generate luminescence because structural defects trap 'free' electrons  
49  
50 44 produced by naturally occurring ionising radiation. Resetting of luminescence systems  
51  
52 45 requires such trapped electrons to be released under stimulation in natural or laboratory  
53  
54 46 settings. Relaxation processes can include recombination at luminescence centres, where a  
55  
56 47 proportion of the energy that is liberated is released as light (Aitken 1985, 1998). Resetting is  
57  
58  
59  
60

1  
2  
3 48 widely considered to be dominated by the effects of heat and light (Wintle & Huntley 1979;  
4  
5 49 Liritzis, 2000), making luminescence a useful tool for dating (cf. Lian & Roberts 2006) or  
6  
7  
8 50 process tracing (e.g. Rink *et al.* 1999; Bateman *et al.* 2007) in geology and geomorphology.  
9  
10 51 Potential as a process tracer in the glacial environment has been demonstrated by Gemmell  
11  
12 52 (1994, 1997), who attributed the substantial residual dose of proglacial stream suspended  
13  
14 53 sediment to the entrainment of sediment from mainly subglacial sources. Resetting of  
15  
16 54 residual dose at the glacier bed as a result of subglacial grinding and crushing has been  
17  
18 55 proposed (e.g. Morozov, 1968; Dreimanis *et al.* 1978; Singhvi *et al.* 1994), but the efficacy  
19  
20 56 of such 'geomechanical resetting' remains controversial (Toyoda *et al.* 2000).  
21  
22  
23  
24

25 57 We set out to examine whether residual dose could be used to elucidate the sources  
26  
27 58 of sediment evacuated by the subglacial drainage system at Haut Glacier d'Arolla,  
28  
29 59 Switzerland (Fig. 1). Firstly, extraglacial and subglacial sediments representing inputs to  
30  
31 60 and outputs from the drainage system were sampled under night-time conditions;  
32  
33 61 extraglacial sediment was sampled at the glacier margin and from glacial streams, whilst  
34  
35 62 subglacial sediment was sampled *in situ* from beneath ~100 m of glacier ice, utilising  
36  
37 63 boreholes drilled through the ice to the glacier bed (see Fig. 1 for drill site location). For  
38  
39 64 reasons given below, residual dose was initially characterised using simple polymineral  
40  
41 65 screening measurements, with full single-aliquot regenerative (i.e. SAR) procedures being  
42  
43 66 undertaken on a subset of samples only. We show that, rather than exhibiting substantial  
44  
45 67 equivalent dose commensurate with a geological irradiation history, the luminescence of the  
46  
47 68 subglacial sample group was substantially reset relative to that of the other major sediment  
48  
49 69 types. Possible reasons for these surprising observations are explored.  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

70 **Field area and sampling method**

71 Haut Glacier d’Arolla (Fig. 1A) is a classic alpine glacier at which sediment transport is  
72 dominated by the subglacial drainage system (Sharp *et al.* 1993; Swift *et al.* 2002). This  
73 system accesses a thin layer of deformable sediment at the ice-bed interface that is produced  
74 by erosion of the underlying bedrock (Hubbard *et al.* 1995; Harbor *et al.* 1997; Fischer &  
75 Hubbard, 1999). The majority of the annual sediment load is evacuated by hydraulically  
76 efficient subglacial channels that evolve in spring and summer (Nienow *et al.* 1998; Swift *et*  
77 *al.* 2002) and in which sediment transport is limited only by the rate of sediment supply  
78 (Swift *et al.* 2005; cf. Alley *et al.* 1997). Nevertheless, a portion of the sediment transported  
79 by subglacial channels is entrained in extraglacial streams, such as those fed by western-  
80 facing cirque glaciers below the Bouquetins ridge (Fig. 1b; Swift *et al.* 2005). Runoff from  
81 glacial sources causes sediment evacuation from the ice-bed interface to peak shortly after  
82 midday; however, runoff from the Bouquetins cirques continues into the evening. The  
83 catchment geology is complex, consisting of amphibolites, granites and gabbros that  
84 represent various stages of the Alpine Orogeny (Fig. 1C).

85 Sediments sampled at night in August 2000 comprised seven samples from the base  
86 of two ~100 m-deep glacial boreholes and 16 extraglacial samples: seven samples from  
87 marginal streams; three surface samples from marginal moraine; and six samples from two  
88 proglacial streams that emerge from the eastern portion of the subglacial drainage system  
89 (Fig. 1A, B). Stream samples comprised suspended sediment obtained by immersing an  
90 opaque sample bottle into a well-mixed section of the flow; moraine samples were scraped  
91 into opaque 35-mm film canisters from exposed sediment surfaces. Borehole sampling was  
92 undertaken using a water sampler modified from the design of Blake & Clarke (1991) (see  
93 Tranter *et al.* 2002). The boreholes had been drilled in mid-July using a hot-water drill

1  
2  
3 94 (ambient drill-tip water temperature  $\sim 50^{\circ}\text{C}$ ) and were sampled  $\sim 30$  days later, after  
4  
5 95 subglacial instrumentation – which had been deployed at the time of drilling – had been  
6  
7 96 removed. The sampler was shaken vigorously at the base of each borehole prior to closure  
8  
9  
10 97 of the sampler *in situ*; samples were protected from light and were stored and transported in  
11  
12 98 opaque polypropylene bottles.  
13  
14

15 99 Drilling and sampling methods do not indicate potential for significant  
16  
17 100 contamination of borehole samples by optically-reset sediment. There is potential to release  
18  
19 101 reset sediment from glacier ice during drilling; however, because debris causes problems  
20  
21 102 during drilling, boreholes were located away from supraglacial and englacial debris  
22  
23 103 accumulations, and, other than the highly conspicuous eastern medial moraine (Fig. 1), no  
24  
25 104 significant debris structures are known to exist in the vicinity of the drill site (see Goodsell  
26  
27 105 *et al.* 2005). Supraglacial and/or englacial streams are another potential source of reset  
28  
29 106 sediment; however, supraglacial runoff is characterised by extremely low sediment  
30  
31 107 concentrations, and boreholes do not act as a focus for runoff from wide areas of the glacier  
32  
33 108 surface. Furthermore, as the basal sediment layer in the vicinity of the drill-site is up to 10  
34  
35 109 cm thick (Hubbard *et al.* 1995; Harbor *et al.* 1997; Fischer & Hubbard 1999), the potential  
36  
37 110 for contamination by reset sediment would have been further reduced by thorough mixing  
38  
39 111 of the basal sediment layer both during drilling and by vigorous shaking of the Nielsen  
40  
41 112 sampler at the base of each borehole when sampling.  
42  
43  
44  
45  
46  
47

48 113 Another potential source of reset sediment is turbid water that down-borehole video  
49  
50 114 has shown to enter boreholes from small englacial channels (e.g. Copland *et al.* 1997).  
51  
52 115 However, such channels appear to be rare at Haut Glacier d’Arolla; the best example to  
53  
54 116 have been observed during borehole-survey was the result of turbid water, comprised of  
55  
56 117 sediment disturbed from the glacier bed, being forced into an englacial channel during  
57  
58  
59  
60

1  
2  
3 118 drilling (Copland *et al.* 1997). Furthermore, Copland *et al.* (1997) concluded that the  
4  
5 119 majority of borehole turbidity appeared to be generated by basal water flow through or  
6  
7  
8 120 above unconsolidated basal sediment at the ice-bed interface. Stone & Clarke (1996) have  
9  
10 121 also reported borehole-observations from temperate glaciers during the melt season that  
11  
12 122 show frequent mobilisation of basal sediment at the ice-bed interface.  
13  
14

### 15 123 **Sample preparation and initial screening results**

16  
17  
18  
19 124 Simple preparation techniques and a simple polymineral single-aliquot multiple-stimulation  
20  
21 125 screening approach (Table 1) were used for all samples on account of the small volume of  
22  
23 126 subglacial sediment acquired using the borehole sampling technique. The samples were  
24  
25  
26 127 prepared by settling in water before washing in a 10% HCl solution for 30 minutes to  
27  
28 128 remove carbonate minerals; no reaction with the HCl solution was observed, and because  
29  
30 129 the samples were devoid of organic material, no further pre-treatments were undertaken.  
31  
32 130 Mineralogical and grain size characteristics (the latter estimated to be 10–100  $\mu\text{m}$ ) were  
33  
34  
35 131 later checked for consistency using an FEI Quanta SEM. All luminescence measurements  
36  
37 132 were made from small quantities of sample dispensed onto 0.25 mm-thick 1 cm-diameter  
38  
39 133 stainless steel discs using a Risø DA15 luminescence reader equipped with a bialkali  
40  
41 134 photomultiplier (ET9235QB) and 9 mm Hoya U340 filter to detect near-UV radiation.  
42  
43 135 Although polymineral luminescence was anticipated to be dominated by feldspar emission,  
44  
45 136 and therefore to exhibit fading (cf. Krbetschek *et al.* 1997), the same multiple-stimulation  
46  
47  
48 137 procedure was used for all measurements.  
49  
50

51  
52 138 The multiple-stimulation screening procedure (Table 1) was applied to two discs per  
53  
54 139 sample and comprised sequential measurement of: (i) Infra-Red-Stimulated Luminescence  
55  
56 140 (IRSL) (60 s stimulation at 60°C with an 830 nm laser diode delivering approximately 240  
57  
58 141 mW cm<sup>-2</sup> to the sample); (ii) post-IR blue Optically Stimulated Luminescence (OSL) (30 s  
59  
60



1  
2  
3 142 stimulation at 125°C with GaN diodes at 470 nm delivering approximately 30 mW cm<sup>-2</sup> to  
4  
5 143 the sample); and (iii) Thermally-stimulated Luminescence (TL) (ambient to 500°C at 5°C s<sup>-1</sup>  
6  
7 144 <sup>1</sup> with a second heating to enable background-subtraction). Background-corrected  
8  
9  
10 145 luminescence signals were then extracted from raw IRSL and OSL shine-down and TL  
11  
12 146 glow-curves as shown in Fig. 2 and used to estimate the Residual Dose ( $D_r$ ) using the  
13  
14 147 simplest form of the single-aliquot regenerative-dose protocol,

15  
16  
17  
18 148 
$$\text{palaeodose} = \frac{L_n}{T_1} \times \frac{T_2}{L_r} \times \text{regenerative dose}, \quad (1)$$
  
19  
20  
21

22  
23 149 where  $L_n$ ,  $T_1$ ,  $L_r$  and  $T_2$  are the background-corrected natural signal, a subsequent test-dose  
24  
25 150 signal, a regenerative dose signal, and its associated test-dose signal, respectively (Table 1;  
26  
27 151 cf. Galbraith 2002). Similar multiple-stimulation procedures have been used in diverse  
28  
29 152 luminescence profiling studies to provide robust diagnoses of sediment transportation and  
30  
31 153 depositional processes (e.g. Sanderson *et al.* 2003, 2007; Burbidge *et al.* 2007; Sanderson &  
32  
33 154 Murphy 2010).

34  
35  
36  
37 155 Fig. 3 shows that initial  $D_r$  estimates reproduced well and covered several orders of  
38  
39 156 magnitude between the major sample groups, exceeding that which could reasonably be  
40  
41 157 expected to have arisen from methodological problems and uncertainties. Notably, although  
42  
43 158 regenerated signals ( $L_r$ ) were uniformly intense (typically around 10<sup>4</sup> counts for all sample  
44  
45 159 groups), subglacial samples yielded low-intensity natural signals ( $L_n$  in Table 1) compared  
46  
47 160 to those in other sample groups (e.g. sample 1277, Fig. 2). Consequently, the subglacial  
48  
49 161 sample group demonstrated substantially lower residual dose than any of the other sample  
50  
51 162 groups, regardless of stimulation method (Table 2). A small number of samples exhibited  
52  
53 163 weak or non-existent natural signals (see caption to Fig. 3), but largely in the case of post-  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 164 IR OSL, which can be attributed to the dominance of emissions from feldspar minerals  
4  
5 165 (predominantly feldspar mineralogy was confirmed by SEM analyses).  
6  
7

8  
9 166 **Further investigation of luminescence characteristics**

10  
11 167 The surprising results and subsequent discussions with peers inspired us to undertake  
12  
13 168 additional work to assess whether unexpectedly low subglacial residual dose could be  
14  
15 169 readily explained by: (1) differences in luminescence behaviour between the subglacial and  
16  
17 170 extraglacial samples; or (2) rogue luminescence behaviour that could cause the subglacial  
18  
19 171 samples to have apparent lower residual doses.  
20  
21  
22

23  
24 172 *Dose response*  
25  
26

27  
28 173 Uncertainties regarding residual dose estimates using the initial screening procedure and the  
29  
30 174 luminescence behaviour of different sample groups were investigated by applying single-  
31  
32 175 aliquot regenerative-dose (SAR) procedures to six key samples (including two subglacial  
33  
34 176 samples). The procedure employed the same polymineral multiple-stimulation procedure  
35  
36 177 (Table 1) with the addition of a range of regenerative doses (from 10 to 1000 Gy) and  
37  
38 178 recuperation and recycling steps; further, the procedure was applied to eight discs per sample,  
39  
40 179 which, following initial data appraisal, enabled mean values to be calculated for each  
41  
42 180 regeneration point belonging to each sample. SAR residual dose estimates were obtained  
43  
44 181 and compared with the initial screening estimates, bearing in mind the potential timing and  
45  
46 182 role of known sensitivity changes (e.g. Wallinga *et al.* 2000, 2001; Blair *et al.* 2005).  
47  
48  
49

50  
51 183 SAR curves (Fig. 4) were supra-linear but all samples demonstrated good SAR  
52  
53 184 characteristics (Table 3) and similar SAR behaviour, although subglacial TL exhibited  
54  
55 185 higher sensitivity than other samples to doses in excess of 100 Gy (Fig. 4C). Recycling and  
56  
57 186 recuperation values for all samples were mostly good (Table 4), with recycling ratios  
58  
59  
60

1  
2  
3 187 typically within the range 0.9–1.1 at  $\pm 1\sigma$ , and only two OSL recuperation values being >5%  
4  
5 188 (subglacial samples 1277 and 1285). Given the polymineral nature of the samples, the SAR  
6  
7 189 characteristics were therefore as good as could be anticipated and SAR  $D_e$  estimates were  
8  
9  
10 190 well-constrained (Table 4) and within saturation limits (cf. Fig. 4). SAR  $D_e$  estimates also  
11  
12 191 compared well with the initial residual dose estimates (Table 4).  
13  
14

15  
16 192 *Shape of the decay curve*  
17

18  
19 193 Consideration was given to whether natural and regenerated signals of certain sample  
20  
21 194 groups exhibited different decay properties that might invalidate SAR approaches. LM-OSL  
22  
23 195 (e.g. Thomas *et al.* 2006) was rejected because changes in decay properties can also arise  
24  
25 196 from differences in sample mineralogy and/or the number of bleaching-dosing cycles to  
26  
27 197 which sediment has been exposed (e.g. Bailey *et al.* 2003; Lukas *et al.* 2007), and our  
28  
29 198 limited experience of applying to feldspar systems indicated that the complex overlapping  
30  
31 199 signal distributions obtained would be extremely difficult to deconvolve. A standard signal  
32  
33 200 analysis approach (cf. Bailey *et al.* 2003) that used existing data sets was therefore  
34  
35 201 employed, comprising analysis of IRSL and OSL signal-decay plots and  $D_e(t)$  plots. The  
36  
37 202 latter were produced using sensitivity-corrected IRSL and OSL signals from successive  
38  
39 203 integration intervals of the raw shine-down curves (Fig. 5).  
40  
41  
42  
43  
44

45 204 Signal-decay plots (Fig. 6) demonstrated no significant differences in the form of  
46  
47 205 natural and regenerated signals for individual samples, and no obvious differences between  
48  
49 206 sample groups; post-IR OSL is characterised by slow decay, indicating that this signal is  
50  
51 207 likely to be dominated by feldspar (or quartz without a fast component).  $D_e(t)$  plots for  
52  
53 208 IRSL signals were either flat or showed a slight decline, whereas the OSL  $D_e(t)$  plots tended  
54  
55 209 to show some increase (Fig. 7). For quartz minerals, it has been suggested that a rise of  $D_e$   
56  
57 210 with integration time occurs in partially-reset samples as a result of better resetting of the  
58  
59  
60

BOR-017-2010.R2

*Luminescence of subglacial sediment*

1  
2  
3 211 fast component relative to the slower components (e.g. Bailey *et al.* 2003). For feldspar  
4  
5 212 minerals, such components have not been identified, and dependency of residual dose on  
6  
7 213 integration period may have other causes (e.g. signal stability). OSL  $D_e(t)$  plots are  
8  
9 214 therefore consistent with resetting of naturally-acquired luminescence signals, but, given  
10  
11 215 our limited knowledge of feldspar signals, no inferences can be made other than that there  
12  
13  
14 216 are no clear differences between the sample groups.

17  
18 217 *Stability of the signal*

19  
20  
21 218 Fading rates were investigated using further aliquots of the six samples previously subjected  
22  
23 219 to SAR analysis (see above). Eight aliquots of each sample were subjected to the same  
24  
25 220 polymineral multiple-stimulation procedure (Table 1); however, the procedure was  
26  
27 221 modified such that four aliquots were stored for 95 days following administration of the  
28  
29 222 regenerative dose, whilst the remaining aliquots were stored prior to administration of the  
30  
31 223 regenerative dose. Measurement of these 'stored' and 'prompt' regenerative doses was then  
32  
33 224 followed by measurement of a 50 Gy test dose, allowing fading to be quantified using the  
34  
35 225 ratio of the sensitivity-corrected 'faded' and 'prompt' signals. The results demonstrate  
36  
37 226 significant fading of regenerated signals (Table 4); nevertheless, fading was generally  
38  
39 227 consistent across all sample groups.

40  
41  
42  
43  
44 228 *Bleaching characteristics*

45  
46  
47  
48 229 Uncertainties concerning the bleaching rates of signals in the different sample groups were  
49  
50 230 addressed by bleaching regenerated doses. Bleaching rates of regenerated IRSL, OSL and  
51  
52 231 TL signals were quantified by exposing aliquots of each sample to 'artificial daylight'  
53  
54 232 fluorescent lighting inside a sealed 'lightbox' for periods of 1 and 8 minutes, and to direct  
55  
56 233 sunlight for a period of 1 minute. Furthermore, the precise form of the bleaching curve was  
57  
58  
59  
60

1  
2  
3 234 investigated by exposing aliquots from two samples (one subglacial and one extraglacial) to  
4  
5 235 'artificial daylight' for periods of up to 32 minutes. The first approach demonstrated mostly  
6  
7 236 consistent rates of bleaching (Table 5). Exposure to the artificial daylight source did appear  
8  
9  
10 237 to bleach subglacial TL more rapidly than for the other sample types, but this was not  
11  
12 238 observed under exposure to direct sunlight, and may therefore reflect unintended heating of  
13  
14 239 the aliquots as a result of the proximity of the fluorescent lighting, or well-known  
15  
16 240 differences between the spectra of fluorescent lighting and sunlight. Bleaching of  
17  
18 241 regenerated signals (e.g. Fig. 8) exhibited an exponential reduction of signal with exposure  
19  
20  
21 242 time that is typical of geological samples.  
22  
23

24  
25 243 *Sensitivity change*  
26

27  
28 244 Residual dose may to some extent reflect sensitivity changes in our samples that cannot be  
29  
30 245 corrected for using normal SAR procedure (e.g. Murray & Wintle 2003). Notably, our  
31  
32 246 multiple-stimulation procedure involves heating aliquots to 500°C prior to administration  
33  
34 247 and measurement of the test dose, which is likely to introduce some sensitivity changes  
35  
36 248 during the first SAR step. Comprehensive dose-recovery tests using a SARA-SAR  
37  
38 249 procedure (as suggested by Wallinga *et al.* 2000) were not possible due to the limited  
39  
40 250 sample material available, and we recommend that additional research be undertaken on the  
41  
42  
43 251 luminescence behaviour of subglacial material from other sites. However, the magnitude of  
44  
45 252 reported effects, which are typically in the range 10–30% (e.g. Wallinga *et al.* 2000; Blair *et*  
46  
47 253 *al.* 2005, Bateman *et al.* 2010), would be insufficient to account for the observed one to two  
48  
49  
50 254 order of magnitude variation of residual dose between sample groups (Fig. 3, Table 2).  
51  
52  
53 255 Furthermore, there are no reasons to suppose that such effects would lead to different  
54  
55 256 behaviour in the subglacial sample group than in any other.  
56  
57  
58  
59  
60

257 **Discussion**258 *Residual doses of the sample groups and their origin*

259 Unexpectedly low subglacial residual dose dominates residual dose variation in samples  
260 obtained at Haut Glacier d'Arolla and is evident even in the difficult-to-reset TL signal (Fig.  
261 3); few extraglacial samples exhibited such low dose, and only in the easy-to-bleach IRSL  
262 and OSL signals (Fig. 3A, B). Also notable is the high residual dose exhibited by samples  
263 of suspended sediment collected from the proglacial stream, which, given the low residual  
264 dose of the subglacial sample group, is not consistent with the expectation that the majority  
265 of sediment transported by such streams is entrained at the ice-bed interface (cf. Gemmill  
266 1994, 1997; Swift *et al.* 2005). However, this expectation may not have been valid at the  
267 time of sampling because periods of falling discharge are generally associated with the  
268 reduced availability of basal sediment (cf. Swift *et al.* 2005), indicating that the majority of  
269 sediment in transport may actually have been extraglacial sediment, sourced from fluvial  
270 erosion of the slopes below the Bouquetins ridge (Fig. 1A, B).

271 A number of previous studies have reported anomalous luminescence behaviour of  
272 samples from glaciated environments, most notably the poor sensitivity of glacial sediment  
273 that arises from poor-intensity signals with weak or absent fast components (e.g. Lukas *et*  
274 *al.* 2007), recuperation of signals after bleaching (e.g. Rhodes & Pownall 1994), or thermal  
275 transfer of signals during SAR procedures (e.g. Rhodes & Bailey 1997). Our analyses have  
276 shown that such problems do not exist in the case of the samples obtained at Haut Glacier  
277 d'Arolla. Furthermore, our analyses indicate consistent luminescence behaviour across all  
278 sample groups and indicate nothing that could reasonably account for the observed one to  
279 two order of magnitude variation in residual dose between the major sample groups. It  
280 follows that we have found no variation in luminescence intensity or behaviour that could

1  
2  
3 281 be ascribed to differences in sample mineralogy or transport/exposure history (cf. Lukas *et*  
4  
5 282 *al.* 2007).

6  
7  
8  
9 283       There is evidence instead that the luminescence of the sediment types sampled at  
10  
11 284 Haut Glacier d'Arolla reflects natural resetting of geologically-accumulated signals. Firstly,  
12  
13 285 extraglacial sample residual dose, which approaches geological saturation levels (cf. Wintle  
14  
15 286 & Murray 2006), is consistent with only partial resetting, such as that resulting from the  
16  
17 287 reworking of glacially-eroded sediments at or near the ice-margin by debris flows and other  
18  
19 288 mass-movement processes. Secondly, although there are many uncertainties regarding the  
20  
21 289 interpretation of the  $D_e(t)$  plots (Fig. 7; see above), rising extraglacial sample OSL  $D_e(t)$  is  
22  
23 290 again consistent with partial resetting, whereas subglacial sample OSL  $D_e(t)$  is almost flat,  
24  
25 291 which is consistent with total resetting (cf. Bailey *et al.* 2003). Thirdly, the relationship of  
26  
27 292 subglacial sample IRSL, OSL and TL residual dose to that of the other sample groups  
28  
29 293 (Table 2), which indicates substantially lower IRSL and OSL residual dose than for the  
30  
31 294 difficult-to-reset TL signal, is consistent with widely-observed bleaching patterns of natural  
32  
33 295 signals as a result of exposure to heat or light (cf. Table 5).

34  
35  
36  
37  
38  
39 296       Assuming subglacial residual dose is indeed a result of natural resetting of near-  
40  
41 297 saturated geological signals, the energy required to have reset such a signal to observed  
42  
43 298 levels can be estimated from rates of bleaching exhibited by regenerated signals when  
44  
45 299 exposed to artificial daylight (Table 5). Knowledge of the signal present in the subglacial  
46  
47 300 bedrock/sediment prior to resetting is also required, but as this is unknown, we substitute  
48  
49 301 this with the mean residual dose exhibited by the other, presumed partially-reset sample  
50  
51 302 groups. By example, the easy-to-bleach subglacial IRSL residual dose is typically 10% of  
52  
53 303 that of the other sample groups (Table 2), which equates to a level of resetting that is  
54  
55 304 produced by approximately 8 minutes of exposure of a regenerated signal to artificial  
56  
57  
58  
59  
60

BOR-017-2010.R2

*Luminescence of subglacial sediment*

1  
2  
3 305 daylight (Table 5). A similar exposure time is arrived at when using the OSL and TL signals  
4  
5 306 (Tables 2, 5). From the irradiance of the artificial source ( $72.92 \text{ W m}^{-2}$ ), it follows that the  
6  
7 307 energy required to reset subglacial signals from levels exhibited by the extraglacial sample  
8  
9 308 groups would be  $\sim 35 \text{ kJ m}^{-2}$ . In terms of exposure to natural light at midday on the glacier  
10  
11 309 surface, when measured irradiance is typically  $\sim 1 \text{ kW m}^{-2}$ ,  $\sim 35 \text{ kJ m}^{-2}$  equates to an  
12  
13 310 exposure time of  $\sim 30$  seconds.  
14  
15  
16  
17

18 311 The above estimate is a minimum estimate of the energy required to have reset  
19  
20 312 subglacial signals to observed values because: (i) extraglacial samples are believed to have  
21  
22 313 been partially-reset and therefore the actual level of signal present in subglacial bedrock or  
23  
24 314 sediment prior to resetting is likely to have been far greater (SAR growth-curves indicate  
25  
26 315 that it may have been  $\sim 1000 \text{ Gy}$ ; Fig. 4); and (ii) resetting is non-linear (Fig. 8), such that  
27  
28 316 the energy required to reduce the luminescence of a sample by a given proportion increases  
29  
30 317 as trapped electrons are released by the resetting process, such that bleaching rates  
31  
32 318 determined from regenerated signals will be significantly greater than for partially-reset  
33  
34 319 natural signals. Nevertheless, this estimate provides a sound and cautious basis from which  
35  
36 320 to assess possible resetting mechanisms.  
37  
38  
39  
40

#### 41 321 *Traditional resetting mechanisms*

42  
43  
44 322 Subglacial sample residual dose cannot be explained by accidental exposure to light or heat  
45  
46 323 since: (i) light sources present during sampling (i.e. head-torch lights and moon light)  
47  
48 324 cannot have delivered the energy required in the time taken to retrieve and bottle the  
49  
50 325 samples; and (ii) drill-water temperatures during borehole drilling were far below the  $200^\circ\text{C}$   
51  
52 326 preheat used during luminescence measurement (B. Hubbard, pers. comm. 2001). Heat  
53  
54 327 generated by friction between clasts, sediment particles and bedrock during glacier sliding  
55  
56 328 or deformation of basal sediment is also negligible. Consequently, potential resetting  
57  
58  
59  
60



1  
2  
3 329 mechanisms are limited to: (i) bleaching of sediment *in situ* by light reaching the glacier bed  
4  
5 330 through open boreholes or through glacier ice; (ii) bleaching of sediment in an extraglacial  
6  
7 331 location prior to re-deposition beneath the glacier; (iii) glacier advance over bleached  
8  
9 332 extraglacial sediment; and (iv) resetting *in situ* as a result of a natural process that does not  
10  
11 333 require heat or light.  
12  
13

14  
15 334 *In situ* bleaching is extremely unlikely because it requires unacceptably low  
16  
17 335 attenuation of light, regardless of whether light is transmitted down boreholes or through  
18  
19 336 glacier ice. In the case of borehole transmission, the Lambert–Beer equation (Grum &  
20  
21 337 Becherer 1979) indicates that, given an ice thickness of ~100 m and mean daily solar  
22  
23 338 irradiance of ~0.3 kW m<sup>-2</sup> (both obtained from field measurements), delivery of 35 kJ m<sup>-2</sup>  
24  
25 339 to the glacier bed via boreholes that were open for 30 days prior to sampling requires  
26  
27 340 attenuation of light in the borehole to be  $\leq 0.12$  m<sup>-1</sup>. Such attenuation rates are unrealistic,  
28  
29 341 given that: (i) typical values for clear water are ~0.2 m<sup>-1</sup>; (ii) boreholes are normally at least  
30  
31 342 partly water-filled (Hubbard *et al.* 1995); (iii) glacier ice has poor reflective properties; and  
32  
33 343 (iv) boreholes have irregular form and ice-wall texture. Furthermore, flushing of sediment  
34  
35 344 between at the glacier bed (e.g. Hubbard *et al.* 1995; Copland *et al.* 1997) indicates that the  
36  
37 345 sampled sediment is unlikely to have been directly beneath the borehole for 30 days.  
38  
39 346 Similar calculations show that the alternative scenario of bleaching via transmission through  
40  
41 347 ice would require ~268 million years, even when reflection of light at the glacier surface is  
42  
43 348 ignored, and a uniform and generous within-ice attenuation coefficient of 0.8 m<sup>-1</sup> is  
44  
45 349 assumed (cf. Grenfell & Maykut 1977; Pegau & Zaneveld 2000).  
46  
47  
48  
49  
50  
51

52  
53 350 Finally, the possibility of extraglacially-bleached sediment existing beneath the  
54  
55 351 glacier is incompatible with current understanding of subglacial processes. Subglacial re-  
56  
57 352 deposition of extraglacially-bleached sediment is extremely unlikely because sediment  
58  
59  
60

BOR-017-2010.R2

Luminescence of subglacial sediment

1  
2  
3 353 transport within subglacial channels, which are occasionally fed by extraglacial streams, is  
4  
5 354 supply-limited (cf. Swift *et al.* 2002, 2005). Sediment can be deposited subglacially when  
6  
7  
8 355 subglacial channels are required to traverse overdeepenings (Alley *et al.* 2003), but the  
9  
10 356 single probable overdeepening at Haut Glacier d’Arolla is not sufficiently deep and does not  
11  
12 357 in any case extend under the drill site (Sharp *et al.* 1993). The alternative scenario of glacier  
13  
14 358 advance over extraglacially-bleached sediment is even more unlikely given the long history  
15  
16 359 of Alpine glacial retreat and the requirement for the overridden sediment to have resisted  
17  
18 360 evacuation by the subglacial drainage system. At Haut Glacier d’Arolla, this system  
19  
20 361 evacuates 2000+ tonnes of sediment per year (Gurnell *et al.* 1992; Swift *et al.* 2002) from a  
21  
22 362 basal sediment layer only ~10 cm thick (Harbor *et al.* 1997), implying spatially-averaged  
23  
24 363 subglacial erosion rates in excess of  $1 \text{ mm a}^{-1}$ , and a mean basal sediment residence time of  
25  
26 364 only 100 years.  
27  
28  
29  
30

31 365 *Alternative resetting mechanisms*  
32  
33

34 366 Calculations of the attenuation of light through ice relate only to absolute intensities of  
35  
36 367 light, whereas it is well-known that shorter-wavelength parts of the spectrum are most  
37  
38 368 attenuated in water (Berger 1990; Bailey *et al.* 2003), resulting in preferential bleaching of  
39  
40 369 feldspar luminescence at water depths beyond those at which effective bleaching of the  
41  
42 370 quartz system can occur, even for turbid water (Sanderson *et al.* 2003, 2007). Since the  
43  
44 371 polymineral aliquots analysed in this study were predominantly composed of feldspar, it is  
45  
46 372 therefore possible that bleaching at the glacier bed could be more effective than anticipated.  
47  
48 373 Without field measurements of the attenuation of different spectra by glacier ice, it is  
49  
50 374 impossible to know just how effective such a resetting mechanism could be. Nevertheless,  
51  
52 375 given that transmission of only a portion of the spectrum would result in a reduction in light  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 376 intensity, and given that the transmitted wavelengths would still undergo at least some  
4  
5 377 attenuation, such a mechanism remains unlikely.  
6  
7

8  
9 378 The absence of plausible resetting mechanisms related to heat or light raises the  
10  
11 379 possibility of more controversial resetting mechanisms. Resetting by subglacial processes  
12  
13 380 has been postulated, particularly the grinding and crushing processes that are responsible for  
14  
15 381 producing and comminuting subglacial debris, because these processes subject individual  
16  
17 382 sediment grains to extremely high stress (cf. Boulton 1974). Various geomechanical  
18  
19 383 resetting mechanisms related to grain stress have been proposed, including: (i) grain  
20  
21 384 fracture, which should result in fewer active luminescence centres that are surrounded by an  
22  
23 385 extended atomic lattice (Toyoda *et al.* 2000); and (ii) the ejection of trapped electrons by  
24  
25 386 stresses imposed on the crystal lattice (Lee & Schwarz 1994) and/or localised frictional  
26  
27 387 heating at grain boundaries (Fukuchi 1989; Lee & Schwarz 1994).  
28  
29  
30  
31

32 388 Since our analyses indicate no substantial differences in the sensitivity of subglacial  
33  
34 389 and extraglacial sample groups of a kind that would indicate a reduction in the number of  
35  
36 390 active luminescence centres, our observations are most consistent with resetting of  
37  
38 391 subglacial luminescence via trapped electron ejection, as envisaged by Lee & Schwarz  
39  
40 392 (1994) and Fukuchi (1989). Although rates of subglacial sediment deformation at Haut  
41  
42 393 Glacier d'Arolla have been suggested to be low in comparison to other similar glaciers  
43  
44 394 (Fischer & Hubbard 1999), the combination of a high annual fine sediment evacuation rate  
45  
46 395 (Swift *et al.* 2002) and a relatively thin basal sediment layer (Harbor *et al.* 1997) indicates a  
47  
48 396 potentially highly erosive subglacial environment in which sedimentary particles are  
49  
50 397 subjected to extremely high stresses. Nevertheless, such processes have also been postulated  
51  
52 398 to induce luminescence (Aitken 1985; Toyoda *et al.* 2000; Zöller *et al.* 2009), and their net  
53  
54 399 effects on luminescence signals remain unknown.  
55  
56  
57  
58  
59  
60

BOR-017-2010.R2

*Luminescence of subglacial sediment*400 *Luminescence as a process tracer in glacial systems*

401 Although this study has indicated unexpected luminescence variation at Haut Glacier  
402 d'Arolla, the results do indicate that luminescence could elucidate glacial sediment  
403 transport pathways. For example, the origin of sediment being evacuated by the subglacial  
404 drainage system could be investigated using a simple two-component mixing-model that  
405 exploits the contrasting residual dose of extraglacial and subglacial sediments.  
406 Nevertheless, uncertainty regarding the nature and efficacy of a subglacial resetting  
407 mechanism means that such studies would not be easy to apply without further investigation  
408 of the luminescence of glacial erosion products. Further studies of subglacial sediments that  
409 have been obtained *in situ* must be paramount (see below), but such samples are logistically  
410 difficult to obtain. Further investigation of diurnal variation in the residual dose of sediment  
411 evacuated by subglacial drainage systems would also be worthwhile (cf. Gemmell 1994,  
412 1997), but this too is logistically difficult because stream samples are very difficult to obtain  
413 under light-free conditions.

414 Further investigation of a possible subglacial resetting processes might include  
415 sampling of a more extensive network of boreholes, since resetting should vary with basal  
416 shear stress, which should be highest where the ice is thickest and is moving fastest, and  
417 sediment transport distance, which should increase downglacier (provided that not all  
418 sediment that is produced by subglacial erosion is at some point evacuated by the subglacial  
419 drainage system). Sampling of boreholes over time should also be undertaken to fully  
420 eliminate resetting as a result of the transmission of light via boreholes and the  
421 contamination of borehole sediment by sediment bleached in englacial and supraglacial  
422 locations. The results of such work might enable the identification of other glaciers with  
423 subglacial conditions that are conducive to resetting, as well as the identification of

1  
2  
3 424 Quaternary sediments that are likely to have experienced transport, and thus resetting, in  
4  
5 425 such environments. Ultimately, such work could enable the dating of subglacially-deposited  
6  
7  
8 426 tills using luminescence-based techniques, as well as the quantification of sediment strain  
9  
10 427 histories and/or residence times in the contemporary subglacial environment.  
11

12  
13 428 Finally, the results of this study indicate some potential to use the luminescence  
14  
15 429 sensitivity to elucidate sediment transport pathways in a way that is similar to that proposed  
16  
17 430 for residual dose (above). Specifically, SAR measurements (Fig. 4) indicate that the TL  
18  
19 431 saturation of subglacial sediment was markedly higher than for the other sediment types,  
20  
21 432 with that  $D_e$  values at 90% of saturation (as indicated by the form of the curves fitted to the  
22  
23 433 SAR measurements) being three times greater than values for other sediment types.  
24  
25  
26 434 However, this feature of the data is not consistent with the anticipated effects of glacial  
27  
28 435 crushing, which might be expected to reduce the saturation point of glacial sediment  
29  
30 436 relative to non-glacial sediment by reducing the number of luminescence centres  
31  
32 437 surrounded by an extended atomic lattice (cf. Lee & Schwarz 1994). Further work is  
33  
34 438 therefore necessary to understand the source of this effect.  
35  
36  
37  
38

### 39 439 **Conclusion**

40  
41  
42 440 This study has shown that the luminescence of subglacial sediment obtained from boreholes  
43  
44 441 drilled to the bed of Haut Glacier d'Arolla through ~100 m of glacier ice appears to have  
45  
46 442 been substantially reset relative to that of extraglacial sediments sampled within the same  
47  
48 443 small catchment. Although further work is required, the results also demonstrate that the  
49  
50 444 observed differences in residual dose cannot readily be explained by differences in the  
51  
52 445 luminescence characteristics or behaviour of the various sample groups. The discussion has  
53  
54 446 further shown that satisfactory process-based explanations related to exposure to heat or  
55  
56 447 light cannot explain observed subglacial sediment residual dose, and we therefore conclude  
57  
58  
59  
60

BOR-017-2010.R2

*Luminescence of subglacial sediment*

1  
2  
3 448 that further work should also investigate alternative resetting processes, including trapped  
4  
5 449 charge ejection as a result of the grinding and crushing that both produces and comminutes  
6  
7 450 sediment in the subglacial environment. Such processes could enable the dating of  
8  
9 451 subglacially-deposited tills using luminescence-based techniques, as well as the  
10  
11 452 quantification of sediment strain histories and/or residence times in the contemporary  
12  
13 453 subglacial environment.  
14  
15  
16

17 454 It is hoped that the need for further investigation will be at least partially fulfilled by  
18  
19 455 a recently-started research project that aims to shear sediment with naturally-acquired  
20  
21 456 luminescence under conditions that are representative of the subglacial environment (Swift  
22  
23 457 *et al.* 2010). Nevertheless, further study of subglacial sediment that has been sampled *in situ*  
24  
25 458 is also required if the nature and efficacy of any such subglacial resetting is to be rigorously  
26  
27 459 quantified and constrained. Such studies are necessary to identify contemporary and  
28  
29 460 Quaternary glacial environments that are conducive to the resetting of subglacial sediment  
30  
31 461 and the associated sediments and landforms that may provide evidence of having been  
32  
33 462 glacially-reset.  
34  
35  
36  
37

38  
39 463 Acknowledgements. DAS acknowledges a University of Glasgow Scholarship and a BP  
40  
41 464 Royal Society of Edinburgh Personal Research Fellowship. Fieldwork by PWN and  
42  
43 465 ICC was supported by the Carnegie Trust for the Universities of Scotland and the  
44  
45 466 University of Glasgow Robertson Bequest. The manuscript was greatly improved as a  
46  
47 467 result of discussions with members of the luminescence community, as well as  
48  
49 468 numerous reviews, and we thank, in no particular order, Alistair Gemmell, Lewis Owen,  
50  
51 469 Jakob Wallinga, Mark Bateman, Bert Roberts, Eddie Rhodes, Chris Burbidge, and one  
52  
53 470 anonymous reviewer for their interest and invaluable recommendations and advice. We  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 471 also thank Anne Sommerville and Lorna Campbell for laboratory assistance, and Tris  
4  
5 472 Irvine-Fynn for his assistance in the field.  
6  
7

8  
9 473 **References**

10  
11 474 Aitken, M.J. 1985: *Thermoluminescence dating*. 351 pp. Academic Press, London.  
12  
13

14 475 Aitken, M.J. 1998: *An introduction to optical dating: The dating of quaternary sediments by*  
15  
16 476 *the use of photon-stimulated luminescence*. 267 pp. Oxford University Press, Oxford.  
17  
18

19  
20 477 Alley, R.B., Cuffey, K.M., Evenson, E.B., Strasser, J.C., Lawson, D.E. & Larson, G.J.  
21  
22 478 1997: How glaciers entrain and transport basal sediment: Physical constraints. *Quaternary*  
23  
24 479 *Science Reviews* **19**, 1017–1038.  
25  
26

27  
28 480 Alley, R.B., Lawson, D.E., Larsen, G.J., Evenson, E.B. & Baker, G.S. 2003: Stabilizing  
29  
30 481 feedbacks in glacier-bed erosion. *Nature* **424**, 758–760.  
31  
32

33 482 Armitage, S.J., Botha, G.A., Duller, G.A.T., Wintle, A.G., Rebêlo, L.P. & Momade, F.J.  
34  
35 483 2006: The Formation and Evolution of the Barrier Islands of Bazaruto and Inhaca,  
36  
37 484 Mozambique. *Geomorphology* **82**, 295–308.  
38  
39

40  
41 485 Bailey, R.M., Singarayer, J.S., Ward, S. & Stokes, S. 2003: Identification of partial resetting  
42  
43 486 using De as a function of illumination time. *Radiation Measurements* **37**, 511–518.  
44  
45

46  
47 487 Bateman M.D., Boulter C., Carr, A.S., Frederick C.D., Wilder, M. & Peter, D. 2007:  
48  
49 488 Preserving the palaeoenvironmental record in drylands: Bioturbation and its significance for  
50  
51 489 luminescence dating derived chronologies. *Sedimentary Geology* **195**, 5–19.  
52  
53

54 490 Bateman, M.D., Murton J.B. & Boulter C. 2010. The source of De variability in periglacial  
55  
56 491 sand wedges: Depositional processes versus measurement issues. *Quaternary Geochronology*  
57  
58 492 **5**, 250–256.  
59  
60

BOR-017-2010.R2

Luminescence of subglacial sediment

- 1  
2  
3 493 Berger, G.W. 1990: Effectiveness of natural zeroing of the thermoluminescence in  
4  
5 494 sediments. *Journal of Geophysical Research* **95B**, 12375–12397.  
6  
7  
8  
9 495 Blair M.W., Yukihara E.G., & McKeever S.W.S. 2005: Experiences with single-aliquot  
10  
11 496 OSL procedures using coarse-grain feldspars. *Radiation Measurements* **39**, 361–374.  
12  
13  
14 497 Blake, E.W. & Clarke, G.K.C. 1991: Subglacial water and sediment samplers. *Journal of*  
15  
16 498 *Glaciology* **37**, 188–190.  
17  
18  
19  
20 499 Boulton, G.S. 1974: Processes and patterns of glacial erosion. In Coates, D.R. (ed.): *Glacial*  
21  
22 500 *Geomorphology (Proceedings of the Fifth Annual Geomorphology Symposia,*  
23  
24 501 *Binghampton)*, 41–87. Allen & Unwin, London.  
25  
26  
27 502 Burbidge, C.I., Sanderson, D.C.W., Housley, R.A., & Allsworth Jones, P. 2007: Survey of  
28  
29 503 Palaeolithic sites by luminescence profiling, a case study from Eastern Europe. *Quaternary*  
30  
31 504 *Geochronology* **2**, 296–302.  
32  
33  
34  
35 505 Copland, L., Harbor, J., Gordon, S. & Sharp, M. 1997: The use of borehole video in  
36  
37 506 investigating the hydrology of a temperate glacier. *Hydrological Processes* **11**, 211–224.  
38  
39  
40  
41 507 Dreimanis, A., Hutt, G., Raukas, A. & Whippey, P.W. 1978: Dating methods of Pleistocene  
42  
43 508 deposits and their problems: I. Thermoluminescence dating. *Geoscience Canada* **5**, 55–60.  
44  
45  
46 509 Fischer, U.H. & Hubbard, B. 1999: Subglacial sediment textures: character and evolution at  
47  
48 510 Haut Glacier d'Arolla, Switzerland. *Annals of Glaciology* **28**, 241–246.  
49  
50  
51  
52 511 Fuchs, M. & Owen, L.A. 2008: Luminescence dating of glacial and associated sediments:  
53  
54 512 review, recommendations and future directions. *Boreas* **37**, 636–659.  
55  
56  
57  
58  
59  
60



BOR-017-2010.R2

Luminescence of subglacial sediment

- 1  
2  
3 513 Fukuchi, T. 1989: Increase of radiation sensitivity of ESR centres by faulting and criteria of  
4  
5 514 fault dates. *Earth and Planetary Science Letters* **94**, 109–122.  
6  
7  
8  
9 515 Galbraith, R.F. 2002: A note on the variance of a background-corrected OSL count. *Ancient*  
10  
11 516 *TL* **20**, 49–51.  
12  
13  
14 517 Gemmell, A.M.D. 1994: Thermoluminescence in suspended sediment of glacier meltwater  
15  
16 518 streams. *Journal of Glaciology* **40**, 158–166.  
17  
18  
19 519 Gemmell, A.M.D. 1997: Fluctuations in the thermoluminescence signal of suspended  
20  
21 520 sediment in an alpine glacial meltwater stream. *Quaternary Science Reviews* **16**, 281–290.  
22  
23  
24  
25 521 Goodsell, B., Hambrey, M.J. & Glasser, N.F. 2005: Debris transport in a temperate valley  
26  
27 522 glacier: Haut Glacier d’Arolla, Valais, Switzerland. *Journal of Glaciology* **51**, 139–146.  
28  
29  
30 523 Grenfell, T.C. & Maykut, G.A. 1977: The optical properties of ice and snow in an Arctic  
31  
32 524 Basin. *Journal of Glaciology* **18**, 445–463.  
33  
34  
35  
36 525 Grum, F. & Becherer, R.J. 1979: *Optical Radiation Measurements, Vol. 1: Radiometry*. 335  
37  
38 526 pp. Academic Press, New York.  
39  
40  
41 527 Gurnell, A.M., Clark, M.J. & Hill, C.T. 1992: Analysis and interpretation of patterns within  
42  
43 528 and between hydroclimatological time series in an Alpine glacier basin. *Earth Surface*  
44  
45 529 *Processes and Landforms* **17**, 821–839.  
46  
47  
48  
49 530 Harbor, J., Sharp, M., Copland, L., Hubbard, B., Nienow, P. & Mair, D. 1997: Influence of  
50  
51 531 subglacial drainage conditions on the velocity distribution within a glacier cross section.  
52  
53 532 *Geology* **25**, 739–742.  
54  
55  
56  
57  
58  
59  
60

BOR-017-2010.R2

Luminescence of subglacial sediment

- 1  
2  
3 533 Hubbard, B.P., Sharp, M.J., Willis, I.C., Nielsen, M.K. & Smart, C.C. 1995: Borehole  
4  
5 534 water-level variation and the structure of the subglacial hydrological system of Haut Glacier  
6  
7 535 d'Arolla, Valais, Switzerland. *Journal of Glaciology* **41**, 572-583.  
8  
9  
10 536 Krbetschek, M.R., Götze, J., Dietrich, A. & Trautmann, T. 1997: Spectral information from  
11  
12 537 minerals relevant for luminescence dating. *Radiation Measurements* **27**, 695–748.  
13  
14  
15 538 Lee, H.K. & Schwarcz, H.P. 1994: Criteria for complete zeroing of ESR signals during  
16  
17 539 faulting of the San Gabriel fault zone, southern California. *Tectonophysics* **235**, 317–337.  
18  
19  
20 540 Lian, O.B. & Roberts, R.G. 2006: Dating the Quaternary: progress in luminescence dating  
21  
22 541 of sediments. *Quaternary Science Reviews* **25**, 2449–2468.  
23  
24  
25 542 Liritzis, I. 2000: Advances in thermo- and opto-luminescence dating of environmental  
26  
27 543 materials (sedimentary deposits). Part II: Applications. *Global Nest: the International*  
28  
29 544 *Journal* **2**, 29–49.  
30  
31  
32 545 Lukas, S., Spencer, J.Q.G., Robinson, R.A.J. & Benn, D.I. 2007: Problems associated with  
33  
34 546 luminescence dating of Late Quaternary glacial sediments in the NW Scottish Highlands.  
35  
36 547 *Quaternary Geochronology* **2**, 243-248.  
37  
38  
39 548 Morozov, G.V. 1968: The relative dating of quaternary Ukrainian sediments by the TL  
40  
41 549 method. In *Proceedings of the VIIIth International Quaternary Association Congress*  
42  
43 550 *(Paris, France, 1969)*. 167 pp. Naukova Dumka, Kiev (United States Geological Survey  
44  
45 551 Library Catalogue No. 208M8280, Washington, DC).  
46  
47  
48 552 Murray, A.S. & Wintle, A.G. 2003: The single aliquot regenerative dose protocol: potential  
49  
50 553 for improvements in reliability. *Radiation Measurements* **37**, 377–381.  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

BOR-017-2010.R2

Luminescence of subglacial sediment

- 1  
2  
3 554 Nienow, P., Sharp, M. & Willis, I. 1998: Seasonal changes in the morphology of the  
4  
5 555 subglacial drainage system, Haut Glacier d'Arolla, Switzerland. *Earth Surface Processes*  
6  
7 556 *and Landforms* **23**, 825–843.
- 8  
9  
10  
11 557 Pegau, W.S. & Zaneveld, J.R.V. 2000: Field measurements of in-ice radiance. *Cold Regions*  
12  
13 558 *Science and Technology* **31**, 33–46.
- 14  
15  
16 559 Rhodes, E.J. & Bailey, R.M. 1997: Thermal transfer effects observed in the luminescence of  
17  
18 560 quartz from recent glaciofluvial sediments. *Quaternary Geochronology (Quaternary*  
19  
20 561 *Science Reviews)* **16**, 291–298.
- 21  
22  
23  
24 562 Rhodes, E.J. & Pownall, L. 1994: Zeroing of the OSL signal in quartz from young  
25  
26 563 glaciofluvial sediments. *Radiation Measurements* **23**, 581–585.
- 27  
28  
29 564 Rink, W.J., Toyoda, S., Rees-Jones, J. & Schwarcz, H.P. 1999: Thermal activation of OSL as  
30  
31 565 a geothermometer for quartz grain heating during fault movements. *Radiation Measurements*  
32  
33 566 **30**, 97–105.
- 34  
35  
36  
37 567 Sanderson, D.C.W., Bishop, P., Stark, M.T. & Spencer, J.Q. 2003: Luminescence dating of  
38  
39 568 Anthropogenically reset sediments from Angkor Borei, Mekong Delta, Cambodia.  
40  
41 569 *Quaternary Science Reviews* **22**, 1111–1122
- 42  
43  
44  
45 570 Sanderson, D.C.W., Bishop, P., Stark, M., Alexander, S. & Penney, D. 2007: Luminescence  
46  
47 571 dating of canal sediments from Angkor Borei, Mekong Delta, Southern Cambodia.  
48  
49 572 *Quaternary Geochronology* **2**, 322–329.
- 50  
51  
52  
53 573 Sanderson, D.C.W. & Murphy, S. 2010: Using simple portable OSL measurements and  
54  
55 574 laboratory characterisation to help understand complex and heterogeneous sediment  
56  
57 575 sequences for luminescence dating. *Quaternary Geochronology* **5**, 299–305.
- 58  
59  
60

BOR-017-2010.R2

Luminescence of subglacial sediment

- 1  
2  
3 576 Sharp, M., Richards, K., Willis, I., Arnold, N., Nienow, P., Lawson, W. & Tison, J.-L. 1993:  
4  
5 577 Geometry, bed topography and drainage system structure of the Haut Glacier d'Arolla,  
6  
7 578 Switzerland. *Earth Surface Processes and Landforms* **18**, 557–571.  
8  
9  
10  
11 579 Singhvi, A.K., Banerjee, D., Pande, K., Gogte, V. & Valdiya, K.S. 1994: Luminescence  
12  
13 580 studies on Neotectonic events in South Central Kumaun Himalaya - A feasibility Study,  
14  
15 581 *Quaternary Science Reviews* **13**, 595–600.  
16  
17  
18  
19 582 Stone, D.B. & Clarke, G.K.C. 1996: In situ measurements of basal water quality and  
20  
21 583 pressure as an indicator of the character of subglacial drainage systems. *Hydrological*  
22  
23 584 *Processes* **10**, 615–628.  
24  
25  
26 585 Swift, D.A., Bateman, M.D. & Piotrowski, J.A. 2008: Geomechanical modification of  
27  
28 586 sediment luminescence. funded by the Danish Agency for Science, Technology and  
29  
30 587 Innovation  
31  
32  
33  
34 588 Swift, D.A., Nienow, P.W. & Hoey, T.B. 2005: Basal sediment evacuation by subglacial  
35  
36 589 meltwater: suspended sediment transport from Haut Glacier d'Arolla, Switzerland. *Earth*  
37  
38 590 *Surface Processes and Landforms* **30**, 867–883.  
39  
40  
41  
42 591 Swift, D.A., Nienow, P.W., Spedding, N. & Hoey, T.B. 2002: Geomorphic implications of  
43  
44 592 subglacial drainage configuration: rates of basal sediment evacuation controlled by seasonal  
45  
46 593 drainage system evolution. *Sedimentary Geology* **149**, 5–19.  
47  
48  
49  
50 594 Swift, D.A., Bateman, M.D. & Piotrowski, J.A. 2010. Can subglacial processes reset the  
51  
52 595 luminescence of basal sediment? European Geophysical Union General Assembly 2010, 2–  
53  
54 596 7 May, Vienna, Austria. *Geophysical Research Abstracts* **12**, EGU2010-13429.  
55  
56  
57  
58  
59  
60

BOR-017-2010.R2

Luminescence of subglacial sediment

- 1  
2  
3 597 Tranter, M., Sharp, M.J., Lamb, H.R., Brown, G.H., Hubbard, B.P. & Willis, I.C. 2002:  
4  
5 598 Geochemical weathering at the bed of Haut Glacier d'Arolla, Switzerland—a new model.  
6  
7 599 *Hydrological Processes* **16**, 959–993.  
8  
9  
10  
11 600 Thomas, P.J., Murray, A.S., Kjær, K.H., Funder, S. & Larsen, E. 2006: Optically Stimulated  
12  
13 601 Luminescence (OSL) dating of glacial sediments from Arctic Russia—depositional  
14  
15 602 bleaching and methodological aspects. *Boreas* **35**, 587–599.  
16  
17  
18  
19 603 Toyoda, S., Rink, W.J., Schwarcz, H.P. & Rees-Jones, J. 2000: Crushing effects on TL and  
20  
21 604 OSL on quartz: Relevance to fault dating. *Radiation Measurements* **32**, 667–672.  
22  
23  
24 605 Wallinga J., Murray A. & Duller, G. 2000: Underestimation of equivalent dose in single-  
25  
26 606 aliquot optical dating of feldspars caused by preheating. *Radiation Measurements* **32**, 691–  
27  
28 607 695.  
29  
30  
31  
32 608 Wallinga, J., Murray, A.S., Duller, G.A.T. & Tornqvist, T.E. 2001: Testing optically  
33  
34 609 stimulated luminescence dating of sand-sized quartz and feldspar from fluvial deposits.  
35  
36 610 *Earth and Planetary Science Letters* **193**, 617–630.  
37  
38  
39  
40 611 Wintle, A.G. & Huntley, D.J., 1979: Thermoluminescence dating of a deep sea ocean core.  
41  
42 612 *Nature* **279**, 710–712.  
43  
44  
45 613 Wintle, A.G. & Murray, A.S. 2006: A review of quartz optically stimulated luminescence  
46  
47 614 characteristics and their relevance in single-aliquot regeneration dating protocols. *Radiation*  
48  
49 615 *Measurements* **41**, 369–391.  
50  
51  
52  
53 616 Zöller, L., Blanchard, H., & McCammon, C. 2009: Can temperature assisted hydrostatic  
54  
55 617 pressure reset the ambient TL of rocks? – A note on the TL of partially heated country rock  
56  
57 618 from volcanic eruptions. *Ancient TL* **27**, 15–22.  
58  
59  
60

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  

## FIGURE CAPTIONS

1  
2 Figure 1. A. Map of Haut Glacier d'Arolla, Switzerland showing sampling locations discussed in  
3 the text. The inset key indicates the number of samples obtained at each location (see  
4 Supplementary Material for a full sample list). B. Photograph looking SE over the glacier. The  
5 approximate location of the drill site, where subglacial sediment was sampled, is indicated by the  
6 filled triangle. Surface sediment was sampled from marginal moraine in the upper glacier basin,  
7 and stream sediments were obtained from two tributaries of a nearby non-glacier-fed marginal  
8 stream and from the eastern subglacial drainage system portal (symbols indicate sampling  
9 locations). Glacier-fed extraglacial streams below Bouquetins ridge (numbered 1 to 4) also enter  
10 the glacial drainage system and emerge from the eastern drainage portal. C. Distribution of major  
11 rock types and sediments in the catchment and surrounding areas (after Tranter *et al.* 2002).

12 Figure 2. Indicative IRSL and OSL shine-down curves and background-subtracted TL glow-  
13 curves measured during read-out of naturally-trapped charge from individual discs prepared from  
14 samples 1277 (subglacial sediment), 1280 (portal stream sediment), 1293 (marginal stream  
15 sediment) and 1296 (surface sediment). IRSL and OSL signals were calculated by subtracting the  
16 underlying background (determined over the last 14.4 s and 7.2 s of observed signal for IRSL and  
17 OSL, respectively) from the initial signal (obtained by integration over the first 4.8 s and 2.4 s of  
18 observed signal for IRSL and OSL, respectively); TL signals were obtained by integration of the  
19 observed signal over the range 300 to 400°C.

20 Figure 3. Initial Residual Dose ( $D_f$ ) estimates obtained using the simple polymineral single-  
21 aliquot multiple-stimulation screening procedure (see text). Two independent determinations of  
22 IRSL, OSL and TL  $D_f$  were obtained for each sample (i.e.  $D_{f1}$  and  $D_{f2}$ ) and these are shown on  
23 separate axes; error bars reflect photon counting statistics (Galbraith 2002) plus an estimated 2%  
24 analytical error (cf. Armitage *et al.* 2006). Subglacial samples are shown as filled triangles; see  
25 Fig. 1 for the key to other sample types.  $D_f$  values with errors that exceeded  $\pm 100\%$ , largely as a

1  
2  
3 26 result of very weak  $L_n$  signals, were treated with caution; hence, one portal stream sample has  
4  
5 27 been removed from (A) and six samples (including four subglacial sediment samples) have been  
6  
7  
8 28 removed from (B). See Supplementary Material for the full dataset.  
9

10  
11 29 Figure 4. Sensitivity-corrected luminescence growth-curves for various samples using a multiple-  
12  
13 30 stimulation single-aliquot regenerative-dose (SAR) procedure (see text); regeneration points are  
14  
15 31 means of eight aliquots per sample. All plots include a recycling point at 50 Gy; zero dose-point  
16  
17 32 values (not shown) and recycling ratios are summarised in Table 4. Fitted curves are fourth-order  
18  
19 33 polynomials that were also used to calculate the SAR  $D_e$  estimates (Table 3); for all curves  
20  
21 34  $R^2 > 0.999$  and the standard deviation of the back-transformed residuals is  $< 3\%$ . Key to lines and  
22  
23 35 symbols for all plots is shown in (A); see Fig. 1A for sample key.  
24  
25  
26

27  
28 36 Figure 5. Integration intervals ( $a-f$ ) used to plot background-corrected IRSL and OSL signal-  
29  
30 37 decay (Fig. 6) and  $D_e(t)$  (Fig. 7) (background obtained from interval  $x$ ).  
31  
32

33  
34 38 Figure 6. Signal-decay plots obtained from IRSL and OSL shine-down curves for various  
35  
36 39 samples: (A) natural IRSL; (B) natural OSL; (C) regenerated IRSL; and (D) regenerated OSL  
37  
38 40 (key to all samples shown in (A)). The plots show sensitivity-corrected luminescence ( $L_X$ ) for  
39  
40 41 successive integration intervals (i.e.  $L_X = L_X/T_X$ , where  $x$  is the integration interval) as a  
41  
42 42 proportion of the sensitivity-corrected initial signal ( $L_A$ ) in interval  $a$  (integration intervals shown  
43  
44 43 in Fig. 5). Values are means of eight aliquots per sample (except for 1279 in (A) and (B), where  
45  
46 44 values are means of seven determinations). Shine-down curves were measured using the multiple-  
47  
48 45 stimulation approach of Table 1.  
49  
50


51  
52 46 Figure 7.  $D_e(t)$  plots ( $D_e = L_n/L_r \times 50$ ) obtained from shine-down curves for various samples: (A)  
53  
54 47 and (B) natural IRSL; (C) and (D) natural OSL (key to all samples shown in (A)). Values are  
55  
56 48 means of eight aliquots per sample; integration intervals are shown in Fig. 5.  
57  
58  
59  
60

1  
2  
3 49 Figure 8. Resetting of regenerated IRSL signals in sample 1285 (subglacial sediment; filled  
4  
5 50 triangles) and 1296 (surface sediment) as a result of exposure to an artificial daylight source. The  
6  
7  
8 51 graph shows the observed signal after bleaching ( $L_b$ ) as a proportion of the observed signal with  
9  
10 52 no bleaching ( $L_u$ ). Symbols are means of two aliquots per sample; errors were calculated as for  
11  
12 53 Fig. 3.  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

For Review Only




1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



172x125mm (600 x 600 DPI)


www.Only



167x55mm (600 x 600 DPI)

Review Only


1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



164x53mm (600 x 600 DPI)

Review Only

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60




248x62mm (600 x 600 DPI)

For Review Only


1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



56x58mm (600 x 600 DPI)


1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



171x127mm (600 x 600 DPI)


Only

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



172x127mm (600 x 600 DPI)

Only



53x54mm (600 x 600 DPI)

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



Table 1: Multiple-stimulation procedure used for initial screening

| Step <sup>1</sup> | Treatment  | Observed <sup>2</sup> |
|-------------------|--|-----------------------|
| 1                 | Preheat (220°C for 30s)  | –                     |
| 2                 | Stimulate IRSL (60s at 60°C)   | $L_n^{IRSL}$          |
| 3                 | Stimulate OSL (30s at 125°C)   | $L_n^{OSL}$           |
| 4                 | Stimulate TL (ambient to 500°C at 5°C s <sup>-1</sup> )              | $L_n^{TL}$            |
| 5                 | Stimulate TL (ambient to 500°C at 5°C s <sup>-1</sup> ) <sup>3</sup> | –                     |
| 6                 | Give test dose, $D_T$ (5 Gy)   | –                     |
| 7                 | Preheat (220°C for 30s)  | –                     |
| 8                 | Stimulate IRSL (60s at 60°C)   | $T_n^{IRSL}$          |
| 9                 | Stimulate OSL (30s at 125°C)   | $T_n^{OSL}$           |
| 10                | Stimulate TL (ambient to 500°C at 5°C s <sup>-1</sup> )              | $T_n^{TL}$            |
| 11                | Stimulate TL (ambient to 500°C at 5°C s <sup>-1</sup> ) <sup>3</sup> | –                     |

<sup>1</sup>Steps 1–11 repeated following a 50 Gy regenerative dose.

<sup>2</sup>Observed signals obtained from raw stimulation curves (see Fig. 2).

<sup>3</sup>Second heating for TL background subtraction.

Table 2: Comparison of  $D_r$  exhibited by each of the sample groups

| Description              | IRSL    |                   | OSL     |                   | TL      |                   |
|--------------------------|---------|-------------------|---------|-------------------|---------|-------------------|
|                          | $D_r^1$ | $D_{rs}/D_{rx}^2$ | $D_r^1$ | $D_{rs}/D_{rx}^2$ | $D_r^1$ | $D_{rs}/D_{rx}^2$ |
| Subglacial sediment      | 12±8.4  | –                 | 2.8±2.0 | –                 | 90±13   | –                 |
| Portal stream sediment   | 512±77  | 0.02              | 292±208 | 0.01              | 329±19  | 0.27              |
| Marginal stream sediment | 151±120 | 0.08              | 131±121 | 0.02              | 287±66  | 0.31              |
| Surface sediment         | 182±135 | 0.07              | 189±147 | 0.02              | 281±52  | 0.32              |

<sup>1</sup>Values are means of the  $D_r$  estimates shown in Fig. 3; errors are  $\pm 1\sigma$ .

<sup>2</sup>Mean subglacial  $D_r$  (i.e.  $D_{rs}$ ) as a fraction of mean  $D_r$  of the other sample types (i.e.  $D_{rx}$ ).

Table 3:  $D_r$  (i.e. initial screening approach) and SAR  $D_e$  for various samples

| Sample | Description              | $D_r^1$ |         |        | $D_e^2$ |         |         |
|--------|--------------------------|---------|---------|--------|---------|---------|---------|
|        |                          | IRSL    | OSL     | TL     | IRSL    | OSL     | TL      |
| 1277   | Subglacial sediment      | 22±13   | 8.7±8.4 | 116±67 | 28±0.1  | 6±0.2   | 102±1.0 |
| 1285   | Subglacial sediment      | 7.0±2.1 | 3.5±4.6 | 70±13  | 11±0.1  | 3±0.1   | 68±0.8  |
| 1279   | Portal stream sediment   | 513±94† | 245±91† | 325±44 | 453±4.7 | 224±6.0 | 397±3.4 |
| 1292   | Marginal stream sediment | 208±42  | 136±33  | 291±23 | 202±2.4 | 135±1.2 | 475±10  |
| 1296   | Surface sediment         | 138±63† | 77±53†  | 294±61 | 126±0.1 | 75±2.4  | 360±3.7 |
| 1298   | Surface sediment         | 294±95† | 161±93  | 325±65 | 260±2.0 | 157±2.6 | 400±4.5 |

<sup>1</sup>Values are means of eight aliquots per sample (unless indicated by †); errors are ±1σ.

<sup>2</sup> $D_e$  interpolated from the corresponding SAR growth curve (Fig. 4) using the mean sensitivity-corrected natural signal ( $L_n/T_n$ ; n=8); ±1σ error has been estimated from the standard error of the regression curve.

†Values are means of seven aliquots per sample, owing to measurement faults.

Table 4: SAR recycling, recuperation and fading characteristics for various samples

| Sample <sup>1</sup> | Mean recycling ratio <sup>2,3</sup> |           |           | Mean recuperated signal (% of $N$ ) <sup>2,4</sup> |           |           | Signal remaining after 95 days <sup>5</sup> |           |           |
|---------------------|-------------------------------------|-----------|-----------|--|-----------|-----------|---|-----------|-----------|
|                     | IRSL                                | OSL       | TL        | IRSL   | OSL       | TL        | IRSL  | OSL       | TL        |
| 1277                | 0.86±0.13                           | 1.09±0.19 | 0.97±0.05 | 0.52±0.88  | 9.23±8.16 | 0.10±0.08 | 0.62±0.11                                   | 0.47±0.16 | 0.58±0.08 |
| 1285                | 0.89±0.07                           | 0.86±0.11 | 0.94±0.05 | 1.78±1.64  | 24.5±23.9 | 0.12±0.08 | 0.57±0.20                                   | 0.51±0.11 | 0.58±0.10 |
| 1279                | 0.92±0.06                           | 1.24±0.44 | 0.89±0.07 | 0.02±0.03  | 0.17±0.26 | 0.02±0.01 | 0.60±0.07                                   | 0.70±0.20 | 0.74±0.09 |
| 1292                | 0.85±0.08                           | 1.08±0.31 | 0.84±0.03 | 0.04±0.03  | 0.30±0.24 | 0.03±0.03 | 0.70±0.08                                   | 0.75±0.05 | 0.89±0.03 |
| 1296                | 0.94±0.07                           | 1.07±0.15 | 0.92±0.05 | 0.03±0.03  | 0.23±0.16 | 0.05±0.06 | 0.58±0.06                                   | 0.61±0.12 | 0.73±0.16 |
| 1298                | 0.95±0.08                           | 1.02±0.28 | 0.94±0.05 | 0.04±0.04  | 0.70±1.07 | 0.03±0.03 | 0.70±0.06                                   | 0.68±0.14 | 0.79±0.08 |

<sup>1</sup>See Table 3 for sample descriptions.

<sup>2</sup>Values are means of eight aliquots per sample; errors are  $\pm 1\sigma$ .

<sup>3</sup>Recycling ratio obtained from the sensitivity-corrected regenerative signals  $R_1$  and  $R_0$  (see text).

<sup>4</sup>The sensitivity-corrected regenerated signal  $R_2$  (zero dose; see text) is expressed as a % of the sensitivity-corrected natural signal ( $L_n/T_n$ ).

<sup>5</sup>Ratio of the mean sensitivity-corrected regenerated signal in four stored discs to the mean prompt signal in four control discs  $\pm 1\sigma$ .

Table 5: Remaining dose after various periods of exposure to different light sources, as a fraction of the 50 Gy original dose

|   | IRSL      |           | OSL       |           | TL        |           |
|---|-----------|-----------|-----------|-----------|-----------|-----------|
|   | 1 min     | 8 mins    | 1 min     | 8 mins    | 1 min     | 8 mins    |
| <i>Artificial daylight</i> <sup>1,2</sup> : |           |           |           |           |           |           |
| Subglacial samples                          | 0.62±0.06 | 0.10±0.02 | 0.80±0.09 | 0.15±0.04 | 0.29±0.02 | 0.22±0.02 |
| Proglacial stream samples                   | 0.64±0.13 | 0.14±0.01 | 0.53±0.19 | 0.14±0.03 | 0.62±0.09 | 0.26±0.03 |
| Marginal stream samples                     | 0.65±0.12 | 0.13±0.02 | 0.69±0.10 | 0.13±0.05 | 0.64±0.09 | 0.26±0.02 |
| Surface sediment samples                    | 0.61±0.17 | 0.14±0.03 | 0.66±0.18 | 0.18±0.07 | 0.69±0.06 | 0.30±0.05 |
| <i>Direct sunlight</i> <sup>1,3</sup> :     |           |           |           |           |           |           |
| Subglacial samples                          | 0.05±0.03 | –         | 0.07±0.04 | –         | 0.66±0.03 | –         |
| Proglacial stream samples                   | 0.06±0.02 | –         | 0.08±0.08 | –         | 0.59±0.04 | –         |
| Marginal stream samples                     | 0.05±0.02 | –         | 0.06±0.04 | –         | 0.56±0.04 | –         |
| Surface sediment samples                    | 0.05±0.02 | –         | 0.03±0.05 | –         | 0.60±0.05 | –         |

<sup>1</sup>Remaining dose calculated as  $L_i/L_u$ , where  $L_i$  is the observed signal after exposure and  $L_u$  is the observed signal with no exposure; values are means for each sample group (the number of samples in each group is shown in Fig. 1A); errors are  $\pm 1\sigma$ .

<sup>2</sup>Irradiance measured using a Molecron PR500 pyroelectric radiometer was approximately  $73 \text{ W m}^{-2}$ .

<sup>3</sup>Undertaken at East Kilbride on 7th March 2005 at midday GMT; measured energy flux was approximately  $1 \text{ kW m}^{-2}$ .