

Geophysical Research Abstracts  
Vol. 18, EGU2016-661, 2016  
EGU General Assembly 2016  
© Author(s) 2015. CC Attribution 3.0 License.



## Reconstructing temperatures in the Maritime Alps, Italy, since the Last Glacial Maximum using cosmogenic noble gas paleothermometry

Marissa Tremblay (1,2), Matteo Spagnolo (3), Adriano Ribolini (4), David Shuster (1,2)

(1) Department of Earth and Planetary Science, University of California, Berkeley, Berkeley, CA, United States (mtremblay@berkeley.edu), (2) Berkeley Geochronology Center, Berkeley, CA, United States, (3) School of Geosciences, University of Aberdeen, Aberdeen, United Kingdom, (4) Dipartimento di Scienze della Terra, University of Pisa, Pisa, Italy

The Gesso Valley, located in the southwestern-most, Maritime portion of the European Alps, contains an exceptionally well-preserved record of glacial advances during the late Pleistocene and Holocene. Detailed geomorphic mapping, geochronology of glacial deposits, and glacier reconstructions indicate that glaciers in this Mediterranean region responded to millennial scale climate variability differently than glaciers in the interior of the European Alps. This suggests that the Mediterranean Sea somehow modulated the climate of this region. However, since glaciers respond to changes in temperature and precipitation, both variables were potentially influenced by proximity to the Sea. To disentangle the competing effects of temperature and precipitation changes on glacier size, we are constraining past temperature variations in the Gesso Valley since the Last Glacial Maximum (LGM) using cosmogenic noble gas paleothermometry. The cosmogenic noble gases  $^3\text{He}$  and  $^{21}\text{Ne}$  experience diffusive loss from common minerals like quartz and feldspars at Earth surface temperatures. Cosmogenic noble gas paleothermometry utilizes this open-system behavior to quantitatively constrain thermal histories of rocks during exposure to cosmic ray particles at the Earth's surface. We will present measurements of cosmogenic  $^3\text{He}$  in quartz sampled from moraines in the Gesso Valley with LGM, Bhl stadial, and Younger Dryas ages. With these  $^3\text{He}$  measurements and experimental data quantifying the diffusion kinetics of  $^3\text{He}$  in quartz, we will provide a preliminary temperature reconstruction for the Gesso Valley since the LGM. Future work on samples from younger moraines in the valley system will be used to fill in details of the more recent temperature history.