



Expedient synthesis of 1,6-anhydro- α -D-galactofuranose, a useful intermediate for glycobiological tools

Luciana Baldoni and Carla Marino*

Full Research Paper

Open Access

Address:

CIHIDECAR-CONICET-UBA, Departamento de Química Orgánica,
Facultad de Ciencias Exactas y Naturales, Universidad de Buenos
Aires, Pabellón II, Ciudad Universitaria, 1428 Buenos Aires,
Argentina, Tel/Fax: +54-11-45763352

Email:

Carla Marino* - cmarino@qo.fcen.uba.ar

* Corresponding author

Keywords:

1,6-anhydro- α -D-Galf; galactofuranosyl iodide; galactofuranosyl
precursor; per-*tert*-butyldimethylsilyl- β -D-galactofuranose

Beilstein J. Org. Chem. **2014**, *10*, 1651–1656.

doi:10.3762/bjoc.10.172

Received: 24 February 2014

Accepted: 26 June 2014

Published: 21 July 2014

Associate Editor: S. Flitsch

© 2014 Baldoni and Marino; licensee Beilstein-Institut.

License and terms: see end of document.

Abstract

A new and efficient three-step procedure for the synthesis of 1,6-anhydro- α -D-galactofuranose is described. The key step involves the formation of the galactofuranosyl iodide by treatment of per-*O*-TBS-D-Galf with TMSI, the selective 6-*O*-desilylation by an excess of TMSI, and the simultaneous nucleophilic attack of the 6-hydroxy group on the anomeric carbon, with the iodide as a good leaving group. This compound is a good precursor for building blocks for the construction of 1 \rightarrow 6 linkages.

Introduction

Anhydro sugars are formed by the intramolecular elimination of a water molecule, with the simultaneous formation of a new heterocyclic ring of different size. They are valuable intermediates not only in carbohydrate synthesis, but also as starting materials for other natural and non-natural complex products and bioactive compounds. Among the glycosans, the anhydro sugars involving the anomeric center in the ring formation, the 1,6-anhydro sugars are the most common and useful building blocks [1,2]. They can play a role in synthetic methodologies aiming at the obtainment of regioselectively functionalized sugars in a few steps, which could give easy access to convenient glycosyl donors and acceptors [3].

Some sugars, for example galactose, can afford not only the pyranosic derivative **1** but also the furanose 1,6-anhydro derivative **2**, both of which may be equipped with [3.2.1] bicyclic skeletons (Figure 1) [1].

A variety of chemical approaches for the synthesis of 1,6-anhydro sugars have been developed [3-10]. Two classes of methods for the synthesis of **2** can be discriminated, the first of which starts from free galactose (D-Gal) and afford mixtures of **1** and **2** and the second starts from a galactofuranose (D-Galf) template conveniently derivatized. Pioneer procedures for the synthesis of **2** involved the pyrolysis of D-Gal under reduced




Figure 1: Possible 1,6-anhydro derivatives for D-galactose.


pressure [11,12] and the acid treatment under heating [13], with the subsequent tedious separation from several byproducts, including the pyranosic analogue **1**. Compound **2** was thus obtained in very low yield. More recently, **2** was obtained in 32% yield by heating with a resin as an acid catalyst. Despite the greater smoothness of the method, byproducts were also formed, rendering the purification difficult [14].

Compound **2** obtained by these procedures was used to afford polymers to explore their possible applications in the field of biochemistry and pharmacology, as their properties differ from those of the corresponding monosaccharides, and they have a high density of functional groups that can be modified to obtain novel materials [14,15]. Benzylated **2** was polymerized under cationic conditions, which afforded a material not completely characterized, presumably formed by β-D-Galf units [15]. Free compound **2**, as well as the D-Manf and D-Glcf analogues, was also polymerized under cationic conditions to yield a hyper-branched polysaccharide with α- and β-linked pyranosidic and furanosidic units [14].

On the other hand, when compound **2** was envisioned as a D-Galf template, the synthesis was devised starting from convenient derivatives of D-Galf in order to avoid the presence of

1. For example, compound **2** was synthesized in the past in six steps from galactose (Scheme 1) [16]. The 1,6-ring closure was produced by the *O*-debenzylation of the 6-hydroxy group of **4** and the nucleophilic attack of this hydroxy group to C-1, promoted by SnCl₄. An optimized synthesis of **2** following this strategy has recently been described with an overall yield of 48% comprising several column chromatography purification steps [17].

The essential role of galactofuranose in the antigenic response of various pathogenic microorganisms [18–20] has triggered the interest for the development of synthetic methods for D-Galf precursors and efficient galactofuranosylation methods [21–25]. D-Galf units have been shown to be *O*-glycosidically linked to other D-Galf units by 1→6 linkages in many natural structures, e.g., in pathogenic *Mycobacteria* and *Aspergillus* spp and others [25–28]. Benzoylated compound **5** is a good precursor of D-Galf derivatives with differentially protected hydroxy groups at position 1 and 6, for example the diacetyl derivative **6** obtained by the acetolysis of **5** (Scheme 1) [16]. In this way, compound **5** would give access to donors in which the 6-position could subsequently be manipulated for the construction of a 1→6 linkage. Based on this strategy, Ning and co-workers synthesized the β-(1→6)-linked hexasaccharide **7** [29], and



Scheme 1: Reported synthesis of 1,6-anhydro-α-D-galactofuranose [16,17].

Kiessling's group developed the synthesis of compounds **8** used for the characterization of GlfT2, one of the two galactofuranosyl transferases involved in the biosynthesis of D-Galf-containing molecules (Figure 2) [30,31].

Our laboratory has long been involved in the development of new galactofuranosyl derivatives and galactofuranosylation methodologies [32]. In this context, we herein report on an efficient three-step synthesis of 1,6-anhydro- α -D-galactofuranose (**2**) from per-*O*-TBS- β -D-galactofuranose (**9**) as a more efficient alternative to existing methods.

Results and Discussion

In the framework of our project for the development of galactofuranosyl derivatives and glycosylation methods, we have reported the synthesis of per-*O*-TBS- β -D-galactofuranose (**9**), a convenient precursor of D-Galf units, and its glycosylation via the in situ generation of galactofuranosyl iodide **10** (Scheme 2) [32–35]. Galactofuranosyl iodides were not previously described, and **10** proved to be useful for the synthesis of several D-Galf-containing molecules (Scheme 2) [32].

The reported procedure consisted in the treatment of compound **9** with 1.2 equiv of TMSI in anhydrous CH_2Cl_2 at 0 °C until the total conversion of **7** into two lower moving products was observed by TLC: the 1-iodo intermediate **10** ($R_f = 0.70$, 10:1 hexane/EtOAc) and 2,3,5,6-tetra-*O*-TBS- α,β -D-galactofuranose ($R_f = 0.54$) formed as a result of the hydrolysis of **10** on the silica gel plate. The addition of simple alcohols or partially protected sugars as acceptors and $\text{EtN}(\text{iPr})_2$ as an acid scavenger led to the complete consumption of both compounds and afforded the corresponding glycosides (Scheme 2) [32]. With an excess of TMSI, a third product ($R_f = 0.62$) was formed, which was not consumed during the reaction and was still present in the product mixture after the work-up. The ^1H NMR spectrum of this product showed a doublet at δ 5.06 with a relatively large $J_{1,2}$ value (4.5 Hz). This signal correlated with a signal at 98.4 ppm in the ^{13}C NMR spectrum, both indicative of the α -configuration. Signals corresponding to C-5 and C-6 showed similar chemical shifts. The signal corresponding to C-6 (δ 65.9) was shifted slightly downfield compared to the same signal in compound **9** (64.7 ppm), while the signal corresponding to C-5 (64.2 ppm) was significantly deshielded






Figure 2: Examples of glycosidic tools synthesized from compound **5** [29–31].



Scheme 2: D-Galactofuranosylation by the glycosyl iodide method [32–35].

(10 ppm) with respect to C-5 of compound **9**. No aglycone signals were observed. In order to elucidate the structure of this compound we *O*-desilylated it by treatment with *n*-Bu₄NF (TBAF) in THF [36]. The product obtained (96%) was faster moving than galactose on TLC (*R_f* = 0.60, 7:1:2 *n*-PrOH/NH₃/H₂O) and showed ¹H and ¹³C NMR spectra coincident with the data reported for 1,6-anhydro- α -D-galactofuranose (**2**) [16,37]. With the objective of optimizing the conditions for glycosylations via iodide **10**, the formation of **12** was suppressed by strict control of the TMSI amount employed.

Taking into consideration how easily compound **12** was obtained and the versatility of anhydro sugars as intermediates for the preparation of biologically important oligosaccharides [3], we decided to investigate the conditions to obtain it as a main product. We reasoned that during the treatment of **9** with TMSI, in addition to the formation of the anomeric iodide, the 6-hydroxy group could be desilylated by the acid medium developed during the iodide formation (**10** \rightarrow **11**). Then, the free 6-hydroxy group could carry out an intramolecular attack of the anomeric carbon, with iodide as a good leaving group, affording the 1,6-anhydro derivative **12** (Scheme 3).



By treatment of **9** with an excess of TMSI (2.25 equiv) in CH₂Cl₂ at room temperature for 5 h, compound **12** was obtained as a single product in 65% yield. Conducting the reaction at low temperature instead (−20 °C), a lower moving product was detected, presumable **11**, which could not be isolated. The use of molecular sieves, which improve the reaction of **10** with alcohols or complex acceptors [32–34], should be avoided in this case as it slows down the reaction. Based on monitoring

the reaction by ¹H NMR (CDCl₃) spectroscopy we observed that the anomeric signal of **9** (5.15 ppm, *J*_{1,2} 2.6 Hz) was slowly transformed into a broad singlet at 6.53 ppm, corresponding to H-1 of **10** [32], followed by the transformation into the anomeric signal of **12** (5.06 ppm, *J*_{1,2} 4.5). However, the deprotection of the 6-hydroxy group of galactofuranosides affects the pattern of H-6 and H-6' in the ¹H NMR spectrum, effectively equalizing them, as was shown before [38]. During the reaction, the pair of double-doublets of H-6 and H-6' (3.68 and 3.54 ppm) of compound **9** [32] were transformed in a double-doublet (3.72 ppm) and an apparent triplet (3.60 ppm), corresponding to the H-6 and H-6' of **12** [32]. In between these two signals an intense doublet corresponding to equivalent H-6,6' (3.58 ppm) of **11** was observed, in accordance with the behavior of other free HO-galactofuranosides [38], which supports the intermediate formation of **11**.

The treatment of **9** with SnCl₄ afforded compound **12**, but in a lower yield due to the *O*-desilylation of another hydroxy group. The addition of BF₃·OEt₂ to recently formed **10** resulted in the formation of compound **12**.

Several factors favor the formation of the bicyclic system of compound **12**, such as the galactose structure itself, the presence of a good leaving group at C-1, and the electron-donating nature of the TBS groups. Thus, while compounds **13** [38] and **14** [39] were prepared by treatment with Lewis acids of fully protected precursors and proved to be stable and therefore useful as synthetic intermediates, attempts to prepare compound **15** by treatment with BF₃·OEt₂ of the persilylated precursor failed and inevitably led to the anhydro derivative **12**. Moreover, whereas treatment with TFA/THF/H₂O 90:5:2.5 of 4-nitrophenyl per-*O*-TBS- α -D-Araf afforded **16**, 4-nitrophenyl per-*O*-TBS- β -D-galactofuranoside did not lead to compound **15** under the same conditions and compound **12** was obtained instead (Figure 3).

The *O*-desilylation of **12** was performed by treatment with *n*-Bu₄F as previously optimized [32–34], affording compound **2** in almost quantitative yield (Scheme 3).

Conclusion

In conclusion, we have described a new and concise procedure for the synthesis of the 1,6-anhydro derivatives **2** and **12**, the key step of which proceeds by a cascade set of three consecutive reactions. The method compares well to existing methods and by avoiding cumbersome steps, such as a benzylation and several column chromatography purifications, is an effective approach. Compounds **2** and **12** represent profitable intermediates to easily access donors and acceptors for the synthesis of Galf-containing molecules as biochemical tools.




Figure 3: Furanosic derivatives with free primary hydroxy group.

Experimental

General methods

Analytical thin-layer chromatography (TLC) were performed on Silica Gel 60 F₂₅₄ (Merck) aluminum supported plates (layer thickness 0.2 mm) with solvent systems given in the text. Visualization of the spots was effected by exposure to UV light and charring with a solution of 10% (v/v) sulfuric acid in EtOH containing 0.5% *p*-anisaldehyde. Column chromatography was carried out with Silica Gel 60 (230–400 mesh, Merck). Optical rotations were measured with a Perkin-Elmer 343 digital polarimeter. Nuclear magnetic resonance (NMR) spectra were recorded with a Bruker AMX 500 spectrometer. Assignments of ¹H and ¹³C were assisted by 2D ¹H COSY and HSQC experiments. High resolution mass spectra (HRMS–ESI⁺) were recorded in a Bruker micrOTOF-Q II spectrometer.

2,3,5-Tri-*O*-tert-butyldimethylsilyl-1,6-anhydro- α -D-galactofuranose (12). A solution of **9** [32] (0.90 g, 1.20 mmol) in anhydrous CH₂Cl₂ (15 mL) was cooled to 0 °C and stirred for 10 min under Ar. Then, iodotrimethylsilane (2.25 equiv, 0.38 mL, 2.70 mmol) was slowly added by using a syringe (10 min) while stirring was continued at 0 °C. The reaction was allowed to reach room temperature (18–25 °C) and stirred until TLC monitoring showed the complete transformation of **9**, first in two products with *R*_f = 0.70 and 0.54 (10:1 hexane–EtOAc), then the transformation of both products in one with *R*_f = 0.62 (5 h). The solution was diluted with CH₂Cl₂ (250 mL), washed with NaHCO₃ (ss) and water, dried (Na₂SO₄), and concentrated. The residue was purified by column chromatography (98.6:1.4 → 2:1, hexane–EtOAc) affording compound **12** as an amorphous solid (0.272 g, 65%). The analytical data of **12** were identical with those described in ref. [32]: [α]_D +42 (*c* 1, CHCl₃); ¹H NMR (500 MHz, CDCl₃) δ 5.05 (d, *J* = 4.5 Hz, 1H, H-1), 4.20 (d, *J* = 1.8 Hz, 1H, H-3), 4.15 (dd, *J* = 1.8, 4.5 Hz, 1H, H-2), 3.97 (ddd, *J* = 4.3, 6.3, 10.5 Hz, 1H, H-5), 3.91 (broad d, *J* = 4.0 Hz, 1H, H-4), 3.72 (ddd, *J* = 1.5, 6.2, 10.5 Hz, 1H, H-6), 3.60 (apparent t, *J* = 10.7 Hz, 1H, H-6'), 0.94–0.86 (SiC(CH₃)₃), 0.12–0.04 (Si(CH₃)₂); ¹³C NMR (125.8 MHz, CDCl₃) δ 98.4 (C-1), 85.3 (C-4), 83.2 (C-2), 77.6 (C-3), 65.9 (C-6), 64.2 (C-5), 25.84, 25.89, 25.6 (SiC(CH₃)₃), 17.9

(SiC(CH₃)₃), –4.49, –4.57, –4.64, –4.68, –4.93, –5.02 (Si(CH₃)₂); Anal. calcd for C₂₄H₅₂O₅Si₃: C, 57.09; H, 10.38; found: C, 56.90; H, 10.52.

1,6-Anhydro- α -D-galactofuranose (2). To a solution of **12** (0.12 g, 0.22 mmol) in freshly distilled THF (7 mL), cooled at 0 °C, (*n*-Bu)₄NF (12 equiv, 2.32 g, 8.88 mmol) was added [36]. The solution was allowed to reach room temperature and then stirring was continued for 3 h until TLC monitoring showed the complete consumption of the starting material. The solution was diluted with water (50 mL), extracted with CH₂Cl₂ (2 × 30 mL), and the aqueous phase was concentrated under vacuum. Purification of the residue by column chromatography (20:1 AcOEt/hexane) afforded **2** (0.036 g, 95%), [α]_D +54 (*c* 1.0, H₂O), lit. [16] [α]_D +54; ¹H NMR (500 MHz, D₂O) δ 5.31 (d, *J* = 4.6 Hz, 1H, H-1), 4.26–4.23 (m, 2H, H-2, H-3), 4.19 (broad d, *J* = 4.2 Hz, 1H, H-4), 4.08–3.99 (m, 2H, H-5, H-6), 3.55 (apparent t, *J* = 10.4 Hz, 1H, H-6'); ¹³C NMR (125.8 MHz, D₂O) δ 98.6 (C-1), 85.2 (C-4), 80.9 (C-2), 75.4 (C-3), 65.6 (C-6), 62.7 (C-5).

Supporting Information

Supporting Information File 1

¹H and ¹³C NMR spectra of compounds **2** and **12**.

[<http://www.beilstein-journals.org/bjoc/content/supplementary/1860-5397-10-172-S1.pdf>]

Acknowledgements

We are indebted to the University of Buenos Aires and the National Research Council of Argentina (CONICET). C.M. is research members of CONICET. L.B. was supported by a fellowship from CONICET.

References

- Černý, M. *Adv. Carbohydr. Chem. Biochem.* **2003**, *58*, 121–198. doi:10.1016/S0065-2318(03)58004-0
- Bols, M. *Carbohydrate Building Blocks*; Wiley: Chichester, 1996.
- Kulkarni, S. S.; Lee, J.-C.; Hung, S.-C. *Curr. Org. Chem.* **2004**, *8*, 475–509. doi:10.2174/1385272043485800

4. Lafont, D.; Boullanger, P.; Cadas, O.; Descotes, G. *Synthesis* **1989**, 191–194. doi:10.1055/s-1989-27192
5. Boons, G.-J.; Isles, S.; Setälä, P. *Synlett* **1995**, 755–756. doi:10.1055/s-1995-5057
6. Skorupowa, E.; Dmochowska, B.; Madaj, J.; Kasprzykowski, F.; Sokolowski, J.; Wiśniewski, A. *J. Carbohydr. Chem.* **1998**, *17*, 49–59. doi:10.1080/07328309808005768
7. Lauer, G.; Oberdofer, F. *Angew. Chem., Int. Ed. Engl.* **1993**, *32*, 272–273. doi:10.1002/anie.199302721
8. Nicolaou, K. C.; Seitz, S. P.; Papahatjis, D. P. *J. Am. Chem. Soc.* **1983**, *105*, 2430–2434. doi:10.1021/ja00346a053
9. Xue, J.; Guo, Z. *Tetrahedron Lett.* **2001**, *42*, 6487–6489. doi:10.1016/S0040-4039(01)01354-5
And references cited therein.
10. Chun, Y.; Yan, S.; Li, X.; Ding, N.; Zhang, W.; Wang, P.; Li, M.; Li, Y. *Tetrahedron Lett.* **2011**, *52*, 6196–6198. doi:10.1016/j.tetlet.2011.09.055
And references therein.
11. Alexander, B. H.; Dimler, R. J.; Mehlretter, C. L. *J. Am. Chem. Soc.* **1951**, *73*, 4658–4659. doi:10.1021/ja01154a048
12. Hann, R. M.; Hudson, C. S. *J. Am. Chem. Soc.* **1941**, *63*, 2241–2242. doi:10.1021/ja01853a061
13. Richtmyer, B. H. *Methods Carbohydr. Chem.* **1963**, *2*, 390–394.
14. Hoai, N. T.; Sasaki, A.; Sasaki, M.; Kaga, H.; Kakuchi, T.; Satoh, T. *Biomacromolecules* **2011**, *12*, 1891–1899. doi:10.1021/bm2002413
15. Lin, J. W.-P.; Schuerch, C. *Macromolecules* **1972**, *5*, 656–657. doi:10.1021/ma60029a026
16. Sakar, S. K.; Choudhury, A. K.; Mukhopadhyay, B.; Roy, N. *J. Carbohydr. Chem.* **1999**, *18*, 1121–1130. doi:10.1080/07328309908544059
17. Sakar, S. K.; Choudhury, A. K.; Hirsch, J.; Roy, N. Synthesis of 1,6-anhydro-2,3,5-tri-O-benzoyl- α -D-galactofuranose. In *Carbohydrate Chemistry. Proven Synthetic methods*; Kováč, P., Ed.; CRC Press: Boca Raton, USA, 2011; Vol. 1, pp 269–278.
18. de Lederkremer, R. M.; Colli, W. *Glycobiology* **1995**, *5*, 547–552. doi:10.1093/glycob/5.6.547
19. Pedersen, L. L.; Turco, S. J. *Cell. Mol. Life Sci.* **2003**, *60*, 259–266. doi:10.1007/s000180300021
20. Marino, C.; de Lederkremer, R. M. Galactose configurations in Nature with emphasis on the biosynthesis of Galactofuranose in glycans. In *Galactose: Structure and Function in Biology and Medicine*; Pomin, V. H., Ed.; Nova Publisher: New York, USA, 2014; pp 107–134.
21. Peltier, P.; Euzen, R.; Daniellou, R.; Nugier-Chauvin, C.; Ferrières, V. *Carbohydr. Res.* **2008**, *343*, 1897–1923. doi:10.1016/j.carres.2008.02.010
22. Richards, M. R.; Lowary, T. L. *ChemBioChem* **2009**, *10*, 1920–1938. doi:10.1002/cbic.200900208
23. Akihiro, I.; Lowary, T. *Trends Glycosci. Glycotechnol.* **2011**, *23*, 134–152. doi:10.4052/tigg.23.134
24. Marino, C.; Baldoni, L. *ChemBioChem* **2014**, *15*, 188–204. doi:10.1002/cbic.201300638
25. Marino, C.; Gallo-Rodriguez, C.; de Lederkremer, R. M. Galactofuranosyl containing glycans: occurrence, synthesis and biochemistry. In *Glycans: Biochemistry, Characterization and Applications*; Mora-Montes, H. M., Ed.; Nova Science Publisher: Boca Raton, USA, 2012; pp 207–268.
26. Crick, D. C.; Mahaprata, S.; Brennan, P. J. *Glycobiology* **2001**, *11*, 107R–118R. doi:10.1093/glycob/11.9.107R
27. Besra, G. S.; Khoo, K.-H.; McNeil, M. R.; Dell, A.; Morris, H. R.; Brennan, P. J. *Biochemistry* **1995**, *34*, 4257–4266. doi:10.1021/bi00013a015
28. Bhamidi, S.; Scherman, M. S.; Rithner, C. D.; Prenni, J. E.; Chatterjee, D.; Koo, K.-H.; McNeil, M. R. *J. Biol. Chem.* **2008**, *283*, 12992–13000. doi:10.1074/jbc.M800222200
29. Zhang, G.; Fu, M.; Ning, J. *Carbohydr. Res.* **2005**, *340*, 155–159. doi:10.1016/j.carres.2004.11.001
30. Splain, R. A.; Kiessling, L. L. *Bioorg. Med. Chem.* **2010**, *18*, 3753–3759. doi:10.1016/j.bmc.2010.04.068
31. Levengood, M. R.; Splain, R. A.; Kiessling, L. L. *J. Am. Chem. Soc.* **2011**, *133*, 12758–12766. doi:10.1021/ja204448t
32. Baldoni, L.; Marino, C. *J. Org. Chem.* **2009**, *74*, 1994–2003. doi:10.1021/jo8025274
33. Baldoni, L.; Stortz, C. A.; Marino, C. *Carbohydr. Res.* **2011**, *346*, 191–196. doi:10.1016/j.carres.2010.11.013
34. Baldoni, L.; Marino, C. *Carbohydr. Res.* **2012**, *362*, 70–78. doi:10.1016/j.carres.2012.08.021
35. Baldoni, L.; Marino, C. *Carbohydr. Res.* **2013**, *374*, 75–81. doi:10.1016/j.carres.2013.03.032
36. Corey, E. J.; Venkateswarlu, A. *J. Am. Chem. Soc.* **1972**, *94*, 6190–6191. doi:10.1021/ja00772a043
37. Köll, P.; Angyal, S. J. *Carbohydr. Res.* **1988**, *179*, 1–5. doi:10.1016/0008-6215(88)84104-1
38. Mariño, K.; Marino, C.; de Lederkremer, R. M. *Anal. Biochem.* **2002**, *301*, 325–328. doi:10.1006/abio.2001.5508
39. Sauvageau, J.; Foster, A. J.; Khan, A. A.; Chee, S. H.; Sims, I. M.; Timmer, M. S. M.; Stocker, B. L. *ChemBioChem* **2012**, *13*, 2416–2424. doi:10.1002/cbic.201200468

License and Terms

This is an Open Access article under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

The license is subject to the *Beilstein Journal of Organic Chemistry* terms and conditions: (<http://www.beilstein-journals.org/bjoc>)

The definitive version of this article is the electronic one which can be found at: [doi:10.3762/bjoc.10.172](https://doi.org/10.3762/bjoc.10.172)