

This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author(s) and source are credited.



ISSN: 2161-1459

Clinical & Experimental Pharmacology

**The International Open Access
Clinical & Experimental Pharmacology**

Executive Editors

Peter Robert Martin

Psychiatry Vanderbilt Psychiatric Hospital, USA

Y. W. Francis Lam

University of Texas, USA

Q. Ping Dou

Pharmacology and Pathology Wayne State University, USA

Karim A. Alkadhi

University of Houston, USA

Aron D. Mosnaim

University of Medicine and Science North Chicago, USA

Available online at: OMICS Publishing Group (www.omicsonline.org)

This article was originally published in a journal by OMICS Publishing Group, and the attached copy is provided by OMICS Publishing Group for the author's benefit and for the benefit of the author's institution, for commercial/research/educational use including without limitation use in instruction at your institution, sending it to specific colleagues that you know, and providing a copy to your institution's administrator.

All other uses, reproduction and distribution, including without limitation commercial reprints, selling or licensing copies or access, or posting on open internet sites, your personal or institution's website or repository, are requested to cite properly.

Digital Object Identifier: <http://dx.doi.org/10.4172/2161-1459.1000e125>



Maternal Separation on Postnatal Stress in Rats Induces Long-lasting Changes on Neurochemical, Behavioural and Genome Responses

Gabriela Beatriz Acosta*

Institute of Pharmacological Research (ININFA), National Research Council of Argentina (CONICET-UBA), School of Pharmacy and Biochemistry, University of Buenos Aires, Junin 956, 5th Floor, C1113AAD, Buenos Aires, Argentina

Early stress in life results in long-lasting effects in many neurotransmitter and neuroendocrine systems of the brain. Maternal Separation (MS) is used an early life stressor that causes profound neurochemical and behavioural changes in the pups that persist into adulthood. Manipulations such as prolonged MS of pups from the dam have been used as an animal model of early-life trauma.

Animal models are quite useful for elucidating the mechanisms underlying abnormalities toward possible treatment strategies for psychiatric, emotional or neurological diseases.

The Central Nervous System (CNS) maintains a degree of adaptive plasticity, which allows adjusting to certain circumstances and modifying innate patterns from neuronal connections. Alterations during development, such as postnatal stress, provoke changes in the stress response of the pups during their whole lives. It is believed that the effects of these events during early life are due to the great plasticity of the developing in CNS. Along the critical period of certain brain

regions, mainly, those related to adverse situations, for example frontal cortex, hippocampus, and Hypothalamic-Pituitary-Adrenal (HPA) axis can develop abnormalities sometimes irreversible and thus alter the emotional process and the stress response along the animal's life.

Early postnatal period and bond between mother and infant seem particularly important in the development and shaping of normal stress response and emotional behavior. Epigenetic modifications, such as DNA methylation, are critical regulators of persistent gene expression changes and may be related to behavioural disorders. However, the mechanisms underlying the long-term effects of environmental stress early in life are not known, they are likely to involve activation of intracellular signalling pathways, leading to modifications of the genome, resulting in changes in gene expression and neural function. The neural modifies underlying the persistent effects of early life separation and stress in rodents remain to be fully elucidated.

***Corresponding author:** Gabriela Beatriz Acosta, Institute of Pharmacological Research (ININFA), National Research Council of Argentina (CONICET-UBA), School of Pharmacy and Biochemistry, University of Buenos Aires, Junin 956, 5th Floor, C1113AAD, Buenos Aires, Argentina, Tel: 4961-5949; E-mail: gcosta@ffyb.uba.ar

Received November 12, 2013; **Accepted** November 14, 2013; **Published** November 21, 2013

Citation: Acosta GB (2013) Maternal Separation on Postnatal Stress in Rats Induces Long-lasting Changes on Neurochemical, Behavioural and Genome Responses. Clin Exp Pharmacol 3: e125. doi:[10.4172/2161-1459.1000e125](http://dx.doi.org/10.4172/2161-1459.1000e125)

Copyright: © 2013 Acosta GB. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.