

Annals of  
**Nutrition &  
Metabolism**

**Power of Programming - The placenta, maternal diet and  
adipose tissue development in the newborn**

Journal:	<i>Annals of Nutrition and Metabolism</i>
Manuscript ID	Draft
mstype:	Review Article
Date Submitted by the Author:	n/a
Complete List of Authors:	Symonds, Michael; The University of Nottingham, Child Health Bloor, Ian; The University of Nottingham, Child Health Ojha, Shalini; The University of Nottingham, Child Health Budge, Helen
Keyword:	Obesity, Reproduction/Pregnancy/Lactation, Infancy and Childhood

SCHOLARONE™  
Manuscripts

Review

1  
2  
3 The placenta, maternal diet and adipose tissue development in the newborn  
4  
5

6  
7 Michael E Symonds<sup>1</sup>, Ian Bloor, Shalini Ojha and Helen Budge  
8  
9

10  
11 Early Life Research Unit, Academic Division of Child Health, Obstetrics & Gynaecology,  
12 and <sup>1</sup>Nottingham Digestive Disease Centre and Biomedical Research Unit, School of  
13 Medicine, Queen's Medical Centre, The University of Nottingham, Nottingham NG7 2UH,  
14 United Kingdom  
15  
16

17  
18  
19 Corresponding author:

20 Professor Michael E. Symonds

21 Academic Division of Child Health Obstetrics & Gynaecology, School of Medicine, Queen's  
22 Medical Centre, The University of Nottingham, Nottingham NG7 2UH, United Kingdom  
23  
24

25 Phone: +44 1158230625 Fax: +44 1158230626.  
26  
27

28  
29 Email: [michael.symonds@nottingham.ac.uk](mailto:michael.symonds@nottingham.ac.uk)  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Abstract****Background:**

A majority of adipose tissue present in the newborn possess the unique mitochondrial protein, uncoupling protein (UCP)1. It is thus highly metabolically active and capable of producing 300 times more heat per unit mass than any other organ in the body. The extent to which maternal obesity and/or an obesogenic diet impacts on placental function thereby resetting the relative distribution of different types of fat in the fetus is unknown.

**Summary:**

Developmentally the majority (if not all) fat in the fetus can be considered as classical brown fat, in which UCP1 is highly abundant. In contrast beige (or recruitable) fat which possess 90% less UCP1 may only appear after birth, as a majority of fat depots undergo a pronounced transformation that is usually accompanied by the loss of UCP1. The extent to which this process can be modulated in a depot specific manner and/or changes in the maternal metabolic environment remain unknown.

**Key Messages:**

An increased understanding of the mechanism by which offspring born to mothers possess excessive adipose tissue could enable sustainable interventions designed to promote the abundance of UCP1 possessing adipocytes. Ultimately this would increase their energy expenditure and improving glucose homeostasis in these individuals.

1  
2  
3  
4  
5 The current challenge to public health with respect to the continued rise in the incidence of  
6 obesity around the world is widely considered to tackling maternal obesity [1]. This is largely  
7 due to epidemiologically based studies together with reviews indicating the potentially  
8 adverse effects of raised maternal body mass index (BMI) on pregnancy outcomes. The  
9 greater prevalence of obesity in pregnant women has occurred concurrently with an increase  
10 in gestational diabetes mellitus (GDM) [2] which can affect up to 14% of all pregnancies in  
11 the US, and around 2–6% of pregnancies in Europe [3, 4]. At the same time the plethora of  
12 reviews are largely based on gross measurements of outcomes such as birthweight and/or  
13 changes in BMI (body mass index) [5] that give comparatively little insight into body  
14 composition and/or metabolic regulation in those offspring. So although it is clear that  
15 women who are grossly obese are at greater risk of not completing a complication free  
16 pregnancy [6] whether this is the case for women who are overweight or moderately obese is  
17 less well established.  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35

36 It should also be noted that in terms of the classification of being overweight, the BMI  
37 category for this has been lowered with time [7] thereby progressively raising the number of  
38 individuals included. At the same time it has been recognised that the relative risk of  
39 metabolic related disease is not simply related to BMI and can vary greatly depending on  
40 ethnicity, social class, age and gender. The extent to which the same classifications and their  
41 relationship with compromised metabolic health apply to women of reproductive age has yet  
42 to be established. Moreover, it appears that excessive weight gain through pregnancy may be  
43 of more concern than BMI per se [6]. In addition, interventions simply trying to reduce the  
44 incidence of large for gestational age (SGA) infants are likely to be unsuccessful as this is  
45 usually an arbitrary classification of the 10% of largest infants rather than an adverse health  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 outcome. It is likely that a combination of raised fat mass together with enhanced postnatal  
4  
5 and later fat growth will be necessary to result in the distinct phenotype that shows metabolic  
6  
7 dysfunction in later life as well as predisposition to produce a similarly large and  
8  
9 disproportionately sized infant.  
10

11  
12  
13  
14 In terms of interventions aimed at improving fetal outcomes with maternal obesity, given  
15  
16 how unsuccessful these are in adults it is clear that we remain a long way off any clear or  
17  
18 effective intervention. Indeed, for most dietary based interventions the usual success rate in  
19  
20 terms of sustained weight loss over several years can be as low as 15%. The challenge in  
21  
22 pregnant women is further amplified by the profound and quite rapid changes in metabolic  
23  
24 regulation that occur in the mother from the time of conception and then through pregnancy.  
25  
26 This occurs as her metabolism adapts to the dual demands of placenta-fetal growth and the  
27  
28 neuro-endocrine adaptations that accompany pregnancy [8]. In the following brief review we  
29  
30 will focus on the different critical windows of development and the potential interaction  
31  
32 between placental and fetal adipose tissue growth, as high-lighted by others [9].  
33  
34  
35  
36  
37

38  
39 In most mammalian species relatively little adipose tissue is laid down in the fetus which may  
40  
41 reflect the higher energetic costs of lipid accretion compared with carbohydrate and protein  
42  
43 together the limited transfer of lipid across the placenta [10, 11]. The notable exception to  
44  
45 this are humans in which significant quantities of lipid can cross the placenta and the  
46  
47 substantial amounts of fat present in term newborns, due in part to large amounts of  
48  
49 subcutaneous fat [12]. This means that there is no obvious animal model for investigating  
50  
51 their interaction and the extent to which modulating maternal diet at different stages of  
52  
53 gestation impacts on both placental function and fetal adiposity [9]. It must also be noted that  
54  
55 there are now three distinct types of fat i.e. brown, white and beige adipose tissue [10]. These  
56  
57  
58  
59  
60

1  
2  
3 may have different embryonic origins although this has yet to be confirmed in any species  
4  
5 other than mice [13]. Furthermore the fetus is maintained within an hypoxic and thermally  
6  
7 clamped environment which will further constrains adipose tissue growth [14]. It is possible  
8  
9 that all fat laid down in the fetus for a majority of large mammalian species is primarily  
10  
11 brown adipose tissue that is characterised as possessing the unique mitochondrial uncoupling  
12  
13 protein (UCP1) [10]. The one exception being the pig which is litter bearing although brown  
14  
15 fat has been found within an isolated depot [15]. When activated through the unmasking of  
16  
17 GDP binding sites the free flow of protons across the mitochondria enables the rapid  
18  
19 production of heat with the need for the production of ATP as required in mitochondria of all  
20  
21 other tissues or organs [16]. This means that once maximally stimulated brown fat can  
22  
23 produce up to 300 times more heat per unit mass than any other tissue in the body. The  
24  
25 contribution of beige fat remains to be fully quantified as it contains around 10% of the  
26  
27 amount of UCP1 as seen in classical brown fat. However, before it acquires this characteristic  
28  
29 the adipose tissue precursors undergo a defined period of proliferation, and as such is located  
30  
31 within a range of diverse depots in the fetus as summarised in Table 1.  
32  
33  
34  
35  
36  
37

### 38 **Adipose tissue development and the modulation effect of maternal diet**

39  
40 The most abundant brown fat depot in humans is located within the supraclavicular (or neck)  
41  
42 region and in contrast to most other depots is retained throughout the life cycle [17, 18].  
43  
44 Despite being first described fifty years ago [19] it is only very recently that a comparable  
45  
46 depot has been found in any other species i.e. sheep [20]. The anatomical location is likely to  
47  
48 be highly significant with respect to its functional role, given that brown fat has the capacity  
49  
50 to ensure that the temperature of blood supplying the brain is maintained. This role will  
51  
52 therefore not only be critical at birth but throughout the life cycle and could explain why  
53  
54 acute stress, and the accompanying rise in cortisol stimulates heat production in  
55  
56  
57  
58  
59  
60

1  
2  
3 supraclavicular brown fat [21]. In contrast, in rodents, at least, glucocorticoids inhibit brown  
4  
5 fat function within the interscapular depot [22].  
6  
7

8  
9  
10 Given the recent discovery of significant amounts of brown fat in the neck region, currently  
11  
12 there are no publications relating to the impact of changes in maternal diet on its growth and  
13  
14 development pre or post-natally. In contrast, the peri-renal depot has been widely investigated  
15  
16 and under conditions in which placental growth is modulated it shows a parallel response,  
17  
18 possibly reflecting the changes in nutrient partitioning to the fetus [20]. Of particular interest  
19  
20 is the finding that under conditions of increased food intake, although fetal growth is  
21  
22 enhanced, relative fetal peri-renal fat mass are actually reduced, but the abundance of UCP1  
23  
24 is raised [23]. Surprisingly this type of model has not been further investigated as it would be  
25  
26 expected to provide further insights into the impact of excess nutrient supply to the fetus and  
27  
28 its impact of both pre and post-natal adiposity.  
29  
30  
31

### 32 33 34 **Future perspectives**

35  
36 Given the rediscovery of brown fat in humans and the acknowledgment that it is a vital organ  
37  
38 implicated in metabolic regulation throughout the life cycle there is an urgent need for more  
39  
40 studies on understanding its early development, especially in large mammals [24]. One  
41  
42 approach currently being adopted is the use of large scale bioinformatics to describe and  
43  
44 identify novel pathways. To date, this has only been undertaken on epicardial adipose tissue  
45  
46 taken from neonates and infants that have undergone heart surgery [25]. These analyses have  
47  
48 further revealed that the transition to infancy is a critical stage for changes in adipose tissue  
49  
50 morphology, that is reflected by a unique pattern of gene expression that included a  
51  
52 significant proportion of thermogenic gene transcripts (~10%). The patterns identified were  
53  
54 specific for each developmental stage, and persisted even after the rebound in abundance of  
55  
56  
57  
58  
59  
60

1  
2  
3 thermogenic genes in later childhood. Using weighted gene co-expression network analysis,  
4  
5 precise anthropometric specific correlations with (corrected) postnatal growth were  
6  
7 identified. These changes in gene expression pathways followed the decline of thermogenic  
8  
9 capacity within the fat depot. Our results also indicated a sequential order of transcriptional  
10  
11 events affecting cellular pathways that could potentially explain the variation in the amount  
12  
13 or activity of BAT in adulthood. In summary, this type of experiment approach could provide  
14  
15 a novel resource to elucidate gene regulatory mechanisms underlying the progressive  
16  
17 development of adipose tissue through the life cycle.  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



## References

- 1 Fitzsimons KJ, Modder J: Setting maternity care standards for women with obesity in pregnancy. *Seminars in Fetal & Neonatal Medicine* 2010;15:100-107.
- 2 Schafer-Graf U: The HAPO Study and its Consequences for the Diagnostic of Gestational diabetes. *Geburtshilfe Und Frauenheilkunde* 2009;69:259-261.
- 3 Dominguez LJ, Martinez-Gonzalez MA, Basterra-Gortari FJ, Gea A, Barbagallo M, Bes-Rastrollo M: Fast Food Consumption and Gestational Diabetes Incidence in the SUN Project. *PloS one* 2014;9:e106627.
- 4 Buckley BS, Harreiter J, Damm P, Corcoy R, Chico A, Simmons D, Vellinga A, Dunne F, Group DCI: Gestational diabetes mellitus in Europe: prevalence, current screening practice and barriers to screening. A review. *Diabetic medicine : a journal of the British Diabetic Association* 2012;29:844-854.
- 5 Godfrey KM, Reynolds RM, Prescott SL, Nyirenda M, Jaddoe VW, Eriksson JG, Broekman BF: Influence of maternal obesity on the long-term health of offspring. *Lancet Diabetes Endocrinol* 2016
- 6 Yu Z, Han S, Zhu J, Sun X, Ji C, Guo X: Pre-pregnancy body mass index in relation to infant birth weight and offspring overweight/obesity: a systematic review and meta-analysis. *PLoS One* 2013;8:e61627.
- 7 Kuczmarski RJ, Flegal KM: Criteria for definition of overweight in transition: background and recommendations for the United States. *Am J Clin Nutr* 2000;72:1074-1081.
- 8 Ojha S, Fainberg HP, Sebert S, Budge H, Symonds ME: Maternal health and eating habits: metabolic consequences and impact on child health. *Trends in molecular medicine* 2015;21:126-133.
- 9 Wankhade UD, Thakali KM, Shankar K: Persistent influence of maternal obesity on offspring health: Mechanisms from animal models and clinical studies. *Mol Cell Endocrinol* 2016;435:7-19.
- 10 Symonds ME, Pope M, Budge H: The ontogeny of brown adipose tissue. *Ann Rev Nutr* 2015;35:295-320.
- 11 Symonds ME: Brown adipose tissue growth and development. *Scientifica* 2013;2013:14. <http://dx.doi.org/10.1155/2013/305763>.
- 12 Widdowson EM: Chemical composition of newly born animals. *Nature* 1950;116:626-628.
- 13 Cypess AM, Haft CR, Laughlin MR, Hu HH: Brown fat in humans: Consensus points and experimental guidelines. *Cell Metab* 2014;20:408-415.
- 14 Symonds ME, Bird JA, Clarke L, Gate JJ, Lomax MA: Nutrition, temperature and homeostasis during perinatal development. *Exp Phys* 1995;80:907-940.
- 15 Mostyn A, Attig L, Larcher T, Dou S, Chavatte-Palmer P, Boukthir M, Gertler A, Djiane J, Symonds ME, Abdennebi-Najar L: UCP1 is present in porcine adipose tissue and is responsive to postnatal leptin. *J Endocrinol* 2014;223:M31-38.
- 16 Cannon B, Nedergaard J: Brown adipose tissue: function and physiological significance. *Phys Rev* 2004;84:277-359.
- 17 Cypess AM, White AP, Vernochet C, Schulz TJ, Xue R, Sass CA, Huang TL, Roberts-Toler C, Weiner LS, Sze C, Chacko AT, Deschamps LN, Herder LM, Truchan N, Glasgow AL, Holman AR, Gavrilu A, Hasselgren PO, Mori MA, Molla M, Tseng YH: Anatomical localization, gene expression profiling and functional characterization of adult human neck brown fat. *Nat Med* 2013;19:635-639.

- 1  
2  
3 18 Symonds ME, Henderson K, Elvidge L, Bosman C, Sharkey D, Perkins AC, Budge  
4 H: Thermal imaging to assess age-related changes of skin temperature within the  
5 supraclavicular region co-locating with brown adipose tissue in healthy children. *J Pediatr*  
6 2012;161:892-898.
- 7 19 Aherne W, Hull D: Brown adipose tissue and heat production in the newborn infant. *J*  
8 *Pathol Bacteriol* 1966;91:223-234.
- 9 20 Symonds ME, Pope M, Sharkey D, Budge H: Adipose tissue and fetal programming.  
10 *Diabetologia* 2012;55:1597-1606.
- 11 21 Robinson LJ, Law JM, Symonds ME, Budge H: Brown adipose tissue activation as  
12 measured by infrared thermography by mild anticipatory psychological stress in lean healthy  
13 females. *Exp Physiol* 2016;101:549-557.
- 14 22 Ramage LE, Akyol M, Fletcher AM, Forsythe J, Nixon M, Carter RN, van Beek EJ,  
15 Morton NM, Walker BR, Stimson RH: Glucocorticoids Acutely Increase Brown Adipose  
16 Tissue Activity in Humans, Revealing Species-Specific Differences in UCP-1 Regulation.  
17 *Cell Metab* 2016;24:130-141.
- 18 23 Budge H, Bispham J, Dandrea J, Evans E, Heasman L, Ingleton PM, Sullivan C,  
19 Wilson V, Stephenson T, Symonds ME: Effect of maternal nutrition on brown adipose tissue  
20 and its prolactin receptor status in the fetal lamb. *Pediatr Res* 2000;47:781-786.
- 21 24 Aldiss P, Davies G, Woods R, Budge H, Sacks HS, Symonds ME: 'Browning' the  
22 cardiac and peri-vascular adipose tissues to modulate cardiovascular risk. *Int J Cardiol*  
23 2016;228:265-274.
- 24 25 Ojha S, Fainberg HP, Wilson V, Pelella G, Castellanos M, May ST, Lotto AA, Sacks  
25 H, Symonds ME, Budge H: Gene pathway development in human epicardial adipose tissue  
26 during early life. *JCI Insight* 2016;1:e87460.
- 27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4  
5 **Figure Title**  
6

7 Figure 1. Summary of the developmental trajectories of the placenta and adipose tissue in  
8  
9 large mammals.  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

For Peer Review

Table 1. Summary of the main fat depots in the fetus and newborn that contain the brown adipose tissue specific uncoupling protein (UCP1).

Anatomical location	Description	Potential function	Nutritionally responsive in utero
Supraclvicular	Located in the neck region surrounding blood vessels	Maintaining the temperature of blood supplying the brain	Not known
Pericardial	Surrounds the heart	Maintaining heart temperature and preventing lipotoxicity	Responds to prevailing nutrient supply in late gestation
Epicardial	Along myocardium, beneath visceral pericardium and pericardial sac. Surrounds coronary arteries	Thermogenesis and related processes	Not known
Perirenal	Surrounds the kidney and adrenal glands	Thermogenesis	Reduced mass with suboptimal nutrition. Increased nutrition promotes UCP1 but not total fat mass

Ann Nutr Metab

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25

Attachment/  
implantation

Rapid placental  
growth

Placental  
adaptation

Proliferation

Expansion and  
Preparation  
for birth

Rapid growth  
and expansion

Embryo

Early gestation

Late gestation

Birth

Post natal


Early organ growth

Organ Maturation

Lactation



Placental  
development



Adipose tissue  
development