

1           **Biomechanical locomotion adaptations on uneven surfaces can be**  
2                   **simulated with a randomly deforming shoe midsole**

3  
4   **Authors and affiliations:**

5   **\*Charlotte Apps**

6   Nottingham Trent University and Li Ning Sports Science Research Center

7   School of Science and Technology, Clifton Campus, Nottingham, United Kingdom,

8   NG11 8NS

9   Phone: +44 (0) 7889622840

10   Email: [charlotte.apps@ntu.ac.uk](mailto:charlotte.apps@ntu.ac.uk)

11  
12   **Thorsten Sterzing**

13   Li Ning Sports Science Research Center

14   Li Ning (China) Sports Goods Co., Ltd, No. 8 Xing Guang 5<sup>th</sup> Street, Beijing, China, 101111

15   Phone: +86 186 11975065

16   E-Mail: [thorsten.sterzing@web.de](mailto:thorsten.sterzing@web.de)

17  
18   **Thomas O'Brien**

19   Liverpool John Moores University

20   School of Sport and Exercise Sciences, Tom Reilly Building, Byrom Street Campus,

21   John Moores University, Liverpool, United Kingdom, L3 3AF

22   Phone: +44 (0)151 231 3179

23   Email: [T.D.O'Brien@ljmu.ac.uk](mailto:T.D.O'Brien@ljmu.ac.uk)

24  
25   **Rui Ding**

26   Li Ning Sports Science Research Center

27   Li Ning (China) Sports Goods Co., Ltd, No. 8 Xing Guang 5<sup>th</sup> Street, Beijing, China, 101111

28   E-Mail: [dingrui@li-ning.com.cn](mailto:dingrui@li-ning.com.cn)

29  
30   **Mark Lake**

31   Liverpool John Moores University

32   School of Sport and Exercise Sciences, Tom Reilly Building, Byrom Street Campus,

33   John Moores University, Liverpool, United Kingdom, L3 3AF

34 Phone: +44 (0)151 904 6260

35 Email: M.J.Lake@ljmu.ac.uk

36

37 \* Corresponding author

38 E-mail: charlotte.apps@ntu.ac.uk (CA)

39

#### 40 **Acknowledgments**

41 The authors would like to thank Ruiya Ma for her help with data collection and participant  
42 recruitment, and all participants for taking part in this research.

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58

59

60

61

62

63 **Biomechanical locomotion adaptations on uneven surfaces can be**  
64 **simulated with a randomly deforming shoe midsole**

65  
66 **ABSTRACT**

67 **Background:** A shoe with unsystematic perturbations, similar to natural uneven terrain, may  
68 offer an enhanced training stimulus over current unstable footwear technologies. This study  
69 compared the instability of a shoe with unpredictably random midsole deformations, an  
70 irregular surface and a control shoe-surface whilst treadmill walking and running.

71 **Methods:** Three-dimensional kinematics and electromyography were recorded of the lower  
72 limb in 18 active males. Gait cycle characteristics, joint angles at initial ground contact and  
73 maximum values during stance, and muscle activations prior to initial contact and during  
74 loading were analysed. Perceived stability, injury-risk and energy consumption were  
75 evaluated. Instability was assessed by movement variability, muscular activations and  
76 subjective ratings.

77 **Results:** Posture alterations at initial contact revealed active adaptations in the irregular  
78 midsole and irregular surface to maintain stability whilst walking and running. Variability of  
79 the gait cycle and lower limb kinematics increased on the irregular surface compared to the  
80 control across locomotion types. Similarly increased variability (coefficient of variation)  
81 were found in the irregular midsole compared to the control for frontal ankle motion (walk:  
82 31.1 and 14.9, run: 28.1 and 11.6), maximum sagittal knee angle (walk: 7.6 and 4.8, run: 2.8  
83 and 2.4), and global gait characteristics during walking only ( $2.1 \pm 0.5$  and  $1.6 \pm 0.3$ ). Tibialis  
84 anterior pre-activation reduced and gastrocnemius activation increased in the irregular  
85 midsole compared to the control across locomotion types. During running, peroneus longus  
86 activation increased in the irregular midsole and irregular surface.

87 **Conclusions:** Results indicate random shoe midsole deformations enhanced instability  
88 relative to the control and simulated certain locomotion adaptations of the irregular surface,  
89 although less pronounced. Thus, a shoe with unpredictable instability revealed potential as a  
90 novel instability-training device.

91  
92 **Keywords:** footwear; instability; kinematics; electromyography; lower-limb  
93

## 94 **1. Introduction**

95 A relatively new concept of training footwear, termed unstable shoes are designed to create  
96 instability with the aim of providing functional benefits, such as increasing muscle  
97 activations and improving balance. Innovative shoe technologies developed for this purpose  
98 include rocker soles, balance pods and midsoles of multiple densities (Price, Smith, Graham-  
99 Smith, & Jones, 2013). The concept behind unstable footwear is similar to traditional  
100 instability training devices, such as Swiss balls, BOSU balls and wobble boards. Such  
101 equipment reduces the base of support causing instability, which can be observed through an  
102 increase of movement variability (Cimadoro, Paizis, Alberti, & Babault, 2013). The  
103 neuromuscular system has to make alterations to maintain stability and regular use is  
104 proposed to enhance balance and train the lower-limb muscles. A limitation of instability  
105 training devices is they are only utilised during restricted, isolated exercises and not during  
106 functional movements. Unstable shoes in contrast, may allow habitual training during  
107 walking, running or aerobic exercises.

108 Increased muscle activation is one of the acute responses of wearing unstable footwear. For  
109 the most frequently tested unstable shoe, Masai Barefoot Technology (MBT), increased  
110 tibialis anterior and peroneus longus activations have been commonly reported whilst  
111 standing (Buchecker, Pfusterschmied, Moser, & Müller, 2012; Nigg, Hintzen, & Ferber, 2006;  
112 Landry, Nigg, & Tecante, 2010) and increased gastrocnemius medialis activations whilst  
113 walking (Price et al., 2013; Romkes, Rudmann, & Brunner, 2006). As would be predicted,  
114 long-term wear strengthens and conditions the ankle muscles (Kaelin, Segesser, & Wasser,  
115 2011). However, other studies report no significant increases in muscle activation whilst  
116 walking (Horsak & Baca, 2013; Nigg et al., 2006; Sacco et al., 2012; Stöggl, Haudum,  
117 Birklbauer, Murrer, & Müller, 2010) or running in unstable shoes (Sobhani et al., 2013).

118 Improved balance is another suggested training effect of regularly wearing unstable footwear.  
119 Increased centre of pressure excursion during static two-legged standing reduced over 6-  
120 weeks in healthy adults aged between 40 to 70 years old (Landry et al., 2010). The authors  
121 suggested this demonstrated improved static balance performance, but results from a dynamic  
122 systems perspective suggest this may not always be the case (van Emmerik, & van Wegen,  
123 2000). A better determinant of the postural system's ability could be assessing reactive  
124 balance after an external perturbation. Females older than 50 years old did improve their  
125 reactive balance over 8-weeks, but not significantly compared to a control group (Ramstrand,  
126 Thuesen, Nielsen, & Rusaw, 2010).

127 The inconsistent findings, particularly during dynamic locomotion may be due to the  
128 different number of participants, amount of pre-exposure to the unstable footwear and  
129 evaluation analyses applied. Another potential reason is the majority of previous research  
130 included active participants who were less likely to be affected by the unstable shoe  
131 instability during locomotion. Perhaps a more challenging unstable shoe construction would  
132 have a more pronounced effect. One such shoe design, Reflex Control has a thin sole bar  
133 along the longitudinal foot axis, compared to the Masai Barefoot Technology (MBT) shoe  
134 that has an anteroposterior sole rocker. Compared to barefoot walking, Reflex Control  
135 increased shank muscle activation, but no effect was found in MBT during walking  
136 (Schiemann, Lohrer, & Nauck, 2015). In addition, reactive balance during one-legged  
137 standing improved after a training program in Reflex Control, but not in MBT (Turbanski,  
138 Lohrer, Nauck, & Schmidtbleicher, 2011).

139 Moreover, although movement variability initially increases whilst walking in MBT shoes,  
140 this variability reduces after a 10-week training period (Stöggl et al., 2010). This suggests  
141 that instability becomes predictable, due to the cyclic repetitions during gait with the same  
142 fixed outsole stimulus of the MBT. Furthermore, Blair, Lake and Sterzing (2013) found  
143 initially increased vastus medialis activation whilst walking in an unstable shoe reduced to a  
144 similar level to a stable shoe after one hour, but tibialis anterior activation further increased.  
145 Trunk acceleration in the unstable shoe also tended to reduce after the hour walking. This  
146 suggests neuromuscular adaptations are learnt quickly and benefits of further training reduce  
147 over time.

148 Uneven natural terrain surfaces may provide a superior training modality by creating a  
149 continually changing and unpredictable instability. Increased muscle activations, a cautious  
150 gait pattern and increased movement variability has been found whilst walking (Gates,  
151 Wilken, Scott, Sinitski, & Dingwell, 2012; Marigold, & Patla, 2008; McAndrew, Dingwell,  
152 & Wilken, 2010; Sterzing, Apps, Ding, & Cheung, 2014a; Thies, Richardson, & Ashton-  
153 Miller, 2005; Voloshina, Kuo, Daley, & Ferris, 2013) and running on irregular surfaces  
154 (Sterzing, Apps, Ding, & Cheung, 2014b; Voloshina & Ferris, 2015). However, irregular  
155 surfaces are often not accessible in urban areas for convenient and frequent use. An  
156 alternative and novel solution would be to develop footwear that causes irregular and  
157 unpredictable instability. Kim and Ashton-Miller (2012) constructed experimental sandals  
158 with medial and lateral flaps in the sole, which could be deployed at random times to assess  
159 response to an unpredictable perturbation. Although the sandals were controlled

160 electronically, which made them unsuitable for use by the general public. Consequently, we  
161 developed a training shoe with random irregular midsole deformations. The purpose of this  
162 study was to investigate the locomotion instability induced by this shoe compared to an  
163 irregular surface during walking and running.

164 Based on previous research, it was hypothesised the irregular midsole and an irregular surface  
165 would provide a similar, higher level of instability compared to a regular shoe-surface. This  
166 would be indicated by an increase in movement variability of the global spatial-temporal gait  
167 cycle characteristics and at the joint level, although this does not necessarily represent loss of  
168 stability. Moreover, there will be postural adjustments and increases in muscle activations to  
169 maintain balance. These hypotheses were applicable to both walking and running.

170

## 171 **2. Methods**

### 172 ***2.1. Participants***

173 Eighteen active male sports science students, who were regular runners participated in this  
174 research (22.7 years  $\pm$  1.7, 177.2 cm  $\pm$  3.8, 69.1 kg  $\pm$  5.7). All participants had been injury  
175 free for at least 6 months prior to testing and had Brannock foot size male US 10.0  $\pm$  0.5 (The  
176 Brannock Device Co., Liverpool, NY, USA). Liverpool John Moores University research  
177 ethics committee approved the study protocol and participants gave their written informed  
178 consent prior to testing.

179 As no previous data were available in the irregular midsole condition, a priori power analysis  
180 was performed on results of a previous study that compared the irregular treadmill surface  
181 condition to the regular treadmill surface (Sterzing et al., 2014a; Sterzing et al., 2014b) in  
182 G\*Power software (Faul, Erdfelder, Lang, & Buchner, 2007). Kinematic variability of  
183 maximum sagittal and frontal ankle and sagittal knee angles during stance phase of walking  
184 and running (as used in this study) were tested. Across results a maximum of 13 participants  
185 were required to obtain an effect size of 0.75 (p value = .05,  $\beta$  = .20). Along with previous  
186 unstable footwear studies, this sample size was deemed appropriate for this study.

### 187 ***2.2. Shoe-surface Conditions***

188 Three shoe-surface conditions were tested on a treadmill during walking and running:

- 189 1. A shoe with irregular midsole deformations and a regular treadmill surface (IM)
- 190 2. A regular shoe midsole and an irregular treadmill surface (IS)

191 3. The regular shoe midsole and regular treadmill surface as a control condition (CC)  
192 Both shoe conditions had the same upper (Li Ning Fengchao TD, Li Ning Co, Beijing, size  
193 male US 10.0) while the two different midsole modifications were attached. The irregular  
194 midsole was created using three highly flexible rubber bags (hardness: 28 Asker C, thickness:  
195 1.5 mm) attached to the shoe upper by Velcro at the rearfoot, midfoot and forefoot at 30%, 30%  
196 and 40% shoe length respectively. The segregation of foot regions is based upon previous  
197 biomechanical research (Cavanagh & Ulbrecht, 1994). The heel to toe offset was 10 mm  
198 unweighted, but due to the deformable bag material this reduced when wearing the IM shoe.  
199 In total, 51 ball bearings (12 mm diameter) and 10 cube shapes (height 15 mm, hardness: 85A  
200 Shore, TPU material) were placed inside the rubber bags and moved freely during swing  
201 phase of the gait cycle, creating a different shoe-surface profile at every ground contact and  
202 thus unpredictable perturbations. The ratio of ball bearings was 15:15:21 and cube shapes  
203 were 4:3:3 inside the rearfoot, midfoot and forefoot bags respectively.

204 The regular shoe midsole condition was developed with the midsole of the original shoe and  
205 used in CC and IS trials. The medio-lateral midsole shape was cut to identical dimensions of  
206 the IM shoe. Aluminium weights (5g) were glued evenly to replicate the weight of the IM  
207 bags (Fig 1). The regular shoe midsole weighed 234g and the irregular midsole shoe weighed  
208 233g. Thus, weight and shape midsole differences were minimised. The heel to toe offset of  
209 the regular midsole was 10 mm.

210

211 \*\*\*Figure 1 near here\*\*\*

212

213 All walking and running trials were performed on a treadmill (Pro XL, Woodway Inc., WI,  
214 USA). The treadmill belt slats were covered with Velcro strips (700mm x 58mm), which  
215 served as the regular surface. The irregular treadmill surface (IS) was created by randomly  
216 fixing 4 types of EVA dome shaped inserts ( $\varnothing$ : 140mm) of different height (10 and 15 mm)  
217 and hardness (40 and 70 Asker C) to the treadmill belt by Velcro attachment (Fig 2), as used  
218 in previous research (Sterzing et al., 2014a, 2014b). To eliminate visual targeting of foot  
219 placements, participants were instructed to look straight ahead. This was monitored by  
220 investigators, ensuring participants could not predict what they were to land on.

221

222 \*\*\*Figure 2 near here\*\*\*

223

### 224 **2.3. Protocol**

225 The treadmill speed was set at 5 km/hr for walking trials, as used in previous unstable  
226 footwear research (Nigg et al., 2006; Stöggl et al., 2010), and 8 km/hr for running trials. The  
227 slow run speed was selected to improve the level of comfort, as previously tested on IS  
228 (Sterzing et al., 2014b; Voloshina & Ferris, 2015). The order of shoe-surface conditions was  
229 arranged so CC trials were always first to avoid potential crossover effects from IM and IS,  
230 whose order was alternated between participants. Walking trials preceded running trials in the  
231 same shoe-surface condition. Before data collection participants were briefed about the  
232 testing conditions. After 60 seconds of walking and running in each shoe-surface condition to  
233 allow participants to get into a regular locomotion rhythm, biomechanical data were collected  
234 for 30 seconds. Surface EMG and lower limb kinematics were recorded synchronously from  
235 the subjects' left leg.

### 236 **2.4. Kinematics**

237 Kinematics were captured by a seven-camera motion analysis system at 300 Hz (Vicon Peak,  
238 Oxford, UK). Reflective markers were attached to the following locations to define the left  
239 thigh, shank and foot segments: The greater trochanter, medial and lateral femoral  
240 epicondyles, the lateral and medial malleoli, on the tip of the shoe and dorsal metatarsal  
241 heads 1 and 5. Tracking markers clusters were attached on the lateral side of the thigh (5  
242 markers) and shank (4 markers), and to the shoe at the proximal posterior, distal posterior and  
243 lateral heel counter. Position and orientation of anatomical markers relative to tracking  
244 markers were determined from a static trial in the anatomical position in the regular shoe only,  
245 similar to the CAST technique (Cappozzo, Catani, Della-Croce, & Leardini, 1995). The same  
246 shoe upper was kept on throughout all trials allowing identical marker placement in all  
247 conditions, ensuring kinematic differences observed cannot be attributed to different marker  
248 location. Utilising a global neutral configuration is advantageous because the absolute  
249 angular differences between midsole conditions can be compared, which are not influenced  
250 by changes in the sole configuration.

251 After digitising, raw marker co-ordinate data were filtered using a low pass fourth order zero-  
252 lag Butterworth filter with cut-off frequencies of 10Hz for walking and 20Hz for running,  
253 based on visual inspection of the power spectrum. Stance phase was determined by ground



254 contact algorithms which matched well against pilot data measurements with a foot switch  
255 placed inside the shoe-conditions on a treadmill and verified with a force plate. Vertical  
256 velocity change of the midpoint between the heel and toe markers identified gait events  
257 during walking (O'Connor, Thorpe, O'Malley, & Vaughan, 2007) and the vertical  
258 acceleration of the heel and tip of shoe markers was used during running (Maiwald, Sterzing,  
259 Mayer, & Milani, 2009). Some kinematic data were not collected successfully due to  
260 technical issues and are excluded from subsequent analyses. Kinematic results are based on  
261 16 participants for walking and 17 for running.

262 Characteristics of the gait cycle were derived from ground contact times. Positive sagittal  
263 knee and ankle angles reflect joint flexion, and positive frontal ankle angle represents  
264 eversion. To show preparatory posture adaptations shoe-surface and joint angles were  
265 calculated at initial contact. We expected the unpredictable instability to have a greater effect  
266 during loading occurring in the first half of stance. Therefore, maximum joint angles and  
267 ankle ranges of motion between initial contact and maximum positive angles during stance  
268 were determined. The single largest ankle inversion angle of all steps between initial ground  
269 contact and maximum eversion angle was recorded to indicate any outliers that were  
270 obscured when looking at the variability through the standard deviation.

## 271 ***2.5. Surface Electromyography***

272 Surface electromyography (EMG) was recorded the left gastrocnemius medialis, peroneus  
273 longus, tibialis anterior, bicep femoris, vastus medialis and vastus lateralis muscle activations  
274 using a wireless telemetric system (TeleMyo DTS, Noraxon Inc., Scottsdale, AZ, USA) at 3  
275 kHz. Pre-gelled bi-polar Ag/AgCl circular electrodes (Tian run, Beijing, China) of 10mm  
276 diameter and inter-electrode spacing of 25mm were positioned according to international  
277 recommendations (SENIAM). To reduce impedance, hair was shaved and skin cleaned with  
278 ethanol. The analogue signal was converted to a digital signal by a 16-bit transmitter data  
279 acquisition system. Certain electrode data contained artefacts and were excluded from  
280 subsequent analyses. After exclusion the number of subjects per muscle for walking and  
281 running respectively contained: gastrocnemius medialis (N=14, 15), peroneus longus (N=12,  
282 13), tibialis anterior (N=9, 10), bicep femoris (N=14, 15), vastus medialis (N=13, 15) and  
283 vastus lateralis (N=11, 16).

284 The EMG data were processed in Visual 3D software together with the kinematic data (C-  
285 Motion, Rockville, MD, USA). The raw signal was digitally band-pass filtered using a bi-

286 directional 4<sup>th</sup> order Butterworth filter with cut-off frequencies of 10 and 300Hz, full wave  
287 rectified and smoothed using an 11-point root mean square moving average filter. In  
288 subsequent analysis, EMG values were normalised to the average peak value of each muscle  
289 during the gait cycle of CC trials of the same locomotion type. The normalised mean value  
290 was calculated in a pre-activation phase (150ms before initial contact) and a loading phase  
291 (from initial contact until maximum knee flexion) to supplement kinematic variables.

## 292 ***2.6. Subjective Perception Assessment***

293 Immediately after biomechanical data collection, subjective perception of the level of  
294 stability, injury risk, and energy consumption were collected while participants were still  
295 walking or running on the treadmill. Prior to data collection, variables were defined to  
296 participants, with the instructor explaining their perceived level of magnitude (low, high)  
297 rather than their interpretation (good, bad) was being assessed. Participants assessed all  
298 variables verbally from a large 9-point Likert scale (1-very low, 3-low, 5-moderate, 7-high  
299 and, 9-very high, with other numbers not denominated) mounted in front of the treadmill,  
300 (Fig 2) (adapted from Au & Goonetilleke, 2007; Lam et al., 2013; Sterzing et al., 2014c).  
301 This method is advantageous because participants can think solely about the perception  
302 variable whilst walking and running (Sterzing et al., 2014a; Sterzing et al., 2014b).

## 303 ***Statistics***

304 All steps ( $41.0 \pm 2.6$  for running and  $28.6 \pm 1.5$  for walking) were analysed to compute the  
305 mean of all variables for each participant. Variability of gait cycle and kinematic variables  
306 were calculated with the coefficient of variation (CV). The CV was calculated by dividing the  
307 standard deviation by the mean and multiplying by 100. The CV can be useful for  
308 determining the relative magnitude of variability when there are differences in mean readings,  
309 but is limited if the mean value is close to zero (James, 2004).

310 All statistical processing was performed in SPSS (v22, SPSS Inc, Chicago, IL, USA).  
311 Normality of data were checked using the Shapiro-Wilk test and visually verified for outliers  
312 with boxplots. Most variables followed parametric assumptions and a one-way repeated  
313 measures ANOVA, with Bonferroni adjusted post hoc tests were applied to define differences  
314 between shoe-surface conditions for walking and running ( $p < .05$ ). The non-parametric  
315 Friedman test with Bonferroni adjusted Wilcoxon post hoc tests were applied to the variables  
316 with outliers ( $p < .05$ ). Missing data were deleted listwise, as it were the only option available  
317 in SPSS. This meant always the same number of participant mean variables were compared.

318

### 319 **3. Results**

#### 320 **3.1. Kinematics**

321 Differences to mean kinematic results were generally small between conditions but consistent  
322 across participants whilst walking (Table 1) and running (Table 2). The gait cycle in IM was  
323 characterised by shorter, thus more frequent steps. Variability of gait was significantly  
324 increased in IS compared to CC, with the difference being greater in running (Fig 3). During  
325 walking IM had rather higher variability similar to IS ( $26 \pm 14\% > CC$ ), whereas during  
326 running IM had rather lower level of variability similar to CC ( $3 \pm 2\% > CC$ ).

327

328 \*\*\*Figure 3 near here\*\*\*

329

330 At initial ground contact, knee flexion increased in IM compared to IS and CC whilst walking  
331 and running. Shoe-surface angle was flattest in IM during walking, and flatter in IM and IS  
332 compared to CC during running. Variability of parameters at initial ground contact tended to  
333 be greatest in IS across participants and locomotion (Fig 4). Ankle angle variability could not  
334 be computed due to mean values ranging around zero. Therefore, the standard deviation is  
335 reported separately in Supplementary Table 1 and 2, to give an indication of ankle angle  
336 variability.

337

338 \*\*\*Figure 4 near here\*\*\*

339

340 During stance, maximum ankle eversion reduced in IM whilst walking and running (Fig 5).  
341 Sagittal ankle range of motion reduced whilst walking and frontal ankle range of motion  
342 reduced whilst running in IM compared to CC and IS. The largest ankle inversion angles  
343 recorded were no different between IM and IS during locomotion. During walking, CC had a  
344 significantly reduced maximum inversion angle compared to IM and IS ( $p = .005$ ; IM =  $11.5$   
345  $\pm 6.1^\circ$ , IS =  $10.1 \pm 7.1$ , CC =  $5.9 \pm 3.1$ ) but no different during running ( $p = .008$ ; IM =  $11.1$   
346  $\pm 4.8$ , IS =  $9.3 \pm 4.7$ , CC =  $8.7 \pm 3.4$ ). Variability of parameters during stance were largely  
347 more variable in IS, with IM having similar variability levels of frontal ankle range of motion

348 (walk: 109% > CC, run 143% > CC) and knee flexion (walk: 60% > CC, run: 19% > CC) (Fig  
349 4, Fig 5) across locomotion.

350

351 \*\*\*Figure 5 near here\*\*\*

352

353 \*\*\*Table 1 near here\*\*\*

354

355 \*\*\*Table 2 near here\*\*\*

356

### 357 **3.2. Electromyography**

358 Electromyography results showed differences mostly occurred in the shank muscles for both  
359 walking (Table 3) and running (Table 4). Tibialis anterior activation significantly reduced  
360 during pre-activation and loading in IM whilst walking compared to CC and IS. During pre-  
361 activation whilst running, tibialis anterior activation significantly reduced in IM and IS  
362 compared to CC. Peroneus longus activation significantly increased during loading in IM and  
363 IS compared to CC, and during pre-activation in IS compared to CC whilst running. The  
364 gastrocnemius medialis had significantly greater pre-activation in IM than CC during walking  
365 and running.

366

367 \*\*\*Table 3 near here\*\*\*

368

369 \*\*\*Table 4 near here\*\*\*

370

### 371 **3.3 Perception**

372 Subjective ratings results showed IM was perceived the least stable, with IS less stable than  
373 CC for walking and running. Injury risk level was perceived greatest in IM and greater in IS  
374 than CC for walking and running. Energy requirement was perceived greater for IM and IS  
375 than CC during walking and running (Table 5).

376

377 \*\*\*Table 5 near here\*\*

378

#### 379 **4. Discussion**

380 This study compared the instability caused by both a shoe and surface exhibiting irregular  
381 perturbations during treadmill walking and running. Biomechanical instability were assessed  
382 by changes in movement variability of the spatial-temporal gait cycle and lower limb  
383 kinematics, as well as, muscle activations. Whether participants could also perceive changes  
384 to instability were also assessed. Results confirmed our hypothesis that the irregular midsole  
385 shoe (IM) and irregular surface (IS) increased biomechanical and subjectively perceived  
386 instability compared to a regular shoe-surface (CC). Similarly increased variability of frontal  
387 ankle motion and maximum knee flexion for both walking and running were found between  
388 IM and IS, indicating a comparable, higher level of instability compared to CC. This suggests  
389 IM could provide an enhanced training shoe to active consumers, over current unstable  
390 footwear technologies, by creating instability in an unpredictable manner similar to IS. Other  
391 adaptations were dependant on the type of locomotion or the different stimuli of IM or IS.

392 Consistent with previous research on uneven surfaces, IM trials triggered increased stride  
393 frequency and reduced step length (Marigold & Patla, 2008; McAndrew et al., 2010;  
394 Voloshina et al., 2013), reduced shoe-surface angle (Marigold & Patla, 2002; Menant et al.,  
395 2008) and increased knee flexion (Gates et al., 2012; Thomas & Derrick, 2003) at initial  
396 contact in both walking and running. Shorter steps and a reduced sagittal shoe-surface angle  
397 reduce the risk of slipping by decreasing the shear forces and consequently reducing the  
398 friction coefficient at the shoe-floor interface (Menant et al., 2008). Increased knee flexion  
399 would help to lower the centre of mass, increasing stability (MacLellan & Patla, 2006). These  
400 active posture adaptations at initial contact in IM suggest a cautious locomotion pattern was  
401 adopted (Menant et al., 2008; Marigold & Patla, 2002). Stability was subjectively perceived  
402 lowest in IM, giving further evidence the level of instability was enough to induce these  
403 cautious posture alterations. Similar cautious kinematic adaptations at initial contact were  
404 found in IS during running, but not walking. This may be due to injury risk of the IS stimuli  
405 being subjectively perceived greater in running than walking, and enough to induce a  
406 cautious gait strategy.

407 The higher maximum ankle inversion across all steps and more variable frontal ankle motion  
408 in IM and IS compared to the control (Fig 5) were caused by the size, shape and hardness of

409 the materials imposed between the shoe-surface interfaces. This may have caused the greater  
410 perceived instability and injury risk. However, this does not mean they were more dangerous  
411 to participants. Increased ankle inversion is not a risk factor for ankle sprain in healthy  
412 participants whilst running (Willems, Witvrouwa, Delbaere, De Cock, & De Clercq, 2005).  
413 Also, the maximum ankle inversion angle was within the normal range of frontal ankle  
414 motion (Ottaviani, Ashton-Miller, Kothari, & Wojtys, 1995). Keeping ankle range of motion  
415 within this safe range is an advantage of the IM shoe compared to a natural irregular terrain  
416 that imposes a greater risk and could cause injury. Thus, the irregular midsoles provide a  
417 similar stimulus to an IS, which is not always available or safe to use, and offer a viable  
418 alternative.

419 The increased gait cycle variability in IM and IS during walking, and IS during running is an  
420 indicator of instability and has been linked to risk of falling (Moe-Nilssen & Helbostad, 2005;  
421 Thies et al., 2005). Previous research also found increased variability of step length and step  
422 time on IS (Gates et al., 2012; Marigold & Patla, 2008; McAndrew et al., 2010; Thies et al.,  
423 2005; Voloshina et al., 2013; Voloshina & Ferris 2015). However, the increased gait cycle  
424 variability does not necessarily represent loss of balance, but rather active alterations to  
425 maintain stability to the unpredictable perturbations, allowing the acquisition of more flexible  
426 locomotion patterns. The reason for variability being higher in IM during walking than  
427 running is related to the reduced shoe-surface angle (walking =  $16.6^\circ$ , running =  $12.4^\circ$ ).  
428 Reducing the angular displacement of the shoe to the ground likely reduced the perturbation  
429 effect whilst running in IM, enabling a more regular locomotion pattern. How to increase the  
430 variability whilst running in IM to a similar level as the IS should be considered in the design  
431 of future prototypes.

432 The increased lower-limb kinematic variability in IS and IM has also been reported  
433 previously on irregular surfaces during walking (Gates et al., 2012; Sterzing et al., 2014a;  
434 Voloshina et al., 2013) and running (Sterzing et al., 2014b; Voloshina & Ferris 2015) and,  
435 walking in unstable shoes (Stöggl et al., 2010). According to Dynamics Systems Theory,  
436 opposed to the more global movement level, increasing variability at the joint/segment level  
437 is associated with functional benefits and not necessarily related with reduced stability (Li,  
438 Haddad, & Hamill, 2005). Performance can be achieved consistently through a variety of  
439 movement pathways, increasing adaptability to perturbations (Davids et al., 2006; Latash,  
440 2012; Wilson, Simpson, van Emmerik, & Hamill, 2008). There is some evidence to suggest  
441 this also reduces the risk of chronic overuse injuries in running because the stresses are

442 spread more evenly over the soft tissues (Hamill, van Emmerik, & Heiderscheit, 1999). In  
443 this respect, we propose IM offers wearers another training benefit, in addition to those  
444 discussed already, of improving the level of this functional joint variability. Whether the level  
445 of functional variability remains high, or reduces to the level of a regular shoe, as reported  
446 previously (Stöggl et al., 2010), warrants further investigation.

447 Electromyography results revealed few common activation strategies to the irregular shoe-  
448 surfaces. One prevalent approach to IS and IM was to increase the peroneus longus activation  
449 during the loading phase of running. The peroneal muscles are the main muscles to provide  
450 eccentric control to protect against lateral ankle sprains (Ashton-Miller, Ottaviani,  
451 Hutchinson, & Wojtys, 1996). Therefore, it appears the increased peroneus longus activation  
452 was a mechanism to control the increased inversion and more variable frontal ankle motion of  
453 IM and IS. With training, this would increase the peroneus muscle strength and reduce the  
454 risk of ankle sprains, as found in conventional unstable shoes (Kaelin et al., 2011). The  
455 perceived risk of injury and energy requirement were lower walking compared to running in  
456 IM and IS, similar to previous research on IS (Sterzing et al., 2014a; Sterzing et al., 2014b).  
457 This may relate to the lack of increased peroneus longus activation during walking in IM and  
458 IS compared to running. However, some participants increased the peroneus longus  
459 activation whilst walking in IM and IS, suggesting individual adaptation strategies for coping  
460 with the constraints occurred, as referred to previously (Apps, Ding, Cheung, & Sterzing,  
461 2014). The other common finding was a reduced tibialis anterior activation on the irregular  
462 shoe and surface conditions, particularly in IM whilst walking. This result supports previous  
463 observations on irregular surfaces (Hettinga, Stefanyshyn, Fairbairn, & Worobets, 2005;  
464 Voloshina et al., 2013), and in unstable shoes (Nigg et al., 2006) and is associated with the  
465 reduced shoe-surface angle at initial contact.

466 This research is subject to certain limitations. The use of set speeds on a treadmill, has been  
467 shown to affect variability compared to when subjects run at their preferred speed (Sekiya,  
468 Nagasaki, Ito, & Furuna, 1997) and overground (Wheat, Milner, & Bartlett, 2004). However,  
469 we do not expect that this would have affected any of the conditions differently and  
470 confounded our conclusions. The time to accommodate to the shoe-surface conditions was  
471 limited to 60 seconds, so the results reported only apply to the acute responses. It is likely  
472 adaptations would change after the initial accommodation period, as previously reported  
473 (Stöggl et al., 2010; Blair et al., 2013). Furthermore, although the irregular treadmill surface  
474 developed did provide continuous unpredictable perturbations, it was limited by the size,

475 hardness and shape of inserts attached and would not have provided the same variety of  
476 perturbations as a natural uneven terrain. In IM trials, participants could perceive the objects  
477 inside the rubber bags under the plantar sole which may have caused the kinematic  
478 adaptations, rather than the instability. Future prototypes should aim to reduce this haptic  
479 sensation.

## 480 **5. Conclusion**

481 In conclusion, we have created a novel shoe that provides continuously random perturbations.  
482 The motivation for developing such a shoe was to have a more challenging stimulus than  
483 existing unstable footwear, thus providing greater functional training benefits. This shoe  
484 successfully increased biomechanical and perceived instability relative to a stable shoe and  
485 simulated certain adaptations of an unpredictable irregular surface during walking and  
486 running. An additional training benefit of the irregular midsole, of increasing the functional  
487 level of joint kinematic variability is proposed, which aligns with the dynamics systems  
488 perspective. Future studies should confirm these suggested training advantages over unstable  
489 shoes, by assessing the adaptability to unpredictable perturbations after regular use.

490

## 491 **References**

492 Apps, C., Ding, R., Cheung, J. T. M., & Sterzing, T. (2014). Individual and generalized lower  
493 limb muscle activity and kinematic adaptations during walking on an unpredictable irregular  
494 surface. *Journal of Foot and Ankle Research*, 7(1), 1. doi: 10.1186/1757-1146-7-s1-a3.

495 Ashton-Miller, J. A., Ottaviani, R. A., Hutchinson, C., & Wojtys, E. M. (1996). What best  
496 protects the inverted weightbearing ankle against further inversion? Evertor muscle strength  
497 compares favorably with shoe height, athletic tape, and three orthoses. *The American Journal  
498 of Sports Medicine*, 24(6), 800-809.

499 Au, E.Y.L., & Goonetilleke, R.S. (2007). A qualitative study on the comfort and fit of ladies'  
500 dress shoes. *Applied Ergonomics*, 38, 687–696.

501 Blair, S. J., Lake, M. J., Sterzing, T., & Cheung, J. T. (2013). Lower extremity biomechanics  
502 following a period of adaptation to wearing unstable shoes. *Footwear Science*, 5(sup1), S139-  
503 S141. doi: 10.1080/19424280.2013.799607.

504 Buchecker, M., Pfusterschmied, J., Moser, S., & Müller, E. (2012). The effect of different  
505 Masai Barefoot Technology (MBT) shoe models on postural balance, lower limb muscle



506 activity and instability assessment. *Footwear Science*, 4(2), 93-100.  
507 doi.org/10.1080/19424280.2012.674560.

508 Cappozzo, A., Catani, F., Della Croce, U., & Leardini, A. (1995). Position and orientation in  
509 space of bones during movement: anatomical frame definition and determination. *Clinical*  
510 *Biomechanics*, 10(4), 171-178. doi:10.1016/0268-0033(95)91394-T.

511 Cavanagh, P. R & Ulbrecht, J. S. (1994). Clinical plantar pressure measurement in diabetes:  
512 rationale and methodology. *The Foot*, 4, 123-135. doi:10.1016/0958-2592(94)90017-5.

513 Cimadoro, G., Paizis, C., Alberti, G., & Babault, N. (2013). Effects of different unstable  
514 supports on EMG activity and balance. *Neuroscience Letters*, 548, 228-232.  
515 doi.org/10.1016/j.neulet.2013.05.025.

516 Davids, K., Bennett, S., & Newell, K. M. (2006). *Movement system variability*. Champaign,  
517 Human Kinetics.

518 Faul, F., Erdfelder, E., Lang, A.-G., & Buchner, A. (2007). G\*Power 3: A flexible statistical  
519 power analysis program for the social, behavioral, and biomedical sciences. *Behavior*  
520 *Research Methods*, 39, 175-191.

521 Forghany, S., Nester, C. J., Richards, B., Hatton, A. L., & Liu, A. (2014). Rollover footwear  
522 affects lower limb biomechanics during walking. *Gait & Posture*, 39(1), 205-212.  
523 doi.org/10.1016/j.gaitpost.2013.07.009.

524 Gates, D. H., Wilken, J. M., Scott, S. J., Sinitski, E. H., & Dingwell, J. B. (2012). Kinematic  
525 strategies for walking across a destabilizing rock surface. *Gait & Posture*, 35(1), 36-42. doi:  
526 10.1016/j.gaitpost.2011.08.001 PMID: 21890361.

527 Hamill, J., van Emmerik, R. E., Heiderscheit, B. C., & Li, L. (1999). A dynamical systems  
528 approach to lower extremity running injuries. *Clinical Biomechanics*, 14(5), 297-308.  
529 doi.org/10.1016/S0268-0033(98)90092-4.

530 Hettinga, B. A., Stefanyshyn, D. J., Fairbairn, J. C., & Worobets, J. T. (2005). Biomechanical  
531 effects of hiking on a non-uniform surface. In *Proceeding. of the 7th Symposium* (pp. 41-42).  
532 Footwear Biomechanics.

533 Horsak, B., & Baca, A. (2013). Effects of toning shoes on lower extremity gait  
534 biomechanics. *Clinical Biomechanics*, 28(3), 344-349.  
535 doi.org/10.1016/j.clinbiomech.2013.01.009.

536 James, C. R. (2004). Considerations of movement variability in biomechanics research. In  
537 Stergiou, N (Eds.) *Innovative analyses of human movement*. Champaign: Human Kinetics.

538 Kaelin, X., Segesser, B., & Wasser, T. (2011). Unstable shoes and rehabilitation. *Footwear*  
539 *Science*, 3(sup1), S85-S86. doi: 10.1080/19424280.2011.575822.

540 Kim, H., & Ashton-Miller, J. A. (2012). A shoe sole-based apparatus and method for  
541 randomly perturbing the stance phase of gait: Test–retest reliability in young adults. *Journal*  
542 *of Biomechanics*, 45(10), 1850-1853. doi: 10.1016/j.jbiomech.2012.05.003.

543 Lam, W. K., Sterzing, T., & Cheung, J. T. M. (2013). Influence of protocol complexity on fit  
544 perception of basketball footwear. *Footwear Science*, 5(3), 155-163.

545 Landry, S. C., Nigg, B. M., & Tecante, K. E. (2010). Standing in an unstable shoe increases  
546 postural sway and muscle activity of selected smaller extrinsic foot muscles. *Gait &*  
547 *Posture*, 32(2), 215-219. doi.org/10.1016/j.gaitpost.2010.04.018.

548 Latash, M. L. (2012). The bliss (not the problem) of motor abundance (not  
549 redundancy). *Experimental Brain Research*, 217(1), 1-5. doi:10.1007/s00221-012-3000-4.

550 Li, L., Haddad, J. M., & Hamill, J. (2005). Stability and variability may respond differently to  
551 changes in walking speed. *Human movement science*, 24(2), 257-267.

552 MacLellan, M. J., & Patla, A. E. (2006). Adaptations of walking pattern on a compliant  
553 surface to regulate dynamic stability. *Experimental Brain Research*, 173(3), 521-530.  
554 doi:10.1007/s00221-006-0399-5.

555 Maiwald, C., Sterzing, T., Mayer, T. A., & Milani, T. L. (2009). Detecting foot-to-ground  
556 contact from kinematic data in running. *Footwear Science*, 1(2), 111-118. doi:  
557 10.1080/19424280903133938.

558 Marigold, D. S., & Patla, A. E. (2002). Strategies for dynamic stability during locomotion on  
559 a slippery surface: effects of prior experience and knowledge. *Journal of*  
560 *Neurophysiology*, 88(1), 339-353. doi: 10.1152/jn.00691.2001.

561 Marigold, D. S., & Patla, A. E. (2008). Age-related changes in gait for multi-surface  
562 terrain. *Gait & Posture*, 27(4), 689-696. doi: 10.1016/j.gaitpost.2007.09.005.

563 McAndrew, P. M., Dingwell, J. B., & Wilken, J. M. (2010). Walking variability during  
564 continuous pseudo-random oscillations of the support surface and visual field. *Journal of*  
565 *Biomechanics*, 43(8), 1470-1475. doi: 10.1016/j.jbiomech.2010.02.003.

566 Menant, J. C., Perry, S. D., Steele, J. R., Menz, H. B., Munro, B. J., & Lord, S. R. (2008).  
567 Effects of shoe characteristics on dynamic stability when walking on even and uneven  
568 surfaces in young and older people. *Archives of Physical Medicine and Rehabilitation*, 89(10),  
569 1970-1976. doi: 10.1016/j.apmr.2008.02.031.

570 Moe-Nilssen, R., & Helbostad, J. L. (2005). Interstride trunk acceleration variability but not  
571 step width variability can differentiate between fit and frail older adults. *Gait &*  
572 *Posture*, 21(2), 164-170. doi: 10.1016/j.gaitpost.2004.01.013.

573 Nigg, B., Hintzen, S., & Ferber, R. (2006). Effect of an unstable shoe construction on lower  
574 extremity gait characteristics. *Clinical Biomechanics*, 21(1), 82-88. doi:  
575 10.1016/j.clinbiomech.2005.08.013.

576 O'Connor, C. M., Thorpe, S. K., O'Malley, M. J., & Vaughan, C. L. (2007). Automatic  
577 detection of gait events using kinematic data. *Gait & Posture*, 25(3), 469-474. doi:  
578 10.1016/j.gaitpost.2006.05.016.

579 Ottaviani, R. A., Ashton-Miller, J. A., Kothari, S. U., & Wojtys, E. M. (1995). Basketball  
580 shoe height and the maximal muscular resistance to applied ankle inversion and eversion  
581 moments. *The American Journal of Sports Medicine*, 23(4), 418-423.

582 Price, C., Smith, L., Graham-Smith, P., & Jones, R. (2013). The effect of unstable sandals on  
583 instability in gait in healthy female subjects. *Gait & Posture*, 38(3), 410-415.  
584 doi.org/10.1016/j.gaitpost.2013.01.003.

585 Ramstrand, N., Thuesen, A. H., Nielsen, D. B., & Rusaw, D. (2010). Effects of an unstable  
586 shoe construction on balance in women aged over 50 years. *Clinical Biomechanics*, 25(5),  
587 455-460. doi.org/10.1016/j.clinbiomech.2010.01.014.

588 Romkes, J., Rudmann, C., & Brunner, R. (2006). Changes in gait and EMG when walking  
589 with the Masai Barefoot Technique. *Clinical Biomechanics*, 21(1), 75-81.  
590 doi.org/10.1016/j.clinbiomech.2005.08.003.

591 Sacco, I. C., Sartor, C. D., Cacciari, L. P., Onodera, A. N., Dinato, R. C., Pantaleão, E., ... &  
592 Yokota, M. (2012). Effect of a rocker non-heeled shoe on EMG and ground reaction forces  
593 during gait without previous training. *Gait & Posture*, 36(2), 312-315.  
594 doi.org/10.1016/j.gaitpost.2012.02.018.

595 Schiemann, S., Lohrer, H., & Nauck, T. (2015). Influence of three different unstable shoe  
596 constructions on EMG-activity during treadmill walking—a cross-sectional study with respect

597 to sensorimotor activation. *Footwear Science*, 7(1), 1-7.  
598 doi.org/10.1080/19424280.2014.939231.

599 Sekiya, N., Nagasaki, H., Ito, H., & Furuna, T. (1997). Optimal walking in terms of  
600 variability in step length. *Journal of Orthopaedic & Sports Physical Therapy*, 26(5), 266-272.

601 Sobhani, S., Hijmans, J., van den Heuvel, E., Zwerver, J., Dekker, R., & Postema, K. (2013).  
602 Biomechanics of slow running and walking with a rocker shoe. *Gait & Posture*, 38(4), 998-  
603 1004. doi.org/10.1016/j.gaitpost.2013.05.008.

604 Sterzing, T., Apps, C., Ding, R., & Cheung, J. T. M. (2014a). Walking on an unpredictable  
605 irregular surface changes lower limb biomechanics and subjective perception compared to  
606 walking on a regular surface. *Journal of Foot and Ankle Research*, 7(1), A81. doi:  
607 10.1186/1757-1146-7-s1-a81.

608 Sterzing, T., Apps, C., Ding, R., & Cheung, J. T. M. (2014b). Running on an unpredictable  
609 irregular surface changes lower limb biomechanics and subjective perception compared to  
610 running on a regular surface. *Journal of Foot and Ankle Research*, 7(1), A80. doi:  
611 10.1186/1757-1146-7-s1-a80.

612 Sterzing, T., Wulf, M., Qin, T. Y., Cheung, J. T. M., & Brauner, T. (2014c). Effect of soccer  
613 shoe ball girth differences on fit perception, agility running and running speed  
614 perception. *Footwear Science*, 6(2), 97-103.

615 Sterzing, T., Cheung, J., & Li, W. (2013). Li Ning Sports Goods Co Ltd, China. Dynamically  
616 unpredictable unstable footwear. Chinese Patent: 2013 1 0084410.4., granted 2015 Feb 2.

617 Stöggl, T., Haudum, A., Birklbauer, J., Murrer, M., & Müller, E. (2010). Short and long term  
618 adaptation of variability during walking using unstable (Mbt) shoes. *Clinical*  
619 *Biomechanics*, 25(8), 816-822. doi: 10.1016/j.clinbiomech.2010.05.012.

620 Thies, S. B., Richardson, J. K., & Ashton-Miller, J. A. (2005). Effects of surface irregularity  
621 and lighting on step variability during gait: A study in healthy young and older women. *Gait*  
622 *& Posture*, 22(1), 26-31. doi: 10.1016/j.gaitpost.2004.06.004.

623 Thomas, J. M., & Derrick, T. R. (2003). Effects of step uncertainty on impact peaks, shock  
624 attenuation, and knee/subtalar synchrony in treadmill running. *Journal of Applied*  
625 *Biomechanics*, 19(1), 60-70.

626 Turbanski, S., Lohrer, H., Nauck, T., & Schmidtbleicher, D. (2011). Training effects of two  
627 different unstable shoe constructions on postural control in static and dynamic testing  
628 situations. *Physical Therapy in Sport*, 12(2), 80-86. doi.org/10.1016/j.ptsp.2011.01.001.

629 Wheat, J. S., Milner, C. E., & Bartlett, R. M. (2004). Kinematic variability during overground  
630 and treadmill running. *Journal of Sports Sciences*, 22(3), 245-246.

631 Willems, T., Witvrouw, E., Delbaere, K., De Cock, A., & De Clercq, D. (2005). Relationship  
632 between gait biomechanics and inversion sprains: a prospective study of risk factors. *Gait &*  
633 *posture*, 21(4), 379-387.

634 Wilson, C., Simpson, S. E., Van Emmerik, R. E., & Hamill, J. (2008). Coordination  
635 variability and skill development in expert triple jumpers. *Sports Biomechanics*, 7(1), 2-9. doi:  
636 10.1080/14763140701682983.

637 Van Emmerik, R. E., & van Wegen, E. E. (2000). On variability and stability in human  
638 movement. *Journal of Applied Biomechanics*, 16(4), 394-406.

639 Voloshina, A. S., Kuo, A. D., Daley, M. A., & Ferris, D. P. (2013). Biomechanics and  
640 energetics of walking on uneven terrain. *Journal of Experimental Biology*, 216(21), 3963-  
641 3970. doi: 10.1242/jeb.081711.

642 Voloshina, A. S., & Ferris, D. P. (2015). Biomechanics and energetics of running on uneven  
643 terrain. *Journal of Experimental Biology*, 218(5), 711-719. doi: 10.1242/jeb.106518.

644 **Table 1. Mean (SD) gait cycle parameters and kinematics during walking across participants.**

<b>Walking</b>	<b>Variable</b>	<b>CC</b>	<b>IM</b>	<b>IS</b>	<b>ANOVA p-value</b>	<b>Post hoc result</b>
Gait cycle	Stance time [secs]	.63 (.04)	.62 (.04)	.65 (.02)	.010	IS > IM
	Swing time [secs]	.38 (.02)	.36 (.02)	.38 (.02)	<.001	IS, CC > IM
	Step length [m]	.87 (.05)	.86 (.05)	.90 (.03)	.010	IS > IM
	Stride frequency [stride/min]	59.4 (3.2)	61.3 (3.1)	58.5 (1.9)	<.001	IM > IS, CC
Kinematics at initial contact	Shoe-surface [°]	24.7 (4.3)	18.6 (4.8)	22.8 (5.0)	.001	CC, IS > IM
	Ankle dorsiflexion [°]	0.9 (3.0)	-1.1 (4.0)	-0.4 (3.6)	.161	---
	Ankle inversion [°]	-3.3 (3.1)	-3.9 (3.0)	-2.5 (3.9)	.028	IM > IS
	Knee flexion [°]	14.5 (5.7)	20.1 (7.1)	16.9 (5.6)	<.001	IM > CC, IS
Kinematics during stance	Ankle dorsiflexion MAX [°]	7.0 (3.1)	8.0 (3.1)	7.8 (3.7)	.248	---
	Ankle eversion MAX [°]	7.3 (2.1)	5.3 (5.5)	8.7 (3.6)	.005	IS > IM
	Sagittal ankle ROM [°]	17.6 (4.5)	12.5 (4.8)	18.7 (4.0)	<.001	CC, IS > IM
	Frontal ankle ROM [°]	10.6 (3.7)	10.6 (3.3)	11.8 (2.2)	.128	---
	Knee flexion MAX [°]	31.2 (7.5)	33.6 (8.4)	32.1 (7.2)	.038	---

645 MAX = maximum, ROM = Range of motion

646

647

648

649

650

651

652

653 **Table 2. Mean (SD) gait cycle parameters and kinematics during running across participants.**

Running	Variable	CC	IM	IS	ANOVA p-value	Post hoc result
Gait cycle	Stance time [secs]	.35 (.02)	.34 (.01)	.34 (.02)	.014	CC > IS
	Swing time [secs]	.39 (.04)	.38 (.04)	.40 (.04)	.018	IS > IM
	Step length [m]	.77 (.04)	.75 (.03)	.75 (.04)	.011	CC > IS
	Stride frequency [stride/min]	82.2 (3.5)	84.3 (4.4)	82.2 (3.8)	.001	IM > CC, IS
Kinematics at initial contact	Shoe-surface [°]	16.4 (2.5)	12.5 (3.0)	12.9 (3.8)	< .001	CC > IM, IS
	Ankle dorsiflexion [°]	6.7 (3.1)	6.1 (0.4)	5.0 (3.9)	.017	CC > IS
	Ankle inversion [°]	-5.7 (3.4)	-6.1 (3.4)	-4.6 (4.4)	.530	---
	Knee flexion [°]	22.2 (4.2)	28 (4.4)	26.9 (4.1)	< .001	IM > IS > CC
Kinematics during stance	Ankle dorsiflexion MAX [°]	13.6 (2.9)	16.2 (4.0)	13.5 (3.5)	< .001	IM > CC, IS
	Ankle eversion MAX [°]	9.2 (3.5)	4.1 (7.5)	9.6 (5.2)	< .001	CC, IS > IM
	Sagittal ankle ROM [°]	16.6 (1.9)	17.0 (2.0)	17.2 (2.3)	.439	---
	Frontal ankle ROM [°]	14.9 (3.0)	11.1 (4.3)	14.4 (2.9)	.001	CC, IS > IM
	Knee flexion MAX [°]	48.6 (4.3)	48.4 (4.8)	49.7 (4.7)	.006	IS > CC, IM

654 MAX = maximum, ROM = Range of motion

655

656 **Table 3: Normalised mean (SD) electromyography data during pre-activation and loading phases**  
 657 **across participants during walking**

Muscle	Phase	CC	IM	IS	ANOVA p-value	Post hoc result
Gastrocnemius Medialis	Pre-activation	1.8 (1.2)	4.4 (3.4)	3.1 (3.2)	.008	IM>CC
	Loading	4.2 (2.1)	5.2 (2.6)	4.2 (1.9)	.263	---
Tibialis Anterior	Pre-activation	18.7 (5.8)	11.8 (5.6)	15.0 (6.9)	.004	CC, IS>IM
	Loading	19.2 (3.8)	9.1 (4.1)	18.2 (6.0)	<.001	CC, IS>IM
Peroneus Longus	Pre-activation	4.7 (2.0)	5.1 (2.6)	6.4 (2.5)	.113	---
	Loading	9.3 (3.7)	14.0 (5.8)	13.5 (5.6)	.062	---
Bicep Femoris	Pre-activation	27.2 (3.6)	23.0 (9.2)	22.9 (5.5)	.005	CC>IS
	Loading	12.1 (4.7)	13.4 (7.4)	12.1 (4.9)	.484	---
Vastus Medialis	Pre-activation	14.4 (6.5)	14.1 (7.6)	13.6 (6.8)	.843	---
	Loading	28.2 (5.4)	28.8 (9.7)	29.9 (8.4)	.699	---
Vastus Lateralis	Pre-activation	10.2 (4.5)	9.0 (5.2)	8.8 (4.7)	.307	---
	Loading	29.1 (5.6)	23.5 (7.5)	23.8 (6.9)	.030	CC>IS

658  
 659  
 660  
 661  
 662  
 663  
 664  
 665  
 666  
 667  
 668  
 669  
 670  
 671  
 672  
 673  
 674  
 675



676  
677

**Table 4: Normalised mean (SD) electromyography data during pre-activation and loading phases across participants during running**

Muscle	Phase	CC	IM	IS	ANOVA p-value	Post hoc result
Gastrocnemius Medialis	Pre-activation	2.3 (1.7)	3.5 (3.2)	2.9 (2.6)	.039	IM>CC
	Loading	21.3 (4.8)	20.8 (6.4)	19.2 (5.6)	.234	---
Tibialis Anterior	Pre-activation	24.1 (3.5)	10.6 (8.2)	12.6 (5.6)	<.001	CC>IM,IS
	Loading	10.4 (4.2)	10.4 (7.0)	15.5 (15.8)	.301	---
Peroneus Longus	Pre-activation	4.3 (1.5)	7.0 (5.2)	6.9 (3.8)	.018	IS>CC
	Loading	24.0 (5.4)	30.8 (10.0)	34.6 (22.2)	.023	IM,IS>CC
Bicep Femoris	Pre-activation	24.3 (5.3)	24.1 (12.2)	21.4 (7.9)	.420	---
	Loading	10.6 (5.2)	10.5 (6.6)	9.9 (3.8)	.803	---
Vastus Medialis	Pre-activation	8.7 (2.9)	8.8 (2.9)	8.8 (2.6)	.963	---
	Loading	31.8 (3.2)	28.3 (6.0)	31.5 (7.3)	.069	---
Vastus Lateralis	Pre-activation	6.6 (3.2)	6.6 (2.4)	6.9 (3.7)	.752	---
	Loading	29.5 (4.9)	26.4 (8.3)	29.6 (15.4)	.144	---

678  
679  
680  
681  
682  
683  
684  
685  
686  
687  
688  
689  
690  
691  
692  
693  
694  
695  
696

697 **Table 5: Subjective perception scores (Mean (SD)) during walking and running across participants**

Variable	Locomotion	CC	IM	IS	ANOVA p-value	Post hoc result
Stability	Walk	5.6 (1.2)	2.9 (1.2)	4.2 (1.4)	<.001	IM<IS<CC
	Run	5.4 (1.6)	2.7 (1.2)	3.8 (1.6)	<.001	IM<IS<CC
Injury risk	Walk	3.2 (1.3)	6.3 (1.1)	5.8 (1.5)	<.001	IM>IS>CC
	Run	3.7 (1.3)	6.8 (1.4)	6.0 (1.6)	<.001	IM>IS>CC
Energy Consumption	Walk	3.1 (1.4)	4.6 (1.5)	4.7 (1.4)	<.001	IM, IS>CC
	Run	4.9 (0.9)	6.5 (1.3)	6.3 (1.4)	<.001	IM, IS>CC

698

699

700

701

702

703

704

705

706

707

708

709

710

711

712

713

714

715

716

717

718

719

720

721 **Fig 1. The regular and irregular shoe midsoles.** The regular midsole (left, top) was removed from  
722 the original shoe upper and cut into same width as IM bags (left, middle), weights attached (left,  
723 bottom). The irregular midsole shoe (right, top), the rubber midsole bags (right middle) with cubes  
724 and ball bearings placed inside (close up: bottom right). © 2013. All rights reserved (Sterzing et al.,  
725 2013 (Li Ning Sports Goods Co. Ltd, China)).

726

727 **Fig 2. The regular and irregular treadmill surface.** The regular treadmill surface covered with strips  
728 of Velcro (top left) and the irregular treadmill surface, created by attaching 4 kinds of EVA inserts to  
729 the belt via Velcro (top right). Data collection of an IS run trial, the large 9-point Likert scale allowed  
730 scores to be taken whilst participants were still on the treadmill (bottom).

731

732 **Fig 3. Variability (CV) of gait cycle parameters across participants.** 1 = significantly greater than CC,  
733 2 = significantly greater than IM, 3 = significantly greater than IS ( $p < .05$ ). Notice IM has higher values  
734 similar to IS during walking and lower values similar to CC during running.

735

736 **Fig 4. Variability (CV) of joint/segment angles at initial contact (IC) and during stance across**  
737 **participants.** ROM = range of motion. 1 = significantly greater than CC, 2 = significantly greater than  
738 IM ( $p < .05$ ).

739

740 **Fig 5. Joint angle plotted against stance phase during walking and running across subjects.**

741 Solid thick lines represent mean sagittal ankle angle (top), frontal ankle angle (middle) and  
742 sagittal knee angle (bottom). CC illustrated by the black line, IM the lighter line and IS the  
743 lightest line (mostly overlaid by CC). Shaded areas (CC, IM) and dotted lines (IS) illustrate  
744 mean intra-subject variability at each percentage of stance phase from 0% at heel-strike to  
745 100% at toe-off.