



The University of Manchester

The University of Manchester Research

# Rumination and psychosis: an experimental, analogue study of the role of perseverative thought processes in voice-hearing

DOI:

10.1080/17522439.2017.1280073

#### **Document Version**

Accepted author manuscript

Link to publication record in Manchester Research Explorer

# Citation for published version (APA):

Hartley, S., Bucci, S., & Morrison, A. (2017). Rumination and psychosis: an experimental, analogue study of the role of perseverative thought processes in voice-hearing. *Psychosis*. https://doi.org/10.1080/17522439.2017.1280073

# Published in:

**Psychosis** 

# Citing this paper

Please note that where the full-text provided on Manchester Research Explorer is the Author Accepted Manuscript or Proof version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version.

## **General rights**


Copyright and moral rights for the publications made accessible in the Research Explorer are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

# Takedown policy

If you believe that this document breaches copyright please refer to the University of Manchester's Takedown Procedures [http://man.ac.uk/04Y6Bo] or contact uml.scholarlycommunications@manchester.ac.uk providing relevant details, so we can investigate your claim.



# **Psychosis**



# Rumination and psychosis: an experimental, analogue study of the role of perseverative thought processes in voice-hearing

Journal:	Psychosis
Manuscript ID	RPSY-2016-0021.R2
Manuscript Type:	Brief Report
Keywords:	Hearing voices, Hallucinations, Affect, Causal explanations, Trauma
Abstract:	We rigorously tested the role of rumination in the development of voice-hearing type experiences. One-hundred and two students watched a video depicting a physical assault and then either ruminated about the contents or were distracted. Participants listened to an anomalous auditory stimulus and recorded any words/ phrases along with distress. Manipulation checks confirmed that the rumination group showed greater perseveration regarding the film content than the distraction group. However, the groups did not significantly differ on number of words recorded, convergence with the video content or distress. These findings indicate that rumination might not necessarily be involved in the development of these experiences or associated distress. The current work should provide an impetus to conduct additional rigorously controlled experimental or prospective work to fully discern the key processes relevant to the experience of voice-hearing following exposure to stressful events.

SCHOLARONE™ Manuscripts

URL: http:/mc.manuscriptcentral.com/rpsy



# <u>Abstract</u>

We rigorously tested the role of rumination in the development of voice-hearing type experiences. One-hundred and two students watched a video depicting a physical assault and then either ruminated about the contents or were distracted. Participants listened to an anomalous auditory stimulus and recorded any words/ phrases along with distress. Manipulation checks confirmed that the rumination group showed greater perseveration regarding the film content than the distraction group. However, the groups did not significantly differ on number of words recorded, convergence with the video content or distress. These findings indicate that rumination might not necessarily be involved in the development of these experiences or associated distress. The current work should provide an impetus to conduct additional rigorously controlled experimental or prospective work to fully discern the key processes relevant to the experience of voice-hearing following exposure to stressful events.


**Keywords:** hearing voices; hallucinations; affect; causal explanations; trauma

Rumination is a type of negative perseverative processing, which maintains depressed mood (Nolen-Hoeksema & Morrow, 1991). Recently, research has demonstrated high levels of rumination in groups experiencing psychosis (Vorontsova, Garety, & Freeman, 2013) and has implicated rumination in delusional ideas and voice-hearing (Hartley, Haddock, Vasconcelos E Sa, Emsley, & Barrowclough, 2014) via increases in thought suppression in response to rumination and subsequent rebound intrusive thoughts that are then misinterpreted as voices (Jones & Fernyhough, 2009). Rumination also maintains post-trauma intrusions (Ehring, Frank, & Ehlers, 2008) and verbal appraisals of stressful material may be relived subsequently as a trauma-related psychotic intrusion. The current study is the first randomised, experimental study of a role for rumination in the development of voice-hearing type experiences. Our aim was to investigate whether experimentally-induced rumination following exposure to stressful video material would be related to hallucination-like experiences, their thematic content and the distress associated with these.

# Method

An opportunity sample of 102 university staff and students who were English-speaking, 18 years old or above, with normal/corrected vision and hearing completed the full procedure described in Figure 1. Potential participants were excluded if they had secondary care psychiatric service history, experience of physical assault or had viewed the video footage previously. The mean age of the sample was 21 years (SD 4.12), comprised 78 females and 24 males, with the majority (n=87, 85%) currently engaged in a psychology-related area of study.

Figure 1: Study procedure



# **Results**

There were no significant differences in demographic or baseline measures between the conditions. The manipulation was successful, generating higher mean levels of perseverative thought in the rumination condition (17.14, SD 7.60) as compared to the distraction condition (14.00, SD 7.90); [t(100)=-2.04, p=.044]. There were no significant differences in the primary outcome of number of eligible words recorded as part of the auditory task [t(100)=.400, p=.690]. There were also no significant differences in the number of overall recordings [t(100)=.502, p=.616], thematic matches [t(100)=.557, p=.579] or average distress [t(100)=.724, p=.471]. In order to confirm that the voice-hearing task operated in the manner expected and previously demonstrated (Feelgood & Rantzen, 1994), the association between the number of recordings and proneness to hallucinate was checked, which showed that the tasks did indeed tap into similar phenomena as outlined in the original study [r(99)=.250; p=.013], in addition to similar average levels of recordings and a broad range of total number of recordings.

# **Conclusions**

Increased ruminative processing did not result in increased levels of voice-hearing experiences, distress or thematic convergence between experiences and the film content. These divergent, non-significant findings are an important contribution to the hitherto cross-sectional literature, which may imply that rumination is engaged in as a reaction to psychotic phenomena that subsequently exacerbates their severity and distress-inducing qualities, rather than being active in their development.

Distraction might not be the most divergent comparator for ruminative processing; an alternative might be to engage in mindful awareness of intrusions. The voice-hearing paradigm is valid and levels of recordings comparable (Feelgood & Rantzen, 1994). Nevertheless, future work might utilise 'high-risk' samples, although this would present ethical issues. The study also rests on the assumption that a continuum model of psychosis is valid.

The current work does not negate a role for rumination in the maintenance of voicehearing and should rather serve as an impetus for further research, with rigorous experimental and prospective methods, alongside consideration of the interpersonal experiences so intrinsic to this kind of event.



### References

Bentall, R. P., & Slade, P. D. (1985). Reliability of a scale measuring disposition towards hallucination: a brief report. *Personality and Individual Differences*, 6(4), 527-529.

Ehring, T., Zetsche, U., Weidacker, K., Wahl, K., Schonfeld, S., & Ehlers, A. (2011). The Perseverative Thinking Questionnaire (PTQ): Validation of a content-independent measure of repetitive negative thinking. *Journal of Behavior Therapy and Experimental Psychiatry*, 42(2), 225-232.

Ehring, T., Frank, S., & Ehlers, A. (2008). The role of rumination and reduced concreteness in the maintenance of posttraumatic stress disorder and depression following trauma. *Cognitive Therapy and Research*, *32*(4), 488-506. doi:10.1007/s10608-006-9089-7

Hartley, S., Haddock, G., Vasconcelos E Sa, D., Emsley, R., & Barrowclough, C. (2014). An experience sampling study of worry and rumination in psychosis. *Psychol Med, 44*(8), 1605-1614. doi:10.1017/S0033291713002080

Jones, S. R., & Fernyhough, C. (2009). Rumination, reflection, intrusive thoughts, and hallucination-proneness: Towards a new model. *Behaviour Research and Therapy*, 47(1), 54-59

Nolen-Hoeksema, S., & Morrow, J. (1991). A prospective study of depression and posttraumatic stress symptoms after a natural disaster: the 1989 Loma Prieta Earthquake. *Journal of personality and social psychology*, 61(1), 115-121.

Vorontsova, N., Garety, P., & Freeman, D. (2013). Cognitive factors maintaining persecutory delusions in psychosis: The contribution of depression. *Journal of abnormal psychology*,122(4), 1121

Zetsche, U., Ehring, T., & Ehlers, A. (2009). The effects of rumination on mood and intrusive memories after exposure to traumatic material: An experimental study. *Journal of Behavior Therapy and Experimental Psychiatry*, 40(4), 499-514. doi:10.1016/j.jbtep.2009.07.001