
This is an electronic reprint of the original article.
This reprint may differ from the original in pagination and typographic detail.

Author(s): Hakonen, Pertti J. & Krusius, M. & Salomaa, M. M. & Simola, J. T.
Title: Comment on 'Nucleation of $^3\text{He-B}$ from the A Phase: A Cosmic-Ray Effect?'
Year: 1985
Version: Final published version

Please cite the original version:

Hakonen, Pertti J. & Krusius, M. & Salomaa, M. M. & Simola, J. T. 1985. Comment on 'Nucleation of $^3\text{He-B}$ from the A Phase: A Cosmic-Ray Effect?'. *Physical Review Letters*. Volume 54, Issue 3. 245-245. ISSN 0031-9007 (printed). DOI: 10.1103/physrevlett.54.245

Rights: © 1985 American Physical Society (APS). This is the accepted version of the following article: Hakonen, Pertti J. & Krusius, M. & Salomaa, M. M. & Simola, J. T. 1985. Comment on 'Nucleation of $^3\text{He-B}$ from the A Phase: A Cosmic-Ray Effect?'. *Physical Review Letters*. Volume 54, Issue 3. 245-245. ISSN 0031-9007 (printed). DOI: 10.1103/physrevlett.54.245, which has been published in final form at <http://journals.aps.org/prl/abstract/10.1103/PhysRevLett.54.245>.

All material supplied via Aaltodoc is protected by copyright and other intellectual property rights, and duplication or sale of all or part of any of the repository collections is not permitted, except that material may be duplicated by you for your research use or educational purposes in electronic or print form. You must obtain permission for any other use. Electronic or print copies may not be offered, whether for sale or otherwise to anyone who is not an authorised user.

Comment on "Nucleation of $^3\text{He-B}$ from the A Phase: A Cosmic-Ray Effect?"

In an interesting Letter, Leggett¹ suggests that the nucleation of $^3\text{He-B}$ from the metastable A phase is a consequence of the passage of a cosmic ray through the sample cell. Here we present new experimental data obtained with a rotating cryostat in NMR experiments² on ^3He . Figure 1(a) shows our data, taken on the $A \rightarrow B$ transition in magnetic fields of 28.4 and 56.9 mT; no field dependence was found between these values. We find that supercooling of the A phase into the B phase is substantial, and there appears to be a threshold temperature ("catastrophe line") $T_{AB}^*(p) \ll T_{AB}(p)$, where the metastable A phase first changes into $^3\text{He-B}$.




FIG. 1. (a) Supercooling of the A phase into the B phase as a function of pressure. (b) Observed nucleation events (histogram) at 29.3 bars are inconsistent with the probability distribution of Ref. 1 with $P(T) \propto (1 - T/T_{AB}^*)^3$. Clearly, only a much more sharply peaked probability at a lower temperature than suggested in Ref. 1 can describe the data; an exponential law for the activation process might be appropriate.

We want to emphasize that in our ^3He cell the $A \rightarrow B$ transition could *only* be observed *during cooling*, never on warmup. The cooling rate varied between 5 and 29 $\mu\text{K}/\text{min}$, but no clear dependence of B -phase nucleation on the cooling rate could be established. When the temperature was stable, the A phase was found to persist, and no $A \rightarrow B$ transition was ever observed in this case. For example, at $p = 29.3$ bars (with $T_{AB} = 0.85 T_c$) and 28.4 mT, the A liquid was cooled as low as $T_{\min} = 0.67 T_c$, and was maintained below $0.7 T_c$ for 2 h; in this run the metastable A phase persisted a total of 8 h, never nucleating the B phase. The temperatures in Fig. 1 are upper-bound estimates for T_{AB}^* . They are determined from NMR susceptibility measurements on platinum powder immersed in the ^3He . A comparison with the A -phase NMR frequency-shift measurement shows that on cooling, the platinum temperature lags behind, at a cooling rate of 20–30 $\mu\text{K}/\text{min}$ by $\leq 0.02 T/T_c$. Even lower liquid temperatures exist in the heat exchanger between the ^3He sample and the refrigerator.

Figure 1(b) represents the data points at 29.3 bars as a histogram and compares them with a fitted distribution of nucleation probability. We conclude that experimentally the nucleation mechanism appears to be active only in a narrow temperature interval near the "catastrophe line" well below the thermodynamic $A \rightarrow B$ transition. An interesting point to note is that on cooling through the $A \rightarrow B$ transition during rotation, the A phase stayed supercooled as long as in the stationary liquid. This is further (indirect) evidence in support of the conclusion³ that continuous vortices are induced by rotation in a bulk liquid sample of $^3\text{He-A}$.

P. J. Hakonen
M. Krusius
M. M. Salomaa
J. T. Simola
Low Temperature Laboratory
Helsinki University of Technology
SF-02150 Espoo 15, Finland

Received 1 October 1984

PACS numbers: 67.50.Fi, 05.70.Fh, 94.40.Vf

¹A. J. Leggett, Phys. Rev. Lett. **53**, 1096 (1984).

²P. J. Hakonen, O. T. Ikkala, S. T. Islander, O. V. Lounasmaa, and G. E. Volovik, J. Low Temp. Phys. **53**, 425 (1983).

³H. K. Seppälä, P. J. Hakonen, M. Krusius, T. Ohmi, M. M. Salomaa, J. T. Simola, and G. E. Volovik, Phys. Rev. Lett. **52**, 1802 (1984).