

Manuscript Number: NEUROPHARM-D-11-00193R1

Title: Kainate postconditioning restores LTP in ischemic hippocampal CA1: onset-dependent second pathophysiological stress

Article Type: Research Paper

Section/Category: Disease

Keywords: 2-vessel occlusion; CA1; kainic acid; LTP; postconditioning; spine density

Corresponding Author: Dr. Tamas Farkas, Ph.D.

Corresponding Author's Institution: University of Szeged

First Author: Dávid Nagy, Ph.D.

Order of Authors: Dávid Nagy, Ph.D.; Kitti Kocsis; János Fuzik; Máté Marosi, Ph.D.; Zsolt Kis, Ph.D.; Vivian I Teichberg, Ph.D., D.Sc.; József Toldi, Ph.D., D.Sc.; Tamas Farkas, Ph.D.

Abstract: Postconditioning can be induced by a broad range of stimuli within minutes to days after an ischemic cerebral insult. A special form is elicited by pharmacological intervention called second pathophysiological stress. The present study aimed to evaluate the effects of low-dose (5 mg/kg) kainate postconditioning with onsets 0, 24 and 48 h after the ischemic insult on the hippocampal synaptic plasticity in a 2-vessel occlusion model in rat. The hippocampal function was tested by LTP measurements of Schaffer collateral-CA1 pyramidal cell synapses in acute slices and the changes in density of Golgi-Cox-stained apical dendritic spines. Postconditioning 0 and 24 h after ischemia was not protective, whereas 48-h-onset postconditioning resulted in the reappearance of a normal spine density (>100,000 spines) 3 days after ischemia, in parallel with the long-term restoration of the damaged LTP function. Similar, but somewhat less effects were observed after 10 days. Our data clearly demonstrate the onset dependence of postconditioning elicited by a subconvulsant dose of kainate treatment in global ischemia, with restoration of the structural plasticity and hippocampal function.

Kainate postconditioning restores LTP in ischemic hippocampal CA1: onset-dependent second pathophysiological stress

Dávid Nagy, Kitti Kocsis, János Fuzik, Máté Marosi, Zsolt Kis, Vivian I. Teichberg, József Toldi
and Tamás Farkas

Corresponding author:

Tamás Farkas Ph.D.

University of Szeged
Department of Physiology, Anatomy and
Neuroscience
Hungary
Szeged, Közép fasor 52.
H-6726

Phone: +36-62-544-381

Fax: +36-62-544-291

email: tfarkas@bio.u-szeged.hu (T. Farkas)

Abstract

Postconditioning can be induced by a broad range of stimuli within minutes to days after an ischemic cerebral insult. A special form is elicited by pharmacological intervention called second pathophysiological stress. The present study aimed to evaluate the effects of low-dose (5 mg/kg) kainate postconditioning with onsets 0, 24 and 48 h after the ischemic insult on the hippocampal synaptic plasticity in a 2-vessel occlusion model in rat. The hippocampal function was tested by LTP measurements of Schaffer collateral-CA1 pyramidal cell synapses in acute slices and the changes in density of Golgi-Cox-stained apical dendritic spines. Postconditioning 0 and 24 h after ischemia was not protective, whereas 48-h-onset postconditioning resulted in the reappearance of a normal spine density (>100,000 spines) 3 days after ischemia, in parallel with the long-term restoration of the damaged LTP function. Similar, but somewhat less effects were observed after 10 days. Our data clearly demonstrate the onset dependence of postconditioning elicited by a subconvulsant dose of kainate treatment in global ischemia, with restoration of the structural plasticity and hippocampal function.

1. Introduction

The cerebral consequences of global hypoperfusion can range from slight changes in synaptic plasticity (Marosi et al., 2009), as seen after 2-vessel occlusion (2VO) (Farkas et al., 2007), to extensive (70-75%) cell death in vulnerable brain areas, including the hippocampal CA1 region (Burda et al., 2006; Sas et al., 2008), after 4-vessel-occlusion (4VO) (Pulsinelli et al., 1983). The cortex and hippocampus are highly sensitive to hypoperfusion (Pulsinelli et al., 1983). 2VO (only the bilateral common carotid arteries are clamped) reduces the blood flow to one-third in rats (Farkas et al., 2007), with only a slight malfunction in synaptic plasticity (Li et al., 2006; Marosi et al., 2009), and no detectable morphological changes at a cellular level (Fluoro JadeB, S-100 and Nissl staining) (Marosi et al., 2009). Through glutamatergic synapses, long-term potentiation (LTP) involves cellular mechanisms of learning and memory (Bliss and Collingridge, 1993; Martin et al., 2000), expressed mainly in dendritic spines in the hippocampus and cortex. Any LTP malfunction in the ischemic hippocampus is paralleled by changes in dendritic spine density through remodeling of spines. Can appropriate intervention normalize both spine density and LTP function?

Numerous neuroprotectants are tested in ischemic animal models with the aims of infarction reductions and neurological status improvement, but few reach the clinic (Hoyte et al., 2004). Glutamate scavenging by oxaloacetate to reduce excitotoxicity in the ischemic brain appears attractive (Teichberg et al., 2009), preserving neurons (Zlotnik et al., 2008) and synaptic plasticity (Marosi et al., 2009; Nagy et al., 2010). Another endogenous strategy is ischemic postconditioning, which has undergone a renaissance during the last 5-10 years of intensive research (reviewed by (Zhao, 2009), after its first description in heart physiology (Sewell et al., 1955). Ischemic postconditioning (Na et al., 1996) in the brain involves several repeated cycles of brief reperfusion and reocclusion of the common carotid artery after the onset of full reperfusion, and is an effective technique (Burda et al., 2006; Wang et al., 2008). Postconditioning can also be elicited by the single

post-ischemic administration of pharmaceuticals (second pathophysiological stress), bradykinin, 3-nitropropionic acid and norepinephrine all successfully preserving the principal cells in the hippocampus after 4VO (Burda et al., 2006; Burda et al., 2005; Danielisova et al., 2008; Danielisova et al., 2009).

Kainic acid (KA), obtained first from the red alga *Digenea simplex*, is a potent agonist of the kainate class of ionotropic glutamatergic receptors, and hence a potent neuroexcitant through excitotoxicity (Olney et al., 1974). High doses (at least 10 mg/kg) produce lesions in vulnerable brain areas (Nadler et al., 1978; Sperk et al., 1983), while physiological activation of kainate receptor helps the central nervous system develop normally, building and maintaining synapses and plasticity (Bortolotto et al., 1999).

We have now tested whether KA treatment is suitable for pharmacological postconditioning after incomplete global ischemia. In particular, we determined the optimal application timepoint (onset dependence) for postconditioning during the 72-h post-ischemic period after 2VO, the effectiveness of postconditioning over a longer post-ischemic period (10 days), and the changes in spine density in the hippocampal CA1 region, in the basal glutamatergic synaptic transmission and in the LTP.

2. Methods

2.1. *Animals and housing conditions*

Male Charles-River rats (200-250 g) of the Wistar strain (N=77) were used. The animals were housed individually in standard plastic cages with free access to food and water. The house of the rats was light-controlled in 12-h cycles and the animals were kept under conditions of constant room temperature (22 ± 1 °C) and humidity. Every effort was made to minimize the number of animals used and their suffering. The principles of animal care (NIH publication No. 85-23), and the protocol for animal care approved by both the Hungarian Health Committee (1998) and the European Communities Council Directive (86/609/EEC) were followed.

2.2. *Surgical and experimental procedures, drug application*

All of the surgical procedures were carried out under deep anesthesia. Before and during the induction of transient cerebral ischemia, the rats were anesthetized with an intraperitoneal (i.p.) injection of 4% chloral-hydrate (Sigma, Germany). The body temperature of the animals was maintained at 37 ± 0.5 °C throughout the procedure with a feedback-regulated heating lamp. The common carotid arteries were exposed via a ventral cervical incision and separated from the vagal nerves. Both arteries were reversibly clamped with non-traumatic aneurysm clips (Aesculap, B. Braun Medical Ltd, Hungary) for a 30-min period (Marosi et al., 2009). The carotid artery blood flow was then reperfused by releasing the clips following the 30-min global hypoperfusion. The surgery was accompanied by a ~70% survival rate following 2VO. The sham-operated control rats underwent the same procedure, but without bilateral common carotid artery occlusion. In the postconditioned groups the reperfusion period was always 72 h (survival time), with the exception of the 10-day-follow-up group (2VO 48KA 10d). For KA treatment (Tocris, UK) the original 8 mg/kg dose (Burda et al., 2009) was lowered to 5 mg/kg i.p. so as to obtain the maximal postconditioning effects (a result of a personal communication with J. Burda). To

gain information concerning the effects of low-dose KA treatment on the normal EEG activity, 3 control and 3 2VO animals were anesthetized (4% chloral-hydrate injection, i.p.) and conventional EEG recordings were made from the frontal and parietal cortices of both hemispheres before and during 5 mg/kg KA treatment, and then for an additional 100 min. No signs of epileptiform activity were observed during these experiments (data not shown). Our observations are in agreement with the experimental data of Hellier et al. 5 mg/kg KA produced seizure activity only if it was administered repeatedly. In the groups in which KA was applied alone, no deaths occurred and no signs of epileptiform activity were detected in the behavior. In the postconditioned groups, the survival rate was lower, and it was worst (~50%) when 2VO immediately preceded KA. No signs of epileptiform activity were observed in the behavior of the animals in the postconditioned groups. These data are presented in Table 1.

2.3. In vitro electrophysiology

With the exception of the 12 animals sacrificed for Golgi-Cox staining, taken randomly from their experimental groups, all animals in each experimental group underwent electrophysiological recordings after different survival times (see Table 1). Coronal slices (400 μm), prepared from the middle part of their hippocampi with a vibratome (Campden Instruments, UK), were kept in an ice-cold artificial cerebrospinal solution (ACSF) composed of (in mM): 130 NaCl, 3.5 KCl, 1 NaH_2PO_4 , 24 NaHCO_3 , 1 CaCl_2 , 3 MgSO_4 and 10 D-glucose (all from Sigma, Germany), saturated with 95% O_2 + 5% CO_2 . The slices then were transferred to a Haas recording chamber and incubated at room temperature for 1 h to allow the slices to recover in the solution used for recording (differing only in that it contained 3 mM CaCl_2 and 1.5 mM MgSO_4). The flow rate of the recording solution was 1.5–2 ml/min and the experiments were performed at 34 °C. In order to obtain orthodromic stimulation of the Schaffer collateral/commissural pathway, the stimulating electrode (either a micropipette pulled from a theta glass or a bipolar concentric stainless steel electrode developed by Neuronelektrod Ltd,

Hungary) was placed in the stratum radiatum between the CA1 and CA2 regions. The stimulus intensity was adjusted to between 30 and 60 μA (constant current, 0.2-ms pulses delivered at 0.05 Hz) to evoke the half-maximal response.

Field excitatory postsynaptic potentials (fEPSPs) were recorded with glass micropipettes with a resistance of 1.5 M Ω filled with ACSF. The recordings were amplified with a neutralized, high-input-impedance preamplifier and filtered (1 Hz-3 kHz). The fEPSPs were digitized (AIF-03, Experimetria Ltd. Hungary), acquired at a sampling rate of 10 kHz, saved to a PC and analyzed off-line with Origin 6.0 software (OriginLab Corporation, USA).

The fEPSPs were monitored for 40-60 min until the amplitudes were generally stable, and their mean value was determined as the 10-min-long baseline. LTP of the Schaffer collateral-CA1 synaptic response was induced by high-frequency stimulation (HFS, 0.2-ms pulses delivered at 100 Hz for 5 s) at 100% intensity of the test stimulus (Marosi et al., 2009; Sas et al., 2008). After the HFS, the fEPSPs were recorded for at least a further 60 min. The basal glutamergic synaptic properties were also tested by means of fEPSP recordings; in IO (input-output) graphs the fEPSP amplitudes are expressed against gradually increasing stimulating currents (0-100 μA , in 5- μA steps) in certain groups. Each slice was subjected to only one measurement (Marosi et al., 2009; Sas et al., 2008); there were 96 recordings for LTP measurements and 18 recordings for IO curves (see Table 1).

2.4. Golgi-Cox staining technique

In order to determine the number of apical dendritic spines in the hippocampal CA1 region in the control, 2VO and KA-postconditioned animals, we used the Golgi techniques (Gibb and Kolb, 1998; Glaser and Van der Loos, 1981). The animals were decapitated, and the brains were rapidly, but carefully removed from the skull. Approximately 3-mm sagittal blocks including the whole hippocampi were placed in Golgi-Cox solution in the dark for 10 days at

room temperature (22-23 °C). This solution is prepared from 3 stock solutions: a 5% solution of potassium dichromate in distilled water (DW), a 5% solution of potassium chromate in DW, and a 5% solution of mercuric chloride in DW (all from Sigma, Germany). The components were diluted and mixed as described by Glaser and Van Der Loos (1981). The blocks were replaced in new solution every 2-3 days. When the impregnation was complete, the blocks were placed into 30% sucrose at 4 °C for at least 2-3 days until they sank. 100- μ m sagittal sections were cut with a vibratome (Leica VT1000S, Leica Biosystems Nussloch GmbH, Germany). Sections were collected in 6% sucrose and mounted on gelatinized (2%) glass slides. The sections were kept wet throughout the sectioning. To ensure that the sections in all certainty stuck on the slides, moist blotting paper was placed onto the sections and downward pressure was exerted via another slide. The blotted slides were stored in a humidity chamber overnight, after which they were placed in a glass staining tray and processed as described by Gibb and Kolb (1997). After the staining and dehydration, the slides were coverslipped with Fluoromount to prevent curling of the sections.

2.5. Quantitative analysis

The clear Golgi sections were studied by light microscopic stereology, using oil immersion objectives. In this study, a total of 20 cells from each of 3 animals were used from each of the 4 experimental groups (control, 2VO, 2VO 48h KA and 2VO 48h KA 10d). Pyramidal neurons from the hippocampal CA1 subfield near the subiculum were studied; in this region, the CA3-proceeding Schaffer collaterals synapse with dendritic spines from the dendritic arbor of pyramidal cells, mostly in the stratum radiatum (Amaral and Witter, 1989). The spine density of the proximal apical dendrite area (100-200 μ m from the soma) was analyzed. The second-order dendritic ramifications are particularly sensitive to plastic changes (Gould et al., 1990; Pyapali and Turner, 1994). One segment 100 μ m in length from a second-order oblique dendrite protruding from its parent apical dendrite was chosen in each examined neuron, for

spine density quantification. The dendrites were selected under a 100X oil-immersion lens and the images (600X) of these apical dendrites were captured through a DP70 camera connected to a light microscope (BX51, Olympus, Tokyo, Japan) and a computer. Serial images were made from each dendrite in the whole of the analyzed segment. The captured multiple photomicrographs from one dendrite were then stacked into one file. To stack the images and determine the density of spines, ImageJ 1.44d software (National Institute of Health, Bethesda, USA) was used. The whole process of measurement of the spine number and density was performed separately by three experimenters blind to the analysis. The total numbers of detected spines are presented in Table 1.

2.6. Statistics

For the statistical analysis of IO curves, the Kruskal-Wallis test was chosen (Fig. 2). For LTP measurements, the fEPSP amplitudes were normalized to the means of the 10-min pre-HFS control data. As normal distribution of the data could not be presumed and the Levene test did not demonstrate equality of variances, a nonparametric test on two independent samples was chosen for the statistical analysis of the LTP data (Mann-Whitney U-test) in Fig. 4A. A P value of ≤ 0.05 was considered significant. SPSS10.0 for Windows software (SPSS Inc., Chicago, IL, USA) was utilized. In the spine density analysis of the data for the various experimental groups, parametric t-tests for two independent samples were used (Fig. 4B).

3. Results

3.1. *In vitro* electrophysiological recordings

The HFS of Schaffer collaterals resulted in a $143.55\% \pm 2.60$ increase in fEPSP amplitudes in the controls (Fig. 1A), which remained stable throughout the 1-h recording period. A 30-min 2VO, 72 h prior to LTP induction, resulted in a lesser increase ($115.75\% \pm 2.19$) in amplitudes (Fig. 1A), with a constant decay in time. These data are in good accordance with our previous results (Marosi et al., 2009). Figure 1B depicts the changes in LTP inducibility and height as a consequence of postconditioning at 0, 24, 48 h and 10 d post-ischemia. KA postconditioning applied 24 h after ischemia (2VO 24h KA) had a worsening effect on the LTP function ($112.87\% \pm 5.51$). When KA postconditioning was applied immediately after 2VO (2VO 0h KA), the fEPSP amplitude increase was least, at only $111.84\% \pm 1.98$, and in this group no post-tetanic potentiation was detected. In both these early postconditioned groups, the amplitudes had reached the pre-HFS levels by the end of the 1-h follow-up period. An LTP function recovery was seen only when KA treatment was applied 48 h after ischemia (delayed postconditioning), clearly demonstrating the onset dependence of the postconditioning (Fig. 1B). In the 2VO 48h KA group, the post-HFS values were close to the control levels ($138.95\% \pm 2.38$) throughout the 1-h follow-up period (Fig. 1B). Since KA treatment alone had a worsening effect on the LTP function at 72 h (Fig. 1D), it was important to test whether the protective effect of KA postconditioning was persistent for a longer time period. In the 2VO 48h 10d group, LTP measurements were made 8 days after postconditioning. As shown in Fig. 1B, the LTP function still existed in the ischemic brain, and the fEPSPs amplitudes were stable throughout the whole of the follow-up period ($128.18\% \pm 3.38$).

In order to exclude the possibility of spontaneous recovery in synaptic plasticity over time, a systematic check was carried out on a day-by-day basis from the first (24 h) to the fourth day (96 h) after 2VO. As indicated in Fig. 1C, stable LTP could not be induced at any examined

timepoint (here the data of the 2VO 72h group are taken from Fig. 1A to facilitate comparison). There was no difference in the overall decrease in synaptic plasticity between the 2VO 48h, 2VO 72h and 2VO 96h groups. However, the 2VO 96h group exhibited a more pronounced decrease in fEPSP amplitudes, which reached the pre-HFS levels during the first 15 min. The reason for this is unclear. The 2VO 24h group displayed the most severe decrease in synaptic plasticity: the fEPSP amplitudes reached the pre-HFS levels after only 30 min. We also tested the temporal effects of single KA treatments at different survival times (Fig. 1D). In the KA 24h group, the amplitudes were around the control values (140-145%) during the first 10-15 min after HFS, but they later began to decrease steadily in time, revealing disturbances in synaptic plasticity. No LTP was observed in this group. In the KA 48h and KA 72h groups, a more severe malfunction was detected, the fEPSP amplitudes reaching the pre-HFS values by 30 min, with a steady decay throughout the 1-h follow-up period. The electrophysiological data presented in Figs. 1A and 1B are summarized in Fig. 4A, where the post-HFS fEPSP amplitudes are box-plotted.

The basal synaptic properties were also tested to evaluate how the various experimental interventions impaired the Schaffer collateral-CA1 synaptic transmission. IO curves were established by plotting the fEPSP amplitude against various intensities of the test pulse, ranging from 0 to 100 μA in 5 μA steps. It is clearly seen in Fig. 2 that there was a marked difference between the IO curves of the control and the 2VO groups, implying that the basal functions of the pyramidal cells and synapses were affected by the 2VO. However, the KA-postconditioned group demonstrated normal IO values (Fig. 2).

3.2. Golgi-Cox impregnation

The Golgi-Cox staining method labeled a subset of neurons in all parts of the hippocampus. There were no differences in overall staining between the 4 studied groups: usually 20-25 fully impregnated CA1 pyramidal cells could be detected per slice. No signs of irregular dendritic swellings were detected in the histologically examined ischemic groups (Fig. 3).

Although different types of spines (e.g. thin, stubby, mushroom, wide, double and branched) (Gonzalez-Burgos et al., 2007) were always clearly seen, our study was restricted merely to the determination of spine density (Fig. 3). In the control group, the spine density was $182.07 \pm 3.25 / 100 \mu\text{m}$ (Figs. 3 and 4B), a value similar to that detected by others ($116.7 \text{ spines} / 50 \mu\text{m}$) (Gonzalez-Burgos et al., 2007); the difference is probably due to slight variations in staining technique. In the 2VO animals, the mild, incomplete ischemia resulted in a marked decrease in spine density ($84.95 \pm 2.23 / 100 \mu\text{m}$) (Fig. 4B). With the aid of prior electrophysiological experiments (LTP measurements), the ideal timepoint for KA-postconditioning was determined, which proved to be 48 h after 2VO. For examination of the consequences of postconditioning as concerns the spine density, measurements were subsequently made only at this single timepoint. As a result of KA-postconditioning, by 48 h after ischemia the spine density had virtually recovered to the control level ($171.96 \pm 2.58 / 100 \mu\text{m}$ vs $182.07 \pm 3.25 / 100 \mu\text{m}$) (Fig. 4B) in parallel with restoration of the synaptic plasticity. The elevated spine density level persisted 10 days after 2VO ($128.25 \pm 1.41 / 100 \mu\text{m}$, Fig. 4B).

4. Discussion

A great wealth of experimental data strengthens the assumption that remodeling and stabilization of the spine synapses of the pyramidal cells are features of the cellular mechanism by which LTP and memory processes occur (Kandel, 2004; Kasai et al., 2003). Pyramidal cells receive almost all their glutamatergic excitatory input through their spine synapses, and changes in spine density and spine synapses can therefore greatly influence the excitability and responsiveness of these cells. During the development of LTP, profound morphological changes in spine morphology occur (Matsuzaki et al., 2004). As a consequence of the changes in dendritic spine morphology, the conduction of postsynaptic potentials (Araya et al., 2006) and the diffusion signal proteins (Hugel et al., 2009) between spines and dendritic shafts undergo alteration. Under pathological circumstances, as in the ischemic brain, the longer dendritic spines (Brown et al., 2008) at the network level can isolate the soma biochemically and electrically from the local excitotoxic effects, preventing the initiation of apoptotic processes. Spines can buffer the elevated Ca^{2+} levels during excitotoxicity (Majewska et al., 2000). The dendritic plasticity in the post-ischemic brain is the early adaptive response by which affected neurons can fend off pathological processes, such as excitotoxicity, increased Ca^{2+} influx (Bano and Nicotera, 2007), and spontaneous spreading depolarizations (Risher et al., 2010). However, incomplete forebrain ischemia, e.g. that caused by 2VO, does not lead to cell death (Marosi et al., 2009); thus, the change in dendritic plasticity alone may be responsible for the reduction of LTP in the CA1 region.

KA is a neuroexcitatory and neurotoxic substance (Olney et al., 1974); its administration locally or systemically induces widespread neurodegeneration in the brain, the most intense being in the hippocampus (Sperk et al., 1985), accompanied by the loss of the principal cells in the CA1, CA3 and hilar regions (Nadler et al., 1978; Sperk et al., 1983). However, the administration of KA also triggers neurogenesis in the dentate gyrus (Gray and Sundstrom, 1998; Jaako et al.,

2009), resulting in aberrant synaptic contacts and neural networks, affecting the hippocampal functioning (Scharfman et al., 2000). Rats receiving KA display a status epilepticus, with recurrent seizures and severe spatial memory and learning impairments (Mikati et al., 1994). Administration of KA at a convulsant dose (e.g. 10 or 50 mg/kg) also induces the expression of proto-oncoproteins, such as Fra, c-Fos and Jun in the rat hippocampus (Le Gal La Salle, 1988; Popovici et al., 1990). These changes in local gene expressions have been suggested to be related to hippocampal neural death and survival after KA-induced seizure activity (Xia et al., 1995). KA also induces the activities of mitogen-activated kinases (MAPKs): extracellular signal-regulated protein kinase (ERK), c-Jun N terminal kinase (JNK) and p38 MAPK (p38) (DeCoster et al., 1998). JNK targets several apoptosis-related proteins, including c-Jun, p53 and Bcl-2, and can activate the caspase cascade response (Haydar et al., 1999). Whether or not a cell survives or undergoes apoptosis is dependent on the activation of these MAPKs (Xia et al., 1995). Calcium/calmodulin-dependent kinase II (CAMKII) regulates a number of cellular functions related to elevated intracellular Ca^{2+} concentrations (Hudmon and Schulman, 2002). CaMKII plays a key role in mediating cell death following hypoxia/hypoglycemia (Hajimohammadreza et al., 1995), and displays an elevated phosphorylated level following KA treatment (Lee et al., 2003). The activation of MAPKs and CAMKII after KA treatment can be blocked by cycloheximide pretreatment, suggesting the need for newly synthesized proteins (Lee et al., 2003). It should be noted that one or more brief, non-harmful seizure episodes (seizure preconditioning) evoked by either low-dose KA (Tanaka et al., 2010) or domoic acid (a KA analog (Hesp et al., 2007)) can dramatically reduce hippocampal damage when given prior to status epilepticus, via a G-protein coupled pathway (PKA and CAMKII). In our experiments, the applied KA concentration was too low to induce epileptiform activity; thus, our results appear to be related not to the pathological over activation, but rather to the physiological activation of kainate receptors.

The kainate type of glutamate receptors also play pivotal roles in the modulation of glutamatergic synaptic transmission and plasticity in the brain (Bortolotto et al., 1999). Kainate receptors have a specific role in synaptic development (Marchal and Mulle, 2004; Tashiro et al., 2003), and can regulate the motility of the filopodia at hippocampal mossy fibers (Tashiro et al., 2003), an early step during synapse formation. The CA1 interneurons containing kainate receptors depolarize the cell soma through ionotropic action (Cossart et al., 1998; Rodriguez-Moreno et al., 2000), while another population of kainate receptors depress evoked inhibitory postsynaptic currents and GABA release through a G-protein-dependent mechanism (Christensen et al., 2004; Clarke et al., 1997; Mulle et al., 2000; Rodriguez-Moreno et al., 2000). In the CA1, the pharmacological activation of Glu5R-containing kainate receptors strongly depresses glutamatergic transmission (Clarke and Collingridge, 2002; Vignes et al., 1998) through the action of the presynaptic receptors via G-protein-coupled signaling (Frerking et al., 2001; Lauri et al., 2006). Moreover, the sustained activation of GluR5 receptors causes a specific and lasting increase in the number of glutamatergic synapses in area CA1, dependent on PKC activation (Vesikansa et al., 2007). It is possible that a similar G-protein-dependent signaling mechanism underlies the kainate receptor-dependent synaptic stabilization in both the CA3 and CA1 areas. However, these receptors are endogenously activated in the developing, but not in the adult hippocampus (Lauri et al., 2006). Our results demonstrate that, while KA treatment hampers synaptic plasticity in the adult hippocampus under normal circumstances, in the ischemic hippocampus KA is able to restore the injured synaptic plasticity within its therapeutic window through restoration of the control level of spine density and normal-like glutamatergic synaptic transmission.

Ischemic postconditioning *per se* cannot be studied separately from the ischemic event, and its underlying mechanisms, at least in part, are common with those seen in connection with kainate receptor activation. Postconditioning enhances Akt phosphorylation and Akt activity (Gao et al., 2008). The MAPK pathways (ERK1/2, p38 and JNK) are also closely related to ischemic

injury and neuronal survival (Sawe et al., 2008). The ERK1/2 phosphorylation level, transiently elevated after reperfusion, is reduced during postconditioning (Gao et al., 2008) in the penumbra. Postconditioning also involves PKC pathways: δ PKC activity usually leads to cell death (Shimohata et al., 2007b), whereas ϵ PKC promotes neuronal survival (Shimohata et al., 2007a). Postconditioning inhibits the increase in level of the cleaved form of δ PKC and improves ϵ PKC phosphorylation (Gao et al., 2008). *De novo* synthesized proteins, e.g. heat shock proteins (Liu et al., 1993), Bcl-2 (Shimazaki et al., 1994), superoxide dismutase (MnSOD) (Bordet et al., 2000) and pro-survival inhibitor-of-apoptosis (Tanaka et al., 2004), contribute to the acquisition of delayed ischemic tolerance. Delayed postconditioning is based on the synthesis of proteins and is able to reverse the ischemia-induced apoptosis-like process of delayed neuronal death 48 h after ischemia.

Our experiments have demonstrated that KA treatment induces reappearance of the normal spine density in the adult hippocampal CA1 subfield after incomplete global ischemia, which results in a marked restoration of experimental LTP induction. The beneficial effects of KA postconditioning on the hippocampal function are long-lasting and can be detected as late as 8 days after KA treatment. However, KA proved most effective on the second post-ischemic day, revealing a strict onset-dependent feature of its application. This is in agreement with previous studies, which involved either ischemic or pharmacological postconditioning with therapeutic windows ranging from 6 h (Ren et al., 2008) up to 2 days (Burda et al., 2006; Burda et al., 2005; Danielisova et al., 2008; Danielisova et al., 2009). Future experiments are needed to clarify the exact mechanism of the KA-mediated protection reported here.

Acknowledgments

This work was supported by TÁMOP-4.2.1/B-09/1/KONV-2010-0005, OTKA K 75628 and GVOP-3.2.1-2004-04-0357/3.0. T.F. was a Bolyai Fellow of the Hungarian Academy of Sciences.

References

- Amaral, D. G., Witter, M. P., 1989. The three-dimensional organization of the hippocampal formation: a review of anatomical data. *Neuroscience* 31, 571-591.
- Araya, R., Jiang, J., Eisenthal, K. B., Yuste, R., 2006. The spine neck filters membrane potentials. *Proc Natl Acad Sci U S A* 103, 17961-17966.
- Bano, D., Nicotera, P., 2007. Ca²⁺ signals and neuronal death in brain ischemia. *Stroke* 38, 674-676.
- Bliss, T. V., Collingridge, G. L., 1993. A synaptic model of memory: long-term potentiation in the hippocampus. *Nature* 361, 31-39.
- Bordet, R., Deplanque, D., Maboudou, P., Puisieux, F., Pu, Q., Robin, E., Martin, A., Bastide, M., Leys, D., Lhermitte, M., Dupuis, B., 2000. Increase in endogenous brain superoxide dismutase as a potential mechanism of lipopolysaccharide-induced brain ischemic tolerance. *J Cereb Blood Flow Metab* 20, 1190-1196.
- Bortolotto, Z. A., Clarke, V. R., Delany, C. M., Parry, M. C., Smolders, I., Vignes, M., Ho, K. H., Miu, P., Brinton, B. T., Fantaske, R., Ogden, A., Gates, M., Ornstein, P. L., Lodge, D., Bleakman, D., Collingridge, G. L., 1999. Kainate receptors are involved in synaptic plasticity. *Nature* 402, 297-301.
- Brown, C. E., Wong, C., Murphy, T. H., 2008. Rapid morphologic plasticity of peri-infarct dendritic spines after focal ischemic stroke. *Stroke* 39, 1286-1291.
- Burda, J., Danielisova, V., Nemethova, M., Gottlieb, M., Kravcukova, P., Domorakova, I., Mechirova, E., Burda, R., 2009. Postconditioning and anticonditioning: possibilities to interfere to evoked apoptosis. *Cell Mol Neurobiol* 29, 821-825.
- Burda, J., Danielisova, V., Nemethova, M., Gottlieb, M., Matiasova, M., Domorakova, I., Mechirova, E., Ferikova, M., Salinas, M., Burda, R., 2006. Delayed postconditioning initiates additive mechanism necessary for survival of selectively vulnerable neurons after transient ischemia in rat brain. *Cell Mol Neurobiol* 26, 1141-1151.
- Burda, J., Matiasova, M., Gottlieb, M., Danielisova, V., Nemethova, M., Garcia, L., Salinas, M., Burda, R., 2005. Evidence for a role of second pathophysiological stress in prevention of delayed neuronal death in the hippocampal CA1 region. *Neurochem Res* 30, 1397-1405.
- Christensen, J. K., Paternain, A. V., Selak, S., Ahring, P. K., Lerma, J., 2004. A mosaic of functional kainate receptors in hippocampal interneurons. *J Neurosci* 24, 8986-8993.
- Clarke, V. R., Ballyk, B. A., Hoo, K. H., Mandelzys, A., Pellizzari, A., Bath, C. P., Thomas, J., Sharpe, E. F., Davies, C. H., Ornstein, P. L., Schoepp, D. D., Kamboj, R. K., Collingridge, G. L., Lodge, D., Bleakman, D., 1997. A hippocampal GluR5 kainate receptor regulating inhibitory synaptic transmission. *Nature* 389, 599-603.
- Clarke, V. R., Collingridge, G. L., 2002. Characterisation of the effects of ATPA, a GLU(K5) receptor selective agonist, on excitatory synaptic transmission in area CA1 of rat hippocampal slices. *Neuropharmacology* 42, 889-902.
- Cossart, R., Esclapez, M., Hirsch, J. C., Bernard, C., Ben-Ari, Y., 1998. GluR5 kainate receptor activation in interneurons increases tonic inhibition of pyramidal cells. *Nat Neurosci* 1, 470-478.
- Danielisova, V., Gottlieb, M., Nemethova, M., Burda, J., 2008. Effects of bradykinin postconditioning on endogenous antioxidant enzyme activity after transient forebrain ischemia in rat. *Neurochem Res* 33, 1057-1064.
- Danielisova, V., Gottlieb, M., Nemethova, M., Kravcukova, P., Domorakova, I., Mechirova, E., Burda, J., 2009. Bradykinin postconditioning protects pyramidal CA1 neurons against delayed neuronal death in rat hippocampus. *Cell Mol Neurobiol* 29, 871-878.
- DeCoster, M. A., Mukherjee, P. K., Davis, R. J., Bazan, N. G., 1998. Platelet-activating factor is a downstream messenger of kainate-induced activation of mitogen-activated protein kinases in primary hippocampal neurons. *J Neurosci Res* 53, 297-303.

- Farkas, E., Luiten, P. G., Bari, F., 2007. Permanent, bilateral common carotid artery occlusion in the rat: a model for chronic cerebral hypoperfusion-related neurodegenerative diseases. *Brain Res Rev* 54, 162-180.
- Frerking, M., Schmitz, D., Zhou, Q., Johansen, J., Nicoll, R. A., 2001. Kainate receptors depress excitatory synaptic transmission at CA3-->CA1 synapses in the hippocampus via a direct presynaptic action. *J Neurosci* 21, 2958-2966.
- Gao, X., Zhang, H., Takahashi, T., Hsieh, J., Liao, J., Steinberg, G. K., Zhao, H., 2008. The Akt signaling pathway contributes to postconditioning's protection against stroke; the protection is associated with the MAPK and PKC pathways. *J Neurochem* 105, 943-955.
- Gibb, R., Kolb, B., 1998. A method for vibratome sectioning of Golgi-Cox stained whole rat brain. *J Neurosci Methods* 79, 1-4.
- Glaser, E. M., Van der Loos, H., 1981. Analysis of thick brain sections by obverse-reverse computer microscopy: application of a new, high clarity Golgi-Nissl stain. *J Neurosci Methods* 4, 117-125.
- Gonzalez-Burgos, I., Letechipia-Vallejo, G., Lopez-Loeza, E., Morali, G., Cervantes, M., 2007. Long-term study of dendritic spines from hippocampal CA1 pyramidal cells, after neuroprotective melatonin treatment following global cerebral ischemia in rats. *Neurosci Lett* 423, 162-166.
- Gould, E., Woolley, C. S., Frankfurt, M., McEwen, B. S., 1990. Gonadal steroids regulate dendritic spine density in hippocampal pyramidal cells in adulthood. *J Neurosci* 10, 1286-1291.
- Gray, W. P., Sundstrom, L. E., 1998. Kainic acid increases the proliferation of granule cell progenitors in the dentate gyrus of the adult rat. *Brain Res* 790, 52-59.
- Hajimohammadreza, I., Probert, A. W., Coughenour, L. L., Borosky, S. A., Marcoux, F. W., Boxer, P. A., Wang, K. K., 1995. A specific inhibitor of calcium/calmodulin-dependent protein kinase-II provides neuroprotection against NMDA- and hypoxia/hypoglycemia-induced cell death. *J Neurosci* 15, 4093-4101.
- Haydar, T. F., Kuan, C. Y., Flavell, R. A., Rakic, P., 1999. The role of cell death in regulating the size and shape of the mammalian forebrain. *Cereb Cortex* 9, 621-626.
- Hellier, J. L., Patrylo, P. R., Dou, P., Nett, M., Rose, G. M., Dudek, F. E., 1999. Assessment of inhibition and epileptiform activity in the septal dentate gyrus of freely behaving rats during the first week after kainate treatment. *J Neurosci* 19, 10053-10064.
- Hesp, B. R., Clarkson, A. N., Sawant, P. M., Kerr, D. S., 2007. Domoic acid preconditioning and seizure induction in young and aged rats. *Epilepsy Res* 76, 103-112.
- Hoyte, L., Kaur, J., Buchan, A. M., 2004. Lost in translation: taking neuroprotection from animal models to clinical trials. *Exp Neurol* 188, 200-204.
- Hudmon, A., Schulman, H., 2002. Neuronal CA2+/calmodulin-dependent protein kinase II: the role of structure and autoregulation in cellular function. *Annu Rev Biochem* 71, 473-510.
- Hugel, S., Abegg, M., de Paola, V., Caroni, P., Gahwiler, B. H., McKinney, R. A., 2009. Dendritic spine morphology determines membrane-associated protein exchange between dendritic shafts and spine heads. *Cereb Cortex* 19, 697-702.
- Jaako, K., Zharkovsky, T., Zharkovsky, A., 2009. Effects of repeated citalopram treatment on kainic acid-induced neurogenesis in adult mouse hippocampus. *Brain Res* 1288, 18-28.
- Kandel, E. R., 2004. The molecular biology of memory storage: a dialog between genes and synapses. *Biosci Rep* 24, 475-522.
- Kasai, H., Matsuzaki, M., Noguchi, J., Yasumatsu, N., Nakahara, H., 2003. Structure-stability-function relationships of dendritic spines. *Trends Neurosci* 26, 360-368.
- Lauri, S. E., Vesikansa, A., Segerstrale, M., Collingridge, G. L., Isaac, J. T., Taira, T., 2006. Functional maturation of CA1 synapses involves activity-dependent loss of tonic kainate receptor-mediated inhibition of glutamate release. *Neuron* 50, 415-429.

- Le Gal La Salle, G., 1988. Long-lasting and sequential increase of c-fos oncoprotein expression in kainic acid-induced status epilepticus. *Neurosci Lett* 88, 127-130.
- Lee, H. K., Choi, S. S., Han, K. J., Han, E. J., Suh, H. W., 2003. Cycloheximide inhibits neurotoxic responses induced by kainic acid in mice. *Brain Res Bull* 61, 99-107.
- Li, Z., Zhou, R., Cui, S., Xie, G., Cai, W., Sokabe, M., Chen, L., 2006. Dehydroepiandrosterone sulfate prevents ischemia-induced impairment of long-term potentiation in rat hippocampal CA1 by up-regulating tyrosine phosphorylation of NMDA receptor. *Neuropharmacology* 51, 958-966.
- Liu, Y., Kato, H., Nakata, N., Kogure, K., 1993. Temporal profile of heat shock protein 70 synthesis in ischemic tolerance induced by preconditioning ischemia in rat hippocampus. *Neuroscience* 56, 921-927.
- Majewska, A., Brown, E., Ross, J., Yuste, R., 2000. Mechanisms of calcium decay kinetics in hippocampal spines: role of spine calcium pumps and calcium diffusion through the spine neck in biochemical compartmentalization. *J Neurosci* 20, 1722-1734.
- Marchal, C., Mulle, C., 2004. Postnatal maturation of mossy fibre excitatory transmission in mouse CA3 pyramidal cells: a potential role for kainate receptors. *J Physiol* 561, 27-37.
- Marosi, M., Fuzik, J., Nagy, D., Rakos, G., Kis, Z., Vecsei, L., Toldi, J., Ruban-Matuzani, A., Teichberg, V. I., Farkas, T., 2009. Oxaloacetate restores the long-term potentiation impaired in rat hippocampus CA1 region by 2-vessel occlusion. *Eur J Pharmacol* 604, 51-57.
- Martin, S. J., Grimwood, P. D., Morris, R. G., 2000. Synaptic plasticity and memory: an evaluation of the hypothesis. *Annu Rev Neurosci* 23, 649-711.
- Matsuzaki, M., Honkura, N., Ellis-Davies, G. C., Kasai, H., 2004. Structural basis of long-term potentiation in single dendritic spines. *Nature* 429, 761-766.
- Mikati, M. A., Holmes, G. L., Chronopoulos, A., Hyde, P., Thurber, S., Gatt, A., Liu, Z., Werner, S., Stafstrom, C. E., 1994. Phenobarbital modifies seizure-related brain injury in the developing brain. *Ann Neurol* 36, 425-433.
- Mulle, C., Sailer, A., Swanson, G. T., Brana, C., O'Gorman, S., Bettler, B., Heinemann, S. F., 2000. Subunit composition of kainate receptors in hippocampal interneurons. *Neuron* 28, 475-484.
- Na, H. S., Kim, Y. I., Yoon, Y. W., Han, H. C., Nahm, S. H., Hong, S. K., 1996. Ventricular premature beat-driven intermittent restoration of coronary blood flow reduces the incidence of reperfusion-induced ventricular fibrillation in a cat model of regional ischemia. *Am Heart J* 132, 78-83.
- Nadler, J. V., Perry, B. W., Cotman, C. W., 1978. Intraventricular kainic acid preferentially destroys hippocampal pyramidal cells. *Nature* 271, 676-677.
- Nagy, D., Knapp, L., Marosi, M., Farkas, T., Kis, Z., Vecsei, L., Teichberg, V. I., Toldi, J., 2010. Effects of Blood Glutamate Scavenging on Cortical Evoked Potentials. *Cell Mol Neurobiol*, PMID: 20607387 [Epub ahead of print].
- Olney, J. W., Rhee, V., Ho, O. L., 1974. Kainic acid: a powerful neurotoxic analogue of glutamate. *Brain Res* 77, 507-512.
- Popovici, T., Represa, A., Crepel, V., Barbin, G., Beaudoin, M., Ben-Ari, Y., 1990. Effects of kainic acid-induced seizures and ischemia on c-fos-like proteins in rat brain. *Brain Res* 536, 183-194.
- Pulsinelli, W. A., Levy, D. E., Duffy, T. E., 1983. Cerebral blood flow in the four-vessel occlusion rat model. *Stroke* 14, 832-834.
- Pyapali, G. K., Turner, D. A., 1994. Denervation-induced dendritic alterations in CA1 pyramidal cells following kainic acid hippocampal lesions in rats. *Brain Res* 652, 279-290.
- Ren, C., Gao, X., Niu, G., Yan, Z., Chen, X., Zhao, H., 2008. Delayed postconditioning protects against focal ischemic brain injury in rats. *PLoS One* 3, e3851.

- Risher, W. C., Ard, D., Yuan, J., Kirov, S. A., 2010. Recurrent spontaneous spreading depolarizations facilitate acute dendritic injury in the ischemic penumbra. *J Neurosci* 30, 9859-9868.
- Rodriguez-Moreno, A., Lopez-Garcia, J. C., Lerma, J., 2000. Two populations of kainate receptors with separate signaling mechanisms in hippocampal interneurons. *Proc Natl Acad Sci U S A* 97, 1293-1298.
- Sas, K., Robotka, H., Rozsa, E., Agoston, M., Szenasi, G., Gigler, G., Marosi, M., Kis, Z., Farkas, T., Vecsei, L., Toldi, J., 2008. Kynurenine diminishes the ischemia-induced histological and electrophysiological deficits in the rat hippocampus. *Neurobiol Dis* 32, 302-308.
- Sawe, N., Steinberg, G., Zhao, H., 2008. Dual roles of the MAPK/ERK1/2 cell signaling pathway after stroke. *J Neurosci Res* 86, 1659-1669.
- Scharfman, H. E., Goodman, J. H., Sollas, A. L., 2000. Granule-like neurons at the hilar/CA3 border after status epilepticus and their synchrony with area CA3 pyramidal cells: functional implications of seizure-induced neurogenesis. *J Neurosci* 20, 6144-6158.
- Sewell, W. H., Koth, D. R., Huggins, C. E., 1955. Ventricular fibrillation in dogs after sudden return of flow to the coronary artery. *Surgery* 38, 1050-1053.
- Shimazaki, K., Ishida, A., Kawai, N., 1994. Increase in bcl-2 oncoprotein and the tolerance to ischemia-induced neuronal death in the gerbil hippocampus. *Neurosci Res* 20, 95-99.
- Shimohata, T., Zhao, H., Steinberg, G. K., 2007a. Epsilon PKC may contribute to the protective effect of hypothermia in a rat focal cerebral ischemia model. *Stroke* 38, 375-380.
- Shimohata, T., Zhao, H., Sung, J. H., Sun, G., Mochly-Rosen, D., Steinberg, G. K., 2007b. Suppression of deltaPKC activation after focal cerebral ischemia contributes to the protective effect of hypothermia. *J Cereb Blood Flow Metab* 27, 1463-1475.
- Sperk, G., Lassmann, H., Baran, H., Kish, S. J., Seitelberger, F., Hornykiewicz, O., 1983. Kainic acid induced seizures: neurochemical and histopathological changes. *Neuroscience* 10, 1301-1315.
- Sperk, G., Lassmann, H., Baran, H., Seitelberger, F., Hornykiewicz, O., 1985. Kainic acid-induced seizures: dose-relationship of behavioural, neurochemical and histopathological changes. *Brain Res* 338, 289-295.
- Tanaka, H., Yokota, H., Jover, T., Cappuccio, I., Calderone, A., Simionescu, M., Bennett, M. V., Zukin, R. S., 2004. Ischemic preconditioning: neuronal survival in the face of caspase-3 activation. *J Neurosci* 24, 2750-2759.
- Tanaka, K., Jimenez-Mateos, E. M., Matsushima, S., Taki, W., Henshall, D. C., 2010. Hippocampal damage after intra-amygdala kainic acid-induced status epilepticus and seizure preconditioning-mediated neuroprotection in SJL mice. *Epilepsy Res* 88, 151-161.
- Tashiro, A., Dunaevsky, A., Blazeski, R., Mason, C. A., Yuste, R., 2003. Bidirectional regulation of hippocampal mossy fiber filopodial motility by kainate receptors: a two-step model of synaptogenesis. *Neuron* 38, 773-784.
- Teichberg, V. I., Cohen-Kashi-Malina, K., Cooper, I., Zlotnik, A., 2009. Homeostasis of glutamate in brain fluids: an accelerated brain-to-blood efflux of excess glutamate is produced by blood glutamate scavenging and offers protection from neuropathologies. *Neuroscience* 158, 301-308.
- Vesikansa, A., Sallert, M., Taira, T., Lauri, S. E., 2007. Activation of kainate receptors controls the number of functional glutamatergic synapses in the area CA1 of rat hippocampus. *J Physiol* 583, 145-157.
- Vignes, M., Clarke, V. R., Parry, M. J., Bleakman, D., Lodge, D., Ornstein, P. L., Collingridge, G. L., 1998. The GluR5 subtype of kainate receptor regulates excitatory synaptic transmission in areas CA1 and CA3 of the rat hippocampus. *Neuropharmacology* 37, 1269-1277.

- Wang, J. Y., Shen, J., Gao, Q., Ye, Z. G., Yang, S. Y., Liang, H. W., Bruce, I. C., Luo, B. Y., Xia, Q., 2008. Ischemic postconditioning protects against global cerebral ischemia/reperfusion-induced injury in rats. *Stroke* 39, 983-990.
- Xia, Z., Dickens, M., Raingeaud, J., Davis, R. J., Greenberg, M. E., 1995. Opposing effects of ERK and JNK-p38 MAP kinases on apoptosis. *Science* 270, 1326-1331.
- Zhao, H., 2009. Ischemic postconditioning as a novel avenue to protect against brain injury after stroke. *J Cereb Blood Flow Metab* 29, 873-885.
- Zlotnik, A., Gurevich, B., Cherniavsky, E., Tkachov, S., Matuzani-Ruban, A., Leon, A., Shapira, Y., Teichberg, V. I., 2008. The contribution of the blood glutamate scavenging activity of pyruvate to its neuroprotective properties in a rat model of closed head injury. *Neurochem Res* 33, 1044-1050.

Legends to figures

Fig. 1.

Post-ischemic kainic acid (KA) administration prevents the global ischemia-induced long-term potentiation (LTP) impairment, but its protective power depends greatly on its application onset. **A:** In the control group, LTP was induced by high-frequency stimulation (HFS). The 2-vessel occlusion (2VO) group did not show stable LTP. **B:** As in the 2VO animals, in the early postconditioned groups (0 and 24 h after 2VO) no stable LTP was induced; on postconditioning with 48-h onset after 2VO, the LTP was near the control level. The beneficial effect of 48-h-onset postconditioning persisted 10 days after 2VO (2VO 48h 10d). **C:** LTP impairment caused by 2VO with different reperfusion intervals (24, 48, 72 and 96 h). Even after 96 h of reperfusion, no stable LTP was detected. There were no significant differences in post-HFS amplitudes between the 48, 72 and 96-h groups, but the LTP function was most severely decreased in the 2VO 24h group. **D:** LTP impairment caused by a single KA treatment with different survival times (24, 48 and 72 h). In the KA 24h group, HFS elicited a robust post-tetanic potentiation, and LTP was around the control level during the first third of the registration period, while in the last two-thirds it decayed steadily. In the KA 48h and KA 72h groups, no LTP was induced, and the fEPSP values fell even below the pre-HFS levels. Dashed lines (values from **A**) in **B**, **C** and **D** denote mean levels of LTP in control animals. Data points are means \pm S.E.M. of normalized amplitudes of fEPSPs.

Fig. 2.

Input-output curves for fEPSP amplitudes in the control group (\square), the 2-vessel occlusion (2VO) group (\bullet) and the group postconditioned with kainic acid (KA \blacklozenge), where the protective effect of postconditioning was maximal (2VO 48 h KA). Data points are means \pm S.E.M. of amplitudes of fEPSPs; N=6 for all groups presented here. Asterisks (*P<0.05 and **P<0.01) denote significant

differences between the data of the 2VO animals and those of the control and postconditioned (2VO 48h KA) groups (Kruskal-Wallis test).

Fig. 3.

Golgi-Cox staining revealing changes in spine density. **A:** A schematic drawing of a CA1 subfield pyramidal neuron, with a small box showing the site of spine density measurement. **B:** Photomicrographs of oblique dendritic segments from hippocampal CA1 pyramidal neurons of control, 2VO and 2VO-postconditioned animals treated with kainic acid (KA) at 48 h post-ischemia (2VO 48h KA). Note the fewer spines on the dendrites in the 2VO than in the control and the 2VO-postconditioned animals. Scale bar for **A:** 100 μm ; for **B:** 20 μm .

Fig. 4.

Changes in LTP and spine density. **A:** Post-ischemic administration of kainic acid (KA) prevents the long-term potentiation (LTP) impairment only within its therapeutic window. Box-plots demonstrate the means \pm S.E.M. of all amplitudes measured throughout the 60-min recording period following HFS. In the intact animals (control), HFS resulted in LTP with amplitudes remaining stable up to the end of the test period. In the 2-vessel occlusion (2VO) animals, however, after HFS the potentiation level was low during the 1-h follow-up period. KA postconditioning prevented the LTP impairment in an onset-dependent manner in the ischemic animals: in the 2VO 0h KA and 2VO 24h KA groups, the fEPSP amplitudes were around or even below the level of the 2VO group. Full protection was seen only when postconditioning was performed on the second post-ischemic day (2VO 48h KA). When the effect of 48-h postconditioning was measured 8 days after KA treatment (2VO 48KA 10d), a stable LTP could still be induced, though with a somewhat smaller amplitude. Asterisks denote significant differences from the control, and # denotes significant differences from the 2VO groups ($^{\#}P<0.05$, $^{**}P<0.001$, *** , $^{####}P<0.001$ Mann-Whitney U-test). **B:** Apical dendritic spine density

analysis revealed that the transient global ischemia induced a reduction in spine density 72 h after the ischemic insult in the 2VO group relative to the controls. In the postconditioned group (treated with KA) 48 h after the ischemic insult, 2VO 48h KA), the spine density was close to the control level. 10 days after the ischemic insult, the spine number elevation due to KA postconditioning was still detected (2VO 48h KA 10d group). In each experimental group, 60 dendrite shafts of 3 animals were analyzed. * $P < 0.05$, *** $P < 0.001$ significant differences from the control group; ### $P < 0.001$ significant differences from the 2VO group (parametric t-tests for two independent samples).

Table 1.

Summary of animal groups, interventions and morphological and electrophysiological observations.


Groups		<i>In vitro</i> electrophysiology		Golgi-Cox staining
		No. of animals / No. of mortalities	LTP recordings / IO curve measurements	No. of animals / No. of cells / No. of detected spines
Control		5/1	8/6	3/60/32,774
Postconditioned	2VO 48h KA	5/1	8/6	3/60/30,954
	2VO 48h KA 10d	6/2	8/0	3/60/23,085
	2VO 0h KA	8/4	8/0	0
	2VO 24h KA	6/1	8/0	0
2VO- only	2VO 24h	6/2	8/0	0
	2VO 48h	5/1	8/0	0
	2VO 72h	6/2	8/6	3/60/15,036
	2VO 96h	6/2	8/0	0
KA- only	KA 24h	4/0	8/0	0
	KA 48h	4/0	8/0	0
	KA 72h	4/0	8/0	0
Total		65/16	96/18	12/240/101,849

In the control and postconditioned groups, the reperfusion period was always 72 h, with the exception of the 2VO 48h KA 10d group, where the survival time was 10 d. In the 2VO-only groups, the survival times were 24, 48, 72 and 96 h. In the KA-only groups, the survival times were 24, 48 and 72 h.


Figure 1

[Click here to download high resolution image](#)

A



C





B**D**

Figure 2

[Click here to download high resolution image](#)





Figure 3
[Click here to download high resolution image](#)

A



control



2VO




2VO 48h KA




Figure 4

[Click here to download high resolution image](#)



B

Spine density (Golgi-Cox)



Highlights

- Postconditioning (PC) can be induced by kainic acid in the post-ischemic hippocampus.
- LTP returns to the control level in the CA1 subfield.
- The structural basis is the restoration of the spines of the CA1 pyramidal cells.
- The optimal timepoint for PC is post-ischemic day 2, with an onset dependence.
- The beneficial effects of PC are long-lasting (≥ 8 days).

Graphical abstract

Postconditioning elicited by a subconvulsant dose of kainic acid (KA) treatment is onset-dependent in global ischemia, resulting in restoration of the structural plasticity and hippocampal function (LTP).

