

## Gastroduodenal involvement in asymptomatic Crohn's disease patients in two areas of emerging disease: Asia and Eastern Europe

Klaudia Farkas<sup>1\*</sup>, Heyson Chan<sup>2\*</sup>, Mariann Rutka<sup>1</sup>, Zoltán Szepes<sup>1</sup>, Ferenc Nagy<sup>1</sup>, László Tizslavicz<sup>3</sup>, Tibor Nyári<sup>4</sup>, Whitney Tang<sup>2</sup>, Grace Wong<sup>2</sup>, Raymond Tang<sup>2</sup>, Angeline Lo<sup>2</sup>, Christina Cheung<sup>2</sup>, Sunny Wong<sup>2</sup>, Rashid Lui<sup>2</sup>, Tamás Molnár<sup>1\*</sup>, Siew C Ng<sup>2\*</sup>

<sup>1</sup>First Department of Medicine, University of Szeged, Szeged, Hungary

<sup>2</sup>Institute of Digestive Disease, Department of Medicine and Therapeutics, the Chinese University of Hong Kong, Hong Kong

<sup>3</sup>Department of Pathology, University of Szeged, Szeged, Hungary

<sup>4</sup>Department of Medical Physics and Informatics, University of Szeged

**\*Contributed equally**

**Short title: Upper GI Crohn's disease in Asia and Eastern Europe**

**Conflict of interests: none**

**Author contributions:** Study design, data collection, supervision of patient selection and manuscript preparation: Siew C Ng, Tamas Molnar, Klaudia Farkas, Heyson Chan; data collection: Klaudia Farkas, Heyson Chan, Mariann Rutka, Zoltán Szepes, Ferenc Nagy, László Tizslavicz, Whitney Tang, Grace Wong, Raymond Tang, Angeline Lo, Christina Cheung, Sunny Wong, Rashid Lui; data collection and manuscript preparation: Siew C Ng, Tamas Molnar, Klaudia Farkas, Heyson Chan; statistical analysis: Tibor Nyári. All authors have approved the final draft submitted.

**Abbreviations:**

**CARD15:** caspase recruitment domain-containing protein 15

**CD:** Crohn's disease

**GI:** gastrointestinal

**H.pylori:** Helicobacter pylori

**IBD:** inflammatory bowel disease

**NOD2:** nucleotide-binding oligomerization domain-containing protein 2

**OGD:** esophagealgastroduodenoscopy

**Co-Corresponding authors:**

**Prof Tamás Molnár** PhD, 1<sup>st</sup> Department of Medicine, University of Szeged, Szeged

H-6720, Korányi fasor 8, Szeged, Hungary

**E-mail:** [molnar.tamas@med.u-szeged.hu](mailto:molnar.tamas@med.u-szeged.hu)

**Tel.:** +36-62-545186, **Fax.:** +36-62-545185

**Prof Siew C Ng**, PhD, Department of Medicine and Therapeutics, Institute of Digestive Disease, The Chinese University of Hong Kong, Hong Kong.

**Email:** [siewchieng@cuhk.edu.hk](mailto:siewchieng@cuhk.edu.hk)

**Tel:** 852 2632 3996

**Fax:** 852 2637 3852

Accepted Manuscript

## Abstract

**Background:** The incidence of Crohn's disease (CD) is increasing in Asia and Eastern Europe. Limited studies have reported on the frequency of upper gastrointestinal (GI) involvement in patients with CD in non-Western countries. This prospective study compared the rate of macroscopic and microscopic upper GI manifestations and *Helicobacter pylori* (*H. pylori*) positivity in asymptomatic CD patients in Asia and Eastern Europe. **Methods:** Consecutive asymptomatic CD patients were prospectively recruited for upper GI endoscopy between 2013 and 2015 in Hong Kong and in Hungary. Endoscopy and biopsy findings were recorded and histology was performed to assess for *H. pylori* and microscopic signs characteristic for CD, using standardized diagnostic criteria. **Results:** One hundred and eighty CD patients (100 Hong Kong; 80 Hungary; 70.6% male; mean age, 38.5 years) and 189 controls (100 Hong Kong; 89 Hungary; 57.7% male; mean age 41 years) were included. Gastroduodenal involvement of CD was significantly higher in Hungary than in Hong Kong (16.5% vs 2.0%,  $p \leq 0.001$ ). *H. pylori* positivity was significantly higher in Hungarian than Chinese CD patients (13.9% vs. 4.0%,  $p \leq 0.001$ ). Granulomas were detected in 1% in Hong Kong and 7.6% in Hungary ( $p \leq 0.001$ ). Chinese CD subjects had a significantly lower *H. pylori* positivity compared with controls (6% vs. 15%;  $p \leq 0.001$ ). **Conclusion:** Upper GI CD was significantly higher in Eastern Europe compared with Asia. The detection of granuloma in Hungary was similar to the literature data, whereas focal gastritis was lower than expected in both cohorts.

**Key words:** Crohn's disease, Asia, Eastern Europe, upper GI tract

## Introduction

The incidence of Crohn's disease (CD) has been increasing in previously low incidence areas, including Asia and Eastern Europe (1-3). Susceptibility genes and disease phenotype may differ in CD patients between the East and the West (4). Male predominance and the absence of nucleotide-binding oligomerization domain-containing protein 2 (NOD2) / caspase recruitment domain-containing protein 15 (CARD15) gene mutation in ileal CD have been reported in CD in Asian population (5). A lower proportion of isolated colonic CD and an increased proportion of ileocolonic CD in Asia and Eastern Europe (2, 4) have been described when compared with Western European CD patients (6, 7). Observational studies have reported a higher proportion of upper GI involvement in Chinese CD patients (8, 9).

Unlike in children with CD whereby upper endoscopy is advocated irrespective of the presence or absence of upper GI symptoms (10), in adults with CD, upper GI endoscopy is not routinely performed in patients who do not have symptoms. Studies included multiple upper GI biopsies on endoscopy have shown that the diagnosis of CD may be missed in 11-29% of the cases (10). In a Hungarian study of pediatric CD, macroscopic lesions have been found on esophagealgastroduodenoscopy (OGD) in 64% of patients with CD. Characteristic lesions for CD were noted in 31% of CD patients, and the diagnostic yield of OGD proved to be 9% of CD patients (11).

Chow et al. have reported that CD patients with upper GI involvement have worse clinical outcomes including increased need of hospitalization and are associated with stricturing and penetrating behavior (9). Therefore, there is a need to assess the frequency of upper GI involvement among CD as they may represent a distinct or worse phenotype.

This prospective study aimed to compare the prevalence of macroscopic and microscopic

upper GI manifestations in asymptomatic CD patients in Asia and Eastern Europe in comparison with non-IBD controls. Secondly, we also compared the rates of *Helicobacter pylori* (*H. pylori*) positivity among CD patients and matched controls.

## Patients and methods

### *Patient recruitment*

Consecutive asymptomatic CD patients were prospectively recruited for upper GI endoscopy between 2013 and 2015 at the Institute of Digestive Disease, the Chinese University of Hong Kong, China and at the 1<sup>st</sup> Department of Medicine, University of Szeged, Hungary. All patients from the Hong Kong cohort were ethnically Chinese. The diagnosis of CD was confirmed based on strict criteria in both centers. Both hospitals have a dedicated IBD in- and outpatient clinic. Only patients with a confirmed diagnosis of CD based on clinical evaluation, endoscopic/radiological appearances, and histology were recruited for the study (12). The Montreal classification was used to classify CD location (ileal (L1), colonic (L2), ileocolonic (L3), and upper GI (L4)) and disease behavior (non-stricturing, non-penetrating (B1), stricturing (B2), and penetrating (B3)) at the time of the diagnosis (13). Date of symptom onset and the diagnosis of IBD were recorded. Data on smoking status, alcohol consumption, concomitant diseases, history of appendectomy, history of tuberculosis (TB), family history of IBD and previous surgeries were collected. OGD and biopsy findings were recorded and histology was performed to assess for *H. pylori*, granuloma, celiac disease and microscopic signs characteristic for CD. Patients enrolled in the study did not receive eradication therapy at least 6 months prior to inclusion.

OGD evaluated the presence of normal mucosa, macroscopic inflammation, erosions, ulcerations, strictures and fistulas. OGD in both center were performed by 2-2 experienced

gastroenterologist-endoscopists. Both centres used Olympus endoscopic equipments: Olympus EXERA II series in Szeged and EXERA III series in Hong-Kong. To reduce inter-observer variations; both teams agreed on predefined criteria on macroscopic definition based on photos and discussion prior to start of study, standardized reporting system was used. A shared common database was used to collect clinical and endoscopic data from both sites based on a standardized reporting system. This system captures macroscopic appearance in the esophagus, stomach and duodenum. Presence of macroscopic inflammation, erosions, ulcerations, strictures and fistulas were recorded. Both centers used the same standardized proforma which is prospectively collected. The endoscopic interpretation was based on agreed and clear definitions prior to study initiation. In each patient, two biopsies were taken from each of the corpus, the antrum and the proximal part of the duodenum for histological assessment by a single GI pathologist blinded to the endoscopic findings. Both pathologists assessed the biopsy samples based on well-defined histopathological criteria for the diagnosis of IBD.

The endoscopic criteria for diagnosis of CD contained discontinuous (skip) inflammatory lesions, mucosal nodularity or "cobblestoned" mucosa, thickened gastric folds, multiple aphthous-like ulcerations and/or linear ulcerations, presence of fistula opening and/or stricture. Anatomical criteria of severity are defined as deep ulcerations eroding the muscle layer (14). On OGD, a) macroscopic inflammation was defined as mucosal hyperemia or erythema, edema or mucosal nodularity; b) aphthous erosion was defined as a superficial mucosal break 2-3 mm in diameter and c) ulcer was defined as a linear or serpiginous mucosal break  $\geq 3$  mm in diameter (15). Histological examination assessed the presence of histological features of chronicity and severity of the inflammation, presence of focally active gastritis and evaluated the presence of *H. pylori*, granuloma and celiac disease. For the

purpose of this study, upper GI CD was defined according to established criteria, defined as macroscopic alteration of the stomach or the duodenum consistent with CD on histology (16). Histological features for CD included the presence of non-caseating granulomas, H. pylori—negative focally enhanced gastritis, crypt abscesses, or lymphoid aggregates (16).

The control group was also recruited prospectively during the same time period. The control group consisted of age- and sex-matched non-IBD participants who underwent OGD for different indications including dyspepsia, epigastric pain and regurgitation.

The primary outcome of this study is differences in upper GI CD between Hungarian and Chinese CD patients. Secondary outcome included H. pylori positivity in CD in Hungary and Hong Kong.

#### *Statistical analyses*

Statistical analysis was performed by an experienced biostatistician. Baseline characteristics, endoscopic findings and histological findings between Asian IBD patients and East European IBD patients were compared.  $\chi^2$  –test were used for categorical variables and Mann – Whitney U-test were used for continuous variables. P values < 0.05 were considered statistically significant.

#### *Ethical statement*

All patients gave informed consent, and the study was approved by the Regional and Institutional Human Medical Biological Research Ethics Committee of the University of Szeged and by the ethics committee of the New Territory East Cluster with site-specific approval from each participating hospital trust. The study was carried out under the

declaration of Helsinki.

## Results

### *Patients with CD*

One hundred and eighty CD patients (100 Hong Kong, China; 80 Szeged, Hungary; 70.6% male; mean age: 38.5 years) were included. The clinical demographics of CD subjects are summarized in Table 1. The mean age of CD patients from Hong Kong and Hungary was 41 and 36 years old ( $p=0.62$ ), respectively. There were more current smokers amongst CD patients in Hungary than in Hong Kong (33.8% vs. 11%;  $p=0.001$ ). History of appendectomy, time of symptom onset and time to diagnosis did not differ between Hungarian and Chinese CD patients (Table 1).

There was no difference in disease location between Hungary and Hong Kong. Non-stricturing, non-penetrating CD was less common in Chinese than in Hungarian patients (41.3% vs. 57%,  $p=0.036$ ). The proportion of subjects with stricturing or penetrating disease did not differ between the two cohorts. There were also no differences in the rates of perianal disease, family history of IBD and surgery (Table 1).

At endoscopy, there was no difference in the presence of macroscopic inflammation in the stomach between the two cohorts (39.2% vs. 31%;  $p=0.213$ ). However, gastric erosions were identified in a significantly higher proportion of Chinese than Hungarian CD patients (2.5% vs. 19%;  $p<0.001$ ). Macroscopic inflammation in the duodenum and duodenal erosions occurred more frequently in Hungarian than in Chinese CD patients (13.9% vs. 2%,  $p\leq 0.001$  and 10.1% vs. 1%,  $p\leq 0.001$ ) (Table 2).



At the time of the study, macroscopic finding of OGD suggested the presence of CD in 24% of the Hungarian cases and 4% of the Chinese cases ( $p \leq 0.001$ ). Histological inflammation suggestive of CD was present in 25.3% of the Hungarian patients and 28% of Chinese patients ( $p = 0.658$ ). Combining the macroscopic and microscopic features, CD was proved to be present in 16.5% vs. 2% of the Hungarian and Chinese patients ( $p \leq 0.001$ ).

Overall 12 patients (6.7%) were re-classified due to OGD. In the Chinese cohort 3 patients without upper GI manifestation at diagnosis were classified to have upper GI involvement after endoscopy. In addition, 3 patients with initial diagnosis of L4 disease had no evidence of upper GI disease on OGD performed during the study thus they were reclassified not having upper GI manifestation of CD. These patients were under immunosuppressive therapy and remained asymptomatic. In the Hungarian cohort 4 patients without the initial diagnosis of L4 disease were classified to have upper GI involvement after endoscopy, while 2 patients with upper GI manifestation at diagnosis were reclassified not to suffer from upper GI disease.

On histological examination chronic inflammation was the most common finding in both groups with predominance of the Hungarian patients (100% vs. 77%,  $p \leq 0.001$ ). Focally active gastritis was present in 2% of the Chinese and 23.8% of the Hungarian patients ( $p \leq 0.001$ ). Granulomas were detected in 1% of the Chinese patients and 7.6% of the Hungarian patients ( $p \leq 0.001$ ). In subjects with macroscopic inflammation, *H. pylori* positivity was found in 33.3% and 7.1% of the Hungarian and the Chinese patients; respectively. Overall *H. pylori* positivity was significantly higher in the Hungarian than in the Chinese CD patients (13.9% vs. 4%,  $p \leq 0.001$ ). 2.5% of the Hungarian and none of the Chinese CD patients were diagnosed with celiac disease ( $p = 0.748$ ). All of the Chinese patients with upper GI manifestation received

maintenance azathioprine therapy. In the Hungarian cohort, 40% of the patients with upper GI CD were on azathioprine, 50% were treated with 5-aminosalicylate and 35% with corticosteroid therapy.

### *Control patients*

One hundred and eighty nine controls (100 Hong Kong; 89 Hungary; 57.7% male; mean age 41 years) were recruited in the study. Table 3 shows the basic demographic data of the control subjects. The main indication for OGD was epigastric discomfort both in Chinese and Hungarian cohorts. Macroscopic inflammation of the esophagus and the stomach was more frequently reported in the Chinese than the Hungarian controls (4.5% vs. 16% and 27% vs. 76%,  $p \leq 0.001$ ), although esophageal and gastric erosions occurred more frequently in the Hungarian than the Chinese patients (24.7% vs. 0% and 10.1% vs. 3%,  $p \leq 0.001$ ). No difference was shown in *H. pylori* positivity between the two control populations (14.6% vs. 15%,  $p = 0.564$ ). Granuloma did not occur on histological examination, and no difference was revealed in the rate of celiac disease. However, none of the patients were diagnosed with celiac disease in Chinese patients (2.2% vs. 0%).

CD subjects in Hong Kong had significantly lower *H. pylori* positivity compared to the matched controls (6% vs. 15%;  $p \leq 0.001$ ), whereas no difference was found in *H. pylori* positivity between the Hungarian CD and control population (13.9 vs. 14.6%;  $p = 0.159$ ) (Table 2 and 3).

### **Discussion**

To our knowledge this is the first study that has prospectively examined and compared the prevalence of macroscopic and microscopic upper GI manifestations and *H. pylori* positivity

in asymptomatic adult CD patients in two populations of different ethnicities and emerging CD incidence. The presence of upper GI CD confirmed by established criteria was 16.5% of the Hungarian patients and 2% of the Hong Kong patients; however, the presence of histological inflammation suggestive of CD did not differ between the Hungarian and Hong Kong CD subjects. When considering endoscopic appearance, gastric erosions occurred more frequently in Hong Kong CD patients, while duodenal inflammation and erosions were more common in Hungarian vs. Chinese CD patients.

These results can be discussed from at least two different views: on one hand, histological criteria of mild upper GI CD are quite accurate and the frequency of upper GI CD is almost the same in both areas, however the endoscopic interpretation of macroscopic lesions of suspected CD is completely different. On the other hand, the presence of macroscopically obvious CD may differ significantly in the two regions. In controls, macroscopic inflammation of the upper GI tract was more frequent in Hong Kong patients and erosions occurred more frequently in the Hungarian patients. No difference was shown in *H. pylori* positivity between the two control populations. However, *H. pylori* infection was more commonly detected in Hong Kong controls vs. CD patients.

The prevalence of gastroduodenal involvement of CD patients varies between 0.5-13% (16-18). In an Italian study upper GI CD was found in approximately 19% of CD subjects being similar to our results (19). Macroscopic and histological findings of CD were detected in 48.7% and 59.7% of CD patients, respectively, and *H. pylori* was found in 8.4% CD subjects which appeared also to be comparable to the present Hungarian results.

Focally active gastritis proved to be low in both groups. According to a recently published review about the prevalence of upper GI CD and its histopathologic changes, focal gastritis

was present in 31% of the patients that is highly similar to that of the Hungarian cohort. The review included 20 studies mainly from European countries or the US. Only 2 studies came from Japan showing that data from Asia are extremely limited. Moreover, since focal gastritis may present in 12% of UC patients and 19% of healthy persons undergoing gastroscopy, it cannot distinguish between CD and UC (20, 21).

CD has been reported with an increasing frequency in some Asian countries during the recent years. The review by Wang et al. summarized proportions of patients with only small intestinal involvement, colonic involvement, or both varying between 28.6-53.8%, 26.9-27%, and 19.2-44.4% in Chinese patients compared to the developed countries with a result of 30%, 15% and 55% in the same locations (22). However, data on upper GI tract involvement in Asian CD patients are generally lacking. The study by Leong et al. revealed 19% of the patients having involvement proximal to the terminal ileum suggesting a higher frequency of upper GI tract CD (8). Notably, in our study we evaluated the involvement of the upper GI tract between the esophagus and the duodenum; therefore data of these two studies are difficult to be compared. The Hungarian cohort of Lovasz BD et al. revealed upper GI tract involvement to be 4.8%, while isolated upper GI tract involvement was only 0.7% (23), which is consistent with our findings that none of the patients had isolated upper GI tract involvement. During our study, 6.7% of the patients were re-classified due to OGD.

Upper GI endoscopy is a regularly performed examination in pediatric IBD and significantly less frequently used in adults suffering from CD. Although the European Crohn's and Colitis Organization guidelines recommend examination of the location and extent of CD in the upper GI tract, irrespective of the findings at ileocolonoscopy, in routine practice, upper endoscopy is mainly performed in CD patients with symptoms (14). As confirming the need

of routine OGD, routine upper endoscopy has been shown to reveal various rates of macroscopic gastroduodenal lesions in 30–75% of CD patients (24, 25). Microscopic inflammation is detected in 24–70% of the patients undergoing upper GI endoscopy (26). Our results also revealed some macroscopic abnormalities in more than one third of the patients with mild chronic inflammation on histology in more than 80% of the patients. The occurrence of epithelioid granulomas, being the histological hallmark of CD, varies between 10–25% (27–29). *H. pylori* infection on histology occurs in about 10% of CD patients (26). In our study, granulomas and *H. pylori* positivity were more common in Hungarian CD patients. Some studies have found a lower prevalence of *H. pylori* infection in IBD patients vs. healthy individuals (30, 31). In the present study, significant difference in *H. pylori* infection between CD patients and controls was only detected in the Chinese population. The meta-analysis by Castaño-Rodríguez et al. also reported a stronger negative association between *H. pylori* infection and IBD in Eastern populations than Western populations. Our consistent findings may be explained by the difference between the *H. pylori* strain in Eastern and Western populations (31, 32).

The incidence of CD lesions in the upper GI tract is frequently underestimated. Although the majority of the macroscopic and microscopic abnormalities were independent from CD, macroscopic findings suspicious of CD differed significantly, while histological evaluation of CD was highly similar in the two ethnics, endoscopy and histology unanimously disclosed CD in 16.5% of the Hungarian and 2% of the Chinese patients of our cohort. This study showed different upper GI tract manifestations in the Hungarian and the Hong Kong cohort; however, histology suggested CD in significantly higher than expected rate in asymptomatic patients.

Since the majority of the asymptomatic patients whose initial diagnosis of upper GI tract involvement has been altered after the study OGD were under immunosuppressive therapy

suggest that adequate immunosuppressive therapy may reverse the upper GI manifestations of CD.

The strength of our study is that to our knowledge the first prospective study assessing upper GI CD in two populations of emerging disease incidence in parallel during the same time period using specific predefined criteria for definition agreed upon prior to start of study. Matched controls were also recruited to assess the *H. pylori* prevalence in two different countries of different *H. pylori* background incidence.

There are also some limitations. There may be inter-observer variation between centers on the macroscopic appearance as photos were not routinely captured during study period. Nonetheless, photos and strict and validated criteria were used to define these lesions. Although central reading for endoscopic mucosal assessment would have provided a more relevant comparison between the two regions, centralized assessment of endoscopic findings usually relates to studies evaluating experimental drugs or novel scoring systems in IBD. Although central reading has been proven to be effective in the assessment of disease activity, it has not been tested in the assessment of upper GI CD. Moreover, video acquisition is not yet the standard of care in clinical endoscopy suites and may require segmental annotation. Moreover, this is a prospective study included data of the real practice collected by trained gastroenterologists as well as endoscopists. Endoscopic interpretation was based on clear definitions, statistical analysis was correct performed by biostatistician. Our controls underwent OGD for certain symptoms so there may be a higher prevalence of finding lesions, asymptomatic controls are more ideal to enroll, but ethically challenging.

In conclusion, we have shown that upper GI lesions are less common among asymptomatic

Chinese vs. Hungarian CD patients based upon our endoscopic criteria but did not differ if considering histological assessment. Gastric erosions are proved to be more common in Hong-Kong, duodenal inflammation was more characteristic for Hungarian CD patients. The detection of granuloma in Hungary was similar to the literature data, whereas focal gastritis was lower than expected in both cohorts.

### **Acknowledgement**

This paper was supported by the János Bolyai Research Scholarship of the Hungarian Academy of Sciences (BO/00632/14/5).

### **References**

1. Thia KT, Loftus EV, Sandborn WJ, Yang S-K. An update on the epidemiology of inflammatory bowel disease in Asia. *Am J Gastroenterol* 2008; 103: 3167–3182
2. Lakatos L, Kiss LS, David G, et al. Incidence, disease phenotype at diagnosis, and early disease course in inflammatory bowel diseases in Western Hungary, 2002–2006. *Inflamm Bowel Dis* 2011; 17: 2558–2565
3. Siew NG, Tang W, Ching JY et al. Incidence and phenotype of inflammatory bowel disease based on results from the Asia-Pacific Crohn's and Colitis Epidemiology Study. *Gastroenterology*, 2013; 145: 158-165
4. Prideaux L, Kamm MA, De Cruz P et al. Inflammatory bowel disease in Asia: A systematic review. *J Gastroenterol and Hepatol*, 2012; 27: 1266-12804
5. Ahuja V, Tandon RK. Inflammatory bowel disease in the Asia-Pacific area: A comparison with developed countries and regional differences. *Journal of Digestive Diseases*, 2010; 11: 134-147
6. Solberg IC, Vatn MH, Høie O, et al. Clinical course in Crohn's disease: results of a

Norwegian population-based ten-year follow-up study. *Clin Gastroenterol Hepatol*, 2007; 5: 1430–1438

7. Henriksen M, Jahnsen J, Lygren I et al. Clinical course in Crohn's disease: results of a five-year population-based follow-up study (the IBSEN study). *Scand J Gastroenterol*, 2007; 42: 602-610

8. Leong RWL, Lau JY, Sung JY. The epidemiology and phenotype of Crohn's disease in the Chinese population. *Inflamm Bowel Dis*, 2004; 10: 646-651

9. Chow DK, Sung JJ, Wu JC et al. Upper gastrointestinal tract phenotype of Crohn's disease is associated with early surgery and further hospitalization. *Inflamm Bowel Dis*, 2009; 15: 551–557

10. IBD Working Group of the European Society for Paediatric Gastroenterology, Hepatology and Nutrition. Inflammatory bowel disease in children and adolescents: recommendations for diagnosis--the Porto criteria. *J Pediatr Gastroenterol Nutr*, 2005; 41: 1-7

11. Kovacs M, Muller KE, Arato A et al. Diagnostic yield of upper endoscopy in paediatric patients with Crohn's disease and ulcerative colitis. Subanalysis of the HUPIR registry. *J Crohns Colitis*, 2012; 6: 86-94

12. Van Assche G, Dignass A, Panes J *et al.* The second European evidence-based consensus on the diagnosis and management of Crohn's disease: definitions and diagnosis. *J Crohns Colitis* 2010; 4: 7 – 27

13. Silverberg MS, Satsangi J, Ahmad T *et al.* Toward an integrated clinical, molecular and serological classification of inflammatory bowel disease: Report of a Working Party of the 2005 Montreal World Congress of Gastroenterology. *Can J Gastroenterol* 2005; 19 (Suppl A): 5 – 36

14. Van Assche G, Axel Dignass A, Panes J, et al. The second European evidence-based consensus on the diagnosis and management of Crohn's disease: definitions and diagnosis.



JCC. 2010; 4: 7–27

15. Yeomans ND, Naesdal J. Systematic review: Ulcer definition in NSAID ulcer prevention trials. *Aliment Pharmacol Ther*, 2008; 27: 465-472

16. Kefalas CH. Gastroduodenal Crohn's disease. *Proc (Bayl Univ Med Cent)*, 2003; 16: 147-151

17. Rutgeerts P, Onette E, Vantrappen G, et al. Crohn's disease of the stomach and duodenum: a clinical study with emphasis on the value of endoscopy and endoscopic biopsies. *Endoscopy*. 1980; 12: 288–294

18. Witte AM, Veenendal RA, Van Hogezaand RA et al. Crohn's disease of the upper gastrointestinal tract: the value of endoscopic examination. *Scand J Gastroenterol Suppl*, 1998; 225: 100-105

19. Annunziata ML, Caviglia R, Papparella LG et al. Upper gastrointestinal involvement of Crohn's disease: A prospective study on the role of upper endoscopy in the diagnostic work-up. *Dig Dis Sci*, 2012; 57: 1618-1623

20. Diaz L, Hernandez-Oquet RE, Deshpande AR et al. Upper gastrointestinal involvement in Crohn Disease: Histopathologic and endoscopic findings. *South Med J*, 2015; 108: 695-700

21. Parente F, Cucino C, Bollani S et al. Focal gastric inflammatory infiltrates in inflammatory bowel disease: prevalence, immunohistochemical characteristics, and diagnostic role. *Am J Gastroenterol*, 2000; 95: 705-711

22. Wang YF, Zhang H, Ouyang Q et al. Clinical manifestations of inflammatory bowel disease: East and West differences, *Journal of Digestive Diseases*, 2007, 8: 121-127

23. Lovasz BD, Lakatos L, Horvath A et al. Evolution of disease phenotype in adult and pediatric onset Crohn's disease in a population-based cohort. *World J Gastroenterol*, 2013; 19: 2217-2226

24. Sawczenko A, Sandhu BK. Presenting features of inflammatory bowel disease in Great

Britain and Ireland. *Arch Dis Child*. 2003; 88: 995–1000

25. Kuriyama M, Kato J, Morimoto N, et al. Specific gastroduodenoscopic findings in Crohn's disease: comparison with findings in patients with ulcerative colitis and gastroesophageal reflux disease. *Dig Liver Dis*. 2008; 40: 468–475

26. Korelitz BI, Waye JD, Kreuning J, et al. Crohn's disease in endoscopic biopsies of the gastric antrum and duodenum. *Am J Gastroenterol*. 1981; 76: 103–109; Cameron DJ. Upper and lower gastrointestinal endoscopy in children and adolescents with Crohn's disease: a prospective study. *J Gastroenterol Hepatol*. 1991; 6: 355–358

27. Wright CL, Riddell RH. Histology of the stomach and duodenum in Crohn's disease. *Am J Surg Pathol*. 1998; 22: 383–390

28. Freeman HJ. Granuloma-positive Crohn's disease. *Can J Gastroenterol*, 2007; 21: 583-587

29. Heresbach D, Alexandre JL, Branger B et al. Frequency and significance of granulomas in a cohort of incident cases of Crohn's disease. *Gut*, 2005; 54: 215-222

30. El-Omar E, Penman I, Cruikshank G, et al. Low prevalence of *Helicobacter pylori* in inflammatory bowel disease: association with sulphasalazine. *Gut*. 1994; 35: 1385–1388

31. Castaño-Rodríguez N, Kaakoush NO, Lee WS et al. Dual role of *Helicobacter* and *Campylobacter* species in IBD: a systematic review and meta-analysis. *Gut*, 2015; epub ahead of print

32. Cook KW, Letley DP, Ingram RJ et al. CCL20/CCR6-mediated migration of regulatory T cells to the *Helicobacter pylori*-infected human gastric mucosa. *Gut*, 2014; 63: 1550–1559

Table 1. The clinical demographics of the Hungarian and Chinese CD patients enrolled in the study

	Hungarian CD patients (n=80)	Chinese CD patients (n=100)	p-value
Gender (male/female)	51/29	76/24	0.74
Mean age at present (years)	36	41	0.602
Mean age at time of symptom onset (years)	28	30	0.568
Mean age at diagnosis (years)	29	33	0.458
Smoking habits			
- Current smoker	27	11	0.001
- Ex smoker	8	14	
- Never smoked	43	75	
History of appendectomy	10	10	0.602
Family history of IBD	3	3	0.465
Previous bowel surgery	25	28	0.864
Location at diagnosis			
-L1 (ileal)	26	27	0.747
-L2 (colonic)	16	21	0.648
-L3 (ileocolonic)	38	51	0.682
-L4 (upper GI)	12	4	0.011
Behaviour at diagnosis			
-B1 (non stricturing non penetrating)	33	57	0.036
-B2 (stricturing)	20	23	0.783
-B3 (penetrating)	23	26	0.132
-Perianal manifestation	22	33	0.400
-H. pylori positivity (%)	13.9	4	≤0.001

Table 2. Endoscopic and histological findings of the Hungarian and Chinese CD patients enrolled in the study

Endoscopic findings of the stomach	Hungarian CD patients (n=80)	Chinese CD patients (n=100)	p-value
Macroscopic inflammation (%)	39.2	31	0.213
Erosions (%)	2.5	19	≤0.001
Ulcers (%)	0	3	0.302
Strictures (%)	0	0	0.452
Fistulation (%)	0	1	0.346

  

Endoscopic findings of the duodenum	Hungarian CD patients (n=80)	Chinese CD patients (n=100)	p-value
Macroscopic inflammation (%)	13.9	2	≤0.001
Erosions (%)	10.1	1	≤0.001
Ulcers (%)	0	0	0.214
Strictures (%)	0	0	0.402
Fistulation (%)	1.3	1	0.426

  

Histological findings	Hungarian CD patients (n=80)	Chinese CD patients (n=100)	p-value
Chronic inflammation (%)	100	77	≤0.001
Severity of the inflammation (%)			
-mild	87.3	71	0.231
-moderate	11.1	14	0.202
-severe	1.6	1	0.110
Focally enhanced gastritis (%)	23.8	2	0.001
Presence of granuloma (%)	7.6	1	≤0.001
H. pylori positivity (%)	13.9	4	≤0.001

Table 3. Demographic data of the Hungarian and Chinese control patients enrolled in the study

	Hungarian controls (n=89)	Chinese controls (n=100)	p-value
Gender (male/female)	41/48	68/32	0.408
Mean age at present (years)	41	41	0.726
Indication of OGD			
-Epigastric discomfort	25	46	
-Dyspepsia	7	25	
-Regurgitation	15	19	
-Abdominal bloating	14	2	
-Dysphagia	4	2	
-History of GI bleeding	1	0	
-Anaemia	7	4	
-Others	4	2	
-H. pylori positivity (%)	14.6	15	0.564