

Meson decay constants from $N_f = 2$ clover fermions

Meinulf Göckeler^a, Roger Horsley^b, Dirk Pleiter^{*c}, Paul E.L. Rakow^d, Gerrit Schierholz^{c,e}, Wolfram Schroers^c, Hinnerk Stüben^f and James M. Zanotti^c

^a *Institut für Theoretische Physik, Universität Regensburg, 93040 Regensburg, Germany*

^b *School of Physics, University of Edinburgh, Edinburgh EH9 3JZ, UK*

^c *John von Neumann-Institut für Computing NIC, Deutsches Elektronen-Synchrotron DESY, 15739 Zeuthen, Germany*

^d *Theoretical Physics Division, Department of Mathematical Sciences, University of Liverpool, Liverpool L69 3BX, UK*

^e *Deutsches Elektronen-Synchrotron DESY, 22603 Hamburg, Germany*

^f *Konrad-Zuse-Zentrum für Informationstechnik Berlin, 14195 Berlin, Germany*

E-mail: dirk.pleiter@desy.de

QCDSF-UKQCD Collaboration

We present recent results for meson decay constants calculated on configurations with two flavours of O(a)-improved Wilson fermions. Non-perturbative renormalisation is applied and quark mass dependencies as well as finite volume and discretisation effects are investigated. In this work we also present the first computation of the coupling of the light vector mesons to the tensor current using dynamical fermions.

XXIIIrd International Symposium on Lattice Field Theory

25-30 July 2005

Trinity College, Dublin, Ireland

*Speaker.

1. Introduction

The theoretical description of decay processes which include light mesons strongly relies on a non-perturbative and model independent determination of the relevant parameters. The experimentally most accurately known decay constant is the pseudo-scalar decay constant, f_{PS} , which can be determined from the leptonic decay of charged pions or kaons. Attempts to compute this quantity from first principles on the lattice is an important validation of the lattice QCD methodology. It allows in particular to test control on the required extrapolations to the infinite volume, continuum and finally chiral limit. For phenomenology much more interesting is a lattice determination of the coupling of the light vector mesons to the vector current (f_V) and the tensor current (f_V^\perp). While the constant f_V can in principle be extracted from experimental studies of, e.g., leptonic decays like $\tau^- \rightarrow V^- \nu_\tau$, the coupling f_V^\perp can only be determined theoretically. It has been pointed out by the authors of [1] that the form factors describing the semi-leptonic B-decay $B \rightarrow \rho l \nu$ strongly depend on these couplings. A precise calculation of these couplings is hence an important step towards a more precise determination of one of the less well known CKM parameters, $|V_{ub}|$.

2. Lattice Details

The configurations with $N_f = 2$ flavours of non-perturbatively improved Wilson fermions and Wilson glue which have been generated by the QCDSF and UKQCD collaborations in recent years now cover a larger range of quark masses and several lattice spacings. The simulations are performed for several values of the gauge coupling β with the lattice spacing varying between 0.07 – 0.11 fm. For each lattice spacing configurations are generated for 3-4 sea quark masses in the range of 590 – 1170 MeV. For additional leverage on the quark mass dependence we performed simulations at a large number of valence quark masses which vary between 470 and 1200 MeV. Also the volumes are varied to explore finite size effects. Finally, a non-perturbative determination of the renormalisation constants is performed to further reduce systematic uncertainties.

Correlation functions for pseudo-scalar and vector mesons are calculated on configurations taken at a distance of 5-10 trajectories using 8-4 different locations of the fermion source. We use binning to obtain an effective distance of 20 trajectories. The size of the bins has little effect on the error, which indicates auto-correlations are small. To improve overlap with the ground state we employ Jacobi smearing.

In order to scale our results at different values of β and κ_{sea} we use the Sommer-scale r_0 , which can be determined with good precision on the lattice. When translating this into more common physical units, we face the problem that the static quark potential is difficult to determine experimentally. We therefore choose to use lattice results for $r_0 m_N$ (where m_N is the nucleon mass) to obtain $r_0 = 0.467$ fm. We typically will use r_0/a at the respective values of β and κ_{sea} . Only when stated explicitly we will use r_0/a extrapolated to the chiral limit [2].

To fully eliminate $O(a)$ discretisation errors, it is not sufficient to improve the action, the operators also need to be improved. To determine the renormalised pseudo-scalar decay constant we therefore have to calculate

$$f_{\text{PS}} = Z_A (1 + b_A a m_{q,\text{sea}}) f_{\text{PS}}^{(0)} \left(1 + c_A \frac{f_{\text{PS}}^{(1)}}{f_{\text{PS}}^{(0)}} \right), \quad (2.1)$$




Figure 1: Comparison of c_A determined from perturbation theory (dotted line, [6]), boosted perturbation theory (dashed line), and non-perturbative methods (solid line, [4]).

where Z_A is the axial vector current renormalisation factor and b_A a coefficient describing its dependency on the bare sea-quark mass $am_{q,\text{sea}} = \frac{1}{2}(1/\kappa_{\text{sea}} - 1/\kappa_{\text{sea}}^c)$. c_A is the improvement coefficient which needs to be chosen such that $O(a)$ discretisation errors cancel.

The unimproved bare decay constant $f_{\text{PS}}^{(0)}$ and the improvement term are computed from the matrix elements $m_{\text{PS}} f_{\text{PS}}^{(0)} = \langle 0|A_4|\text{PS}\rangle$ and $m_{\text{PS}} a f_{\text{PS}}^{(1)} = \langle 0|a\partial_4 P|\text{PS}\rangle$. While the renormalisation constant Z_A [3, 4] (we use [3]) and the critical value of the hopping parameter κ_{sea}^c [5] have been determined non-perturbatively, the coefficient b_A is only known perturbatively [6]. As the expansion in the bare gauge coupling $g^2 = 6/\beta$ is known to have bad convergence properties, we use the boosted coupling $g^{\overline{\text{MS}}}(\mu)$ at the scale $\mu = 1/a$. This coupling is computed from the renormalisation group equations with $\mu/\Lambda^{\overline{\text{MS}}} = (r_0/a)/(r_0\Lambda^{\overline{\text{MS}}})$, taking r_0/a in the chiral limit and $r_0\Lambda^{\overline{\text{MS}}} = 0.617(40)(21)$ [2]. To assess the differences between perturbation theory, boosted perturbation theory and non-perturbative methods, we compare the results for c_A . As can be seen from Fig. 1 the values from boosted perturbation theory are much closer to the non-perturbative results. However, there is still a significant discrepancy between the two results.

Similarly, we define the renormalised and improved vector decay constant f_V :

$$\frac{1}{f_V} = Z_V (1 + b_V am_{q,\text{sea}}) f_V^{(0)} \left(1 + c_V \frac{f_V^{(1)}}{f_V^{(0)}} \right), \quad (2.2)$$

where $e(\lambda)_i m_V^2 f_V^{(0)} = \langle 0|V_i|V, \lambda\rangle$ and $e(\lambda)_i m_V^2 a f_V^{(1)} = \langle 0|a\partial_4 T_{i4}|V, \lambda\rangle$ with $e(\lambda)_i$ being the polarization vector. The renormalisation constant Z_V and the coefficient b_V have been computed non-perturbatively [7], but we have to rely on a perturbative calculation of c_V [6].

Finally, we define the renormalised and improved constant for the coupling of the vector meson to the tensor current, f_V^\perp :

$$f_V^\perp(\mu) = Z_T^\mathcal{S}(\mu) (1 + b_T am_{q,\text{sea}}) \left(f_V^{\perp(0)} + c_T f_V^{\perp(1)} \right), \quad (2.3)$$

where $e(\lambda)_i m_V f_V^{\perp(0)} = \langle 0|T_{\mu\nu}|V, \lambda\rangle$ and $e(\lambda)_i m_V a f_V^{\perp(1)} = \langle 0|a(\delta_{\mu\rho}\partial_\rho V_\nu - \delta_{\nu\rho}\partial_\rho V_\mu)|V, \lambda\rangle$. For the coefficients b_T and c_T only perturbative results are available [6]. The QCDSF collaboration has calculated the renormalisation constant (which depends on both, a scheme \mathcal{S} and a scale μ) non-perturbatively in the RI'-MOM scheme [7]. We use the 4-loop and 3-loop results for the β -function and anomalous dimension to translate our results into the $\overline{\text{MS}}$ scheme at $\mu = 2 \text{ GeV}$ [8].

3. Results

The results which are obtained for different values of the gauge coupling $\beta = 5.20, 5.25, 5.29$ and 5.40 can be well parametrised by the following ansatz:

$$r_0 f_x(a, r_0 m_{\text{PS,sea}}, r_0 m_{\text{PS,val}}) = r_0 F_x(a) \left[1 + \alpha_0 (r_0 m_{\text{PS,sea}})^2 + \alpha_1 (r_0 m_{\text{PS,val}})^2 + \beta_0 (r_0 m_{\text{PS,sea}})^4 + \beta_1 r_0^4 m_{\text{PS,sea}}^2 m_{\text{PS,val}}^2 + \beta_2 (r_0 m_{\text{PS,val}})^4 \right]. \quad (3.1)$$

Here $am_{\text{PS,sea}}$ denotes the mass of the pseudo-scalar meson for $\kappa_{\text{sea}} = \kappa_{\text{val}}$, while for $am_{\text{PS,val}}$ the valence and sea quark mass might be different. Given the small number of sea quark masses, we fixed the value of $\beta_0 = 0$. In case of f_V^\perp we find the valence quark mass dependency to be linear in $(am_{\text{PS,val}})^2$ and therefore we also fix $\beta_2 = 0$. The results of our fits to this ansatz are shown in Fig. 2, 3 and 4. As can be seen from these figures, discretisation effects seem to be small, in particular compared to the statistical errors. For f_{PS} and f_V^\perp we therefore fit the results in the chiral limit to a constant. Only in case of $1/f_V$ we find much better agreement when fitting to a polynomial linear in $(a/r_0)^2$.

For $\beta = 5.29$ and $\kappa_{\text{sea}} = 0.13550, 0.13590$ we determine the decay constants for three different volumes $L^3 \times T = 12 \times 32, 16 \times 32$ and 24×48 , which corresponds to $L \approx 1.0, 1.3, 2.0$ fm. We find the pseudo-scalar decay constant f_{PS} to be shifted by $2 - 6\%$ when increasing the spatial extent from 1.3 to 2.0 fm. This seems to be larger than one would expect from the results of chiral perturbation theory [9]. No significant finite size effects have been observed for the vector couplings f_V and f_V^\perp .

In the chiral limit, our result for the pseudo-scalar decay constant is $131(7)$ MeV, where only the statistical error has been taken into account. An extrapolation to the physical pion mass gives a slightly larger value of $133(7)$ MeV, which still agrees very well with the experimental result $f_\pi^{\text{exp}} = 130.7$ MeV. Chiral logarithms predicted by chiral perturbation theory, which have not




Figure 2: The upper plot displays the result for $r_0 f_{\text{PS}}$ at $\beta = 5.29$. The lines show the result of a fit to Eq. (3.1). Solid lines are used for the valence quark mass dependency, a dotted for the sea quark mass dependency. The lower plot shows $r_0 f_{\text{PS}}$ as a function of $(a/r_0)^2$ together with a fit to a constant. Both r_0/a and $a f_{\text{PS}}$ have been extrapolated to the chiral limit.




Figure 3: Similar to Fig. 2, but for $1/f_V$. An ansatz linear in a^2 was used for the continuum extrapolation.

been taken into account here, would shift this result down. Extrapolating our results to the kaon mass gives us an estimate of $f_K \approx 155$ MeV. This again agrees well with the experimental result $f_K^{\text{exp}} = 159.8$ MeV. It should be noted however that our estimate for f_K is based on simulations with degenerate quark masses. For the rho we find $1/f_\rho = 0.333(12)$. This is slightly larger than the experimental estimate $1/f_\rho^{\text{exp}} \approx 0.27$ obtained from the branching ratio $\text{BR}(\tau \rightarrow \rho^- \nu_\tau)$. Extrapolating our results for the tensor coupling gives $f_V^{\perp \text{RGI}} = 183(4)$ MeV or, after conversion to the $\overline{\text{MS}}$ -scheme, $f_V^{\perp \overline{\text{MS}}}(2 \text{ GeV}) = 168(3)$ MeV. This result is larger compared to previous results from QCD sum rules [10] and from lattice simulations employing the quenched approximation [1, 11], which are in the range of 140 – 150 MeV.




Figure 4: Similar to Fig. 2, but for $f_V^{\perp \text{RGI}}$.

Acknowledgements

The numerical calculations have been performed on the APEmille at NIC/DESY (Zeuthen), the Hitachi SR8000 at LRZ (Munich) and on the T3E at EPCC (Edinburgh) [12]. This work is supported in part by the DFG (Forschergruppe Gitter-Hadronen-Phänomenologie) and by the EU Integrated Infrastructure Initiative under contract number RII3-CT-2004-506078. We would like to thank A. Irving for providing updated results for r_0/a prior to publication.

References

- [1] D. Becirevic et al., *Coupling of the light vector meson to the vector and to the tensor current*, *JHEP* **0305** (2003) 007 [arXiv:hep-lat/0301020].
- [2] M. Göckeler et al. [QCDSF Collaboration], *A determination of the Lambda parameter from full lattice QCD*, [arXiv:hep-ph/0502212].
- [3] M. Göckeler et al. [QCDSF Collaboration], *The axial charge of the nucleon on the lattice and in chiral perturbation theory*, these proceedings, PoS(LAT2005)349.
- [4] M. Della Morte, R. Hoffmann and R. Sommer, *Non-perturbative improvement of the axial current for dynamical Wilson fermions*, *JHEP* **0503** (2005) 029 [arXiv:hep-lat/0503003]; M. Della Morte et al. [Alpha Collaboration], *Non-perturbative renormalization of the axial current with dynamical Wilson fermions*, *JHEP* **0507** (2005) 007 [arXiv:hep-lat/0505026].
- [5] M. Göckeler et al. [QCDSF Collaboration], *Determination of light and strange quark masses from full lattice QCD*, [arXiv:hep-ph/0409312].
- [6] S. Sint and P. Weisz, *Further results on $O(a)$ improved lattice QCD to one-loop order of perturbation theory*, *Nucl. Phys.* **B502** (1997) 251 [arXiv:hep-lat/9704001]; S. Sint and P. Weisz, *Further one-loop results in $O(a)$ improved lattice QCD*, *Nucl. Phys. Proc. Suppl.* **63** (1998) 856 [arXiv:hep-lat/9709096].
- [7] T. Bakeyev et al. [QCDSF-UKQCD Collaboration], *Non-perturbative renormalisation and improvement of the local vector current for quenched and unquenched Wilson fermions*, *Phys. Lett.* **B 580** (2004) 197 [arXiv:hep-lat/0305014]; M. Göckeler et al. [QCDSF Collaboration], in preparation.
- [8] J. A. M. Vermaseren, S. A. Larin and T. van Ritbergen, *The 4-loop quark mass anomalous dimension and the invariant quark mass*, *Phys. Lett.* **B 405** (1997) 327; J. A. Gracey, *Three loop \overline{MS} tensor current anomalous dimension in QCD*, *Phys. Lett.* **B 488** (2000) 175.
- [9] G. Colangelo, S. Dürr and C. Haefeli, *Finite volume effects for meson masses and decay constants*, *Nucl. Phys.* **B721** (2005) 136 [arXiv:hep-lat/0503014].
- [10] P. Ball and V. M. Braun, *The ρ Meson Light-Cone Distribution Amplitudes of Leading Twist Revisited*, *Phys. Rev.* **D54** (1996) 2182 [arXiv:hep-ph/9602323].
- [11] S. Capitani et al. [QCDSF Collaboration], *Towards a lattice calculation of $\Delta(q)$ and $\delta(q)$* , *Nucl. Phys. Proc. Suppl.* **79** (1999) 548 [arXiv:hep-ph/9905573]; Göckeler et al. [QCDSF Collaboration], in preparation; V. M. Braun et al., *A lattice calculation of vector meson couplings to the vector and tensor currents using chirally improved fermions*, *Phys. Rev.* **D68** (2003) 054501 [arXiv:hep-lat/0306006].
- [12] C.R. Allton et al. [UKQCD Collaboration], *Effects of non-perturbatively improved dynamical fermions in QCD at fixed lattice spacing*, *Phys. Rev.* **D65** (2002) 054502 [hep-lat/0107021].