

## Generalized Parton Distributions from Lattice QCD

M. Gökeler,<sup>1,2</sup> R. Horsley,<sup>3</sup> D. Pleiter,<sup>4</sup> P. E. L. Rakow,<sup>5</sup> A. Schäfer,<sup>2</sup> G. Schierholz,<sup>4,6</sup> and W. Schroers<sup>7</sup>

(QCDSF Collaboration)

<sup>1</sup>*Institut für Theoretische Physik, Universität Leipzig, D-04109 Leipzig, Germany*

<sup>2</sup>*Institut für Theoretische Physik, Universität Regensburg, D-93040 Regensburg, Germany*

<sup>3</sup>*School of Physics, University of Edinburgh, Edinburgh EH9 3JZ, United Kingdom*

<sup>4</sup>*John von Neumann-Institut für Computing NIC, Deutsches Elektronen-Synchrotron DESY, D-15738 Zeuthen, Germany*

<sup>5</sup>*Theoretical Physics Division, Department of Mathematical Sciences, University of Liverpool, Liverpool L69 3BX, United Kingdom*

<sup>6</sup>*Deutsches Elektronen-Synchrotron DESY, D-22603 Hamburg, Germany*

<sup>7</sup>*Center for Theoretical Physics, Massachusetts Institute of Technology, Cambridge, Massachusetts 02139, USA*  
(Received 13 May 2003; published 29 January 2004)

We perform a quenched lattice calculation of the first moment of twist-two generalized parton distribution functions of the proton, and assess the total quark (spin and orbital angular momentum) contribution to the spin of the proton.

DOI: 10.1103/PhysRevLett.92.042002

PACS numbers: 12.38.Gc, 13.60.Fz

Generalized parton distributions [1] (GPDs) provide a deeper understanding of the internal structure of hadrons in terms of quarks and gluons. While ordinary parton distributions measure the probability  $|\psi(x)|^2$  of finding a parton with fractional momentum  $x$  in the hadron, GPDs describe the coherence of two different hadron wave functions  $\psi^\dagger(x + \xi/2)\psi(x - \xi/2)$ , one where the parton carries fractional momentum  $x + \xi/2$  and one where this fraction is  $x - \xi/2$ , from which information about parton-parton correlation functions can be deduced. As a consequence, GPDs depend on the momentum transfer  $\Delta^2$  between the initial and final hadron, which provides further information on the transverse location of quarks and gluons [2]. Spatial images of hadrons can thus be obtained, where the resolution is determined by the virtuality  $Q^2$  of the incoming photon. Last, but not least, GPDs allow us to isolate the contribution of the quark orbital angular momentum to the spin of hadrons. Lattice QCD is the only known method that is able to compute moments of GPDs from first principles.

We will restrict ourselves to the GPDs  $H_q$  and  $E_q$  of the nucleon, where  $q = u, d, \dots$  denotes the flavor of the struck quark. We will not consider the gluon sector here. The lowest, zeroth moments of  $H_q$  and  $E_q$  are given by

the Dirac and Pauli form factors:

$$\int_{-1}^1 dx H_q(x, \xi, \Delta^2) = F_1^q(\Delta^2), \quad (1)$$

$$\int_{-1}^1 dx E_q(x, \xi, \Delta^2) = F_2^q(\Delta^2). \quad (2)$$

Both form factors have been computed on the lattice in a similar calculation [3] to the present one and found to be well described by a dipole ansatz

$$F_{1,2}^q(\Delta^2) = F_{1,2}^q(0)/(1 - \Delta^2/M_{1,2}^2)^2 \quad (3)$$

for sufficiently small (and accessible) momenta, with dipole masses  $M_{1,2}$  of the order of the  $\rho$ ,  $\omega$  mass, when extrapolated to the physical pion mass.

The first moments of  $H_q$  and  $E_q$  are of the form [1]

$$\int_{-1}^1 dx x H_q(x, \xi, \Delta^2) = A_2^q(\Delta^2) + \xi^2 C_2^q(\Delta^2), \quad (4)$$

$$\int_{-1}^1 dx x E_q(x, \xi, \Delta^2) = B_2^q(\Delta^2) - \xi^2 C_2^q(\Delta^2), \quad (5)$$

where  $A_2^q(\Delta^2)$ ,  $B_2^q(\Delta^2)$ , and  $C_2^q(\Delta^2)$  are generalized form factors, which are given by the nucleon matrix elements of the energy-momentum tensor (EMT):

$$\begin{aligned} \langle p' | \mathcal{O}_{\{\mu\nu\}}^q | p \rangle &\equiv \frac{i}{2} \langle p' | \bar{q} \gamma_{\{\mu} \vec{D}_{\nu\}} q | p \rangle \\ &= A_2^q(\Delta^2) \bar{u}(p') \gamma_{\{\mu} \bar{p}_{\nu\}} u(p) - B_2^q(\Delta^2) \frac{i}{2m_N} \bar{u}(p') \Delta^\alpha \sigma_{\alpha\{\mu} \bar{p}_{\nu\}} u(p) + C_2^q(\Delta^2) \frac{1}{m_N} \bar{u}(p') u(p) \Delta_{\{\mu} \Delta_{\nu\}}. \end{aligned} \quad (6)$$

Here  $m_N$  denotes the nucleon mass,  $\bar{p} = \frac{1}{2}(p' + p)$ ,  $\Delta = p' - p$ , and curly brackets refer to symmetrization of indices and subtraction of traces. The EMT has twist two and spin two. It is assumed to be renormalized at the scale  $\mu$ , which makes  $A_2^q(\Delta^2)$ ,  $B_2^q(\Delta^2)$ , and  $C_2^q(\Delta^2)$

scale and scheme dependent. For the classification of states of definite  $J^{PC}$  contributing to (6) in the  $t$ -channel, see [4]. The so-called skewedness parameter  $\xi$  is defined by  $\xi = -n \cdot \Delta$ , where  $n$  is a lightlike vector

with  $n \cdot \bar{p} = 1$ , and bounded by  $|\xi| \leq 2\sqrt{\Delta^2/(\Delta^2 - 4m_N^2)}$ . In the forward limit,  $\Delta^2 \rightarrow 0$ , we have

$$A_2^q(0) = \langle x_q \rangle \equiv \int_0^1 dx x(q_{\uparrow}(x) + q_{\downarrow}(x)), \quad (7)$$

where  $q_{\uparrow(\downarrow)}(x)$  are the usual quark distributions with spin parallel (antiparallel) to the spin of the nucleon. Furthermore, one derives [5]

$$\frac{1}{2}(A_2^q(0) + B_2^q(0)) = J_q, \quad (8)$$

where  $J_q$  is the angular momentum of the  $q$  quark, and  $J = \sum_q J_q$  is the total angular momentum of the nucleon carried by the quarks. The angular momentum decomposes, in a gauge invariant way, into two pieces:

$$J_q = L_q + S_q, \quad (9)$$

where  $L_q$  is the orbital angular momentum and

$$S_q = \frac{1}{2}\Delta q \equiv \frac{1}{2}\int_0^1 dx (q_{\uparrow}(x) - q_{\downarrow}(x)) \quad (10)$$

is the spin of the quark. We know  $\Delta q$  from separate calculations [6,7], so that  $L_q$  can be computed from (8).

In this Letter, we perform a quenched lattice calculation of the generalized form factors  $A_2^q(\Delta^2)$ ,  $B_2^q(\Delta^2)$ , and  $C_2^q(\Delta^2)$ . The quenched approximation neglects fluctuations of virtual quark-antiquark pairs from the Dirac sea. The nonforward matrix elements (6) are computed from ratios of three- and two-point functions following [3]. Further details are given in [8]. To keep cutoff effects small, we use nonperturbatively  $\mathcal{O}(a)$  improved Wilson




FIG. 1. The generalized form factors  $A_2^u$ ,  $B_2^u$ , and  $C_2^u$  at  $\kappa = 0.1333$ , together with the dipole fit and the extrapolated values at  $\Delta^2 = 0$  ( $\square$ ).




FIG. 2. The generalized form factors  $A_2^d$ ,  $B_2^d$ , and  $C_2^d$  at  $\kappa = 0.1333$ , together with the dipole fit and the extrapolated values at  $\Delta^2 = 0$  ( $\square$ ).

TABLE I. Parameters of the dipole fit. In the bottom row we give the parameters extrapolated to the physical pion mass.

$\kappa$	$M$ [GeV]	$A_2^u(0)$	$B_2^u(0)$	$C_2^u(0)$	$A_2^d(0)$	$B_2^d(0)$	$C_2^d(0)$
0.1324	1.69(05)	0.419(07)	0.344(028)	-0.084(26)	0.188(04)	-0.281(20)	-0.071(15)
0.1333	1.58(06)	0.415(10)	0.334(044)	-0.101(35)	0.176(05)	-0.260(29)	-0.073(19)
0.1342	1.41(10)	0.404(19)	0.357(117)	-0.117(70)	0.158(10)	-0.265(80)	-0.067(35)
	1.11(20)	0.400(22)	0.334(113)	-0.134(81)	0.147(11)	-0.232(77)	-0.071(42)

fermions. The calculation is done on  $16^3 32$  lattices at  $\beta = 6.0$  and for three different hopping parameters,  $\kappa = 0.1324, 0.1333, \text{ and } 0.1342$ , which allows us to extrapolate our results to the chiral limit. Using  $r_0 = 0.5$  fm to set the scale, which results in the inverse lattice spacing  $1/a = 2.12$  GeV, the corresponding pion masses are 1070, 870, and 640 MeV. If we use the nucleon mass extrapolated to the chiral limit to set the scale, the pion masses are 930, 760, and 550 MeV, and  $1/a = 1.84$  GeV. The corresponding nucleon masses and the choice of nucleon momenta  $p, p'$  can be inferred from [3]. For the EMT we consider two sets of (Euclidean) operators:

$$\frac{1}{\sqrt{2}}(\mathcal{O}_{\mu\nu} + \mathcal{O}_{\nu\mu}), \quad 1 \leq \mu < \nu \leq 4, \quad (11)$$

and

$$\frac{1}{2}(\mathcal{O}_{11} + \mathcal{O}_{22} - \mathcal{O}_{33} - \mathcal{O}_{44}), \quad (12)$$

$$\frac{1}{\sqrt{2}}(\mathcal{O}_{33} - \mathcal{O}_{44}), \quad \frac{1}{\sqrt{2}}(\mathcal{O}_{11} - \mathcal{O}_{22}).$$

Each set transforms irreducibly under the hypercubic group. The operators (11) and (12) are renormalized multiplicatively,  $\mathcal{O}(\mu) = Z(a\mu) \mathcal{O}(a)$ , with renormalization constants [6]  $Z_{v_{2a}}$  and  $Z_{v_{2b}}$ , respectively. The renormalization constants are computed nonperturbatively [9] following [10]. We obtain  $Z_{v_{2a}}^{\overline{MS}}(2 \text{ GeV}) = 1.10$  and  $Z_{v_{2b}}^{\overline{MS}}(2 \text{ GeV}) = 1.09$ . The following results refer to the  $\overline{MS}$  scheme at the renormalization scale  $\mu = 2$  GeV.

In Figs. 1 and 2 we show the generalized form factors  $A_2^u(\Delta^2)$ ,  $B_2^u(\Delta^2)$ ,  $C_2^u(\Delta^2)$ , and  $A_2^d(\Delta^2)$ ,  $B_2^d(\Delta^2)$ ,  $C_2^d(\Delta^2)$  of the proton for  $\kappa = 0.1333$ . Data points with larger errors are not shown here but are included in the fit. The corresponding form factors of the neutron are obtained by interchanging  $u$  and  $d$ . Similarly good results are found for  $\kappa = 0.1324$  and  $0.1342$ . The generalized form factors can be well described by the dipole ansatz

$$A_2^q(\Delta^2) = A_2^q(0)/(1 - \Delta^2/M^2)^2, \quad (13)$$

and similarly for  $B_2^q$  and  $C_2^q$ . Fits of  $A_2^u(\Delta^2)$  and  $A_2^d(\Delta^2)$  give the same dipole mass  $M$  within errors. The dipole masses obtained from separate fits of  $B_2^u(\Delta^2)$ ,  $B_2^d(\Delta^2)$ ,  $C_2^u(\Delta^2)$ , and  $C_2^d(\Delta^2)$  are found to be consistent with that value. We therefore have decided to fit our data by a common dipole mass  $M$ . Our data do not favor a monopole behavior. The results of the fits are shown in

Table I. For a reliable extrapolation to  $\Delta^2 = 0$  we find it important to cover a wide enough range of  $\Delta^2$  values. This may be the reason why our dipole masses turn out to be systematically larger than those found in a previous calculation [11].

In Fig. 3 we show the dipole mass  $M$  as a function of the pion mass. The mass values appear to lie on a straight line, as was observed already in the case of the nucleon form factors [3]. A linear extrapolation in  $m_\pi$  to the physical pion mass gives  $M = 1.1(2)$  GeV. This value is close to the physical masses of the  $f_2, a_2$  mesons, which supports the hypothesis of tensor meson dominance. A quadratic extrapolation in  $m_\pi$  leads to  $M = 1.3(1)$  GeV. The form factor data  $A_2^q(0)$ ,  $B_2^q(0)$ , and  $C_2^q(0)$  show little variation with the quark mass and are extrapolated quadratically in  $m_\pi$  to the physical pion mass. The results are shown in the bottom row of Table I. It should be stressed that all quantities refer (at best) to valence quark distributions, because sea quark effects have been neglected. In unquenched simulations there are also quarkline disconnected contributions. For an estimate, see [11].

If the dipole behavior (3), (13) continues to hold for the higher moments as well, and if we assume that the dipole masses continue to grow in a Regge-like fashion, we would obtain

$$\int_{-1}^1 dx x^n H_q(x, 0, \Delta^2) = \langle x^n \rangle / (1 - \Delta^2/M_{n+1}^2)^2, \quad (14)$$

with  $M_l^2 = \text{const} + l/\alpha'$ ,  $\alpha'$  being the slope of the Regge trajectory. This would mean that with increasing momentum transfer  $|\Delta^2|$  the lower moments of  $H_q(x, 0, \Delta^2)$  are




FIG. 3. The dipole mass  $M$  as a function of  $m_\pi$ , together with a linear extrapolation to the physical pion mass ( $\square$ ).




FIG. 4. The total angular momentum  $J$ , together with a quadratic extrapolation to the physical pion mass ( $\square$ ).

suppressed more than the higher ones, so that the observed peak in  $H_q(x, 0, 0) = q_1(x) + q_1(x)$  around  $x \approx 0.2$  is shifted towards the higher values of  $x$ . As a result, the  $\Delta^2$  dependence cannot be factorized in a simple way, as is sometimes assumed. Knowing  $\langle x_q^n \rangle$ , we can reconstruct  $H_q(x, 0, \Delta^2)$  from (14) by inverse Mellin transform. The  $\xi$  dependence of both  $H_q$  and  $E_q$  appears to be rather weak, based on our knowledge of the first two moments, and in the isovector channel (corresponding to proton-neutron or  $u$ - $d$  matrix elements) it largely cancels out.

In Fig. 4 we show the total angular momentum  $J = J_u + J_d$ . The dependence on the pion mass is rather flat, as expected [12]. The errors are due to the relatively large statistical errors of  $B_2^u$  and  $B_2^d$  and the fact that  $B_2^u$  and  $B_2^d$  cancel to a large extent. In Table II we give our results for  $J$ , and separately for  $J_q$  and  $S_q$ , extrapolated quadratically (linearly in  $m_\pi^2$ ) to the physical pion mass. The numbers for  $S_q$  refer to our latest results [9], computed from the nonperturbatively improved axial vector current with nonperturbative renormalization factors. It turns out that the total angular momentum  $J$  carried by the quarks amounts to  $\approx 70\%$  of the spin of the (quenched) proton, leaving a contribution of  $\approx 30\%$  for the gluons. The major contribution is given by the  $u$  quark, while the contribution of the  $d$  quark is found to be negligible, which hints at strong pairing effects. Our result for  $J$  is somewhat smaller than that of [11,13]. We are able to compute  $L_q$  now. The total orbital angular momentum of the quarks turns out to be consistent with zero:

$$L \equiv L_u + L_d = 0.03(7). \quad (15)$$

This indicates that (at virtuality  $\mu = 2$  GeV) the parton's transverse momentum in the (quenched) proton is small. A similar conclusion can be drawn from our earlier finding [14] of a small twist-three contribution  $d_2$  to the second moment of the polarized structure function  $g_2$ .

The generalized form factors  $C_2^q(\Delta^2)$  contribute to the beam charge asymmetry of deeply virtual Compton scattering. We obtain a rather small value:  $C_2^u(0) + C_2^d(0) =$

TABLE II. The total angular momentum and its individual contributions, extrapolated to the physical pion mass.

$J$	$J_u$	$J_d$	$S_u$	$S_d$
0.33(7)	0.37(6)	-0.04(4)	0.42(1)	-0.12(1)

-0.2(1). This result is to be compared with the value -0.8 obtained in the chiral quark-soliton model at  $\mu \approx 0.6$  GeV [15]. For a discussion, see also [16].

As far as one can compare, quenched and unquenched results agree surprisingly well, and we do not expect to find significant differences here either. For a recent study of quenching artifacts, as well as cutoff effects, see [17].

This work is supported by DFG and under the Feodor Lynen program. Discussions with P. Hägler, J. Negele, D. Renner, and C. Weiss are acknowledged. The numerical calculations have been performed at NIC (Jülich).

- [1] D. Müller *et al.*, Fortschr. Phys. **42**, 101 (1994); X. Ji, Phys. Rev. Lett. **78**, 610 (1997); A.V. Radyushkin, Phys. Rev. D **56**, 5524 (1997); M. Diehl *et al.*, Phys. Lett. B **411**, 193 (1997); X. Ji, J. Phys. G **24**, 1181 (1998); J. Blümlein, B. Geyer, and D. Robaschik, Nucl. Phys. **B560**, 283 (1999).
- [2] M. Diehl, Eur. Phys. J. C **25**, 223 (2002); M. Burkardt, Int. J. Mod. Phys. A **18**, 173 (2003).
- [3] M. Göckeler *et al.*, hep-lat/0303019.
- [4] X. Ji and R. F. Lebed, Phys. Rev. D **63**, 076005 (2001).
- [5] X. Ji, Phys. Rev. Lett. **78**, 610 (1997).
- [6] M. Göckeler *et al.*, Phys. Rev. D **53**, 2317 (1996).
- [7] M. Fukugita *et al.*, Phys. Rev. Lett. **75**, 2092 (1995); S. J. Dong, J.-F. Lagaë, and K. F. Liu, Phys. Rev. Lett. **75**, 2096 (1995); S. Capitani *et al.*, Nucl. Phys. (Proc. Suppl.) **B79**, 548 (1999); S. Güsken *et al.*, Phys. Rev. D **59**, 114502 (1999); D. Dolgov *et al.*, Phys. Rev. D **66**, 034506 (2002).
- [8] P. Hägler *et al.*, Phys. Rev. D **68**, 034505 (2003).
- [9] M. Göckeler *et al.* (to be published).
- [10] G. Martinelli *et al.*, Nucl. Phys. **B445**, 81 (1995); M. Göckeler *et al.*, Nucl. Phys. **B544**, 699 (1999).
- [11] N. Mathur *et al.*, Phys. Rev. D **62**, 114504 (2000).
- [12] J.-W. Chen and X. Ji, Phys. Rev. Lett. **88**, 052003 (2002).
- [13] V. Gadiyak, X. Ji, and C. Jung, Phys. Rev. D **65**, 094510 (2002); see, however, W. Wilcox, Phys. Rev. D **66**, 017502 (2002).
- [14] M. Göckeler *et al.*, Phys. Rev. D **63**, 074506 (2001).
- [15] N. Kivel, M.V. Polyakov, and M. Vanderhaeghen, Phys. Rev. D **63**, 114014 (2001); V.Yu. Petrov *et al.*, Phys. Rev. D **57**, 4325 (1998).
- [16] A.V. Belitsky and D. Müller, Nucl. Phys. **A711**, 118 (2002); F. Ellinghaus, Nucl. Phys. **A711**, 171 (2002); A. Freund, M. McDermott, and M. Strikman, Phys. Rev. D **67**, 036001 (2003).
- [17] T. Bakeyev *et al.*, hep-lat/0311017.