



Brain atrophy in seizure-free temporal lobe epilepsy: implications for predicting pharmacoresistance

Journal:	<i>Epilepsia</i>
Manuscript ID	Draft
Manuscript Type:	Letter
Date Submitted by the Author:	n/a
Complete List of Authors:	Keller, Simon; University of Liverpool, Molecular and Clinical Pharmacology; King's College London, Department of Clinical Neuroscience
Key Words:	Prognosis, Imaging, Outcome, Atrophy

SCHOLARONE™
Manuscripts

Only

**Brain atrophy in seizure-free temporal lobe epilepsy:
implications for predicting pharmacoresistance**

Simon S. Keller

Department of Molecular and Clinical Pharmacology, Institute of Translational
Medicine, University of Liverpool
Department of Neuroradiology, Walton Centre NHS Foundation Trust
Department of Basic and Clinical Neuroscience, Institute of Psychiatry,
Psychology & Neuroscience, King's College London

Dr. Simon S. Keller
Department of Molecular and Clinical Pharmacology
Institute of Translational Medicine
University of Liverpool
Clinical Sciences Centre
Lower Lane
Liverpool, L9 7LJ
simon.keller@liverpool.ac.uk
Tel: +44 (0)151 529 5461

To the Editors:

I read with interest the recent manuscript by Alvim et al.¹ entitled “Progression of gray matter atrophy in seizure-free patients with temporal lobe epilepsy”. Using voxel-based morphometry (VBM) applied to conventional 3D T1-weighted images, the authors report widespread extrahippocampal atrophy in patients with well-controlled temporal lobe seizures, and that progressive atrophy occurs over time in the absence of recurrent seizures. As the authors point out, this suggests that “extrahippocampal atrophy has multifactorial causes and cannot be defined as the origin of the pharmacoresistance. Although seizures might be responsible to some extent for this atrophy, this probably does not influence the AED response” (pg 8). This work could be interpreted to indicate that more widespread brain atrophy is not necessarily a reflection of a seizure disorder that is less amenable to pharmacological or surgical therapy. This idea is not entirely consistent with the authors’ previous interesting papers, which indicate – using the same VBM techniques – that pharmacological² and surgical³ control of temporal lobe seizures is less likely in patients with more widespread atrophy.

TLE is a network disorder. Diffusion tensor imaging (DTI) studies are increasingly reporting that seizure freedom after temporal lobe surgery is less likely when deficits are observed in temporal lobe pathways.^{4, 5} Given the current paucity of data, it is currently unknown whether pharmacoresistance can be predicted using similar approaches in patients with a new diagnosis of TLE. Magnetic resonance spectroscopy may offer important insights.⁶

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

A crucial point is that the development of robust, individualized, imaging prognostic markers of treatment outcome in TLE is more likely to be achieved using imaging approaches that provide biological information that exceeds what is afforded by conventional structural MRI scans, which are typically acquired in context of the clinical evaluation of patients. The volumetric, thickness and surface area information afforded by conventional scans constrain inferences to macroscopic brain morphology. There is no doubt that these approaches have furthered our understanding of pathological brain changes in focal epilepsy. However, the ability to develop highly sensitive and robust prognostic imaging measures of treatment outcome – and a detailed understanding of the mechanisms underlying pharmacoresistance and postoperative outcome - are more likely to be achieved using imaging techniques that probe neurobiology on a microscopic scale. Diffusion approaches are not without limitation, but do represent an important development. Techniques that probe neuroinflammation may represent an important pathway,⁷ but there is of yet no evidence indicating that MRI can be used to directly, accurately and validly determine neuroinflammation in epilepsy. In large scale imaging studies (i.e. >100 patients), imaging modalities are invariably restricted to what is collected for clinical purposes. New advanced imaging approaches that may have the sensitivity to discriminate between individual patients who will or will not achieve seizure remission are likely to be administered in context of research projects in smaller groups of patients, which limits the generalizability of findings. It may therefore be important to consider a coordinated effort across institutions to

1
2
3 harmonize advanced imaging protocols to identify non-invasive markers of
4
5 seizure control.
6
7
8

9 10 **Acknowledgement**

11 SSK is funded by a UK Medical Research Council grant (Grant Number
12
13 MR/K023152/1).
14
15
16

17 18 **Disclosure**

19
20 The author has no conflicts of interest to disclose. I confirm that I have read
21
22 the Journal's position on issues involved in ethical publication and affirm that
23
24 this report is consistent with those guidelines.
25
26
27

28 29 **References**

- 30
31
32 1. Alvim MK, Coan AC, Campos BM, et al. Progression of gray matter atrophy
33 in seizure-free patients with temporal lobe epilepsy. *Epilepsia*. 2016 Feb 11.
34 2. Bilevicius E, Yasuda CL, Silva MS, et al. Antiepileptic drug response in
35 temporal lobe epilepsy: a clinical and MRI morphometry study. *Neurology*. 2010
36 Nov 9;75:1695-701.
37 3. Yasuda CL, Valise C, Saude AV, et al. Dynamic changes in white and gray
38 matter volume are associated with outcome of surgical treatment in temporal
39 lobe epilepsy. *Neuroimage*. 2010 Jan 1;49:71-9.
40 4. Keller SS, Richardson MP, Schoene-Bake JC, et al. Thalamotemporal
41 alteration and postoperative seizures in temporal lobe epilepsy. *Ann Neurol*.
42 2015 May;77:760-74.
43 5. Bonilha L, Jensen JH, Baker N, et al. The brain connectome as a
44 personalized biomarker of seizure outcomes after temporal lobectomy.
45 *Neurology*. 2015 May 5;84:1846-53.
46 6. Campos BA, Yasuda CL, Castellano G, et al. Proton MRS may predict AED
47 response in patients with TLE. *Epilepsia*. 2010 May;51:783-8.
48 7. Feindel KW. Can we develop pathology-specific MRI contrast for "MR-
49 negative" epilepsy? *Epilepsia*. 2013 May;54 Suppl 2:71-4.
50
51
52
53
54
55
56
57
58
59
60