

Vector Reachability Problem in $SL(2, \mathbb{Z})$ *

Igor Potapov¹ and Pavel Semukhin²

1 Department of Computer Science, University of Liverpool, United Kingdom
potapov@liverpool.ac.uk

2 Department of Computer Science, University of Liverpool, United Kingdom
semukhin@liverpool.ac.uk

Abstract

The decision problems on matrices were intensively studied for many decades as matrix products play an essential role in the representation of various computational processes. However, many computational problems for matrix semigroups are inherently difficult to solve even for problems in low dimensions and most matrix semigroup problems become undecidable in general starting from dimension three or four.

This paper solves two open problems about the decidability of the vector reachability problem over a finitely generated semigroup of matrices from $SL(2, \mathbb{Z})$ and the point to point reachability (over rational numbers) for fractional linear transformations, where associated matrices are from $SL(2, \mathbb{Z})$. The approach to solving reachability problems is based on the characterization of reachability paths between points which is followed by the translation of numerical problems on matrices into computational and combinatorial problems on words and formal languages. We also give a geometric interpretation of reachability paths and extend the decidability results to matrix products represented by arbitrary labelled directed graphs. Finally, we will use this technique to prove that a special case of the scalar reachability problem is decidable.

1998 ACM Subject Classification F.2.1 Numerical Algorithms and Problems, F.1.1 Models of Computation

Keywords and phrases Matrix Semigroup, Vector Reachability Problem, Special Linear Group, Linear Fractional Transformation, Automata and Formal Languages

Digital Object Identifier 10.4230/LIPIcs.MFCS.2016.85

1 Introduction

Decision problems on matrices were intensively studied from 1947 when A. Markov showed the connection between classical computations and problems for matrix semigroups [24]. Moreover matrix products play an essential role in the representation of various computational processes, i.e., linear recurrent sequences [18, 27, 28], arithmetic circuits [14], hybrid and dynamical systems [26, 3], probabilistic and quantum automata [7], stochastic games, broadcast protocols [13], optical systems, etc. New algorithms for solving reachability problems in matrix semigroups can be incorporated into software verification tools and used for analysis of mathematical models in physics, chemistry, biology, ecology, and economics.

However, many computational problems for matrix semigroups are inherently difficult to solve even when the problems are considered in dimension two, and most of these problems become undecidable in general starting from dimension three or four. Examples of such problems are the Membership problem (including the special cases of the Mortality

* This work was supported by EPSRC grant “Reachability problems for words, matrices and maps” (EP/M00077X/1).



© Igor Potapov and Pavel Semukhin;
licensed under Creative Commons License CC-BY

41st International Symposium on Mathematical Foundations of Computer Science (MFCS 2016).

Editors: Piotr Faliszewski, Anca Muscholl, and Rolf Niedermeier; Article No. 85; pp. 85:1–85:14

Leibniz International Proceedings in Informatics



LIPICs Schloss Dagstuhl – Leibniz-Zentrum für Informatik, Dagstuhl Publishing, Germany

and Identity problems), vector reachability, scalar reachability, freeness problem and the emptiness problem of matrix semigroups intersection [6]. All above problems are tightly connected, including three central problems:

- **The membership problem:** Let $S = \langle G \rangle$ be a semigroup generated by a finite set G of $n \times n$ matrices. Determine whether a given matrix M belongs to S , that is, determine whether there exists a sequence of matrices M_1, M_2, \dots, M_k in G such that $M = M_1 \cdot M_2 \cdot \dots \cdot M_k$
- **The vector reachability problem:** Let \mathbf{x} and \mathbf{y} be two vectors and S be a given finitely generated semigroup of $n \times n$ matrices. Determine whether there is a matrix $M \in S$ such that $M\mathbf{x} = \mathbf{y}$.
- **The scalar reachability problem:** Let \mathbf{x} and \mathbf{y} be two vectors, λ be a scalar, and S be a given finitely generated semigroup of $n \times n$ matrices. Determine whether there is a matrix $M \in S$ such that $\mathbf{x}^\top M\mathbf{y} = \lambda$.

The vector reachability problem can be seen as a parameterized version of the membership problem, where some elements of a matrix M are either independent variables or variables linked by some equations. In contrast to the original membership problem, where all values of M are constants, in vector reachability we may have an infinite set of matrices that can transform a vector \mathbf{x} to \mathbf{y} . Thus the decidability results for the membership cannot be directly applied to the vector reachability problem.

The scalar reachability can be viewed as a vector to hyperplane reachability problem. Indeed, we can rewrite the equation $\mathbf{x}^\top M\mathbf{y} = \lambda$ as a system of two equations: $M\mathbf{y} = \mathbf{z}$ and $\mathbf{x}^\top \mathbf{z} = \lambda$. So, the question becomes if there is a matrix $M \in S$ that maps a given vector \mathbf{y} to a vector \mathbf{z} that lies on a hyperplane $\mathbf{x}^\top \mathbf{z} = \lambda$. Because there are infinitely many vectors on a hyperplane, decidability of the scalar reachability problem does not follow directly from the decidability of the vector reachability problem.

Most of the problems such as membership, vector reachability and freeness are undecidable for 3×3 integer matrices. The undecidability proofs in matrix semigroups are mainly based on various techniques and methods of embedding universal computations into three and four dimensional matrices and their products. The case of dimension two is the most intriguing one since there is some evidence that if these problems are undecidable, then this cannot be proved using a construction similar to the one used for dimensions 3 and 4. In particular, there is no injective semigroup morphism from pairs of words over any finite alphabet (with at least two elements) into complex 2×2 matrices [8], which means that the coding of independent pairs of words in 2×2 complex matrices is impossible and the exact encoding of the Post Correspondence Problem or a computation of a Turing Machine cannot be used directly for proving undecidability in 2×2 matrix semigroups over \mathbb{Z} , \mathbb{Q} or \mathbb{C} . The only undecidability result in dimension two for the vector reachability and the membership problems has been shown in the case of 2×2 matrices over quaternions [4].

The main hypothesis is that problems for 2×2 matrix semigroups over integers, rationals or complex numbers could be decidable, but not much is known about the status of these problems. Recently, there was some progress on the *Membership problem*, which was shown to be decidable in $SL(2, \mathbb{Z})$, and the *Identity problem*, which was shown to be decidable in $\mathbb{Z}^{2 \times 2}$ [11]. Later the decidability of the *Freeness problem* (that is, to decide whether each element can be expressed uniquely as a product of generating matrices) was shown for $SL(2, \mathbb{Z})$ [9]. On the other hand, the Mortality, Identity and vector reachability problems were shown to be at least NP-hard for $SL(2, \mathbb{Z})$ in [5, 6], but for the modular group the membership was shown to be decidable in polynomial time by Gurevich and Schupp [16].

The algorithmic properties of $\text{SL}(2, \mathbb{Z})$ are important in the context of many fundamental problems in hyperbolic geometry [34, 10, 12], dynamical systems [29], Lorenz/modular knots [22], braid groups [30], particle physics, high energy physics [33], M/string theories [15], ray tracing analysis, music theory [25] and can lead to further decidability results in $\mathbb{Z}^{2 \times 2}$ using matrix presentation in the Smith normal form.

This paper solves two open problems about the decidability of the vector reachability problem for finitely generated semigroups of matrices from $\text{SL}(2, \mathbb{Z})$ and the point to point reachability (over rational numbers) for fractional linear transformations $f_M(x) = \frac{ax+b}{cx+d}$, where the associated matrix $M = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$ belongs to $\text{SL}(2, \mathbb{Z})$. The approach to solving these reachability problems for 2×2 matrix semigroups is based on the analysis of reachability paths between vectors or points. This analysis is then used to translate the numerical reachability problems into computational problems on words and regular languages. We also present several extensions of our main results, give a geometric interpretation of reachability paths, and use this technique to solve a special case of the scalar reachability problem.

The decidability proof of the vector reachability problem in dimension two presented in this paper is the first nontrivial new result for solving vector reachability problems since 1996 when it was shown that the problem is decidable for any commutative matrix semigroup in any dimension [1] and for a special case of non-commuting matrices [20]. On the other hand, in the general case of non-commuting matrices the problem is known to be undecidable already for integer matrices in dimension three [17].

2 Preliminaries

The integers and rationals are denoted by \mathbb{Z} and \mathbb{Q} , respectively, and $\text{SL}(2, \mathbb{Z})$ is a group of 2×2 integer matrices with determinant 1. The notation $a \mid b$ means that a divides b , and $a \nmid b$ means that a does not divide b , when a and b are integer numbers.

► **Definition 1.** With each matrix $M = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \in \text{SL}(2, \mathbb{Z})$ we associate a fractional linear map (also called Möbius transformation) $f_M : \mathbb{Q} \rightarrow \mathbb{Q}$ defined as $f_M(x) = \frac{ax+b}{cx+d}$. This definition can be extended to $f : \mathbb{Q} \cup \{\infty\} \rightarrow \mathbb{Q} \cup \{\infty\}$ in a natural way by setting $f_M(\infty) = \frac{a}{c}$ if $c \neq 0$, $f_M(\infty) = \infty$ if $c = 0$, and $f_M(x) = \infty$ if $cx + d = 0$.

Note that we have $f_{M_1} \circ f_{M_2} = f_{M_1 M_2}$ for any matrices M_1 and M_2 .

Let M_1, \dots, M_n be a finite collection of matrices. Then $\langle M_1, \dots, M_n \rangle$ denotes the multiplicative semigroup (including the identity matrix) generated by M_1, \dots, M_n .

► **Definition 2.** The *vector reachability problem* in $\text{SL}(2, \mathbb{Z})$ is defined as follows: Given two vectors \mathbf{x} and \mathbf{y} with integer coefficients and a finite collection of matrices M_1, \dots, M_n from $\text{SL}(2, \mathbb{Z})$, decide whether there exists a matrix $M \in \langle M_1, \dots, M_n \rangle$ such that $M\mathbf{x} = \mathbf{y}$.

► **Definition 3.** The *reachability problem by fractional linear transformations* in $\text{SL}(2, \mathbb{Z})$ is defined as follows: Given two rational numbers x and y and a finite collection of matrices M_1, \dots, M_n from $\text{SL}(2, \mathbb{Z})$, decide whether there exists a matrix $M \in \langle M_1, \dots, M_n \rangle$ such that $f_M(x) = y$.

The main result of our paper is that the vector reachability problem and the reachability problem by fractional linear transformations for $\text{SL}(2, \mathbb{Z})$ are decidable (Theorem 14). Both proofs follow the same pattern. We will use the fact that any matrix M from $\text{SL}(2, \mathbb{Z})$ can

be expressed as product of matrices $S = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}$ and $R = \begin{bmatrix} 0 & -1 \\ 1 & 1 \end{bmatrix}$. So we can represent any $M \in SL(2, \mathbb{Z})$ by a word w in the alphabet $\Sigma = \{S, R\}$.

The main idea of the proof is to show that the solution set of the equation $M\mathbf{x} = \mathbf{y}$ has the form $\left\{ B \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}^t C : t \in \mathbb{Z} \right\}$, where B and C are some matrices from $SL(2, \mathbb{Z})$ that can be computed in PTIME from \mathbf{x}, \mathbf{y} (Theorem 8). Similarly, the solution set of the equation $f_M(x) = y$ can be presented as a union of two sets of such form (Theorem 10). After translating matrices into words, these sets become regular languages. On the other hand, the language that corresponds to the semigroup $\langle M_1, \dots, M_n \rangle$ is also regular. Indeed, if M_i corresponds to the word w_i , then the semigroup $\langle M_1, \dots, M_n \rangle$ translates into the language $(w_1 + \dots + w_n)^*$. The last step of the proof is to show that the emptiness problem of the intersection of two such languages is decidable (Proposition 13).

Here is a more detailed description of our proofs. Let $M = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$, $\mathbf{x} = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$ and $\mathbf{y} = \begin{bmatrix} y_1 \\ y_2 \end{bmatrix}$. To show that the equation $M\mathbf{x} = \mathbf{y}$ defines a regular language we must solve the following system of three equations in four unknown variables:

$$x_1a + x_2b = y_1 \qquad x_1c + x_2d = y_2 \qquad ad - bc = 1$$

Choosing b as a free parameter, we can reduce it to the following system of linear congruence equations:

$$x_2b \equiv y_1 \pmod{x_1} \qquad y_2b \equiv -x_1 \pmod{y_1} \qquad x_2y_2b \equiv y_1y_2 - x_1x_2 \pmod{x_1y_1}$$

It can be shown that the above system either has no solutions or it has a solution of the form $b = b_1t + b_2$, where $t \in \mathbb{Z}$, and hence all coefficients of the matrix M are linear functions of t . In Proposition 7 we will show that such matrices can be written in the form $M = B \begin{bmatrix} 1 & k \\ 0 & 1 \end{bmatrix}^t C$, where B, C are some matrices from $SL(2, \mathbb{Z})$, k is a fixed integer number and $t \in \mathbb{Z}$ is a free parameter. After that it is not hard to see that such solution translates into a regular language.

We will use a similar approach to prove that the equation $f_M(x) = y$ also defines a regular language. In fact, we will do it by showing that the solution set of $f_M(x) = y$ is equal to the union of the solution sets of the equations $M\mathbf{x} = \mathbf{y}$ and $M\mathbf{x} = -\mathbf{y}$ for suitable vectors \mathbf{x} and \mathbf{y} .

The final step is to show that there is an algorithm that decides whether the intersection of two regular subsets of $SL(2, \mathbb{Z})$ is empty or not. Our idea relies on the fact that the intersection of two regular languages is regular, and that the emptiness problem for regular languages is decidable. The problem here is that we cannot apply these facts directly because for each matrix $M \in SL(2, \mathbb{Z})$ there are infinitely many words $w \in \{S, R\}^*$ that correspond to M , and only some of them may appear in the given language. However there is only one *reduced* word that corresponds to M , that is, the word that does not have a substring of the form SS or RRR . So, our solution is to take any automaton \mathcal{A} and turn it into a new automaton $\tilde{\mathcal{A}}$ that accepts the same language as \mathcal{A} plus all reduced words w that correspond to non-reduced words w' accepted by \mathcal{A} .

The construction of the automaton $\tilde{\mathcal{A}}$ was inspired by a similar construction from [11]. Note that in $SL(2, \mathbb{Z})$ we have an equality $S^2 = R^3 = -I$. Thus to construct $\tilde{\mathcal{A}}$ we add to \mathcal{A} a new ε -transition from a state q_1 to a state q_2 if there is a run of \mathcal{A} from q_1 to q_2 labelled

by SS or RRR . We will apply this procedure iteratively until no new ε -transitions can be added. However we need to keep track of sign changes when we add new ε -transitions. To achieve this we will use *signed automata*, which are slight modifications of the usual finite automata but they take into account such sign changes.

Now to solve the emptiness problem for the intersection of two regular languages L_1 and L_2 , we take the signed automata \mathcal{A}_1 and \mathcal{A}_2 that accept L_1 and L_2 , respectively, and construct new automata $\tilde{\mathcal{A}}_1$ and $\tilde{\mathcal{A}}_2$ as described above. After that we can check whether $L(\tilde{\mathcal{A}}_1) \cap L(\tilde{\mathcal{A}}_2) \neq \emptyset$.

In the Section 4 we will show how to extend these decidability results to arbitrary regular subsets of $\text{SL}(2, \mathbb{Z})$, i.e., subsets that are defined by finite automata. Using this technique we will show how to algorithmically solve the equation $M_1^{x_1} \cdots M_k^{x_k} \mathbf{x} = N_1^{y_1} \cdots N_l^{y_l} \mathbf{y}$, where \mathbf{x}, \mathbf{y} are given vectors from $\mathbb{Z} \times \mathbb{Z}$, the matrices M_1, \dots, M_k and N_1, \dots, N_l are from $\text{SL}(2, \mathbb{Z})$, and x_1, \dots, x_k and y_1, \dots, y_l are unknown non-negative integers. Furthermore, we will show how to apply this method to prove that a special case of the scalar reachability problem is decidable.

All missing proofs can be found in the extended version of this paper, which is available online on arXiv.org [31].

3 Main results

The characterization of the solution set of the equation $M\mathbf{x} = \mathbf{y}$ given in Theorem 8 will follow from Propositions 5 and 7. First, we prove one simple lemma which we will use several times in our arguments.

► **Lemma 4.** Let $\mathbf{x} = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$ and $\mathbf{y} = \begin{bmatrix} y_1 \\ y_2 \end{bmatrix}$ be vectors from $\mathbb{Z} \times \mathbb{Z}$ and M be a matrix from $\text{SL}(2, \mathbb{Z})$ such that $M\mathbf{x} = \mathbf{y}$. Then $\gcd(x_1, x_2) = \gcd(y_1, y_2)$.

Proof. Take any $k \in \mathbb{Z}$ such that $k \mid x_1, x_2$ and let $M = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$. Then from $M\mathbf{x} = \mathbf{y}$ we have $y_1 = ax_1 + bx_2$ and $y_2 = cx_1 + dx_2$. Thus $k \mid y_1, y_2$. Now since $M \in \text{SL}(2, \mathbb{Z})$, M^{-1} is also in $\text{SL}(2, \mathbb{Z})$, and $M\mathbf{x} = \mathbf{y}$ is equivalent to $M^{-1}\mathbf{y} = \mathbf{x}$. So, if $k \in \mathbb{Z}$ is any number such that $k \mid y_1, y_2$, then $k \mid x_1, x_2$. Therefore, $\gcd(x_1, x_2) = \gcd(y_1, y_2)$. ◀

► **Proposition 5.** Let $\mathbf{x} = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$ and $\mathbf{y} = \begin{bmatrix} y_1 \\ y_2 \end{bmatrix}$ be two vectors from $\mathbb{Z} \times \mathbb{Z}$, such that \mathbf{x} is not equal to the zero vector $\mathbf{0}$, and consider the matrix equation $M\mathbf{x} = \mathbf{y}$, where M is an unknown matrix from $\text{SL}(2, \mathbb{Z})$. Then either this equation does not have a solution or all its solutions are given by $M = tA_1 + A_2$, where t is any integer number, A_1, A_2 are some matrices from $\mathbb{Z}^{2 \times 2}$ such that A_1 is a nonzero matrix. Moreover, there is a polynomial time algorithm that determines whether such an equation has a solution and if so, finds it.

Proof. See Section A of the Appendix in [31]. ◀

For the next proposition we will need the following theorem about the Smith normal form of a matrix.

► **Theorem 6** (Smith normal form [19]). For any nonzero matrix $A \in \mathbb{Z}^{2 \times 2}$, there are matrices B, C from $\text{SL}(2, \mathbb{Z})$ such that $A = B \begin{bmatrix} t_1 & 0 \\ 0 & t_2 \end{bmatrix} C$ for some $t_1, t_2 \in \mathbb{Z}$ such that $t_1 \neq 0$ and $t_1 \mid t_2$. Moreover, B, C, t_1, t_2 can be computed in polynomial time.

► **Proposition 7.** *Let A_1 and A_2 be matrices from $\mathbb{Z}^{2 \times 2}$ such that A_1 is a nonzero matrix and, for every $t \in \mathbb{Z}$, we have $tA_1 + A_2 \in SL(2, \mathbb{Z})$. Then there are matrices B and C from $SL(2, \mathbb{Z})$ and $k \in \mathbb{Z}$ such that*

$$tA_1 + A_2 = BT^{kt}C \text{ for every } t \in \mathbb{Z},$$

where $T = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix} \in SL(2, \mathbb{Z})$. Moreover, B , C , and k can be computed in polynomial time.

Proof. Let $A_1 = \begin{bmatrix} a_1 & b_1 \\ c_1 & d_1 \end{bmatrix}$ and $A_2 = \begin{bmatrix} a_2 & b_2 \\ c_2 & d_2 \end{bmatrix}$. By the assumption, for every $t \in \mathbb{Z}$,

$$\begin{vmatrix} a_1t + a_2 & b_1t + b_2 \\ c_1t + c_2 & d_1t + d_2 \end{vmatrix} = 1. \text{ That is } (a_1t + a_2)(d_1t + d_2) - (b_1t + b_2)(c_1t + c_2) = 1 \text{ or}$$

$$(a_1d_1 - b_1c_1)t^2 + (a_1d_2 + a_2d_1 - b_1c_2 - b_2c_1)t + a_2d_2 - b_2c_2 = 1 \text{ for all } t \in \mathbb{Z}.$$

Therefore, $a_1d_1 - b_1c_1 = 0$, $a_1d_2 + a_2d_1 - b_1c_2 - b_2c_1 = 0$, and $a_2d_2 - b_2c_2 = 1$. In particular, $\det(A_1) = 0$ and $\det(A_2) = 1$.

By Theorem 6, there are matrices $F, G \in SL(2, \mathbb{Z})$ such that $A_1 = F \begin{bmatrix} k & 0 \\ 0 & l \end{bmatrix} G$ for some $k, l \in \mathbb{Z}$ such that $k \mid l$. Since $\det(A_1) = 0$ we have that $kl = 0$. However if $k = 0$ and $l = 0$, then A_1 is equal to the zero matrix, contrary to the assumption. Hence we must have that $k \neq 0$ and $l = 0$.

Now $F^{-1}(tA_1 + A_2)G^{-1} = \begin{bmatrix} kt + a & b \\ c & d \end{bmatrix}$, for some $a, b, c, d \in \mathbb{Z}$. Note that since $\det(F) = \det(G) = \det(tA_1 + A_2) = 1$, we have $\begin{vmatrix} kt + a & b \\ c & d \end{vmatrix} = dkt + ad - bc = 1$ for every $t \in \mathbb{Z}$.

Hence $dk = 0$ and so $d = 0$. Substituting $d = 0$ in the above equation, we obtain $bc = -1$. Since b and c are integers, there are only two possibilities: $b = 1, c = -1$, or $b = -1, c = 1$.

So the above matrix actually looks like $F^{-1}(tA_1 + A_2)G^{-1} = \begin{bmatrix} kt + a & \mp 1 \\ \pm 1 & 0 \end{bmatrix}$. Therefore,

$$T^{-c(kt+a)}F^{-1}(tA_1 + A_2)G^{-1} = D, \text{ where } c = \pm 1 \text{ and } D = \begin{bmatrix} 0 & \mp 1 \\ \pm 1 & 0 \end{bmatrix} \in SL(2, \mathbb{Z}). \text{ Hence}$$

$tA_1 + A_2 = FT^{(ck)t}T^{ca}DG$. Note that F and $T^{ca}DG$ are in $SL(2, \mathbb{Z})$. This completes the proof. The bound on complexity follows from the fact that F and G can be computed in PTIME by Theorem 6. ◀

As a corollary of Propositions 5 and 7 we obtain the following theorem.

► **Theorem 8.** *Let $\mathbf{x} = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$ and $\mathbf{y} = \begin{bmatrix} y_1 \\ y_2 \end{bmatrix}$ be vectors from $\mathbb{Z} \times \mathbb{Z}$ such that $\mathbf{x} \neq \mathbf{0}$, and consider the matrix equation $M\mathbf{x} = \mathbf{y}$, where M is an unknown matrix from $SL(2, \mathbb{Z})$. Then either this equation does not have a solution or all its solutions are given by the following formula $M = B \begin{bmatrix} 1 & k \\ 0 & 1 \end{bmatrix}^t C$, where $t \in \mathbb{Z}$.*

In the above expression B and C are some matrices from $SL(2, \mathbb{Z})$, and k is an integer number. Moreover, there is a polynomial time algorithm that determines whether such an equation has a solution and if so, finds the suitable matrices B , C and the integer k .

In Section 4 we will give a geometric interpretation of reachability paths (Figure 1 and Proposition 16), using which we can prove the following corollary.¹ The proof itself can be

¹ Even though we use Corollary 9 in the proofs of Theorem 10 and Proposition 18, it is not essential there for proving decidability. Namely, all references to Corollary 9 in these proofs can be replaced by references to Theorem 8, at the same time replacing T with T^k where appropriate.

found in Section D of the Appendix in [31].

► **Corollary 9.** *The value of the parameter k in Theorem 8 is equal to 1.*

Theorem 8 provides us with a characterization of the matrices $M \in \text{SL}(2, \mathbb{Z})$ that map vector \mathbf{x} to vector \mathbf{y} . This characterization will be used later to prove the decidability of the vector reachability problem. We now give a similar characterization of the matrices $M \in \text{SL}(2, \mathbb{Z})$ for which the fractional linear transformation f_M maps a number x to number y . In fact, we will do this by reducing the problem to finding the solutions of the equation $M\mathbf{x} = \mathbf{y}$ which we discussed above.

► **Theorem 10.** *Let x and y be rational numbers and let $\mathcal{F}(x, y)$ be the following set of matrices from $\text{SL}(2, \mathbb{Z})$:*

$$\mathcal{F}(x, y) = \{M \in \text{SL}(2, \mathbb{Z}) : f_M(x) = y\}.$$

Then $\mathcal{F}(x, y) = \mathcal{F}_1(x, y) \cup \mathcal{F}_2(x, y)$, where each $\mathcal{F}_i(x, y)$ is either empty or has the form

$$\mathcal{F}_i(x, y) = \{B_i T^t C_i : t \in \mathbb{Z}\},$$

where B_i and C_i are some matrices from $\text{SL}(2, \mathbb{Z})$. Moreover, there is a polynomial time algorithm that determines whether each $\mathcal{F}_i(x, y)$ is empty or not and in the latter case finds corresponding matrices B_i and C_i .

Proof. Let us write the numbers x and y as $x = \frac{x_1}{x_2}$ and $y = \frac{y_1}{y_2}$, where we assume that $\gcd(x_1, x_2) = \gcd(y_1, y_2) = 1$. Consider the equation $f_M(x) = y$, where $M = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$ is an unknown matrix from $\text{SL}(2, \mathbb{Z})$. We can rewrite it as

$$\frac{a \frac{x_1}{x_2} + b}{c \frac{x_1}{x_2} + d} = \frac{y_1}{y_2} \quad \text{or} \quad \frac{ax_1 + bx_2}{cx_1 + dx_2} = \frac{y_1}{y_2}. \quad (1)$$

Consider the vectors $\mathbf{x} = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$, $\mathbf{y} = \begin{bmatrix} y_1 \\ y_2 \end{bmatrix}$, and $\mathbf{z} = \begin{bmatrix} z_1 \\ z_2 \end{bmatrix}$, where \mathbf{z} is the vector with coordinates $z_1 = ax_1 + bx_2$ and $z_2 = cx_1 + dx_2$. So we have that $\mathbf{z} = M\mathbf{x}$. In this notation Equation (1) is equivalent to the fact that vector $\mathbf{z} = M\mathbf{x}$ belongs to the set $\{k\mathbf{y} : k \in \mathbb{Z}\}$.

Recall that $\gcd(x_1, x_2) = 1$ and hence, by Lemma 4, we also have that $\gcd(z_1, z_2) = 1$. Thus if $\mathbf{z} = k\mathbf{y}$ for some $k \in \mathbb{Z}$, then we must have that $k = \pm 1$. In other words, we showed that Equation (1) is equivalent to two matrix equations: $M\mathbf{x} = \mathbf{y}$ and $M\mathbf{x} = -\mathbf{y}$. So we have that $\mathcal{F}(x, y) = \mathcal{F}_1(x, y) \cup \mathcal{F}_2(x, y)$, where

$$\mathcal{F}_1(x, y) = \{M \in \text{SL}(2, \mathbb{Z}) : M\mathbf{x} = \mathbf{y}\} \quad \text{and} \quad \mathcal{F}_2(x, y) = \{M \in \text{SL}(2, \mathbb{Z}) : M\mathbf{x} = -\mathbf{y}\}.$$

Note that $\mathbf{x} \neq \mathbf{0}$ because $x_2 \neq 0$. Hence by Theorem 8 and Corollary 9, each $\mathcal{F}_i(x, y)$ is either empty or has the form $\mathcal{F}_i(x, y) = \{B_i T^t C_i : t \in \mathbb{Z}\}$ for some B_i and C_i from $\text{SL}(2, \mathbb{Z})$ which can be computed in polynomial time. ◀

Now we will use signed automata to prove that the emptiness problem for the intersection of two regular subsets of $\text{SL}(2, \mathbb{Z})$ is decidable.

Consider an alphabet $\Sigma = \{S, R\}$ consisting of two symbols S and R and define the mapping $\varphi : \Sigma \rightarrow \text{SL}(2, \mathbb{Z})$ as follows: $\varphi(S) = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}$ and $\varphi(R) = \begin{bmatrix} 0 & -1 \\ 1 & 1 \end{bmatrix}$. We can extend this mapping to the morphism $\varphi : \Sigma^* \rightarrow \text{SL}(2, \mathbb{Z})$ in a natural way. The matrices $\varphi(S)$ and $\varphi(R)$ are in fact generators of $\text{SL}(2, \mathbb{Z})$, so φ is surjective. We call a word $w \in \Sigma^*$ *reduced* if it does not have substrings of the form SS or RRR . In our proof we will make use of the following well-known fact.

► **Theorem 11** ([21, 23, 32]). *For every $M \in \text{SL}(2, \mathbb{Z})$, there exists a unique reduced word $w \in \Sigma^*$ such that either $M = \varphi(w)$ or $M = -\varphi(w)$.*

► **Definition 12.** A *signed automaton* $\mathcal{A} = (\Sigma, Q, I, \Delta, F^+, F^-)$ is a (non-deterministic) finite automaton whose final states are divided into two (not necessarily disjoint) subsets F^+ and F^- .

A *signed language* accepted by a signed automaton \mathcal{A} is a pair $L(\mathcal{A}) = (L(\mathcal{A})^+, L(\mathcal{A})^-)$, where $L(\mathcal{A})^+$ and $L(\mathcal{A})^-$ consists of the words $w \in \Sigma^*$ for which there is a run of \mathcal{A} that ends in the set F^+ or F^- , respectively. Note that we do not assume that $L(\mathcal{A})^+$ and $L(\mathcal{A})^-$ are disjoint.

Let $L = (L^+, L^-)$ be a signed language, then we define a regular subset of $\text{SL}(2, \mathbb{Z})$ corresponding to this language as $\varphi(L) = \{\varphi(w) : w \in L^+\} \cup \{-\varphi(w) : w \in L^-\}$.

The following proposition is an important ingredient of our main results.

► **Proposition 13.** *There is an algorithm that for any given regular signed languages L_1 and L_2 over the alphabet Σ , decides whether $\varphi(L_1) \cap \varphi(L_2)$ is empty or not.*

Proof. See Section B of the Appendix in [31]. ◀

We are now ready to prove our main results.

► **Theorem 14.** *The vector reachability problem and the reachability problem by fractional linear transformations in $\text{SL}(2, \mathbb{Z})$ are decidable.*

Proof. Suppose M_1, \dots, M_n is a given finite collection of matrices from $\text{SL}(2, \mathbb{Z})$. Let $w_1, \dots, w_n \in \Sigma^*$ be some words, not necessarily reduced, such that $M_i = \varphi(w_i)$, for $i = 1, \dots, n$. Define the language $\mathcal{L}_{\text{semigr}}$ that corresponds to the semigroup $\langle M_1, \dots, M_n \rangle$ as $\mathcal{L}_{\text{semigr}} = (w_1 + w_2 + \dots + w_n)^*$.

Recall that in the vector reachability problem we are given two vectors \mathbf{x} and \mathbf{y} from $\mathbb{Z} \times \mathbb{Z}$, and we ask if there is a matrix $M \in \langle M_1, \dots, M_n \rangle$ such that $M\mathbf{x} = \mathbf{y}$. We want to construct a regular language $\mathcal{L}_{\mathbf{x}, \mathbf{y}}^{\text{vrp}}$ that corresponds to these matrices.

If $\mathbf{x} = \mathbf{0}$ and $\mathbf{y} \neq \mathbf{0}$, then we set $\mathcal{L}_{\mathbf{x}, \mathbf{y}}^{\text{vrp}} = \emptyset$ because in this case the equation $M\mathbf{x} = \mathbf{y}$ does not have a solution. On the other hand, if $\mathbf{x} = \mathbf{0}$ and $\mathbf{y} = \mathbf{0}$, then we set $\mathcal{L}_{\mathbf{x}, \mathbf{y}}^{\text{vrp}} = \{S, R\}^*$ because any matrix $M \in \text{SL}(2, \mathbb{Z})$ satisfies the equation $M\mathbf{0} = \mathbf{0}$.

Now assume that $\mathbf{x} \neq \mathbf{0}$. Then by Theorem 8, the matrix equation $M\mathbf{x} = \mathbf{y}$ either has no solution, or its solution has the form $\{BT^tC : t \in \mathbb{Z}\}$, where $T = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$, and B and C are some matrices from $\text{SL}(2, \mathbb{Z})$. Moreover, B and C can be computed from \mathbf{x} and \mathbf{y} in PTIME. In the case when $M\mathbf{x} = \mathbf{y}$ has no solution, we set $\mathcal{L}_{\mathbf{x}, \mathbf{y}}^{\text{vrp}} = \emptyset$. If the solution set is non-empty, then we can rewrite it as

$$\{BT^tC : t \in \mathbb{Z}\} = \{BT^tC : t \geq 0\} \cup \{BT^{-t}C : t \geq 0\}.$$

Let u and v be words from Σ^* such that $B = \varphi(u)$ and $C = \varphi(v)$. It is easy to check that $T = \varphi(S^3R)$ and $T^{-1} = \varphi(R^5S)$. Hence $\mathcal{L}_{\mathbf{x}, \mathbf{y}}^{\text{vrp}} = u(S^3R)^*v + u(R^5S)^*v$ is a regular language that describes the solutions of the equation $M\mathbf{x} = \mathbf{y}$ in $\text{SL}(2, \mathbb{Z})$.

In a similar way we can construct a regular language $\mathcal{L}_{x, y}^{\text{flt}}$ that corresponds to the reachability problem by fractional linear transformations from x to y . By Theorem 10, the set $\mathcal{F}(x, y)$ of matrices from $\text{SL}(2, \mathbb{Z})$ that satisfy the equation $f_M(x) = y$ is equal to $\mathcal{F}(x, y) = \mathcal{F}_1(x, y) \cup \mathcal{F}_2(x, y)$, where each $\mathcal{F}_i(x, y)$ is either empty or has the form $\mathcal{F}_i(x, y) = \{B_iT^tC_i : t \in \mathbb{Z}\}$, where T is as above, and B_i and C_i are some matrices from $\text{SL}(2, \mathbb{Z})$. All these matrices can be computed in PTIME from x and y .

We define $\mathcal{L}_{x,y}^{\text{flt}}$ as the union $\mathcal{L}_{x,y}^{\text{flt}} = \mathcal{L}_1 \cup \mathcal{L}_2$ of two regular languages \mathcal{L}_1 and \mathcal{L}_2 . If $\mathcal{F}_i(x, y)$ is empty, then we set $\mathcal{L}_i = \emptyset$. Otherwise, let u_i and v_i be words from Σ^* such that $B_i = \varphi(u_i)$ and $C_i = \varphi(v_i)$. Then we can define \mathcal{L}_i as $\mathcal{L}_i = u_i(S^3R)^*v_i + u_i(R^5S)^*v_i$. Thus we defined a regular language $\mathcal{L}_{x,y}^{\text{flt}}$ that corresponds the solution set of the equation $f_M(x) = y$ in $\text{SL}(2, \mathbb{Z})$.

We remind that in Proposition 13 we work with signed languages. Therefore, in what follows we convert every regular language L that we have constructed so far into a corresponding signed language (L, \emptyset) .

Finally, the vector reachability problem for \mathbf{x} and \mathbf{y} has a solution if and only if

$$\varphi((\mathcal{L}_{\mathbf{x},\mathbf{y}}^{\text{vfp}}, \emptyset)) \cap \varphi((\mathcal{L}_{\text{semigr}}, \emptyset)) \neq \emptyset.$$

Similarly, the reachability problem by fractional linear transformations for x and y has a solution if and only if

$$\varphi((\mathcal{L}_{x,y}^{\text{flt}}, \emptyset)) \cap \varphi((\mathcal{L}_{\text{semigr}}, \emptyset)) \neq \emptyset.$$

By Proposition 13 these questions are algorithmically decidable. \blacktriangleleft

A characterization of the matrices M from $\text{SL}(2, \mathbb{Z})$ that satisfy the equation $M\mathbf{x} = \mathbf{y}$, which is given in Theorem 8, can be computed in polynomial time. However the overall complexity of the algorithm is EXPTIME if the entries of the matrices are given in binary presentation. This is due to the fact that a reduced word w that corresponds to a given matrix M , i.e., such that $M = \pm\varphi(w)$, has length exponential in the binary presentation of M . So computing symbolic presentations of given matrices and constructing automata for the languages $\mathcal{L}_{\text{semigr}}$, $\mathcal{L}_{\mathbf{x},\mathbf{y}}^{\text{vfp}}$ and $\mathcal{L}_{x,y}^{\text{flt}}$ takes exponential time. The next steps of the algorithm take only polynomial time in the size of these automata. However the PTIME algorithm for computing all mappings from \mathbf{x} to \mathbf{y} could be combined with the result of Gurevich and Schupp [16] to produce a polynomial time algorithm for the vector reachability problem over the modular group.

4 Geometric interpretation and extensions

Consider a semigroup generated by matrices M_1, \dots, M_n from $\text{SL}(2, \mathbb{Z})$. As we showed above, this semigroup can be described by a regular language which we called $\mathcal{L}_{\text{semigr}}$. It's not hard to see that the proof of Theorem 14 remains valid if we replace $\mathcal{L}_{\text{semigr}}$ by any other regular language, that is, a language defined by a finite automaton or a labelled transition system.

► **Proposition 15.** *Suppose that we are given a finite collection of matrices M_1, \dots, M_n from $\text{SL}(2, \mathbb{Z})$ and a regular language $L \subseteq \{1, \dots, n\}^*$. Consider the following generalized reachability problems:*

- **Generalized vector reachability problem.** *Given two vectors \mathbf{x} and \mathbf{y} with integer coefficients, decide whether there exists a word $i_1 \dots i_k$ from the language L such that $M_{i_1} \dots M_{i_k} \mathbf{x} = \mathbf{y}$.*
- **Generalized reachability problem by fractional linear transformations.** *Given two rational numbers x and y , decide whether there exists a word $i_1 \dots i_k$ from L such that $f_{M_{i_1} \dots M_{i_k}}(x) = y$.*

Then the above generalized reachability problems are decidable.

Proof. The proof of this proposition is similar to the proof of Theorem 14. Namely, it follows from the fact that a regular language L defines a regular subset in $\text{SL}(2, \mathbb{Z})$ and Proposition 13, where we proved that the emptiness problem for the intersection of two regular subsets in $\text{SL}(2, \mathbb{Z})$ is decidable. \blacktriangleleft

As an application of Proposition 15 let us consider the follow matrix equation

$$M_1^{x_1} \dots M_k^{x_k} \mathbf{x} = N_1^{y_1} \dots N_l^{y_l} \mathbf{y}, \tag{2}$$

where x_1, \dots, x_k and y_1, \dots, y_l are non-negative integers. In [1] it was proved that if M_1, \dots, M_k and N_1, \dots, N_l are commuting $n \times n$ matrices over algebraic numbers and \mathbf{x}, \mathbf{y} are vectors with algebraic coefficients, then it is decidable in polynomial time whether the Equation (2) has a solution. On the other hand, in [2] it was shown that there is no algorithm for solving the equation $M_1^{x_1} \dots M_k^{x_k} = Z$, where M_1, \dots, M_k are integer $n \times n$ matrices and Z is the zero matrix. Using the construction of Kronecker (or tensor) product of matrices, it is possible to show that the above-mentioned result implies that Equation (2) is algorithmically undecidable in general for non-commuting integer matrices M_1, \dots, M_k and N_1, \dots, N_l .

However with the help of Proposition 15 we can algorithmically solve Equation (2) in the case when M_1, \dots, M_k and N_1, \dots, N_l are matrices from $SL(2, \mathbb{Z})$ and the vectors \mathbf{x}, \mathbf{y} have integer coefficients. Indeed, since the matrices from $SL(2, \mathbb{Z})$ are invertible, we can rewrite (2) as $(N_l^{-1})^{y_l} \dots (N_1^{-1})^{y_1} M_1^{x_1} \dots M_k^{x_k} \mathbf{x} = \mathbf{y}$. It is not hard to see that $\{(N_l^{-1})^{y_l} \dots (N_1^{-1})^{y_1} M_1^{x_1} \dots M_k^{x_k} : x_1, \dots, x_k, y_1, \dots, y_l \in \mathbb{N} \cup \{0\}\}$ is a regular subset of $SL(2, \mathbb{Z})$, and hence the problem is decidable. Using the same idea we can algorithmically solve Equation (2) also in the case when x_1, \dots, x_k and y_1, \dots, y_l are arbitrary integers and the matrices are from $SL(2, \mathbb{Z})$.

In the rest of this section we will give a geometric interpretation of both reachability problems (Figure 1), which we will use later to solve a special case of the scalar reachability problem (Proposition 18).

► **Proposition 16.** *According to Theorem 8, the set of matrices M from $SL(2, \mathbb{Z})$ that transform a vector $\mathbf{x} = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$ to a vector $\mathbf{y} = \begin{bmatrix} y_1 \\ y_2 \end{bmatrix}$ has the form $\mathcal{F} = \{BT^{kt}C : t \in \mathbb{Z}\}$.*

Consider the equation $BT^{kt}C\mathbf{x} = \mathbf{y}$ and let us make the following change of variables: $\mathbf{u} = C\mathbf{x}$ and $\mathbf{v} = B^{-1}\mathbf{y}$: $\mathbf{x} \xrightarrow{C} \mathbf{u} \xrightarrow{T^{kt}} \mathbf{v} \xrightarrow{B} \mathbf{y}$. Then $\mathbf{u} = \mathbf{v} = \begin{bmatrix} d \\ 0 \end{bmatrix}$, where $|d| = \gcd(x_1, x_2) = \gcd(y_1, y_2)$.

Proof. In the new notations, the equation $BT^{kt}C\mathbf{x} = \mathbf{y}$ can be written as $T^{kt}\mathbf{u} = \mathbf{v}$, and this equality holds for any $t \in \mathbb{Z}$. Now let $\mathbf{u} = \begin{bmatrix} u_1 \\ u_2 \end{bmatrix}$ and $\mathbf{v} = \begin{bmatrix} v_1 \\ v_2 \end{bmatrix}$. Hence we have

$\begin{bmatrix} 1 & kt \\ 0 & 1 \end{bmatrix} \begin{bmatrix} u_1 \\ u_2 \end{bmatrix} = \begin{bmatrix} v_1 \\ v_2 \end{bmatrix}$, which is equivalent to $u_2 = v_2$ and $u_1 + kt u_2 = v_1$, for any $t \in \mathbb{Z}$. So, we must have $u_2 = v_2 = 0$ and hence $u_1 = v_1$.

Therefore, the vectors \mathbf{u} and \mathbf{v} have the form $\mathbf{u} = \mathbf{v} = \begin{bmatrix} d \\ 0 \end{bmatrix}$ for some $d \in \mathbb{Z}$. Moreover, since $\mathbf{u} = C\mathbf{x}$, we obtain from Lemma 4 that $|d| = \gcd(x_1, x_2) = \gcd(y_1, y_2)$. ◀

We can give the following geometric interpretation of the transformation $BT^tC\mathbf{x} = \mathbf{y}$: first, we apply C to \mathbf{x} and arrive at $\mathbf{u} = \begin{bmatrix} d \\ 0 \end{bmatrix}$, then we loop at \mathbf{u} for t many times using T , and finally apply B to move from \mathbf{u} to \mathbf{y} (see Figure 1 on the left).

Similarly, we have the following geometric interpretation of the fractional linear transformation $y = f_{BT^tC}(x) = f_B \circ f_{T^t} \circ f_C(x)$: first it maps x to ∞ using f_C , then loops at ∞ for t many times using f_T , and finally maps ∞ to y using f_B (see Figure 1 on the right).

We now show how to apply the geometric interpretation of the vector reachability problem to solve a special case of the scalar reachability problem.




Figure 1 Geometric interpretation of the linear transformation $\mathbf{y} = BT^tC\mathbf{x}$ (left) and of the fractional linear transformation $y = f_{BT^tC}(x)$ (right).

► **Definition 17.** The *scalar reachability problem* in $SL(2, \mathbb{Z})$ is stated as follows: Let $[z_1, z_2]$ and $\begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$ be vectors from $\mathbb{Z} \times \mathbb{Z}$ and let λ be an integer number. We are also given a finite collection of matrices M_1, \dots, M_n from $SL(2, \mathbb{Z})$. The question is to decide whether there exists a matrix $M \in \langle M_1, \dots, M_n \rangle$ which satisfies the equation $[z_1, z_2] M \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \lambda$.

We will consider a special case of this problem when $z_2 = 1$ and $\lambda = 1$. Our proof relies on the characterization from Theorem 8 and Corollary 9 and on Proposition 13 in which we showed that the emptiness problem for the intersection of two regular subsets in $SL(2, \mathbb{Z})$ is decidable.

► **Proposition 18.** *Suppose that the above equation has the form*

$$[a, 1] M \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = 1, \tag{3}$$

where a, x_1 and x_2 are some integer numbers. Then this special case of the scalar reachability problem is decidable.

Proof. The general idea of the proof is the same as in Theorem 14, that is, we will show that the set of matrices $M \in SL(2, \mathbb{Z})$ that satisfy Equation (3) can be described by a regular language. First, let us consider a geometric interpretation of this problem. We can rewrite Equation (3) as a system of two equations: $M \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} y_1 \\ y_2 \end{bmatrix}$ and $ay_1 + y_2 = 1$. So, M satisfies Equation (3) if and only if it maps a fixed vector $\mathbf{x} = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$ to some vector $\mathbf{y} = \begin{bmatrix} y_1 \\ y_2 \end{bmatrix}$ that lies on the line L described by the equation $ay_1 + y_2 = 1$. In other words, we have a *vector to line reachability problem* for the line L that is defined by the equation $ay_1 + y_2 = 1$.

Note that if a vector \mathbf{y} lies of the line $ay_1 + y_2 = 1$, then $\gcd(y_1, y_2) = 1$. Hence by Lemma 4, Equation (3) has a solution only if $\gcd(x_1, x_2) = 1$. So, from now on we assume that $\gcd(x_1, x_2) = 1$.

By Corollary 9, any $M \in SL(2, \mathbb{Z})$ that maps \mathbf{x} to a vector \mathbf{y} on the line L has the form $M = BT^tC$, where B and C are some matrices from $SL(2, \mathbb{Z})$ and $t \in \mathbb{Z}$. Geometrically, the transformation $\mathbf{y} = BT^tC\mathbf{x}$ goes via the point $(1, 0)$ as shown in Figure 2.

Note that the matrices B and C above depend on the vector \mathbf{y} as a parameter. Here we prove a useful lemma which will imply that we can choose only one matrix C that maps \mathbf{x} to $\begin{bmatrix} 1 \\ 0 \end{bmatrix}$ independently of the vector \mathbf{y} .


► **Lemma 19.** Let $\mathbf{x} = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$ and $\mathbf{y} = \begin{bmatrix} y_1 \\ y_2 \end{bmatrix}$ be any vectors from $\mathbb{Z} \times \mathbb{Z}$ such that $\gcd(x_1, x_2) = \gcd(y_1, y_2) = d$. Let d_1 and d_2 be any integer numbers with $|d_1| = |d_2| = d$ and let A_1, B_1 and A_2, B_2 by any matrices from $SL(2, \mathbb{Z})$ such that $B_i \mathbf{x} = \begin{bmatrix} d_i \\ 0 \end{bmatrix}$ and $A_i \begin{bmatrix} d_i \\ 0 \end{bmatrix} = \mathbf{y}$, for $i = 1, 2$. Then $\{A_1 T^t B_1 : t \in \mathbb{Z}\} = \{A_2 T^t B_2 : t \in \mathbb{Z}\}$. In other words, the following diagrams define the same set of matrices that map \mathbf{x} to \mathbf{y} .

$$\mathbf{x} \xrightarrow{B_1} \begin{bmatrix} d_1 \\ 0 \end{bmatrix} \xrightarrow{A_1} \mathbf{y} \qquad \mathbf{x} \xrightarrow{B_2} \begin{bmatrix} d_2 \\ 0 \end{bmatrix} \xrightarrow{A_2} \mathbf{y}$$

$\uparrow T^t$
 $\uparrow T^t$

Proof. See Section C of the Appendix in [31]. ◀

By Lemma 19, we can choose any matrix C from $SL(2, \mathbb{Z})$ that maps a vector \mathbf{x} to the vector $\begin{bmatrix} 1 \\ 0 \end{bmatrix}$, and for each \mathbf{y} on the line L we can choose any matrix $B_{\mathbf{y}}$ that maps $\begin{bmatrix} 1 \\ 0 \end{bmatrix}$ to the vector \mathbf{y} . Then the solution of Equation (3) will be described by the following set $\mathcal{F} = \{B_{\mathbf{y}} T^t C : \mathbf{y} \in L \text{ and } t \in \mathbb{Z}\}$. Figure 2 gives geometric interpretation of this solution.



■ **Figure 2** Geometric interpretation of the scalar reachability problem.

We need to choose $B_{\mathbf{y}}$ in such a way that \mathcal{F} becomes a regular set. Let $\mathbf{y} = \begin{bmatrix} y_1 \\ y_2 \end{bmatrix} \in L$, then we have $ay_1 + y_2 = 1$. As one can check, if we let $B_{\mathbf{y}} = \begin{bmatrix} y_1 & -1 \\ -ay_1 + 1 & a \end{bmatrix}$ then $B_{\mathbf{y}} \in SL(2, \mathbb{Z})$ and $B_{\mathbf{y}} \begin{bmatrix} 1 \\ 0 \end{bmatrix} = \mathbf{y}$.

Since every entry of $B_{\mathbf{y}}$ is a linear function of y_1 , we obtain by Proposition 7 that $B_{\mathbf{y}} = AT^{ky_1}D$, where A and D are some matrices from $SL(2, \mathbb{Z})$ and k is some integer number (in fact, one can show that $k = 1$). Finally, we can write all solutions of Equation (3) as $\mathcal{F} = \{AT^{ky_1}DT^tC : y_1 \in \mathbb{Z} \text{ and } t \in \mathbb{Z}\}$. This is clearly a regular set and, therefore, the scalar reachability problem is decidable. ◀

References

- 1 László Babai, Robert Beals, Jin-yi Cai, Gábor Ivanyos, and Eugene M. Luks. Multiplicative equations over commuting matrices. In *Proceedings of the Seventh Annual ACM-SIAM Symposium on Discrete Algorithms*, SODA '96, pages 498–507, Philadelphia, PA, USA, 1996. Society for Industrial and Applied Mathematics.

- 2 Paul Bell, Vesa Halava, Tero Harju, Juhani Karhumäki, and Igor Potapov. Matrix equations and Hilbert's tenth problem. *Internat. J. Algebra Comput.*, 18(8):1231–1241, 2008.
- 3 Paul Bell and Igor Potapov. On undecidability bounds for matrix decision problems. *Theoretical Computer Science*, 391(1-2):3–13, 2008.
- 4 Paul Bell and Igor Potapov. Reachability problems in quaternion matrix and rotation semigroups. *Information and Computation*, 206(11):1353–1361, 2008.
- 5 Paul C. Bell, Mika Hirvensalo, and Igor Potapov. Mortality for 2x2 matrices is NP-hard. In Branislav Rován, Vladimiro Sassone, and Peter Widmayer, editors, *Mathematical Foundations of Computer Science 2012*, volume 7464 of *Lecture Notes in Computer Science*, pages 148–159. Springer Berlin Heidelberg, 2012.
- 6 Paul C. Bell and Igor Potapov. On the computational complexity of matrix semigroup problems. *Fundam. Inf.*, 116(1-4):1–13, January 2012.
- 7 Vincent D. Blondel, Emmanuel Jeandel, Pascal Koiran, and Natacha Portier. Decidable and undecidable problems about quantum automata. *SIAM J. Comput.*, 34(6):1464–1473, June 2005.
- 8 Julien Cassaigne, Tero Harju, and Juhani Karhumäki. On the undecidability of freeness of matrix semigroups. *International Journal of Algebra and Computation*, 09(03n04):295–305, 1999. URL: <http://www.worldscientific.com/doi/pdf/10.1142/S0218196799000199>.
- 9 Julien Cassaigne and François Nicolas. On the decidability of semigroup freeness. *RAIRO - Theor. Inf. and Applic.*, 46(3):355–399, 2012.
- 10 Fernando Chamizo. Non-euclidean visibility problems. *Proceedings of the Indian Academy of Sciences - Mathematical Sciences*, 116(2):147–160, 2006.
- 11 Christian Choffrut and Juhani Karhumäki. Some decision problems on integer matrices. *RAIRO-Theor. Inf. Appl.*, 39(1):125–131, 2005.
- 12 J. Elstrodt, F. Grunewald, and J. Mennicke. Arithmetic applications of the hyperbolic lattice point theorem. *Proc. London Math. Soc.*, s3-57:pp.239–283, 1988.
- 13 J. Esparza, A. Finkel, and R. Mayr. On the verification of broadcast protocols. In *Logic in Computer Science, 1999. Proceedings. 14th Symposium on*, pages 352–359, 1999.
- 14 Esther Galby, Joël Ouaknine, and James Worrell. On Matrix Powering in Low Dimensions. In Ernst W. Mayr and Nicolas Ollinger, editors, *32nd International Symposium on Theoretical Aspects of Computer Science (STACS 2015)*, volume 30 of *Leibniz International Proceedings in Informatics (LIPIcs)*, pages 329–340, Dagstuhl, Germany, 2015. Schloss Dagstuhl–Leibniz-Zentrum fuer Informatik.
- 15 M. P. García del Moral, I. Martín, J. M. Peña, and A. Restuccia. $Sl(2, z)$ symmetries, supermembranes and symplectic torus bundles. *Journal of High Energy Physics*, 2011(9):1–12, 2011.
- 16 Yuri Gurevich and Paul Schupp. Membership problem for the modular group. *SIAM J. Comput.*, 37(2):425–459, May 2007.
- 17 Vesa Halava, Tero Harju, and Mika Hirvensalo. Undecidability bounds for integer matrices using Claus instances. *International Journal of Foundations of Computer Science*, 18(05):931–948, 2007. URL: <http://www.worldscientific.com/doi/pdf/10.1142/S0129054107005066>.
- 18 Vesa Halava, Tero Harju, Mika Hirvensalo, and Juhani Karhumäki. Skolem's problem - on the border between decidability and undecidability. Technical Report 683, Turku Centre for Computer Science, 2005.
- 19 Ravindran Kannan and Achim Bachem. Polynomial algorithms for computing the Smith and Hermite normal forms of an integer matrix. *SIAM J. Comput.*, 8(4):499–507, 1979.
- 20 Alexei Lisitsa and Igor Potapov. Membership and reachability problems for row-monomial transformations. In *Mathematical Foundations of Computer Science 2004, 29th Interna-*

- tional Symposium, MFCS 2004, Prague, Czech Republic, August 22-27, 2004, Proceedings*, pages 623–634, 2004.
- 21 Roger C. Lyndon and Paul E. Schupp. *Combinatorial group theory*. Springer-Verlag, Berlin-New York, 1977. *Ergebnisse der Mathematik und ihrer Grenzgebiete, Band 89*.
 - 22 Dana Mackenzie. A new twist in knot theory. *What's Happening in the Mathematical Sciences*, Volume 7, 2009.
 - 23 Wilhelm Magnus, Abraham Karrass, and Donald Solitar. *Combinatorial group theory*. Dover Publications, Inc., New York, revised edition, 1976. Presentations of groups in terms of generators and relations.
 - 24 A. Markov. On certain insoluble problems concerning matrices. *Doklady Akad. Nauk SSSR*, 57(6):539–542, June 1947.
 - 25 Thomas Noll. Musical intervals and special linear transformations. *Journal of Mathematics and Music: Mathematical and Computational Approaches to Music Theory, Analysis, Composition and Performance*, vol.1, 2007.
 - 26 Joël Ouaknine, João Sousa Pinto, and James Worrell. On termination of integer linear loops. In *Proceedings of the Twenty-Sixth Annual ACM-SIAM Symposium on Discrete Algorithms*, SODA '15, pages 957–969. SIAM, 2015.
 - 27 Joël Ouaknine and James Worrell. On the positivity problem for simple linear recurrence sequences. In *Automata, Languages, and Programming - 41st International Colloquium, ICALP 2014, Copenhagen, Denmark, July 8-11, 2014, Proceedings, Part II*, pages 318–329, 2014.
 - 28 Joël Ouaknine and James Worrell. Ultimate positivity is decidable for simple linear recurrence sequences. In *Automata, Languages, and Programming - 41st International Colloquium, ICALP 2014, Copenhagen, Denmark, July 8-11, 2014, Proceedings, Part II*, pages 330–341, 2014.
 - 29 Leonid Polterovich and Zeev Rudnick. Stable mixing for cat maps and quasi-morphisms of the modular group. *Ergodic Theory and Dynamical Systems*, 24:609–619, 4 2004.
 - 30 Igor Potapov. Composition problems for braids. In *33rd International Conference on Foundations of Software Technology and Theoretical Computer Science*, volume 24 of *LIPICs. Leibniz Int. Proc. Inform.*, pages 175–187. Schloss Dagstuhl. Leibniz-Zent. Inform., Wadern, 2013.
 - 31 Igor Potapov and Pavel Semukhin. Vector reachability problem in $SL(2, \mathbb{Z})$. *CoRR*, abs/1510.03227, 2015. URL: <http://arxiv.org/abs/1510.03227>.
 - 32 Robert A. Rankin. *Modular forms and functions*. Cambridge University Press, Cambridge-New York-Melbourne, 1977.
 - 33 Edward Witten. $SL(2, \mathbb{Z})$ action on three-dimensional conformal field theories with abelian symmetry. In *From fields to strings: circumnavigating theoretical physics. Vol. 2*, pages 1173–1200. World Sci. Publ., Singapore, 2005.
 - 34 Don Zagier. Elliptic modular forms and their applications. In Kristian Ranestad, editor, *The 1-2-3 of Modular Forms*, Universitext, pages 1–103. Springer Berlin Heidelberg, 2008.