

EUROPEAN ORGANIZATION FOR NUCLEAR RESEARCH (CERN)

CERN-EP-2016-048  
LHCb-PAPER-2016-005  
June 16, 2016

# Search for violations of Lorentz invariance and $CPT$ symmetry in $B_{(s)}^0$ mixing

The LHCb collaboration<sup>†</sup>

## Abstract

Violations of  $CPT$  symmetry and Lorentz invariance are searched for by studying interference effects in  $B^0$  mixing and in  $B_s^0$  mixing. Samples of  $B^0 \rightarrow J/\psi K_s^0$  and  $B_s^0 \rightarrow J/\psi K^+ K^-$  decays are recorded by the LHCb detector in proton–proton collisions at centre-of-mass energies of 7 and 8 TeV, corresponding to an integrated luminosity of  $3 \text{ fb}^{-1}$ . No periodic variations of the particle–antiparticle mass differences are found, consistent with Lorentz invariance and  $CPT$  symmetry. Results are expressed in terms of the Standard Model Extension parameter  $\Delta a_\mu$  with precisions of  $\mathcal{O}(10^{-15})$  and  $\mathcal{O}(10^{-14})$  GeV for the  $B^0$  and  $B_s^0$  systems, respectively. With no assumption on Lorentz (non-)invariance, the  $CPT$ -violating parameter  $z$  in the  $B_s^0$  system is measured for the first time and found to be  $\text{Re}(z) = -0.022 \pm 0.033 \pm 0.005$  and  $\text{Im}(z) = 0.004 \pm 0.011 \pm 0.002$ , where the first uncertainties are statistical and the second systematic.

Published in Phys. Rev. Lett. 116 (2016) 241601

© CERN on behalf of the LHCb collaboration, licence CC-BY-4.0.

---

<sup>†</sup>Authors are listed at the end of this Letter.



Lorentz invariance and the combination of charge conjugation, spatial inversion and time reversal ( $CPT$ ) are exact symmetries in the Standard Model (SM) of particle physics, and are deeply connected in any quantum field theory [1]. Quantum theories that aim to describe Planck-scale physics, such as string theory, might break these fundamental symmetries [2]. Present-day experiments are many orders of magnitude away from the Planck energy scale of  $\sim 10^{19}$  GeV, however, small effects at low energy might still be observable. Interference effects in the mixing of neutral mesons are sensitive to violations of  $CPT$  symmetry, and therefore may provide a window to the quantum gravity scale [3]. Such effects can be quantified in a low-energy, effective field theory, as done in the Standard Model Extension (SME) [4, 5]. In this framework, terms that explicitly break Lorentz and  $CPT$  symmetry are added to the SM Lagrangian to describe the couplings between particles and (hypothetical) uniform tensor fields. These fields would acquire non-zero vacuum expectation values when these symmetries are spontaneously broken in the underlying theory. The SME couplings are expected to be suppressed by powers of the Planck scale [6]. In the SME, the  $CPT$ -violating parameters that can be measured in neutral meson systems also break Lorentz symmetry. The amount of  $CPT$  violation depends on the direction of motion and on the boost of the particle. The SME parameters for the  $B^0$  and  $B_s^0$  systems can be best measured with a time-dependent analysis of the decay channels  $B^0 \rightarrow J/\psi K_s^0$  and  $B_s^0 \rightarrow J/\psi K^+ K^-$ , using the four-velocity of the  $B$  mesons [7]. The notation  $B$  refers to either  $B^0$  or  $B_s^0$  and the inclusion of charge-conjugate processes is implied throughout this Letter. These parameters have been measured previously, albeit with less sensitive decay modes [7], by the BaBar collaboration for the  $B^0$  system [8], and by the D0 collaboration for the  $B_s^0$  system [9].

The LHCb detector is a single-arm forward spectrometer described in detail in Refs. [10, 11]. Simulated events are produced using the software described in Refs. [12–16]. The data used in this analysis correspond to an integrated luminosity of  $3 \text{ fb}^{-1}$ , taken at the LHC at proton–proton centre-of-mass energies of 7 and 8 TeV. The selection of both decay channels is the same as used in Refs. [17] and [18]. The  $J/\psi$  meson is reconstructed in the dimuon channel and the  $K_s^0$  meson in the  $\pi^+\pi^-$  final state.

Interference effects from  $CPT$  violation can be incorporated generically in the time evolution of a neutral  $B$  meson system, described by the Schrödinger equation  $i\partial_t\Psi = \hat{H}\Psi$ . The effective  $2 \times 2$  Hamiltonian is written as  $\hat{H} = \hat{M} - i\hat{\Gamma}/2$  [19]. Diagonalisation gives a heavy-mass eigenstate  $|B_H\rangle$  and a light-mass eigenstate  $|B_L\rangle$  with masses  $m_{H,L}$  and decay widths  $\Gamma_{H,L}$ . The differences between the eigenvalues are defined as  $\Delta m \equiv m_H - m_L$  and  $\Delta\Gamma \equiv \Gamma_L - \Gamma_H$ . The differences between the diagonal matrix elements of the effective Hamiltonian are defined as  $\delta m \equiv M_{11} - M_{22}$  and  $\delta\Gamma \equiv \Gamma_{11} - \Gamma_{22}$ . Any difference between the mass or lifetime of particles and antiparticles (*i.e.*, a non-zero  $\delta m$  or  $\delta\Gamma$ ) would be a sign of  $CPT$  violation, and is characterised by

$$z = \frac{\delta m - i\delta\Gamma/2}{\Delta m + i\Delta\Gamma/2}, \quad (1)$$

and the mass eigenstates are given by  $|B_{H,L}\rangle = p\sqrt{1 \pm z}|B\rangle \mp q\sqrt{1 \mp z}|\bar{B}\rangle$ . Owing to the smallness of the  $B$  mixing parameters  $\Delta m$  and  $\Delta\Gamma$  in the denominator,  $z$  is highly

sensitive to  $CPT$ -violating effects.

Considering only contributions to first order in  $z$ , the decay rate to a  $CP$  eigenstate  $f$  as a function of the  $B$  proper decay time  $t$  becomes

$$\begin{aligned} \frac{d\Gamma_f}{dt} \propto e^{-\Gamma t} & \left[ [1 + \zeta D_f \mathcal{R}e(z) - S_f \mathcal{I}m(z)] \cosh(\Delta\Gamma t/2) \right. \\ & + [D_f + \mathcal{R}e(z)(C_f + \zeta)] \sinh(\Delta\Gamma t/2) \\ & + \zeta [C_f - D_f \mathcal{R}e(z) + \zeta S_f \mathcal{I}m(z)] \cos(\Delta m t) \\ & \left. - \zeta [S_f - \mathcal{I}m(z)(C_f + \zeta)] \sin(\Delta m t) \right], \end{aligned} \quad (2)$$

where  $\Gamma \equiv (\Gamma_{11} + \Gamma_{22})/2$ ,  $\zeta = +1(-1)$  for an initial  $|B\rangle$  ( $|\bar{B}\rangle$ ) state and the following definitions are introduced:

$$C_f \equiv \frac{1 - |\lambda_f|^2}{1 + |\lambda_f|^2}, \quad S_f \equiv \frac{2\mathcal{I}m(\lambda_f)}{1 + |\lambda_f|^2}, \quad D_f \equiv -\frac{2\mathcal{R}e(\lambda_f)}{1 + |\lambda_f|^2}, \quad \lambda_f \equiv \frac{q \bar{A}_f}{p A_f}, \quad (3)$$

with  $A_f$  and  $\bar{A}_f$  the direct decay amplitudes of a  $|B\rangle$  or  $|\bar{B}\rangle$  state to the eigenstate  $f$ .

For the decay  $B^0 \rightarrow J/\psi K_s^0$ , the final state is  $CP$  odd, corresponding to  $CP$  eigenvalue  $\eta_f = -1$ . In the SM,  $\arg(\lambda_{J/\psi K_s^0}) = \pi - 2\beta$ , where  $\beta$  is defined in terms of elements of the CKM matrix as  $\beta \equiv \arg[-(V_{cd}V_{cb}^*)/(V_{td}V_{tb}^*)]$ . Furthermore, in the  $B^0$  system, the approximation  $\Delta\Gamma_d = 0$  is made, as supported by experimental data [20].

The decay  $B_s^0 \rightarrow J/\psi K^+ K^-$  is similar to  $B^0 \rightarrow J/\psi K_s^0$ , but the decay width difference  $\Delta\Gamma_s$  cannot be ignored [20]. Another important difference is that the  $K^+ K^-$  system mostly originates from the  $\phi(1020)$  resonance, giving the  $K^+ K^-$  pair an orbital angular momentum  $L = 1$  (P wave). Since the  $J/\psi \phi$  final state consists of two vector mesons, its orbital angular momentum can be  $L \in \{0, 1, 2\}$  for the polarisation states  $f \in \{0, \perp, \parallel\}$ , respectively, with corresponding  $CP$  eigenvalues  $\eta_f = (-1)^L$ . The  $K^+ K^-$  system has a small S-wave contribution [18], which results in another  $L = 1$  component for the  $J/\psi K^+ K^-$  final state. These four polarisation states can be separated statistically in the helicity formalism [21], using the three decay angles between the final-state particles. The corresponding weak phases,  $\arg(\lambda_{J/\psi K^+ K^-}) = L\pi - \phi_s$ , can, in the SM, be expressed in terms of CKM matrix elements,  $\phi_s = -2\beta_s \equiv -2\arg[-(V_{ts}V_{tb}^*)/(V_{cs}V_{cb}^*)]$ . The decay rate has to be modified compared to Eq. 2 to include the angular dependence. It becomes a sum over all ten combinations of the four helicity amplitudes,

$$\frac{d^4\Gamma_{J/\psi K^+ K^-}}{dt d\vec{\Omega}} \propto \sum_{k=1}^{10} h_k(t) f_k(\vec{\Omega}), \quad (4)$$

where  $f_k(\vec{\Omega})$  are angular functions, given in Ref. [21], and  $h_k(t)$  are products of the

Table 1: Time-dependent functions  $h_k(t)$  in Eq. 4.

$k$	$h_k(t)$	$k$	$h_k(t)$
1	$ A_0(t) ^2$	6	$\mathcal{I}m(A_0^*(t)A_\perp(t))$
2	$ A_\parallel(t) ^2$	7	$ A_S(t) ^2$
3	$ A_\perp(t) ^2$	8	$\mathcal{R}e(A_S^*(t)A_\parallel(t))$
4	$\mathcal{I}m(A_\parallel^*(t)A_\perp(t))$	9	$\mathcal{I}m(A_S^*(t)A_\perp(t))$
5	$\mathcal{R}e(A_0^*(t)A_\parallel(t))$	10	$\mathcal{R}e(A_S^*(t)A_0(t))$

Table 2: Definition of the coefficients in Eq. 5. The following definitions are used:  $\eta^+ \equiv (1 + \eta_l \eta_m)/2$ ,  $\eta^- \equiv (1 - \eta_l \eta_m)/2$ ,  $\eta^{\mathcal{I}m} \equiv i(\eta_l - \eta_m)/2$ ,  $\eta^{\mathcal{R}e} \equiv (\eta_l + \eta_m)/2$ . Furthermore,  $\zeta^+ \equiv (\zeta)^{\eta^+}$ , and  $\zeta^- \equiv (\zeta)^{\eta^-}$ , such that  $\zeta^\pm = 1$  if  $\eta^\pm = 0$  and  $\zeta^\pm = \zeta$  otherwise.

$$\begin{aligned}
 a_k &= (\eta^+ + \eta^- C_f) + \zeta \mathcal{R}e(z)(\eta^{\mathcal{R}e} D_f + \eta^{\mathcal{I}m} S_f) \\
 &\quad + \mathcal{I}m(z)(\eta^{\mathcal{I}m} D_f - \eta^{\mathcal{R}e} S_f) \\
 b_k &= (\eta^{\mathcal{R}e} D_f + \eta^{\mathcal{I}m} S_f) + \mathcal{R}e(z)(\zeta^+ + \zeta^- C_f) \\
 c_k &= \zeta(\eta^- + \eta^+ C_f) - \zeta \mathcal{R}e(z)(\eta^{\mathcal{R}e} D_f + \eta^{\mathcal{I}m} S_f) \\
 &\quad - \mathcal{I}m(z)(\eta^{\mathcal{I}m} D_f - \eta^{\mathcal{R}e} S_f) \\
 d_k &= \zeta(\eta^{\mathcal{I}m} D_f - \eta^{\mathcal{R}e} S_f) + \mathcal{I}m(z)(\zeta^+ + \zeta^- C_f)
 \end{aligned}$$

amplitudes as listed in Table 1. The time dependence of  $h_k(t)$  is given by

$$\begin{aligned}
 A_l^*(t)A_m(t) &= \frac{A_l^*(0)A_m(0)e^{-\Gamma_s t}}{1 + \zeta C_f} \\
 &\quad \left[ a_k \cosh(\Delta\Gamma_s t/2) + b_k \sinh(\Delta\Gamma_s t/2) \right. \\
 &\quad \left. + c_k \cos(\Delta m_s t) + d_k \sin(\Delta m_s t) \right], \tag{5}
 \end{aligned}$$

with the coefficients listed in Table 2.

In the SME, the dimensionless parameter  $z$  is not a constant. It depends on the four-velocity  $\beta^\mu = (\gamma, \gamma\vec{\beta})$  of the neutral meson as [22, 23]

$$z = \frac{\beta^\mu \Delta a_\mu}{\Delta m + i\Delta\Gamma/2}, \tag{6}$$

thereby breaking Lorentz invariance. The SME parameter  $\Delta a_\mu$  describes the difference between the couplings of the valence quarks, within the neutral meson, with the Lorentz-violating fields [22]. Therefore,  $B^0$  and  $B_s^0$  mesons can have different values of  $\Delta a_\mu$ . Since  $\Delta a_\mu$  is real [24], it follows that  $\mathcal{R}e(z)\Delta\Gamma = -2\mathcal{I}m(z)\Delta m$ . For  $B$  mesons,  $\Delta m \gg \Delta\Gamma$ , and so  $\mathcal{I}m(z)$  is two orders of magnitude smaller than  $\mathcal{R}e(z)$ , and can be ignored in the measurements of  $\Delta a_\mu$ . The average boost of  $B$  mesons in the acceptance of LHCb is  $\langle\gamma\beta\rangle \approx 20$ . It follows from Eq. 6 that this large boost results in a high sensitivity to  $\Delta a_\mu$  [7].

To measure  $\Delta a_\mu$ , the meson direction needs to be determined in an absolute reference frame. Such a frame can be defined with respect to fixed stars [24]. In this frame, the  $Z$ -axis points north along the Earth's rotation axis, the  $X$ -axis points from the Sun to the vernal equinox on 1 January 2000 (J2000 epoch) and the  $Y$ -axis completes the right-handed coordinate system. The latitude of the LHCb interaction point is  $46.2414^\circ\text{N}$ , the longitude is  $6.0963^\circ\text{E}$ , and the angle of the beam east of north is  $236.296^\circ$ . The beam axis is inclined with respect to the geodetic plane by  $3.601$  mrad, pointing slightly upwards. The timekeeping is obtained from the LHC machine with a time stamp,  $t_{\text{LHC}}$ , in UTC microseconds since January 1<sup>st</sup>, 1970, 00:00:00 UTC. The time, spatial coordinates and angles have negligible uncertainties and are used to define the rotation from the coordinate system of LHCb to the absolute reference frame. For mesons travelling along the beam axis,  $\mathcal{R}e(z)$  can be expressed as,

$$\begin{aligned} \mathcal{R}e(z) &= \frac{\Delta m}{\Delta m^2 + \Delta\Gamma^2/4} \beta^\mu \Delta a_\mu \\ &\approx \frac{\gamma}{\Delta m} \left[ \Delta a_0 + \cos(\chi) \Delta a_Z + \sin(\chi) [\Delta a_Y \sin(\Omega \hat{t}) + \Delta a_X \cos(\Omega \hat{t})] \right], \end{aligned} \quad (7)$$

where  $|\vec{\beta}|$  is set to unity,  $\Delta a^{X,Y,Z} = -\Delta a_{X,Y,Z}$ , and  $\chi = 112.4^\circ$  is the angle between the beam axis and the rotational axis of the Earth. The time dependence results from the Earth's rotation, giving a periodicity with sidereal frequency  $\Omega$ . The sidereal phase at  $t_{\text{LHC}} = 0$  is found to be  $\hat{t} = (2.8126 \pm 0.0014)$  hr. The  $B$  mesons are emitted at an average angle of about  $5^\circ$  from the beam axis. This means that the LHCb detector is mostly sensitive to the linear combination  $\Delta a_{\parallel} \equiv \Delta a_0 + \cos(\chi) \Delta a_Z = \Delta a_0 - 0.38 \Delta a_Z$ , while there is a much weaker sensitivity to the orthogonal parameter,  $\Delta a_{\perp} = 0.38 \Delta a_0 + \Delta a_Z$ , coming from the smaller transverse component of the  $B$  velocity. Both  $\Delta a_{\parallel}$  and  $\Delta a_{\perp}$  are measured and the correlation between them is negligible.

Unbinned likelihood fits are applied to the decay-time distributions using Eqs. 2 and 4. To obtain the SME parameters, the sidereal variation of  $\mathcal{R}e(z)$  is taken into account by including in the fits the LHC time and the three-momentum of the reconstructed  $B$  candidate. For the  $B_s^0$  sample, the fits are performed to the full angular distribution.

In the invariant mass distributions of the  $B$  candidates, the background is mostly combinatorial. For both decay channels, this background is statistically subtracted using the *sPlot* technique [25], which allows to project out the signal component by weighting each event depending on the mass of the  $B$  candidate. The mass models are the same as in Refs. [17, 18]. The correlation between the shape of the invariant mass distribution and the  $B$  momentum or, for the  $B_s^0$  sample, the decay angles, leads to a small systematic bias for both samples. This effect is included in the systematic uncertainty. In the  $B_s^0 \rightarrow J/\psi K^+ K^-$  sample, there is a small contribution coming from misidentified  $B^0 \rightarrow J/\psi K^+ \pi^-$  and  $A_b^0 \rightarrow J/\psi p K^-$  decays. This background contribution is statistically removed by adding simulated decays with negative weights. A systematic uncertainty is assigned to account for the uncertainty on the size and shape of this background.

The description of the detection efficiency as a function of the decay time, the decay-time resolution model and the flavour tagging (to distinguish initial  $B$  and  $\bar{B}$  mesons) are

the same as in Refs. [17] and [18] for the  $B^0 \rightarrow J/\psi K_s^0$  and  $B_s^0 \rightarrow J/\psi K^+ K^-$  samples, respectively. This description includes the dilution of the asymmetry due to wrong decisions of the flavour tagging method. The decay-time resolution model and tagging calibration do not lead to a systematic bias in the final result. A possible wrong assignment of the primary interaction vertex (PV) to the  $B$  candidate gives a small bias in the  $\Delta a_{\perp}^{B^0}$  parameter, which is included in the systematic uncertainty. The inefficiency at high decay times, caused by the reconstruction algorithms, is described by an exponential function. For the  $B^0$  sample, this function is obtained from simulation and does not lead to a systematic bias in the result. For the  $B_s^0$  sample, the exponential function is obtained from a data-driven method. The change in the final result when using the correction procedure from Ref. [18] is taken as a systematic uncertainty.

The production asymmetry between  $B^0$  and  $\bar{B}^0$  mesons is included in the modelling of the decay rates, and is taken from Refs. [26,27]. The corresponding uncertainties are included in the statistical uncertainty, while a possible momentum dependence of the production asymmetry is considered as a systematic uncertainty. The  $B_s^0$  production asymmetry does not affect the fit to the  $B_s^0$  sample, since the fast  $B_s^0$  oscillations wash out this effect and since the decay rates for  $B_s^0$  and  $\bar{B}_s^0$  tags are normalised separately.

In the fit to the  $B^0$  sample, the correlation between  $\mathcal{R}e(z^{B^0})$  and  $C_{J/\psi K_s^0}$  is too large to allow determination of  $\mathcal{R}e(z^{B^0})$  without making assumptions about the value of  $C_{J/\psi K_s^0}$  [7]. On the other hand, to determine  $\Delta a_{\mu}^{B^0}$ , the averages  $C_{J/\psi K_s^0} = 0.005 \pm 0.020$  and  $S_{J/\psi K_s^0} = 0.676 \pm 0.021$  [19] as measured by the BaBar and Belle collaborations can be used in the fit. Since the boost of the  $B^0$  mesons is about 40 times lower in these experiments, these values are hardly affected by possible Lorentz violation in the SME. The value of  $D_{J/\psi K_s^0}$  is by definition  $\sqrt{1 - S_{J/\psi K_s^0}^2 - C_{J/\psi K_s^0}^2}$ . The uncertainties on these external input values are propagated as systematic uncertainties on  $\Delta a_{\mu}^{B^0}$ . The mass difference,  $\Delta m_d = 0.510 \pm 0.003 \text{ ps}^{-1}$  [19], is allowed to vary in the fit within its uncertainty using a Gaussian constraint. Setting  $\Delta\Gamma_d = 0.007 \text{ ps}^{-1}$ , which corresponds to the experimental uncertainty [20], leads to a small change in  $\Delta a_{\parallel}^{B^0}$ , which is included in the systematic uncertainty. The  $B^0$  lifetime is allowed to vary freely in the fit.

In the fit to the  $B_s^0$  sample, the correlation between  $\mathcal{R}e(z^{B_s^0})$  and  $C_{J/\psi K^+ K^-}$  is small owing to the additional interference terms from the helicity amplitudes, the non-zero  $\Delta\Gamma_s$  and the faster  $B_s^0 - \bar{B}_s^0$  oscillations. For this reason, the same parameters as in Ref. [18] are varied freely in the fit, in addition to either  $\Delta a_{\mu}$  or  $z$ . The detection efficiency is also described as a function of the decay angles. The shape of this angular acceptance is obtained from simulation. The simulated events are weighted to match the kinematic distributions in data. The uncertainty due to the limited number of simulated events and the full effect of correcting for the kinematic distributions in data are added to the systematic uncertainty. Systematic effects due to the decay-angle resolution are negligibly small. The fit to the  $B_s^0$  sample is performed simultaneously in bins of the  $K^+ K^-$  invariant mass [18]. Each bin has a different interference between the P- and S-wave amplitudes. This effect is included in the fit and no systematic biases are observed.

An overview of the systematic uncertainties is given in Table 3. For the  $B^0$  mixing,

Table 3: Systematic uncertainties on  $\Delta a_\mu$  for  $B^0$  mixing and on  $\Delta a_\mu$  and  $z$  for  $B_s^0$  mixing. Contributions marked with – are negligible.

<b><math>B^0</math> mixing</b>	$\Delta a_\parallel$	$\Delta a_\perp$	$\Delta a_{X,Y}$
Source	[ $\times 10^{-15}$ GeV]		
Mass correlation	–	–	0.04
Wrong PV assignment	–	1	–
Production asymmetry	0.28	1	0.05
External input $C_f, S_f$	0.46	4	0.28
Decay width difference	0.07	–	–
Neutral kaon asymmetry	–	1	–
Quadratic sum	0.54	4	0.29

<b><math>B_s^0</math> mixing</b>	$\Delta a_\parallel$	$\Delta a_\perp$	$\Delta a_{X,Y}$	$\mathcal{R}e(z)$	$\mathcal{I}m(z)$
Source	[ $\times 10^{-14}$ GeV]				
Mass correlation	0.10	3	0.24	0.001	0.002
Peaking background	0.14	3	0.15	0.003	–
Decay-time acceptance	0.30	7	0.65	–	0.001
Angular acceptance	0.07	–	–	0.002	0.001
Quadratic sum	0.36	8	0.71	0.003	0.002

the largest contribution comes from the uncertainty on the external parameters  $C_f$  and  $S_f$ . A small systematic bias is observed in  $\Delta a_\perp^{B^0}$  due to the momentum dependence of the cross-sections of neutral kaons in the detector material. For the  $B_s^0$  mixing, the largest contribution comes from the description of the decay-time acceptance. Effects from the correlation between the mass and decay time and from the accuracy of the length scale and momentum scale of the detector are found to be negligible.

The components of the SME parameter  $\Delta a_\mu$  for  $B^0$  mixing, obtained from the fit to the sample of selected  $B^0 \rightarrow J/\psi K_s^0$  candidates, are

$$\begin{aligned}
 \Delta a_\parallel^{B^0} &= (-0.10 \pm 0.82 \text{ (stat)} \pm 0.54 \text{ (syst)}) \times 10^{-15} \text{ GeV} , \\
 \Delta a_\perp^{B^0} &= (-0.20 \pm 0.22 \text{ (stat)} \pm 0.04 \text{ (syst)}) \times 10^{-13} \text{ GeV} , \\
 \Delta a_X^{B^0} &= (+1.97 \pm 1.30 \text{ (stat)} \pm 0.29 \text{ (syst)}) \times 10^{-15} \text{ GeV} , \\
 \Delta a_Y^{B^0} &= (+0.44 \pm 1.26 \text{ (stat)} \pm 0.29 \text{ (syst)}) \times 10^{-15} \text{ GeV} ,
 \end{aligned}$$






Figure 1: Values of  $\mathcal{R}e(z)$  obtained from fits in bins of sidereal phase for (top) the  $B^0$  sample and (bottom) the  $B_s^0$  sample. The solid line shows the variation of  $\mathcal{R}e(z)$  from the  $\Delta a_\mu$  fits, using the average  $B$  momentum.

and the corresponding numbers for  $B_s^0$  mixing, using  $B_s^0 \rightarrow J/\psi K^+ K^-$  candidates, are

$$\begin{aligned} \Delta a_{\parallel}^{B_s^0} &= (-0.89 \pm 1.41 \text{ (stat)} \pm 0.36 \text{ (syst)}) \times 10^{-14} \text{ GeV} , \\ \Delta a_{\perp}^{B_s^0} &= (-0.47 \pm 0.39 \text{ (stat)} \pm 0.08 \text{ (syst)}) \times 10^{-12} \text{ GeV} , \\ \Delta a_X^{B_s^0} &= (+1.01 \pm 2.08 \text{ (stat)} \pm 0.71 \text{ (syst)}) \times 10^{-14} \text{ GeV} , \\ \Delta a_Y^{B_s^0} &= (-3.83 \pm 2.09 \text{ (stat)} \pm 0.71 \text{ (syst)}) \times 10^{-14} \text{ GeV} . \end{aligned}$$

Figure 1 shows the result of fits of  $\mathcal{R}e(z)$  in bins of the sidereal phase for both samples. For the  $B^0$  sample, the external constraints on  $C_{J/\psi K_S^0}$  and  $S_{J/\psi K_S^0}$  are again used. No sidereal variation is observed. Independently of any assumption of Lorentz violation, the complex  $CPT$ -violating parameter  $z$  in the  $B_s^0$  system is found to be

$$\begin{aligned} \mathcal{R}e(z^{B_s^0}) &= -0.022 \pm 0.033 \text{ (stat)} \pm 0.003 \text{ (syst)} , \\ \mathcal{I}m(z^{B_s^0}) &= 0.004 \pm 0.011 \text{ (stat)} \pm 0.002 \text{ (syst)} . \end{aligned}$$

Since the SME fits consider only one specific frequency, *i.e.* the sidereal frequency, a wide range of frequencies is scanned by means of the periodogram method. A periodogram gives the spectral power  $P(\nu)$  of a frequency  $\nu$  in a signal sampled at discrete, not necessarily equidistant, times. In this analysis, the Lomb-Scargle periodogram [28] is used, as in the BaBar measurement of  $\Delta a_\mu^{B^0}$  [8].




Figure 2: Periodograms for (top) the  $B^0$  and (bottom) the  $B_s^0$  sample. The insets show a zoom around the solar (red dashed line) and sidereal (blue solid line) frequencies, which have been made by highly oversampling the frequencies in this narrow range.

The periodogram is determined for the term in the decay rates proportional to  $e^{-\Gamma t} \text{Re}(z)$ . Since negative weights cannot be used in the periodogram, the  $B$  mass windows are narrowed to  $5260 < m_{J/\psi K_s^0} < 5300 \text{ MeV}/c^2$  and  $5350 < m_{J/\psi K^+ K^-} < 5390 \text{ MeV}/c^2$  compared to those used in the fits [17, 18]. In total about 5200 frequencies are scanned in a wide range around the sidereal frequency, from  $0.03 \text{ solar-day}^{-1}$  to  $2.10 \text{ solar-day}^{-1}$ . The number of frequencies oversamples the number of independent frequencies by roughly a factor of two, thereby avoiding any undersampling [29]. As the data are unevenly sampled, the false-alarm probability is determined from simulation [29], where the time stamps are taken from data.

The two periodograms are shown in Fig. 2. No significant peaks are found. For the  $B^0$  periodogram, the highest peak  $P(\nu_{\max}) = 8.09$  is found at a frequency of  $1.5507 \text{ solar-day}^{-1}$  and has a false-alarm probability of 0.57. There are 2707 (1559) sampled frequencies with a larger spectral power than the peak at the sidereal (solar) frequency. For the  $B_s^0$  periodogram the highest peak  $P(\nu_{\max}) = 10.85$  is found at a frequency of  $1.3301 \text{ solar-day}^{-1}$  and has a false-alarm probability of 0.06. There are 3386 (2356) frequencies with a larger spectral power than the sidereal (solar) peak. The absence of any signal in the SME fits is confirmed by the absence of significant peaks at the sidereal frequency.

The results presented here are consistent with  $CPT$  symmetry and Lorentz invariance. The measurement of  $\Delta a_\mu^{B^0}$  is an improvement in precision of about three orders of magnitude compared to the one from the BaBar collaboration [8] when the SM prediction

$\Delta\Gamma_d = -0.0027 \text{ ps}^{-1}$  [30] is used to scale their result. The measurement of  $\Delta a_\mu^{B_s^0}$  is an order of magnitude more precise than the one from the D0 collaboration [9] (note the different definition,  $\Delta a_\perp \equiv \sqrt{\Delta a_X^2 + \Delta a_Y^2}$ , in Ref. [9]). The measurement of  $z^{B_s^0}$  is the first direct measurement of this quantity.

## Acknowledgements

We express our gratitude to our colleagues in the CERN accelerator departments for the excellent performance of the LHC. We thank the technical and administrative staff at the LHCb institutes. We acknowledge support from CERN and from the national agencies: CAPES, CNPq, FAPERJ and FINEP (Brazil); NSFC (China); CNRS/IN2P3 (France); BMBF, DFG and MPG (Germany); INFN (Italy); FOM and NWO (The Netherlands); MNiSW and NCN (Poland); MEN/IFA (Romania); MinES and FANO (Russia); MinECo (Spain); SNSF and SER (Switzerland); NASU (Ukraine); STFC (United Kingdom); NSF (USA). We acknowledge the computing resources that are provided by CERN, IN2P3 (France), KIT and DESY (Germany), INFN (Italy), SURF (The Netherlands), PIC (Spain), GridPP (United Kingdom), RRCKI and Yandex LLC (Russia), CSCS (Switzerland), IFIN-HH (Romania), CBPF (Brazil), PL-GRID (Poland) and OSC (USA). We are indebted to the communities behind the multiple open source software packages on which we depend. Individual groups or members have received support from AvH Foundation (Germany), EPLANET, Marie Skłodowska-Curie Actions and ERC (European Union), Conseil Général de Haute-Savoie, Labex ENIGMASS and OCEVU, Région Auvergne (France), RFBR and Yandex LLC (Russia), GVA, XuntaGal and GENCAT (Spain), Herchel Smith Fund, The Royal Society, Royal Commission for the Exhibition of 1851 and the Leverhulme Trust (United Kingdom).

## References

- [1] O. W. Greenberg, *CPT violation implies violation of Lorentz invariance*, Phys. Rev. Lett. **89** (2002) 231602, [arXiv:hep-ph/0201258](#).
- [2] V. A. Kostelecký and S. Samuel, *Spontaneous breaking of Lorentz symmetry in string theory*, Phys. Rev. **D39** (1989) 683.
- [3] S. Liberati, *Tests of Lorentz invariance: A 2013 update*, Class. Quant. Grav. **30** (2013) 133001, [arXiv:1304.5795](#).
- [4] D. Colladay and V. A. Kostelecký, *CPT violation and the standard model*, Phys. Rev. **D55** (1997) 6760, [arXiv:hep-ph/9703464](#).
- [5] D. Colladay and V. A. Kostelecký, *Lorentz violating extension of the standard model*, Phys. Rev. **D58** (1998) 116002, [arXiv:hep-ph/9809521](#).

- [6] V. A. Kostelecký and R. Potting, *CPT and strings*, Nucl. Phys. **B359** (1991) 545; V. A. Kostelecký and R. Potting, *CPT, strings, and meson factories*, Phys. Rev. **D51** (1995) 3923, [arXiv:hep-ph/9501341](#).
- [7] J. van Tilburg and M. van Veghel, *Status and prospects for CPT and Lorentz invariance violation searches in neutral meson mixing*, Phys. Lett. **B742** (2015) 236, [arXiv:1407.1269](#).
- [8] BaBar collaboration, B. Aubert *et al.*, *Search for CPT and Lorentz violation in  $B^0$ - $\bar{B}^0$  oscillations with dilepton events*, Phys. Rev. Lett. **100** (2008) 131802, [arXiv:0711.2713](#).
- [9] D0 collaboration, V. M. Abazov *et al.*, *Search for violation of CPT and Lorentz invariance in  $B_s^0$  meson oscillations*, Phys. Rev. Lett. **115** (2015) 161601, [arXiv:1506.04123](#), [Erratum: Phys. Rev. Lett. **116** (2016) 019901].
- [10] LHCb collaboration, A. A. Alves Jr. *et al.*, *The LHCb detector at the LHC*, JINST **3** (2008) S08005.
- [11] LHCb collaboration, R. Aaij *et al.*, *LHCb detector performance*, Int. J. Mod. Phys. **A30** (2015) 1530022, [arXiv:1412.6352](#).
- [12] T. Sjöstrand, S. Mrenna, and P. Skands, *PYTHIA 6.4 physics and manual*, JHEP **05** (2006) 026, [arXiv:hep-ph/0603175](#); T. Sjöstrand, S. Mrenna, and P. Skands, *A brief introduction to PYTHIA 8.1*, Comput. Phys. Commun. **178** (2008) 852, [arXiv:0710.3820](#).
- [13] I. Belyaev *et al.*, *Handling of the generation of primary events in Gauss, the LHCb simulation framework*, J. Phys. Conf. Ser. **331** (2011) 032047.
- [14] D. J. Lange, *The EvtGen particle decay simulation package*, Nucl. Instrum. Meth. **A462** (2001) 152.
- [15] Geant4 collaboration, J. Allison *et al.*, *Geant4 developments and applications*, IEEE Trans. Nucl. Sci. **53** (2006) 270; Geant4 collaboration, S. Agostinelli *et al.*, *Geant4: A simulation toolkit*, Nucl. Instrum. Meth. **A506** (2003) 250.
- [16] M. Clemencic *et al.*, *The LHCb simulation application, Gauss: Design, evolution and experience*, J. Phys. Conf. Ser. **331** (2011) 032023.
- [17] LHCb collaboration, R. Aaij *et al.*, *Measurement of CP violation in  $B^0 \rightarrow J/\psi K_S^0$  decays*, Phys. Rev. Lett. **115** (2015) 031601, [arXiv:1503.07089](#).
- [18] LHCb collaboration, R. Aaij *et al.*, *Precision measurement of CP violation in  $B_s^0 \rightarrow J/\psi K^+ K^-$  decays*, Phys. Rev. Lett. **114** (2015) 041801, [arXiv:1411.3104](#).
- [19] Particle Data Group, K. A. Olive *et al.*, *Review of particle physics*, Chin. Phys. **C38** (2014) 090001.

- [20] Heavy Flavor Averaging Group, Y. Amhis *et al.*, *Averages of  $b$ -hadron,  $c$ -hadron, and  $\tau$ -lepton properties as of summer 2014*, arXiv:1412.7515, updated results and plots available at <http://www.slac.stanford.edu/xorg/hfag/>.
- [21] LHCb collaboration, R. Aaij *et al.*, *Measurement of CP violation and the  $B_s^0$  meson decay width difference with  $B_s^0 \rightarrow J/\psi K^+ K^-$  and  $B_s^0 \rightarrow J/\psi \pi^+ \pi^-$  decays*, Phys. Rev. **D87** (2013) 112010, arXiv:1304.2600.
- [22] V. A. Kostelecký, *Sensitivity of CPT tests with neutral mesons*, Phys. Rev. Lett. **80** (1998) 1818, arXiv:hep-ph/9809572.
- [23] V. A. Kostelecký, *CPT, T, and Lorentz violation in neutral-meson oscillations*, Phys. Rev. **D64** (2001) 076001, arXiv:hep-ph/0104120.
- [24] V. A. Kostelecký, *Signals for CPT and Lorentz violation in neutral meson oscillations*, Phys. Rev. **D61** (1999) 016002, arXiv:hep-ph/9909554.
- [25] M. Pivk and F. R. Le Diberder, *sPlot: A statistical tool to unfold data distributions*, Nucl. Instrum. Meth. **A555** (2005) 356, arXiv:physics/0402083.
- [26] LHCb collaboration, R. Aaij *et al.*, *Measurement of the  $\bar{B}^0 - B^0$  and  $\bar{B}_s^0 - B_s^0$  production asymmetries in  $pp$  collisions at  $\sqrt{s} = 7$  TeV*, Phys. Lett. **B739** (2014) 218, arXiv:1408.0275.
- [27] LHCb collaboration, R. Aaij *et al.*, *Measurement of the semileptonic CP asymmetry in  $B^0 - \bar{B}^0$  mixing*, Phys. Rev. Lett. **114** (2015) 041601, arXiv:1409.8586.
- [28] J. D. Scargle, *Studies in astronomical time series analysis: II. Statistical aspects of spectral analysis of unevenly spaced data*, Astrophys. J. **263** (1982) 835.
- [29] J. H. Horne and S. L. Baliunas, *A prescription for period analysis of unevenly sampled time series*, Astrophys. J. **302** (1986) 757.
- [30] A. Lenz and U. Nierste, *Theoretical update of  $B_s^0 - \bar{B}_s^0$  mixing*, JHEP **06** (2007) 072, arXiv:hep-ph/0612167.

## LHCb collaboration

R. Aaij<sup>39</sup>, C. Abellán Beteta<sup>41</sup>, B. Adeva<sup>38</sup>, M. Adinolfi<sup>47</sup>, Z. Ajaltouni<sup>5</sup>, S. Akar<sup>6</sup>, J. Albrecht<sup>10</sup>, F. Alessio<sup>39</sup>, M. Alexander<sup>52</sup>, S. Ali<sup>42</sup>, G. Alkhazov<sup>31</sup>, P. Alvarez Cartelle<sup>54</sup>, A.A. Alves Jr<sup>58</sup>, S. Amato<sup>2</sup>, S. Amerio<sup>23</sup>, Y. Amhis<sup>7</sup>, L. An<sup>3,40</sup>, L. Anderlini<sup>18</sup>, G. Andreassi<sup>40</sup>, M. Andreotti<sup>17,g</sup>, J.E. Andrews<sup>59</sup>, R.B. Appleby<sup>55</sup>, O. Aquines Gutierrez<sup>11</sup>, F. Archilli<sup>39</sup>, P. d'Argent<sup>12</sup>, A. Artamonov<sup>36</sup>, M. Artuso<sup>60</sup>, E. Aslanides<sup>6</sup>, G. Auriemma<sup>26,n</sup>, M. Baalouch<sup>5</sup>, S. Bachmann<sup>12</sup>, J.J. Back<sup>49</sup>, A. Badalov<sup>37</sup>, C. Baesso<sup>61</sup>, S. Baker<sup>54</sup>, W. Baldini<sup>17</sup>, R.J. Barlow<sup>55</sup>, C. Barschel<sup>39</sup>, S. Barsuk<sup>7</sup>, W. Barter<sup>39</sup>, V. Batozskaya<sup>29</sup>, V. Battista<sup>40</sup>, A. Bay<sup>40</sup>, L. Beaucourt<sup>4</sup>, J. Beddow<sup>52</sup>, F. Bedeschi<sup>24</sup>, I. Bediaga<sup>1</sup>, L.J. Bel<sup>42</sup>, V. Bellec<sup>40</sup>, N. Belloli<sup>21,k</sup>, I. Belyaev<sup>32</sup>, E. Ben-Haim<sup>8</sup>, G. Bencivenni<sup>19</sup>, S. Benson<sup>39</sup>, J. Benton<sup>47</sup>, A. Berezhnoy<sup>33</sup>, R. Bernet<sup>41</sup>, A. Bertolin<sup>23</sup>, F. Betti<sup>15</sup>, M.-O. Bettler<sup>39</sup>, M. van Beuzekom<sup>42</sup>, S. Bifani<sup>46</sup>, P. Billoir<sup>8</sup>, T. Bird<sup>55</sup>, A. Birnkraut<sup>10</sup>, A. Bizzeti<sup>18,i</sup>, T. Blake<sup>49</sup>, F. Blanc<sup>40</sup>, J. Blouw<sup>11</sup>, S. Blusk<sup>60</sup>, V. Bocci<sup>26</sup>, A. Bondar<sup>35</sup>, N. Bondar<sup>31,39</sup>, W. Bonivento<sup>16</sup>, A. Borgheresi<sup>21,k</sup>, S. Borghi<sup>55</sup>, M. Borisyak<sup>67</sup>, M. Borsato<sup>38</sup>, M. Boubdir<sup>9</sup>, T.J.V. Bowcock<sup>53</sup>, E. Bowen<sup>41</sup>, C. Bozzi<sup>17,39</sup>, S. Braun<sup>12</sup>, M. Britsch<sup>12</sup>, T. Britton<sup>60</sup>, J. Brodzicka<sup>55</sup>, E. Buchanan<sup>47</sup>, C. Burr<sup>55</sup>, A. Bursche<sup>2</sup>, J. Buytaert<sup>39</sup>, S. Cadeddu<sup>16</sup>, R. Calabrese<sup>17,g</sup>, M. Calvi<sup>21,k</sup>, M. Calvo Gomez<sup>37,p</sup>, P. Campana<sup>19</sup>, D. Campora Perez<sup>39</sup>, L. Capriotti<sup>55</sup>, A. Carbone<sup>15,e</sup>, G. Carboni<sup>25,l</sup>, R. Cardinale<sup>20,j</sup>, A. Cardini<sup>16</sup>, P. Carniti<sup>21,k</sup>, L. Carson<sup>51</sup>, K. Carvalho Akiba<sup>2</sup>, G. Casse<sup>53</sup>, L. Cassina<sup>21,k</sup>, L. Castillo Garcia<sup>40</sup>, M. Cattaneo<sup>39</sup>, Ch. Cauet<sup>10</sup>, G. Cavallero<sup>20</sup>, R. Cenci<sup>24,t</sup>, M. Charles<sup>8</sup>, Ph. Charpentier<sup>39</sup>, G. Chatzikonstantinidis<sup>46</sup>, M. Chefdeville<sup>4</sup>, S. Chen<sup>55</sup>, S.-F. Cheung<sup>56</sup>, M. Chrzaszcz<sup>41,27</sup>, X. Cid Vidal<sup>39</sup>, G. Ciezarek<sup>42</sup>, P.E.L. Clarke<sup>51</sup>, M. Clemencic<sup>39</sup>, H.V. Cliff<sup>48</sup>, J. Closier<sup>39</sup>, V. Coco<sup>58</sup>, J. Cogan<sup>6</sup>, E. Cogneras<sup>5</sup>, V. Cogoni<sup>16,f</sup>, L. Cojocariu<sup>30</sup>, G. Collazuol<sup>23,r</sup>, P. Collins<sup>39</sup>, A. Comerma-Montells<sup>12</sup>, A. Contu<sup>39</sup>, A. Cook<sup>47</sup>, M. Coombes<sup>47</sup>, S. Coquereau<sup>8</sup>, G. Corti<sup>39</sup>, M. Corvo<sup>17,g</sup>, B. Couturier<sup>39</sup>, G.A. Cowan<sup>51</sup>, D.C. Craik<sup>51</sup>, A. Crocombe<sup>49</sup>, M. Cruz Torres<sup>61</sup>, S. Cunliffe<sup>54</sup>, R. Currie<sup>54</sup>, C. D'Ambrosio<sup>39</sup>, E. Dall'Occo<sup>42</sup>, J. Dalseno<sup>47</sup>, P.N.Y. David<sup>42</sup>, A. Davis<sup>58</sup>, O. De Aguiar Francisco<sup>2</sup>, K. De Bruyn<sup>6</sup>, S. De Capua<sup>55</sup>, M. De Cian<sup>12</sup>, J.M. De Miranda<sup>1</sup>, L. De Paula<sup>2</sup>, P. De Simone<sup>19</sup>, C.-T. Dean<sup>52</sup>, D. Decamp<sup>4</sup>, M. Deckenhoff<sup>10</sup>, L. Del Buono<sup>8</sup>, N. Déleage<sup>4</sup>, M. Demmer<sup>10</sup>, D. Derkach<sup>67</sup>, O. Deschamps<sup>5</sup>, F. Dettori<sup>39</sup>, B. Dey<sup>22</sup>, A. Di Canto<sup>39</sup>, F. Di Ruscio<sup>25</sup>, H. Dijkstra<sup>39</sup>, F. Dordei<sup>39</sup>, M. Dorigo<sup>40</sup>, A. Dosil Suárez<sup>38</sup>, A. Dovbnya<sup>44</sup>, K. Dreimanis<sup>53</sup>, L. Dufour<sup>42</sup>, G. Dujany<sup>55</sup>, K. Dungs<sup>39</sup>, P. Durante<sup>39</sup>, R. Dzhelyadin<sup>36</sup>, A. Dziurda<sup>27</sup>, A. Dzyuba<sup>31</sup>, S. Easo<sup>50,39</sup>, U. Egede<sup>54</sup>, V. Egorychev<sup>32</sup>, S. Eidelman<sup>35</sup>, S. Eisenhardt<sup>51</sup>, U. Eitschberger<sup>10</sup>, R. Ekelhof<sup>10</sup>, L. Eklund<sup>52</sup>, I. El Rifai<sup>5</sup>, Ch. Elsasser<sup>41</sup>, S. Ely<sup>60</sup>, S. Esen<sup>12</sup>, H.M. Evans<sup>48</sup>, T. Evans<sup>56</sup>, A. Falabella<sup>15</sup>, C. Färber<sup>39</sup>, N. Farley<sup>46</sup>, S. Farry<sup>53</sup>, R. Fay<sup>53</sup>, D. Fazzini<sup>21,k</sup>, D. Ferguson<sup>51</sup>, V. Fernandez Albor<sup>38</sup>, F. Ferrari<sup>15</sup>, F. Ferreira Rodrigues<sup>1</sup>, M. Ferro-Luzzi<sup>39</sup>, S. Filippov<sup>34</sup>, M. Fiore<sup>17,g</sup>, M. Fiorini<sup>17,g</sup>, M. Firlej<sup>28</sup>, C. Fitzpatrick<sup>40</sup>, T. Fiutowski<sup>28</sup>, F. Fleuret<sup>7,b</sup>, K. Fohl<sup>39</sup>, M. Fontana<sup>16</sup>, F. Fontanelli<sup>20,j</sup>, D. C. Forshaw<sup>60</sup>, R. Forty<sup>39</sup>, M. Frank<sup>39</sup>, C. Frei<sup>39</sup>, M. Frosini<sup>18</sup>, J. Fu<sup>22</sup>, E. Furfaro<sup>25,l</sup>, A. Gallas Torreira<sup>38</sup>, D. Galli<sup>15,e</sup>, S. Gallorini<sup>23</sup>, S. Gambetta<sup>51</sup>, M. Gandelman<sup>2</sup>, P. Gandini<sup>56</sup>, Y. Gao<sup>3</sup>, J. García Pardiñas<sup>38</sup>, J. Garra Tico<sup>48</sup>, L. Garrido<sup>37</sup>, P.J. Garsed<sup>48</sup>, D. Gascon<sup>37</sup>, C. Gaspar<sup>39</sup>, L. Gavardi<sup>10</sup>, G. Gazzoni<sup>5</sup>, D. Gerick<sup>12</sup>, E. Gersabeck<sup>12</sup>, M. Gersabeck<sup>55</sup>, T. Gershon<sup>49</sup>, Ph. Ghez<sup>4</sup>, S. Gianì<sup>40</sup>, V. Gibson<sup>48</sup>, O.G. Girard<sup>40</sup>, L. Giubega<sup>30</sup>, V.V. Gligorov<sup>39</sup>, C. Göbel<sup>61</sup>, D. Golubkov<sup>32</sup>, A. Golutvin<sup>54,39</sup>, A. Gomes<sup>1,a</sup>, C. Gotti<sup>21,k</sup>, M. Grabalosa Gándara<sup>5</sup>, R. Graciani Diaz<sup>37</sup>, L.A. Granado Cardoso<sup>39</sup>, E. Graugés<sup>37</sup>, E. Graverini<sup>41</sup>, G. Graziani<sup>18</sup>, A. Grecu<sup>30</sup>, P. Griffith<sup>46</sup>, L. Grillo<sup>12</sup>, O. Grünberg<sup>65</sup>, E. Gushchin<sup>34</sup>, Yu. Guz<sup>36,39</sup>, T. Gys<sup>39</sup>, T. Hadavizadeh<sup>56</sup>,

C. Hadjivasiliou<sup>60</sup>, G. Haefeli<sup>40</sup>, C. Haen<sup>39</sup>, S.C. Haines<sup>48</sup>, S. Hall<sup>54</sup>, B. Hamilton<sup>59</sup>, X. Han<sup>12</sup>,  
 S. Hansmann-Menzemer<sup>12</sup>, N. Harnew<sup>56</sup>, S.T. Harnew<sup>47</sup>, J. Harrison<sup>55</sup>, J. He<sup>39</sup>, T. Head<sup>40</sup>,  
 A. Heister<sup>9</sup>, K. Hennessy<sup>53</sup>, P. Henrard<sup>5</sup>, L. Henry<sup>8</sup>, J.A. Hernando Morata<sup>38</sup>,  
 E. van Herwijnen<sup>39</sup>, M. Heß<sup>65</sup>, A. Hicheur<sup>2</sup>, D. Hill<sup>56</sup>, M. Hoballah<sup>5</sup>, C. Hombach<sup>55</sup>,  
 L. Hongming<sup>40</sup>, W. Hulsbergen<sup>42</sup>, T. Humair<sup>54</sup>, M. Hushchyn<sup>67</sup>, N. Hussain<sup>56</sup>, D. Hutchcroft<sup>53</sup>,  
 M. Idzik<sup>28</sup>, P. Ilten<sup>57</sup>, R. Jacobsson<sup>39</sup>, A. Jaeger<sup>12</sup>, J. Jalocha<sup>56</sup>, E. Jans<sup>42</sup>, A. Jawahery<sup>59</sup>,  
 M. John<sup>56</sup>, D. Johnson<sup>39</sup>, C.R. Jones<sup>48</sup>, C. Joram<sup>39</sup>, B. Jost<sup>39</sup>, N. Jurik<sup>60</sup>, S. Kandybei<sup>44</sup>,  
 W. Kanso<sup>6</sup>, M. Karacson<sup>39</sup>, T.M. Karbach<sup>39,†</sup>, S. Karodia<sup>52</sup>, M. Kecke<sup>12</sup>, M. Kelsey<sup>60</sup>,  
 I.R. Kenyon<sup>46</sup>, M. Kenzie<sup>39</sup>, T. Ketel<sup>43</sup>, E. Khairullin<sup>67</sup>, B. Khanji<sup>21,39,k</sup>, C. Khurewathanakul<sup>40</sup>,  
 T. Kirn<sup>9</sup>, S. Klaver<sup>55</sup>, K. Klimaszewski<sup>29</sup>, M. Kolpin<sup>12</sup>, I. Komarov<sup>40</sup>, R.F. Koopman<sup>43</sup>,  
 P. Koppenburg<sup>42,39</sup>, M. Kozeiha<sup>5</sup>, L. Kravchuk<sup>34</sup>, K. Kreplin<sup>12</sup>, M. Krepes<sup>49</sup>, P. Krokovny<sup>35</sup>,  
 F. Kruse<sup>10</sup>, W. Krzemien<sup>29</sup>, W. Kucewicz<sup>27,o</sup>, M. Kucharczyk<sup>27</sup>, V. Kudryavtsev<sup>35</sup>, A.  
 K. Kuonen<sup>40</sup>, K. Kurek<sup>29</sup>, T. Kvaratskheliya<sup>32</sup>, D. Lacarrere<sup>39</sup>, G. Lafferty<sup>55,39</sup>, A. Lai<sup>16</sup>,  
 D. Lambert<sup>51</sup>, G. Lanfranchi<sup>19</sup>, C. Langenbruch<sup>49</sup>, B. Langhans<sup>39</sup>, T. Latham<sup>49</sup>, C. Lazzeroni<sup>46</sup>,  
 R. Le Gac<sup>6</sup>, J. van Leerdam<sup>42</sup>, J.-P. Lees<sup>4</sup>, R. Lefèvre<sup>5</sup>, A. Leflat<sup>33,39</sup>, J. Lefrançois<sup>7</sup>,  
 E. Lemos Cid<sup>38</sup>, O. Leroy<sup>6</sup>, T. Lesiak<sup>27</sup>, B. Leverington<sup>12</sup>, Y. Li<sup>7</sup>, T. Likhomanenko<sup>67,66</sup>,  
 R. Lindner<sup>39</sup>, C. Linn<sup>39</sup>, F. Lionetto<sup>41</sup>, B. Liu<sup>16</sup>, X. Liu<sup>3</sup>, D. Loh<sup>49</sup>, I. Longstaff<sup>52</sup>, J.H. Lopes<sup>2</sup>,  
 D. Lucchesi<sup>23,r</sup>, M. Lucio Martinez<sup>38</sup>, H. Luo<sup>51</sup>, A. Lupato<sup>23</sup>, E. Luppi<sup>17,g</sup>, O. Lupton<sup>56</sup>,  
 N. Lusardi<sup>22</sup>, A. Lusiani<sup>24</sup>, X. Lyu<sup>62</sup>, F. Machefert<sup>7</sup>, F. Maciuc<sup>30</sup>, O. Maev<sup>31</sup>, K. Maguire<sup>55</sup>,  
 S. Malde<sup>56</sup>, A. Malinin<sup>66</sup>, G. Manca<sup>7</sup>, G. Mancinelli<sup>6</sup>, P. Manning<sup>60</sup>, A. Mapelli<sup>39</sup>, J. Maratas<sup>5</sup>,  
 J.F. Marchand<sup>4</sup>, U. Marconi<sup>15</sup>, C. Marin Benito<sup>37</sup>, P. Marino<sup>24,t</sup>, J. Marks<sup>12</sup>, G. Martellotti<sup>26</sup>,  
 M. Martin<sup>6</sup>, M. Martinelli<sup>40</sup>, D. Martinez Santos<sup>38</sup>, F. Martinez Vidal<sup>68</sup>, D. Martins Tostes<sup>2</sup>,  
 L.M. Massacrier<sup>7</sup>, A. Massafferri<sup>1</sup>, R. Matev<sup>39</sup>, A. Mathad<sup>49</sup>, Z. Mathe<sup>39</sup>, C. Matteuzzi<sup>21</sup>,  
 A. Mauri<sup>41</sup>, B. Maurin<sup>40</sup>, A. Mazurov<sup>46</sup>, M. McCann<sup>54</sup>, J. McCarthy<sup>46</sup>, A. McNab<sup>55</sup>,  
 R. McNulty<sup>13</sup>, B. Meadows<sup>58</sup>, F. Meier<sup>10</sup>, M. Meissner<sup>12</sup>, D. Melnychuk<sup>29</sup>, M. Merk<sup>42</sup>,  
 A Merli<sup>22,u</sup>, E. Michielin<sup>23</sup>, D.A. Milanese<sup>64</sup>, M.-N. Minard<sup>4</sup>, D.S. Mitzel<sup>12</sup>,  
 J. Molina Rodriguez<sup>61</sup>, I.A. Monroy<sup>64</sup>, S. Monteil<sup>5</sup>, M. Morandin<sup>23</sup>, P. Morawski<sup>28</sup>, A. Mordà<sup>6</sup>,  
 M.J. Morello<sup>24,t</sup>, J. Moron<sup>28</sup>, A.B. Morris<sup>51</sup>, R. Mountain<sup>60</sup>, F. Muheim<sup>51</sup>, D. Müller<sup>55</sup>,  
 J. Müller<sup>10</sup>, K. Müller<sup>41</sup>, V. Müller<sup>10</sup>, M. Mussini<sup>15</sup>, B. Muster<sup>40</sup>, P. Naik<sup>47</sup>, T. Nakada<sup>40</sup>,  
 R. Nandakumar<sup>50</sup>, A. Nandi<sup>56</sup>, I. Nasteva<sup>2</sup>, M. Needham<sup>51</sup>, N. Neri<sup>22</sup>, S. Neubert<sup>12</sup>,  
 N. Neufeld<sup>39</sup>, M. Neuner<sup>12</sup>, A.D. Nguyen<sup>40</sup>, C. Nguyen-Mau<sup>40,q</sup>, V. Niess<sup>5</sup>, S. Nieswand<sup>9</sup>,  
 R. Niet<sup>10</sup>, N. Nikitin<sup>33</sup>, T. Nikodem<sup>12</sup>, A. Novoselov<sup>36</sup>, D.P. O’Hanlon<sup>49</sup>,  
 A. Oblakowska-Mucha<sup>28</sup>, V. Obraztsov<sup>36</sup>, S. Ogilvy<sup>52</sup>, O. Okhrimenko<sup>45</sup>, R. Oldeman<sup>16,48,f</sup>,  
 C.J.G. Onderwater<sup>69</sup>, B. Osorio Rodrigues<sup>1</sup>, J.M. Otalora Goicochea<sup>2</sup>, A. Otto<sup>39</sup>, P. Owen<sup>54</sup>,  
 A. Oyanguren<sup>68</sup>, A. Palano<sup>14,d</sup>, F. Palombo<sup>22,u</sup>, M. Palutan<sup>19</sup>, J. Panman<sup>39</sup>, A. Papanestis<sup>50</sup>,  
 M. Pappagallo<sup>52</sup>, L.L. Pappalardo<sup>17,g</sup>, C. Pappenheimer<sup>58</sup>, W. Parker<sup>59</sup>, C. Parkes<sup>55</sup>,  
 G. Passaleva<sup>18</sup>, G.D. Patel<sup>53</sup>, M. Patel<sup>54</sup>, C. Patrignani<sup>20,j</sup>, A. Pearce<sup>55,50</sup>, A. Pellegrino<sup>42</sup>,  
 G. Penso<sup>26,m</sup>, M. Pepe Altarelli<sup>39</sup>, S. Perazzini<sup>15,e</sup>, P. Perret<sup>5</sup>, L. Pescatore<sup>46</sup>, K. Petridis<sup>47</sup>,  
 A. Petrolini<sup>20,j</sup>, M. Petruzzo<sup>22</sup>, E. Picatoste Olloqui<sup>37</sup>, B. Pietrzyk<sup>4</sup>, M. Piekies<sup>27</sup>, D. Pinci<sup>26</sup>,  
 A. Pistone<sup>20</sup>, A. Piucci<sup>12</sup>, S. Playfer<sup>51</sup>, M. Plo Casasus<sup>38</sup>, T. Poikela<sup>39</sup>, F. Polci<sup>8</sup>,  
 A. Poluektov<sup>49,35</sup>, I. Polyakov<sup>32</sup>, E. Polcarpo<sup>2</sup>, A. Popov<sup>36</sup>, D. Popov<sup>11,39</sup>, B. Popovici<sup>30</sup>,  
 C. Potterat<sup>2</sup>, E. Price<sup>47</sup>, J.D. Price<sup>53</sup>, J. Prisciandaro<sup>38</sup>, A. Pritchard<sup>53</sup>, C. Prouve<sup>47</sup>,  
 V. Pugatch<sup>45</sup>, A. Puig Navarro<sup>40</sup>, G. Punzi<sup>24,s</sup>, W. Qian<sup>56</sup>, R. Quagliani<sup>7,47</sup>, B. Rachwal<sup>27</sup>,  
 J.H. Rademacker<sup>47</sup>, M. Rama<sup>24</sup>, M. Ramos Pernas<sup>38</sup>, M.S. Rangel<sup>2</sup>, I. Raniuk<sup>44</sup>, G. Raven<sup>43</sup>,  
 F. Redi<sup>54</sup>, S. Reichert<sup>55</sup>, A.C. dos Reis<sup>1</sup>, V. Renaudin<sup>7</sup>, S. Ricciardi<sup>50</sup>, S. Richards<sup>47</sup>, M. Rihl<sup>39</sup>,  
 K. Rinnert<sup>53,39</sup>, V. Rives Molina<sup>37</sup>, P. Robbe<sup>7</sup>, A.B. Rodrigues<sup>1</sup>, E. Rodrigues<sup>55</sup>,

J.A. Rodriguez Lopez<sup>64</sup>, P. Rodriguez Perez<sup>55</sup>, A. Rogozhnikov<sup>67</sup>, S. Roiser<sup>39</sup>, V. Romanovsky<sup>36</sup>, A. Romero Vidal<sup>38</sup>, J. W. Ronayne<sup>13</sup>, M. Rotondo<sup>23</sup>, T. Ruf<sup>39</sup>, P. Ruiz Valls<sup>68</sup>, J.J. Saborido Silva<sup>38</sup>, N. Sagidova<sup>31</sup>, B. Saitta<sup>16,f</sup>, V. Salustino Guimaraes<sup>2</sup>, C. Sanchez Mayordomo<sup>68</sup>, B. Sanmartin Sedes<sup>38</sup>, R. Santacesaria<sup>26</sup>, C. Santamarina Rios<sup>38</sup>, M. Santimaria<sup>19</sup>, E. Santovetti<sup>25,l</sup>, A. Sarti<sup>19,m</sup>, C. Satriano<sup>26,n</sup>, A. Satta<sup>25</sup>, D.M. Saunders<sup>47</sup>, D. Savrina<sup>32,33</sup>, S. Schael<sup>9</sup>, M. Schiller<sup>39</sup>, H. Schindler<sup>39</sup>, M. Schlupp<sup>10</sup>, M. Schmelling<sup>11</sup>, T. Schmelzer<sup>10</sup>, B. Schmidt<sup>39</sup>, O. Schneider<sup>40</sup>, A. Schopper<sup>39</sup>, M. Schubiger<sup>40</sup>, M.-H. Schune<sup>7</sup>, R. Schwemmer<sup>39</sup>, B. Sciascia<sup>19</sup>, A. Sciubba<sup>26,m</sup>, A. Semennikov<sup>32</sup>, A. Sergi<sup>46</sup>, N. Serra<sup>41</sup>, J. Serrano<sup>6</sup>, L. Sestini<sup>23</sup>, P. Seyfert<sup>21</sup>, M. Shapkin<sup>36</sup>, I. Shapoval<sup>17,44,g</sup>, Y. Shcheglov<sup>31</sup>, T. Shears<sup>53</sup>, L. Shekhtman<sup>35</sup>, V. Shevchenko<sup>66</sup>, A. Shires<sup>10</sup>, B.G. Siddi<sup>17</sup>, R. Silva Coutinho<sup>41</sup>, L. Silva de Oliveira<sup>2</sup>, G. Simi<sup>23,s</sup>, M. Sirendi<sup>48</sup>, N. Skidmore<sup>47</sup>, T. Skwarnicki<sup>60</sup>, E. Smith<sup>54</sup>, I.T. Smith<sup>51</sup>, J. Smith<sup>48</sup>, M. Smith<sup>55</sup>, H. Snoek<sup>42</sup>, M.D. Sokoloff<sup>58</sup>, F.J.P. Soler<sup>52</sup>, F. Soomro<sup>40</sup>, D. Souza<sup>47</sup>, B. Souza De Paula<sup>2</sup>, B. Spaan<sup>10</sup>, P. Spradlin<sup>52</sup>, S. Sridharan<sup>39</sup>, F. Stagni<sup>39</sup>, M. Stahl<sup>12</sup>, S. Stahl<sup>39</sup>, S. Stefkova<sup>54</sup>, O. Steinkamp<sup>41</sup>, O. Stenyakin<sup>36</sup>, S. Stevenson<sup>56</sup>, S. Stoica<sup>30</sup>, S. Stone<sup>60</sup>, B. Storaci<sup>41</sup>, S. Stracka<sup>24,t</sup>, M. Straticiuc<sup>30</sup>, U. Straumann<sup>41</sup>, L. Sun<sup>58</sup>, W. Sutcliffe<sup>54</sup>, K. Swientek<sup>28</sup>, S. Swientek<sup>10</sup>, V. Syropoulos<sup>43</sup>, M. Szczekowski<sup>29</sup>, T. Szumlak<sup>28</sup>, S. T'Jampens<sup>4</sup>, A. Tayduganov<sup>6</sup>, T. Tekampe<sup>10</sup>, G. Tellarini<sup>17,g</sup>, F. Teubert<sup>39</sup>, C. Thomas<sup>56</sup>, E. Thomas<sup>39</sup>, J. van Tilburg<sup>42</sup>, V. Tisserand<sup>4</sup>, M. Tobin<sup>40</sup>, S. Tolk<sup>43</sup>, L. Tomassetti<sup>17,g</sup>, D. Tonelli<sup>39</sup>, S. Topp-Joergensen<sup>56</sup>, E. Tournefier<sup>4</sup>, S. Tourneur<sup>40</sup>, K. Trabelsi<sup>40</sup>, M. Traill<sup>52</sup>, M.T. Tran<sup>40</sup>, M. Tresch<sup>41</sup>, A. Trisovic<sup>39</sup>, A. Tsaregorodtsev<sup>6</sup>, P. Tsopelas<sup>42</sup>, N. Tuning<sup>42,39</sup>, A. Ukleja<sup>29</sup>, A. Ustyuzhanin<sup>67,66</sup>, U. Uwer<sup>12</sup>, C. Vacca<sup>16,39,f</sup>, V. Vagnoni<sup>15</sup>, S. Valat<sup>39</sup>, G. Valenti<sup>15</sup>, A. Vallier<sup>7</sup>, R. Vazquez Gomez<sup>19</sup>, P. Vazquez Regueiro<sup>38</sup>, C. Vázquez Sierra<sup>38</sup>, S. Vecchi<sup>17</sup>, M. van Veghel<sup>42</sup>, J.J. Velthuis<sup>47</sup>, M. Veltri<sup>18,h</sup>, G. Veneziano<sup>40</sup>, M. Vesterinen<sup>12</sup>, B. Viaud<sup>7</sup>, D. Vieira<sup>2</sup>, M. Vieites Diaz<sup>38</sup>, X. Vilasis-Cardona<sup>37,p</sup>, V. Volkov<sup>33</sup>, A. Vollhardt<sup>41</sup>, D. Voong<sup>47</sup>, A. Vorobyev<sup>31</sup>, V. Vorobyev<sup>35</sup>, C. Voß<sup>65</sup>, J.A. de Vries<sup>42</sup>, R. Waldi<sup>65</sup>, C. Wallace<sup>49</sup>, R. Wallace<sup>13</sup>, J. Walsh<sup>24</sup>, J. Wang<sup>60</sup>, D.R. Ward<sup>48</sup>, N.K. Watson<sup>46</sup>, D. Websdale<sup>54</sup>, A. Weiden<sup>41</sup>, M. Whitehead<sup>39</sup>, J. Wicht<sup>49</sup>, G. Wilkinson<sup>56,39</sup>, M. Wilkinson<sup>60</sup>, M. Williams<sup>39</sup>, M.P. Williams<sup>46</sup>, M. Williams<sup>57</sup>, T. Williams<sup>46</sup>, F.F. Wilson<sup>50</sup>, J. Wimberley<sup>59</sup>, J. Wishahi<sup>10</sup>, W. Wislicki<sup>29</sup>, M. Witek<sup>27</sup>, G. Wormser<sup>7</sup>, S.A. Wotton<sup>48</sup>, K. Wraight<sup>52</sup>, S. Wright<sup>48</sup>, K. Wyllie<sup>39</sup>, Y. Xie<sup>63</sup>, Z. Xu<sup>40</sup>, Z. Yang<sup>3</sup>, H. Yin<sup>63</sup>, J. Yu<sup>63</sup>, X. Yuan<sup>35</sup>, O. Yushchenko<sup>36</sup>, M. Zangoli<sup>15</sup>, M. Zavertyaev<sup>11,c</sup>, L. Zhang<sup>3</sup>, Y. Zhang<sup>3</sup>, A. Zhelezov<sup>12</sup>, Y. Zheng<sup>62</sup>, A. Zhokhov<sup>32</sup>, L. Zhong<sup>3</sup>, V. Zhukov<sup>9</sup>, S. Zucchelli<sup>15</sup>.

<sup>1</sup>Centro Brasileiro de Pesquisas Físicas (CBPF), Rio de Janeiro, Brazil

<sup>2</sup>Universidade Federal do Rio de Janeiro (UFRJ), Rio de Janeiro, Brazil

<sup>3</sup>Center for High Energy Physics, Tsinghua University, Beijing, China

<sup>4</sup>LAPP, Université Savoie Mont-Blanc, CNRS/IN2P3, Annecy-Le-Vieux, France

<sup>5</sup>Clermont Université, Université Blaise Pascal, CNRS/IN2P3, LPC, Clermont-Ferrand, France

<sup>6</sup>CPPM, Aix-Marseille Université, CNRS/IN2P3, Marseille, France

<sup>7</sup>LAL, Université Paris-Sud, CNRS/IN2P3, Orsay, France

<sup>8</sup>LPNHE, Université Pierre et Marie Curie, Université Paris Diderot, CNRS/IN2P3, Paris, France

<sup>9</sup>I. Physikalisches Institut, RWTH Aachen University, Aachen, Germany

<sup>10</sup>Fakultät Physik, Technische Universität Dortmund, Dortmund, Germany

<sup>11</sup>Max-Planck-Institut für Kernphysik (MPIK), Heidelberg, Germany

<sup>12</sup>Physikalisches Institut, Ruprecht-Karls-Universität Heidelberg, Heidelberg, Germany

<sup>13</sup>School of Physics, University College Dublin, Dublin, Ireland

<sup>14</sup>Sezione INFN di Bari, Bari, Italy

<sup>15</sup>Sezione INFN di Bologna, Bologna, Italy



- <sup>16</sup> *Sezione INFN di Cagliari, Cagliari, Italy*
- <sup>17</sup> *Sezione INFN di Ferrara, Ferrara, Italy*
- <sup>18</sup> *Sezione INFN di Firenze, Firenze, Italy*
- <sup>19</sup> *Laboratori Nazionali dell'INFN di Frascati, Frascati, Italy*
- <sup>20</sup> *Sezione INFN di Genova, Genova, Italy*
- <sup>21</sup> *Sezione INFN di Milano Bicocca, Milano, Italy*
- <sup>22</sup> *Sezione INFN di Milano, Milano, Italy*
- <sup>23</sup> *Sezione INFN di Padova, Padova, Italy*
- <sup>24</sup> *Sezione INFN di Pisa, Pisa, Italy*
- <sup>25</sup> *Sezione INFN di Roma Tor Vergata, Roma, Italy*
- <sup>26</sup> *Sezione INFN di Roma La Sapienza, Roma, Italy*
- <sup>27</sup> *Henryk Niewodniczanski Institute of Nuclear Physics Polish Academy of Sciences, Kraków, Poland*
- <sup>28</sup> *AGH - University of Science and Technology, Faculty of Physics and Applied Computer Science, Kraków, Poland*
- <sup>29</sup> *National Center for Nuclear Research (NCBJ), Warsaw, Poland*
- <sup>30</sup> *Horia Hulubei National Institute of Physics and Nuclear Engineering, Bucharest-Magurele, Romania*
- <sup>31</sup> *Petersburg Nuclear Physics Institute (PNPI), Gatchina, Russia*
- <sup>32</sup> *Institute of Theoretical and Experimental Physics (ITEP), Moscow, Russia*
- <sup>33</sup> *Institute of Nuclear Physics, Moscow State University (SINP MSU), Moscow, Russia*
- <sup>34</sup> *Institute for Nuclear Research of the Russian Academy of Sciences (INR RAN), Moscow, Russia*
- <sup>35</sup> *Budker Institute of Nuclear Physics (SB RAS) and Novosibirsk State University, Novosibirsk, Russia*
- <sup>36</sup> *Institute for High Energy Physics (IHEP), Protvino, Russia*
- <sup>37</sup> *Universitat de Barcelona, Barcelona, Spain*
- <sup>38</sup> *Universidad de Santiago de Compostela, Santiago de Compostela, Spain*
- <sup>39</sup> *European Organization for Nuclear Research (CERN), Geneva, Switzerland*
- <sup>40</sup> *Ecole Polytechnique Fédérale de Lausanne (EPFL), Lausanne, Switzerland*
- <sup>41</sup> *Physik-Institut, Universität Zürich, Zürich, Switzerland*
- <sup>42</sup> *Nikhef National Institute for Subatomic Physics, Amsterdam, The Netherlands*
- <sup>43</sup> *Nikhef National Institute for Subatomic Physics and VU University Amsterdam, Amsterdam, The Netherlands*
- <sup>44</sup> *NSC Kharkiv Institute of Physics and Technology (NSC KIPT), Kharkiv, Ukraine*
- <sup>45</sup> *Institute for Nuclear Research of the National Academy of Sciences (KINR), Kyiv, Ukraine*
- <sup>46</sup> *University of Birmingham, Birmingham, United Kingdom*
- <sup>47</sup> *H.H. Wills Physics Laboratory, University of Bristol, Bristol, United Kingdom*
- <sup>48</sup> *Cavendish Laboratory, University of Cambridge, Cambridge, United Kingdom*
- <sup>49</sup> *Department of Physics, University of Warwick, Coventry, United Kingdom*
- <sup>50</sup> *STFC Rutherford Appleton Laboratory, Didcot, United Kingdom*
- <sup>51</sup> *School of Physics and Astronomy, University of Edinburgh, Edinburgh, United Kingdom*
- <sup>52</sup> *School of Physics and Astronomy, University of Glasgow, Glasgow, United Kingdom*
- <sup>53</sup> *Oliver Lodge Laboratory, University of Liverpool, Liverpool, United Kingdom*
- <sup>54</sup> *Imperial College London, London, United Kingdom*
- <sup>55</sup> *School of Physics and Astronomy, University of Manchester, Manchester, United Kingdom*
- <sup>56</sup> *Department of Physics, University of Oxford, Oxford, United Kingdom*
- <sup>57</sup> *Massachusetts Institute of Technology, Cambridge, MA, United States*
- <sup>58</sup> *University of Cincinnati, Cincinnati, OH, United States*
- <sup>59</sup> *University of Maryland, College Park, MD, United States*
- <sup>60</sup> *Syracuse University, Syracuse, NY, United States*
- <sup>61</sup> *Pontifícia Universidade Católica do Rio de Janeiro (PUC-Rio), Rio de Janeiro, Brazil, associated to <sup>2</sup>*
- <sup>62</sup> *University of Chinese Academy of Sciences, Beijing, China, associated to <sup>3</sup>*
- <sup>63</sup> *Institute of Particle Physics, Central China Normal University, Wuhan, Hubei, China, associated to <sup>3</sup>*
- <sup>64</sup> *Departamento de Física, Universidad Nacional de Colombia, Bogota, Colombia, associated to <sup>8</sup>*
- <sup>65</sup> *Institut für Physik, Universität Rostock, Rostock, Germany, associated to <sup>12</sup>*

- <sup>66</sup> *National Research Centre Kurchatov Institute, Moscow, Russia, associated to* <sup>32</sup>
- <sup>67</sup> *Yandex School of Data Analysis, Moscow, Russia, associated to* <sup>32</sup>
- <sup>68</sup> *Instituto de Fisica Corpuscular (IFIC), Universitat de Valencia-CSIC, Valencia, Spain, associated to* <sup>37</sup>
- <sup>69</sup> *Van Swinderen Institute, University of Groningen, Groningen, The Netherlands, associated to* <sup>42</sup>
- <sup>a</sup> *Universidade Federal do Triângulo Mineiro (UFMT), Uberaba-MG, Brazil*
- <sup>b</sup> *Laboratoire Leprince-Ringuet, Palaiseau, France*
- <sup>c</sup> *P.N. Lebedev Physical Institute, Russian Academy of Science (LPI RAS), Moscow, Russia*
- <sup>d</sup> *Università di Bari, Bari, Italy*
- <sup>e</sup> *Università di Bologna, Bologna, Italy*
- <sup>f</sup> *Università di Cagliari, Cagliari, Italy*
- <sup>g</sup> *Università di Ferrara, Ferrara, Italy*
- <sup>h</sup> *Università di Urbino, Urbino, Italy*
- <sup>i</sup> *Università di Modena e Reggio Emilia, Modena, Italy*
- <sup>j</sup> *Università di Genova, Genova, Italy*
- <sup>k</sup> *Università di Milano Bicocca, Milano, Italy*
- <sup>l</sup> *Università di Roma Tor Vergata, Roma, Italy*
- <sup>m</sup> *Università di Roma La Sapienza, Roma, Italy*
- <sup>n</sup> *Università della Basilicata, Potenza, Italy*
- <sup>o</sup> *AGH - University of Science and Technology, Faculty of Computer Science, Electronics and Telecommunications, Kraków, Poland*
- <sup>p</sup> *LIFAELS, La Salle, Universitat Ramon Llull, Barcelona, Spain*
- <sup>q</sup> *Hanoi University of Science, Hanoi, Viet Nam*
- <sup>r</sup> *Università di Padova, Padova, Italy*
- <sup>s</sup> *Università di Pisa, Pisa, Italy*
- <sup>t</sup> *Scuola Normale Superiore, Pisa, Italy*
- <sup>u</sup> *Università degli Studi di Milano, Milano, Italy*
- <sup>†</sup> *Deceased*