

Additional observations of *Spiniferites alaskensis* from topotype material

Fabienne Marret^{1*}, Kenneth Mertens²

¹Department of Geography and Planning, School of Environmental Sciences, University of Liverpool, Liverpool, L69 7ZT, UK

²Ifremer, LER BO, Station de Biologie Marine, Place de la Croix, BP40537, F-29185 Concarneau Cedex, France

Here we present new observations of *Spiniferites alaskensis*, a relatively rare species described from the Eemian of the Gulf of Alaska. We show that the species shows a gonyaulacacean tabulation: Po, 4', 6'', 6c, ?s, 6''', 1p, 1'''''. The surface is finely granulate to scabrate. The species bears characteristic processes: these are exclusively gonal, membranous, perforated and end distally in platforms with stumpy ends. We provide more detail as how this species compares to closely related species belonging to the genus *Spiniferites*.

Keywords: Gulf of Alaska, chorate, *Spiniferites*, processes, Eemian, ODP

1. Introduction

Spiniferites alaskensis Marret et al. 2001 ex Marret in Fensome and Williams 2004 was first identified in the North East Pacific Ocean, in the Eemian interval that was recovered during the ODP Leg 145, site 887B (54°21' N – 148°23' W, 3647 m water depth) (Marret et al. 2001). The name was validated by Fensome & Williams (2004) because Marret et al. (2001) did not indicate which of the illustrations represented the holotype. The original description of this taxon mentioned a chorate cyst of ovoid shape with an apical boss, thin cyst wall, a finely granulate surface, gonal and broad terminally trifurcate processes, low sutural septa between the processes and a gonyaulacacean tabulation. This taxon was described as differing from other known *Spiniferites* only by the shape of the processes and their pointed-end termination.

New observations have been carried out, using a combination of light-transmitted and scanning electron microscopy, enabling to fully characterise the morphology of this relatively unknown species.

2. Material and methods

Permanent slides made from residues prepared by Marret et al. (2001) from the type locality (ODP Leg 145 core 887B, section 2H5 at 65 cm in section, Gulf of Alaska) were examined using a light microscope at Geotop, Montreal, Canada (Leica DMR equipped with a Leica DFC490 digital camera). Single specimens were picked under an inverted microscope with a micropipette and observed using a scanning electron microscope (Hitachi S-3400N SEM) at Geotop, Montreal, Canada. Kofoid's nomenclature is used to designate the plates.

3. Results

Division DINOFLAGELLATA (Bütschli 1885) Fensome et al. 1993, emend. Adl et al. 2005
Class DINOPHYCEAE Pascher 1914
Subclass PERIDINIPHYCIDAE Fensome et al. 1993
Order GONYAULACALES Taylor 1980

¹ Corresponding author. Email: f.marret@liverpool.ac.uk

48 Suborder Gonyaulacineae autonym
49 Family Gonyaulacaceae Lindemann 1928
50 Subfamily Gonyaulacoideae autonym
51 Genus *Spiniferites* Mantell 1850, emend. Sarjeant 1970
52 *Spiniferites alaskensis* Marret et al. 2001 ex Marret in Fensome and Williams 2004
53 Plate 1, Figures 1–10; Plate 2, Figures 1–6.

54 **Synonymy.** None.

55 **Holotype.** Marret et al. 2001, their Plate 1, Figures 7–9.

56 **Locus typicus.** ODP core 887B, section 2H5 at 65 cm in section, Gulf of Alaska.

57 **Stratum typicum.** Sediments deposited during isotope stage 5e in the Gulf of Alaska (ODP
58 Leg 145, site 887) northeastern North Pacific.

59 **Etymology.** Named *alaskensis* from its type locality in the Gulf of Alaska, NE Pacific
60 (Marret et al. 2001).

61 **Distinguishing characters.** Ovoid to pear shaped cysts with an apical boss (Plate 1, Figures
62 4–6) that touches all the apical plates (Plate 1, Figure 1). The cyst wall is thin (less than 1
63 μm) and has a finely granulate to scabrate wall surface, which is confirmed by SEM
64 observation (Plate 2, Figures 1–4). This surface texture continues on the bases of processes
65 and the bases of the connecting crests of the processes (Plate 2, Figure 2), but not on the more
66 distal part, which is smooth to shagreenate (Plate 2, Figures 5–6). Processes are exclusively
67 gonial, stout, membranous, with perforated polygonal platforms with stumpy distal ends (Plate
68 2, Figures 5–6). Processes at the junctions of the apical plates are shorter compared to others,
69 and those along 1'''' are the longest. The processes do not bear granules, as opposed to the
70 sutural septa connecting the processes. These sutural septa define a gonyaulacacean
71 tabulation: Po, 4', 6'', 6c, ?s, 6''', 1p, 1'''' (Plate 1, Figures 1, 3). The tabulation is expressed
72 with a generally low sutural arrangement, with slightly undulating sutures (Plate 1, Figure 7),
73 except being high where they connect cingular processes (arrow "a"), between 1', 4' and as
74 (arrow "b"), between 6''' and 1'''' (arrow "c"), and between 1p and 1'''' (arrow "d") (Plate
75 1, Figure 10). Observation of the epicyst shows an arrangement of four apical and six
76 precingular plates (Plate 1, Figure 1), and for the hypocyst, six postcingular plates although
77 1''' is not well expressed, 1p, and an asymmetrical 1'''''. The cingular arrangement shows six
78 plates, with a typical displacement of one cingular width, without overhang. In the sulcal
79 area, a pronounced suture is observed above ps and below as. The other sulcal plates are not
80 well reflected. The archeopyle has a pentagonal shape, corresponding to the third precingular
81 plate (3''), with rounded angles, and is reduced and free (Plate 1, Figure 8).

82 **Dimensions.** Central body diameter length 26.3 (31.4) 36.8 μm and width 23.6 (29.3) 31.5
83 μm with length of processes 7.5 (10.1) 12.5 μm (Marret et al. 2001). Number of specimen
84 measured: 11

85 **Biological affiliation.** Unknown.

86 **Comparison to other taxa.** *Spiniferites alaskensis* shows some similarities to a number of
87 *Spiniferites* species, such as *Spiniferites ludhamensis* Head 1996 which has a similar shape
88 and stout processes, but *Spiniferites ludhamensis* has hollow processes, and the cyst has a
89 thicker wall and no apical boss (Head 1996). *Spiniferites ristingensis* Head 2007 and
90 *Spiniferites delicatus* Reid 1974 also have membranous processes with petaloid distal ends.
91 However, *Spiniferites ristingensis* has small, densely distributed blisters and hollow
92 undulations over the surface (comparable to "bubble-wrap") (Head 2007). *Spiniferites*
93 *delicatus* has a granular surface, with high sutural crests (Reid 1974). *Spiniferites bentorii*
94 (Rossignol 1964) Wall & Dale 1970 also has a pear shaped body, with an apical boss, but has
95 processes often with claustra at their base and these processes are not membranous.
96 *Spiniferites belerius* Reid 1974 can also have an oval to pear shaped body, but has a shorter
97 apical boss, its wall surface is not as granular and there is a larger cingulum displacement,

98 and has box-like processes (Reid 1974). *Spiniferites alaskensis* also differs from *Spiniferites*
99 *falcipediis* Warny & Wrenn 1997, as the latter has hollow processes, no apical boss and high
100 crests between the antapical processes (Warny & Wrenn 1997). Lastly, *S. alaskensis* is
101 distinguished from *Spiniferites lazus* Reid 1974 as the latter has a clearly elongated ovoidal
102 shape and claustra at the base of the processes (Reid 1974).

103

104 **Acknowledgements**

105 We like to thank Anne de Vernal for use of microscope facilities at GEOTOP. Maryse Henry
106 and Raynald Lapointe are acknowledged for help with SEM preparations at GEOTOP. We
107 also are very grateful for the constructive comments from reviewers Laurent Londeix and
108 André Rochon that helped to improve the manuscript.

109

110 **References**

111

112 Adl SM, Simpson AGB, Farmer MA, Andersen RA, Anderson OR, Barta JR, Bowser SS,
113 Brugerolle G, Fensome RA, Fredericq S, James TY, Karpov S, Kugrens P, Krug J, Lane CE,
114 Lewis LA, Lodge J, Lynn DH, Mann DG, McCourt RM, Mendoza L, Moestrup Ø, Mozley-
115 Standridge SE, Nerad TA, Shearer CA, Smirnov AV, Spiegel FW, Taylor M.F.J.R. 2005. The
116 new higher level classification of Eukaryotes with emphasis on the taxonomy of Protists. *J*
117 *Eukaryot Microbiol.* 52:399–451.

118

119 Bütschli O. 1885. II. Abtheilung: Mastigophora. In: Dr. H.G. Bronn's Klassen und
120 Ordnungen des Thier-Reichs, wissenschaftlich dargestellt in Wort und Bild. Ester Band
121 Protozoa. Leipzig & Heidelberg: C.F. Winter'sche Verlagsbuchhandlung; p. 865–1088.

122

123 Fensome RA, Taylor FJR, Norris G, Sarjeant WAS, Wharton DI, Williams GL. 1993. A
124 classification of fossil and living dinoflagellates. *Micropaleontology Special Publication 7*: 1–
125 245.

126

127 Fensome RA, Williams GL. 2004. The Lentin and Williams Index of fossil dinoflagellates
128 2004 Edition. American Association of Stratigraphic Palynologists, Contributions Series 42.

129

130 Head MJ. 1996. Late Cenozoic dinoflagellates from the Royal Society borehole at Ludham,
131 Norfolk, eastern England. *J Paleol.* 70: 543–570.

132

133 Head MJ. 2007. Last Interglacial (Eemian) hydrographic conditions in the southwestern
134 Baltic Sea based on dinoflagellate cysts from Ristinge Klint, Denmark. *Geological Magazine*
135 144:987–1013.

136

137 Lindemann E. 1928. Abteilung Peridineae (Dinoflagellatae). In Engler A & Prantl K [Eds.]
138 Die Natürlichen Pflanzenfamilien nebst ihren Gattungen und wichtigeren Arten insbesondere
139 den Nutz-pflanzen. Zweite stark vermehrte und verbesserte Auflage herausgegeben von A.
140 Engler. 2 Band. Wilhelm Engelmann, Leipzig; pp. 3–104.

141

142 Mantell, GA. 1850: A Pictorial Atlas of Fossil Remains Consisting of Coloured Illustrations
143 Selected from Parkinson's "Organic Remains of a Former World", and Artis's "Antediluvian
144 Phytology". xii+207 p., 74 pl.; Henry G. Bohn, London, U.K.

145

- 146 Marret F, de Vernal A, Pedersen TF, McDonald D. 2001. Middle Pleistocene to Holocene
147 palynostratigraphy of Ocean Drilling Program Site 887 in the Gulf of Alaska, northeastern
148 North Pacific. *Can J Earth Sci.* 38:373–386.
- 149
- 150 Pascher A. 1914. Über Flagellaten und Algen. *Berichte der Deutschen Botanischen*
151 *Gesellschaft* 32: 136–160.
- 152
- 153 Reid PC. 1974. Gonyaulacacean dinoflagellate cysts from the British Isles. *Nova Hedwigia*
154 25:579–637.
- 155
- 156 Rossignol M. 1964. Hystrichosphères du Quaternaire en Méditerranée orientale, dans les
157 sédiments Pléistocènes et les boues marines actuelles. *Revue de micropaléontologie* 7:83–99.
- 158
- 159 Sarjeant WAS. 1970. The genus *Spiniferites* Mantell, 1850 (Dinophyceae). *Grana* 10:74–78.
- 160
- 161 Taylor F. 1980. On dinoflagellate evolution. *BioSystems* 13: 65–108.
- 162
- 163 Wall D, Dale B. 1970. Living Hystrichosphaerid Dinoflagellate Spores from Bermuda and
164 Puerto Rico. *Micropaleontology* 16:47–58.
- 165
- 166 Warny SA, Wrenn JH. 1997. New species of dinoflagellate cysts from the Bou Regreg core: a
167 Miocene-Pliocene boundary section on the Atlantic coast of Morocco. *Rev Palaeobot Palynol*
168 96:281–304.
- 169
- 170

171 Plate captions

172

173 Plate 1. Light microscope and SEM micrographs of topotype specimens of *Spiniferites*
174 *alaskensis*. Figures 1–9. High focus to low focus of single specimen. 1. Shows epicyst with
175 four apical plates and six precingular plates, stumpy distal ends of the processes are clearly
176 visible; the suture between 1' and 4' is faintly visible under light microscope. 2. Cross-
177 section in polar view. 3. Hypocyst showing four of the six postcingular plates, one posterior
178 intercalary plate (1p) and one large, asymmetrical antapical plate. 4. Mid-focus of pear-
179 shaped specimen. 5–6. Slightly higher than median focus to median focus of ovoid shaped
180 specimen. 7. Specimen showing morphology and distribution of exclusively gonial processes
181 with operculum still in place. 8. Specimen showing reduced archeopyle with rounded angles.
182 9. View of epicysta showing processes distribution and sutural crests. 10. View of hypocyst
183 showing sulcus and high crests between some of the processes (a,b,c,d). All scale bars = 10
184 μm .

185

186 Plate 2. Topotype specimens of *Spiniferites alaskensis*. 1–4. SEM micrographs and light
187 micrographs of wall texture, showing scabrate to granulate central body surface and
188 shagreenate septa and processes. 5–6. SEM micrographs of process structure showing stumpy
189 distal ends and low crests connecting the processes. All scale bars = 10 μm .



