

Manuscript version: Published Version

The version presented in WRAP is the published version (Version of Record).

Persistent WRAP URL:

<http://wrap.warwick.ac.uk/87737>

How to cite:

The repository item page linked to above, will contain details on accessing citation guidance from the publisher.

Copyright and reuse:

The Warwick Research Archive Portal (WRAP) makes this work by researchers of the University of Warwick available open access under the following conditions.

Copyright © and all moral rights to the version of the paper presented here belong to the individual author(s) and/or other copyright owners. To the extent reasonable and practicable the material made available in WRAP has been checked for eligibility before being made available.

Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge. Provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

Publisher's statement:

Please refer to the repository item page, publisher's statement section, for further information.

For more information, please contact the WRAP Team at: wrap@warwick.ac.uk

Spectral gap for the stochastic quantization equation on the 2-dimensional torus

Pavlos Tsatsoulis and Hendrik Weber

University of Warwick, Coventry, UK. E-mail: p.tsatsoulis@warwick.ac.uk; hendrik.weber@warwick.ac.uk

Received 27 September 2016; revised 24 March 2017; accepted 10 April 2017

Abstract. We study the long time behavior of the stochastic quantization equation. Extending recent results by Mourrat and Weber (Global well-posedness of the dynamic ϕ^4 in the plane (2015) Preprint) we first establish a strong non-linear dissipative bound that gives control of moments of solutions at all positive times independent of the initial datum. We then establish that solutions give rise to a Markov process whose transition semigroup satisfies the strong Feller property. Following arguments by Chouk and Friz (Support theorem for a singular SPDE: the case of gPAM (2016) Preprint) we also prove a support theorem for the laws of the solutions. Finally all of these results are combined to show that the transition semigroup satisfies the Doeblin criterion which implies exponential convergence to equilibrium.

Along the way we give a simple direct proof of the Markov property of solutions and an independent argument for the existence of an invariant measure using the Krylov–Bogoliubov existence theorem. Our method makes no use of the reversibility of the dynamics or the explicit knowledge of the invariant measure and it is therefore in principle applicable to situations where these are not available, e.g. the vector-valued case.

Résumé. Nous étudions le comportement sur le long terme de l'équation de quantification stochastique. Dans la continuité de récents résultats par Mourrat et Weber (Global well-posedness of the dynamic ϕ^4 in the plane (2015) Preprint), nous établissons en premier lieu une borne dissipative forte non-linéaire qui contrôle les moments des solutions, pour tout choix de temps, indépendamment des conditions initiales. Nous prouvons ensuite que les solutions génèrent un processus Markovien dont le semigroupe satisfait la propriété de Feller forte. Nous obtenons également un théorème pour le support des lois des solutions grâce à des arguments adaptés de Chouk et Friz (Support theorem for a singular SPDE: the case of gPAM (2016) Preprint). Enfin, en combinant tous ces résultats, nous montrons que le semigroupe de transition satisfait le critère de Doeblin, ce qui entraîne une convergence exponentielle vers l'équilibre.

Nous obtenons également au passage une preuve directe de la propriété de Markov pour les solutions, ainsi qu'un argument indépendant pour l'existence de mesures invariantes en utilisant le théorème d'existence de Krylov–Bogoliubov. Notre méthode n'utilise pas le caractère réversible de la dynamique ni la connaissance explicite de la mesure invariante, et peut donc en théorie s'appliquer dans des cas où ces propriétés ne sont pas connues, par exemple le cas vectoriel.

MSC: 37A25; 60H15; 81T08

Keywords: Singular SPDEs; Strong Feller property; Support theorem; Exponential mixing

1. Introduction

We consider the stochastic quantization equation on the 2-dimensional torus \mathbb{T}^2 given by

$$\begin{cases} \partial_t X = \Delta X - X - \sum_{k=0}^n a_k : X^k : + \xi, & \text{in } \mathbb{R}_+ \times \mathbb{T}^2, \\ X(0, \cdot) = x, & \text{on } \mathbb{T}^2, \end{cases} \quad (1.1)$$

where n is odd, $a_n > 0$, ξ is a Gaussian space time white noise and x is a distribution of suitably negative regularity. Here X^k stands for the k th Wick power of X (see Section 2 for its definition). This equation was first proposed by Parisi and Wu (see [19]) as a natural reversible dynamics for the Φ_2^{n+1} measure which is given by

$$\nu(dX) \propto \exp \left\{ -2 \int_{\mathbb{T}^2} \sum_{k=0}^n \frac{a_k}{k+1} : X^{k+1} : (z) dz \right\} \mu(dX), \quad (1.2)$$

where μ is the law of a massive Gaussian free field.

The interpretation and construction of solutions for (1.1) remained a challenge for many years with important contributions by Jona-Lasinio and Mitter in [13] (solution of a modified equation via Girsanov's transformation) and Albeverio and Röckner in [1] (construction of solutions using the theory of Dirichlet forms). In [5] Da Prato and Debussche proposed a simple transformation of (1.1) which allowed them to prove local in time existence of strong solutions for any initial datum x of suitable (negative) regularity and non-explosion for x in a set of measure one with respect to (1.2). Recently Mourrat and Weber [15] obtained global in time solutions on the full space for any initial datum of suitable regularity by following a similar strategy. In [21] Röckner et al. identified these solutions with the solutions obtained via Dirichlet forms.

The aim of this paper is to establish exponential convergence to equilibrium for solutions of (1.1). Building on the analysis in [15] and using a simple comparison test for non-linear ordinary differential equations we establish a strong dissipative bound for the solutions. We then prove the strong Feller property for the Markov semigroup generated by the solution generalizing the method in [12, Section 4.2]. Although for convenience we make (moderate) use of global in time existence which follows from the strong dissipative bounds derived before, this part of the analysis could also be implemented using only local existence (see Remark 5.9); the linearized dynamics of Galerkin approximations are controlled by combining a localization via stopping times and the small-time bounds obtained from the local existence theory. We furthermore establish a support theorem in the spirit of [4]. Finally, we combine all of these ingredients to show that the associated Markov semigroup satisfies the Doeblin criterion which implies exponential convergence to the unique invariant measure uniformly over the state space.

All steps are implemented for general odd n except for the support theorem which we only show in the case $n = 3$. The reason for this restriction is explained in Remark 6.2. We expect however that a support theorem for (1.1) holds true for all odd n and that such a result could be combined with the results of this paper to generalize Theorem 6.5 to the case of an arbitrary odd n .

Along the way we give independent proofs of the Markov property for the dynamics as well as existence of the invariant measure. The Markov property was already established previously in [21] based on the identification of the dynamics with the solutions constructed via Dirichlet form. The same paper [21] also established that (1.2) is a reversible (and in particular invariant) measure for the dynamics. We stress that our approach completely circumvents the theory of Dirichlet forms and uses neither the symmetry of the process nor the explicit form of the invariant measure. We therefore expect that our methods could be applied in situations where the reversibility is absent and where there is no explicit representation of the invariant measure, for example in situations where X is vector rather than scalar valued.

Finally, we would like to mention two independent works on a similar subject – one [22] published very recently and one [10] about to appear. In [22] the authors establish that (1.2) is the unique invariant measure for the dynamics and that the transition probabilities converge to this invariant measure. Their method is based on the asymptotic coupling technique from [11] and relies on the bounds from [15]. This analysis does however not include the strong Feller property or the support theorem and does not imply exponential convergence to equilibrium. In the forthcoming article [10] the authors present a general method to establish the strong Feller property, for solutions of SPDE solved in the framework of the theory of regularity structures. As an example this method is implemented for the dynamic Φ_3^4 model. We expect that their method can also treat the case of (1.1) but at first glance it only implies continuity of the associated Markov semigroup with respect to the total variational norm, whereas Theorem 5.10 implies Hölder continuity with respect to this norm.

1.1. Outline

In Section 2 we introduce some notation for Wick powers and their approximations. The results in this section are essentially contained in [5] and [15] and the purpose of the section is mostly to fix notation. In Section 3 we first

briefly sketch the construction of solutions to (1.1) including a short time bound and a stability result which are used in Section 5. We then prove the strong dissipative bound which is independent of the initial condition, improving on the bounds obtained in [15]. In Section 4 we prove the Markov property for the solution using a simple factorization argument as in [6] and we furthermore prove existence of invariant measures based on the bounds obtained in Section 3. The strong Feller property for the associated Markov semigroup is shown in Section 5. Finally, in Section 6 we prove a support theorem for (1.1) in the case of $n = 3$ which we combine with the results of the previous sections to prove exponential mixing.

1.2. Notation

Let \mathbb{T}^d be the d -dimensional torus of size 1 (throughout the article $d = 2$). We denote by $C^\infty(\mathbb{R}^d)$ and $C^\infty(\mathbb{T}^d)$ the space of real-valued smooth functions over \mathbb{R}^d and \mathbb{T}^d respectively as well as by $\mathcal{S}'(\mathbb{T}^d)$ the dual space of Schwarz distributions acting on $C^\infty(\mathbb{T}^d)$. We furthermore denote by $L^p(\mathbb{T}^d)$ the space of p -integrable functions on \mathbb{T}^d , endowed with the norm

$$\|f\|_{L^p} := \left(\int_{\mathbb{T}^d} |f(z)|^p dz \right)^{\frac{1}{p}}.$$

Although we only deal with spaces of real-valued functions, we prefer to work with the orthonormal basis $\{e_m\}_{m \in \mathbb{Z}^d}$ of trigonometric functions

$$e_m(z) := e^{2\pi i m \cdot z},$$

for $z \in \mathbb{T}^d$. Thus some complex-valued functions appear and we write

$$\langle f, g \rangle = \int_{\mathbb{T}^d} f(z) \overline{g(z)} dz$$

for their inner product. In this notation, for $f \in L^2(\mathbb{T}^d)$, the m th Fourier coefficient is given by

$$\hat{f}(m) := \langle f, e_m \rangle$$

and since f is real-valued we have the symmetry condition

$$\hat{f}(-m) = \overline{\hat{f}(m)}, \tag{1.3}$$

for any $m \in \mathbb{Z}^d$. For $f \in \mathcal{S}'(\mathbb{T}^d)$ we define the m th Fourier coefficient as

$$\hat{f}(m) := \langle f, \cos(2\pi i m \cdot) \rangle + i \langle f, \sin(2\pi i m \cdot) \rangle,$$

with the convention that $\langle f, \cdot \rangle$ stands for the action of f on $C^\infty(\mathbb{T}^d)$.

For $\zeta \in \mathbb{R}^d$ and $r > 0$ we denote by $B(\zeta, r)$ the ball of radius r centered at ζ . We consider the annulus $\mathcal{A} = B(0, \frac{8}{3}) \setminus B(0, \frac{3}{4})$ and a dyadic partition of unity $(\chi_\kappa)_{\kappa \geq -1}$ such that

- (i) $\chi_{-1} = \tilde{\chi}$ and $\chi_\kappa = \chi(\cdot/2^\kappa)$, $\kappa \geq 0$, for two radial functions $\tilde{\chi}, \chi \in C^\infty(\mathbb{R}^d)$.
- (ii) $\text{supp } \tilde{\chi} \subset B(0, \frac{4}{3})$ and $\text{supp } \chi \subset \mathcal{A}$.
- (iii) $\tilde{\chi}(\zeta) + \sum_{\kappa=0}^\infty \chi(\zeta/2^\kappa) = 1$, for all $\zeta \in \mathbb{R}^d$.

We furthermore let

$$\mathcal{A}_{2^\kappa} := 2^\kappa \mathcal{A}, \quad \kappa \geq 0.$$

Notice that $\text{supp } \chi_\kappa \subset \mathcal{A}_{2^\kappa}$, for every $\kappa \geq 0$. We also keep the convention that $\mathcal{A}_{2^{-1}} = B(0, \frac{4}{3})$. The existence of such a dyadic partition of unity is given by [2, Proposition 2.10].

For a function $f \in C^\infty(\mathbb{T}^d)$ we define the κ th Littlewood–Paley block as

$$\delta_\kappa f(z) := \sum_{m \in \mathbb{Z}^d} \chi_\kappa(m) \hat{f}(m) e^{2\pi i m \cdot z}, \quad \kappa \geq -1. \quad (1.4)$$

Sometimes it is convenient to write (1.4) as $\delta_\kappa f = \eta_\kappa * f$, $\kappa \geq -1$, where

$$\eta_\kappa * f(\cdot) = \int_{\mathbb{T}^d} \eta_\kappa(\cdot - z) f(z) \, dz,$$

and

$$\eta_\kappa(z) := \sum_{m \in \mathbb{Z}^d} \chi_\kappa(m) e^{2\pi i m \cdot z}.$$

For $\alpha \in \mathbb{R}$ and $p, q \in [1, \infty]$ we define the non-homogeneous periodic Besov norm (see [2, Section 2.7]),

$$\|f\|_{\mathcal{B}_{p,q}^\alpha} := \left\| (2^{\alpha\kappa} \|\delta_\kappa f\|_{L^p})_{\kappa \geq -1} \right\|_{\ell^q}. \quad (1.5)$$

The Besov space $\mathcal{B}_{p,q}^\alpha$ is defined as the completion of $C^\infty(\mathbb{T}^d)$ with respect to the norm (1.5). We are mostly interested in the Besov space $\mathcal{B}_{\infty,\infty}^\alpha$ which from now on we denote by \mathcal{C}^α . Note that for $p = q = \infty$ our definition of Besov spaces differs from the standard definition as the set of those distributions for which (1.5) is finite. Our convention has the advantage that all Besov spaces are separable. Some basic properties of Besov spaces are collected in Appendix A.

Throughout the article we fix $\alpha_0 \in (0, \frac{1}{n})$ (we measure the regularity of the initial condition in $\mathcal{C}^{-\alpha_0}$) as well as $\beta > 0$ (the regularity of the remainder v defined in Section 3.1) and $\gamma > 0$ (the rate of blowup of the $\|v_t\|_{\mathcal{C}^\beta}$ for t close to 0, see e.g. Theorem 3.3) such that

$$\gamma < \frac{1}{n}, \quad \frac{\beta + \alpha_0}{2} < \gamma. \quad (1.6)$$

For an arbitrary $\alpha \in (0, 1)$ we let

$$C^{n,-\alpha}(0; T) := C([0, T]; \mathcal{C}^{-\alpha}) \times C((0, T]; \mathcal{C}^{-\alpha})^{n-1} \quad (1.7)$$

and denote by $\underline{Z} = (Z^{(1)}, Z^{(2)}, \dots, Z^{(n)})$ a generic $C^{n,-\alpha}(0; T)$ -valued vector. For $\alpha' > 0$ we also define

$$\|\underline{Z}\|_{\alpha'; T} := \max_{k=1,2,\dots,n} \left\{ \sup_{0 \leq t \leq T} t^{(k-1)\alpha'} \|Z_t^{(k)}\|_{\mathcal{C}^{-\alpha}} \right\}.$$

Throughout the whole article C denotes a positive constant which might differ from line to line but we make explicit the dependence on different parameters where necessary. Furthermore, through the proofs of our statements, in cases where we do not want to keep track of the various constants in the inequalities we use \lesssim instead of $\leq C$. Finally, we use $a \vee b$ and $a \wedge b$ to denote the maximum and the minimum of a and b .

2. Preliminaries

In this section we present the necessary stochastic tools to handle (1.1). In Section 2.1 we introduce the stochastic heat equation along with its Wick powers in terms of abstract iterated stochastic integrals in the spirit of [18, Chapter 1]. In Section 2.2 we describe how these iterated stochastic integrals arise as limits of powers of solutions to finite dimensional approximations after renormalization.

2.1. The stochastic heat equation and its Wick powers

Let ξ be a space-time white noise on $\mathbb{R} \times \mathbb{T}^2$ (see Appendix B) on some probability space $(\Omega, \mathcal{F}, \mathbb{P})$, which is fixed from now on. We set

$$\tilde{\mathcal{F}}_t = \sigma(\{\xi(\phi) : \phi|_{(t, +\infty) \times \mathbb{T}^2} \equiv 0, \phi \in L^2(\mathbb{R} \times \mathbb{T}^2)\}), \tag{2.1}$$

for $t > -\infty$ and denote by $(\mathcal{F}_t)_{t > -\infty}$ the usual augmentation (as in [20, Chapter 1.4]) of the filtration $(\tilde{\mathcal{F}}_t)_{t > -\infty}$.

Consider the stochastic heat equation with zero initial condition at time $s \in (-\infty, \infty)$

$$\begin{cases} \partial_t \mathfrak{I}_{s,t} = \Delta \mathfrak{I}_{s,t} - \mathfrak{I}_{s,t} + \xi, & \text{in } (s, \infty) \times \mathbb{T}^2, \\ \mathfrak{I}_{s,s} = 0, & \text{on } \mathbb{T}^2. \end{cases} \tag{2.2}$$

There are several ways to give a meaning to this equation. We simply use Duhamel’s principle (see [8, Section 2.3]) as a definition and set for every $\phi \in C^\infty(\mathbb{T}^2)$ and $s < t$

$$\mathfrak{I}_{s,t}(\phi) := \int_s^t \int_{\mathbb{T}^2} \langle \phi, H(t-r, z-\cdot) \rangle \xi(dr, dz), \tag{2.3}$$

where $H(r, \cdot)$, $r \in \mathbb{R} \setminus \{0\}$, stands for the periodic heat kernel on $L^2(\mathbb{T}^2)$ given by

$$H(r, z) := \sum_{m \in \mathbb{Z}^2} e^{-(1+4\pi^2|m|^2)r} e_m(z), \tag{2.4}$$

for all $z \in \mathbb{T}^2$. We furthermore let

$$S(t) := e^{-t} e^{t\Delta}$$

be the semigroup associated to the generator $\Delta - 1$ in $L^2(\mathbb{T}^2)$, i.e. the convolution operator with respect to the space variable $z \in \mathbb{T}^2$ with the kernel $H(t, \cdot)$.

The integral in (2.3) is a stochastic integral (see Appendix B for definitions) and for fixed $s < t$, $\mathfrak{I}_{s,t}$ is a family of Gaussian random variables indexed by $C^\infty(\mathbb{T}^2)$.

Since it is more convenient to work with stationary processes we extend definition (2.3) for $s = -\infty$. For $\phi \in C^\infty(\mathbb{T}^2)$, $n \geq 2$ and $t > -\infty$ we also consider the multiple stochastic integral (see Appendix B) given by

$$\mathfrak{W}_{-\infty,t}^{\circlearrowleft n}(\phi) := \int_{\{(-\infty, t] \times \mathbb{T}^2\}^n} \left\langle \phi, \prod_{k=1}^n H(t-s_k, z_k-\cdot) \right\rangle \xi \left(\bigotimes_{k=1}^n ds_k, \bigotimes_{k=1}^n dz_k \right). \tag{2.5}$$

We call $\mathfrak{W}_{-\infty,\cdot}^{\circlearrowleft n}$ the n th Wick power of $\mathfrak{I}_{-\infty,\cdot}$ and we recall that for every $n \geq 1$ and $\phi \in C^\infty(\mathbb{T}^2)$, $\mathfrak{W}_{-\infty,\cdot}^{\circlearrowleft n}(\phi)$ is an element in the n th homogeneous Wiener chaos (see Appendix B for definitions). We furthermore point out that $\mathfrak{W}_{-\infty,\cdot}^{\circlearrowleft n}(\phi)$ is stationary, for every $\phi \in C^\infty(\mathbb{T}^2)$.

The next theorem collects the optimal regularity properties of the processes $\{\mathfrak{W}_{-\infty,\cdot}^{\circlearrowleft n}\}$, $n \geq 1$ and is very similar to the bounds originally derived in [5, Lemma 3.2]. The precise statement is a consequence of the Kolmogorov-type criterion [15, Lemma 5.2, Lemma 5.3] and the proof follows similar lines to the one of [15, Theorem 5.4].

Theorem 2.1. *Let $p \geq 2$. For every $n \geq 1$ and $t_0 > -\infty$, the process $\mathfrak{W}_{-\infty,t_0+}^{\circlearrowleft n}$ admits a modification $\widetilde{\mathfrak{W}}_{-\infty,t_0+}^{\circlearrowleft n}$ such that $\widetilde{\mathfrak{W}}_{-\infty,t_0+}^{\circlearrowleft n} \in C([0, T]; C^{-\alpha})$, for every $T > 0$ and $\alpha > 0$. Furthermore, there exists $\theta \equiv \theta(\alpha) \in (0, 1) > 0$ and $C \equiv C(T, \alpha, p)$ such that*

$$\mathbb{E} \sup_{s,t \in [0, T]} \frac{\|\widetilde{\mathfrak{W}}_{-\infty,t_0+t}^{\circlearrowleft n} - \widetilde{\mathfrak{W}}_{-\infty,t_0+s}^{\circlearrowleft n}\|_{C^{-\alpha}}^p}{|t-s|^{p\theta}} \leq C. \tag{2.6}$$

For notational convenience we always refer to $\widetilde{\mathfrak{W}}_{-\infty,\cdot}^{\circlearrowleft n}$ as $\mathfrak{W}_{-\infty,\cdot}^{\circlearrowleft n}$.

Proof. See Appendix D. □

Notice that for every $t \geq s$ we have that

$$\mathfrak{I}_{s,t} = \mathfrak{I}_{-\infty,t} - S(t-s)\mathfrak{I}_{-\infty,s}.$$

It is then reasonable to define (see also [15, p. 34] for equivalent definitions) the n th shifted Wick power of $\mathfrak{I}_{s,t}$, $t > s > -\infty$, as

$$\mathfrak{W}_{s,t}^{(n)} := \sum_{k=0}^n \binom{n}{k} (-1)^k (S(t-s)\mathfrak{I}_{-\infty,s})^k \mathfrak{W}_{-\infty,t}^{(n-k)}. \quad (2.7)$$

Here and below we use the convention $\mathfrak{W}_{s,t}^{(k)} \equiv 1$ for $k = 0$ and any $-\infty \leq s < t$. We furthermore point out that the n th shifted Wick power is not an element of the n th homogeneous Wiener chaos (see Appendix B for definitions). We refer the reader to Proposition 2.3 below for a natural approximation of the objects defined in (2.7).

At this point we would like to mention that one might work directly with $\mathfrak{W}_{-\infty,\cdot}^{(n)}$ instead of introducing (2.7) (see for example [5] and [9]). This alternative approach has the advantage that the diagrams are stationary in time. However, we prefer to work with (2.7) (as in [15]) because when proving the Markov property (see Section 4.1) we use heavily that $\mathfrak{W}_{s,t}^{(n)}$ is independent of \mathcal{F}_s for any $s < t$ (see Proposition 2.3). A slight disadvantage of our convention is the logarithmic divergence of $\mathfrak{W}_{s,t}^{(n)}$ as $t \downarrow s$ (see (2.8)).

The next proposition uses the regularization property of the heat semigroup (see Proposition A.5) to show that for every $t > s$ and $n \geq 2$, $\mathfrak{W}_{s,t}^{(n)}$ is a well-defined element in a Besov space of negative regularity close to 0.

Proposition 2.2. *Let $p \geq 2$ and $T > 0$. For every $s_0 > -\infty$, $\alpha \in (0, 1)$ and $\alpha' > 0$ there exist $\theta \equiv \theta(\alpha, \alpha') > 0$ and $C \equiv C(T, \alpha, \alpha', p, n)$ such that*

$$\mathbb{E} \sup_{0 \leq s \leq t} (s^{(n-1)\alpha'} p \|\mathfrak{W}_{s_0, s_0+s}^{(n)}\|_{\mathcal{C}^{-\alpha}}^p) \leq Ct^{p\theta}, \quad (2.8)$$

for every $t \leq T$.

Proof. We show (2.8) for $s_0 = 0$.

Let $\bar{\alpha} < \alpha \wedge \frac{2}{3}\alpha'$ and $V(s) = S(s)(-\mathfrak{I}_{-\infty,0})$. Using (A.1) as well as Propositions A.7 and A.5 we have that

$$\|V(s)^n\|_{\mathcal{C}^{-\alpha}} \lesssim \|V(s)\|_{\mathcal{C}^{2\bar{\alpha}}}^{n-1} \|V(s)\|_{\mathcal{C}^{-\bar{\alpha}}} \lesssim s^{-(n-1)\frac{3}{2}\bar{\alpha}} \|\mathfrak{I}_{-\infty,0}\|_{\mathcal{C}^{-\bar{\alpha}}}^n.$$

In a similar way, for $k \notin \{0, n\}$, we have that

$$\|V(s)^k \mathfrak{W}_{-\infty,s}^{(n-k)}\|_{\mathcal{C}^{-\alpha}} \lesssim s^{-k\frac{3}{2}\bar{\alpha}} \|\mathfrak{I}_{-\infty,0}\|_{\mathcal{C}^{-\bar{\alpha}}}^k \|\mathfrak{W}_{-\infty,s}^{(n-k)}\|_{\mathcal{C}^{-\bar{\alpha}}}.$$

Thus

$$\|\mathfrak{W}_{0,s}^{(n)}\|_{\mathcal{C}^{-\alpha}} \lesssim s^{-(n-1)\frac{3}{2}\bar{\alpha}} \|\mathfrak{I}_{-\infty,0}\|_{\mathcal{C}^{-\bar{\alpha}}}^n + \sum_{k=0}^{n-1} \binom{n}{k} s^{-k\frac{3}{2}\bar{\alpha}} \|\mathfrak{I}_{-\infty,0}\|_{\mathcal{C}^{-\bar{\alpha}}}^k \|\mathfrak{W}_{-\infty,s}^{(n-k)}\|_{\mathcal{C}^{-\bar{\alpha}}}.$$

Hence

$$\begin{aligned} \mathbb{E} \sup_{0 \leq s \leq t} s^{(n-1)\alpha'} p \|\mathfrak{W}_{0,s}^{(n)}\|_{\mathcal{C}^{-\alpha}}^p &\lesssim t^{(n-1)(\alpha' - \frac{3}{2}\bar{\alpha})p} \mathbb{E} \|\mathfrak{I}_{-\infty,0}\|_{\mathcal{C}^{-\bar{\alpha}}}^{np} \\ &+ \sum_{k=0}^{n-1} \binom{n}{k} t^{((n-1)\alpha' - k\frac{3}{2}\bar{\alpha})p} (\mathbb{E} \|\mathfrak{I}_{-\infty,0}\|_{\mathcal{C}^{-\bar{\alpha}}}^{2kp})^{\frac{1}{2}} \left(\mathbb{E} \sup_{0 \leq s \leq t} \|\mathfrak{W}_{-\infty,s}^{(n-k)}\|_{\mathcal{C}^{-\bar{\alpha}}}^{2p} \right)^{\frac{1}{2}}, \end{aligned}$$

where we use a Cauchy–Schwarz inequality in the last line. Combining with (2.6) we finally obtain (2.8). □

2.2. Finite dimensional approximations

Let $\rho_\varepsilon(z) = \sum_{|m| < \frac{1}{\varepsilon}} e_m(z)$ and define a finite dimensional approximation of $\mathfrak{I}_{s,t}$ by

$$\mathfrak{I}_{s,t}^\varepsilon := \mathfrak{I}_{s,t}(\rho_\varepsilon(z \cdot)).$$

We introduce the renormalization constant

$$\mathfrak{N}^\varepsilon := \|\mathbf{1}_{[0,\infty)} H_\varepsilon\|_{L^2(\mathbb{R} \times \mathbb{T}^2)}^2, \tag{2.9}$$

where $H_\varepsilon(r, z) = (H(r, \cdot) * \rho_\varepsilon)(z)$ noting that $\mathfrak{N}^\varepsilon \sim \log \varepsilon^{-1}$ as $\varepsilon \rightarrow 0^+$. For any integer $n \geq 1$ and $s \geq -\infty$ we define

$$\mathfrak{H}_{s,t}^{(n)\varepsilon} := \mathcal{H}_n(\mathfrak{I}_{s,t}^\varepsilon, \mathfrak{N}^\varepsilon),$$

where $\mathcal{H}_n(X, C)$, $X, C \in \mathbb{R}$, stands for the n th Hermite polynomial given by the recursive formula

$$\begin{cases} \mathcal{H}_{-1}(X, C) = 0, & \mathcal{H}_0(X, C) = 1, \\ \mathcal{H}_n(X, C) = X\mathcal{H}_{n-1}(X, C) - (n-1)C\mathcal{H}_{n-2}(X, C). \end{cases} \tag{2.10}$$

The first three Hermite polynomials are given by $\mathcal{H}_1(X, C) = X$, $\mathcal{H}_2(X, C) = X^2 - C$, $\mathcal{H}_3(X, C) = X^3 - 3CX$.

Proposition 2.3. *Let $\alpha, \alpha' > 0$. Then for every $n \geq 1$ and $p \geq 2$ we have that*

$$\begin{aligned} \lim_{\varepsilon \rightarrow 0^+} \mathbb{E} \sup_{0 \leq t \leq T} \left\| \mathfrak{H}_{-\infty, s+t}^{(n)\varepsilon} - \mathfrak{H}_{-\infty, s+t}^{(n)} \right\|_{C^{-\alpha}}^p &= 0, \\ \lim_{\varepsilon \rightarrow 0^+} \mathbb{E} \sup_{0 \leq t \leq T} t^{(n-1)\alpha' p} \left\| \mathfrak{H}_{s, s+t}^{(n)\varepsilon} - \mathfrak{H}_{s, s+t}^{(n)} \right\|_{C^{-\alpha}}^p &= 0, \end{aligned}$$

for every $s > -\infty$. In particular, $\mathfrak{H}_{s, s+}^{(n)}$ is independent of \mathcal{F}_s and for $s_1, s_2 \neq -\infty$, $\mathfrak{H}_{s_1, s_1+}^{(n)} \stackrel{\text{law}}{=} \mathfrak{H}_{s_2, s_2+}^{(n)}$.

Proof. See Appendix E. □

An immediate consequence of the above proposition is the following corollary which we later use in Section 4 to prove the Markov property.

Corollary 2.4. *For every $n \geq 1$ and $t, h > 0$ the following identity holds \mathbb{P} -almost surely,*

$$\mathfrak{H}_{0, t+h}^{(n)} = \sum_{k=0}^n \binom{n}{k} (S(h)\mathfrak{I}_{0,t})^k \mathfrak{H}_{t, t+h}^{(n-k)}. \tag{2.11}$$

Proof. It suffices to check (2.11) for $\mathfrak{H}_{0, t+h}^{(n)\varepsilon}$. The result then follows from the previous proposition. □

3. Solving the equation

3.1. Analysis of the problem

We are interested in solving the following renormalized stochastic partial differential equation,

$$\begin{cases} \partial_t X = \Delta X - X - \sum_{k=0}^n a_k : X^k : + \xi, & \text{in } \mathbb{R}_+ \times \mathbb{T}^2, \\ X(0, \cdot) = x, & \text{on } \mathbb{T}^2, \end{cases} \tag{3.1}$$

where X^k stands for the k th Wick power of X and $x \in C^{-\alpha_0}$. Motivated by the Da Prato–Debussche method (see [5]) we search for solutions to (3.1) by writing $X = \mathfrak{I}_{0,\cdot} + v$, where $\mathfrak{I}_{0,\cdot}$ is the solution to (2.2) and the remainder v is a mild solution of the following random partial differential equation,

$$\begin{cases} \partial_t v = \Delta v - v - \sum_{k=0}^n a_k \sum_{j=0}^k \binom{k}{j} v^j \mathfrak{I}_{0,\cdot}^{k-j}, \\ v(0, \cdot) = x. \end{cases} \quad (3.2)$$

Remark 3.1. In [15] $\mathfrak{I}_{0,\cdot}$ is started from x and consequently there (3.2) is solved with zero initial condition. Our approach of starting $\mathfrak{I}_{0,\cdot}$ from 0 and the remainder v from x has the advantage that the strong non-linear damping in (3.2) acts directly on the initial condition, yielding a strong dissipative bound for v that is independent of x (see Proposition 3.7).

We can rewrite (3.2) as

$$\begin{cases} \partial_t v = \Delta v - v - \sum_{j=0}^n v^j Z^{(n-j)}, \\ v(0, \cdot) = x, \end{cases} \quad (3.3)$$

where

$$Z^{(n-j)} = \sum_{k=j}^n a_k \binom{k}{j} \mathfrak{I}_{0,\cdot}^{k-j},$$

for all $0 \leq j \leq n-1$ and $Z^{(0)} = a_n$. For the rest of this section \underline{Z} always denotes a vector of this form. This convention will not be used in the other sections.

Notice that for every $\alpha \in (0, 1)$, $\underline{Z} \in C^{n, -\alpha}(0; T)$ (see (1.7) for the definition of the space), for every $T > 0$, and by (2.8) for every $\alpha' > 0$ there exists $\theta > 0$ such that

$$\mathbb{E} \|\underline{Z}\|_{\alpha; \alpha'; t}^p \leq C t^{p\theta} \quad (3.4)$$

for every $t \leq T$, $p \geq 2$.

We now fix $\alpha < \alpha_0$ small enough (the precise value is fixed below in the proof of Theorem 3.3) and $\underline{Z} \in C^{n, -\alpha}(0; T)$, for every $T > 0$, and a norm $\|\cdot\|_{\alpha; \alpha'; T}$, for some $\alpha' > 0$ but still sufficiently small. We furthermore let

$$F(v, \underline{Z}) := \sum_{j=0}^n v^j Z^{(n-j)}. \quad (3.5)$$

3.2. Short time existence

We are interested in solutions to the PDE problem (3.3).

Definition 3.2. Let $T > 0$ and $x \in C^{-\alpha_0}$. We say that a function v is a mild solution of (3.3) up to time T if $v \in C((0, T]; C^\beta)$ and

$$v_t = S(t)x - \int_0^t S(t-s)F(v_s, \underline{Z}_s) ds, \quad (3.6)$$

for every $t \leq T$.

The next theorem implies the existence of local in time solutions to (3.3).

Theorem 3.3 ([5, Proposition 4.4], [15, Theorem 6.2]). *Let $x \in C^{-\alpha_0}$ and $R > 0$ such that $\|x\|_{C^{-\alpha_0}} \leq R$. Then for every $\beta, \gamma > 0$ satisfying (1.6) and $T > 0$ there exists $T^* \equiv T^*(R, \|\underline{Z}\|_{\alpha; \alpha'; T}) \leq T$ such that (3.3) has a unique mild solution on $[0, T^*]$ and*

$$\sup_{0 \leq s \leq T^*} s^\gamma \|v_s\|_{C^\beta} \leq 1.$$

If we furthermore assume that $\|\underline{Z}\|_{\alpha; \alpha'; T} \leq 1$, then there exists $\theta > 0$ and a constant $C > 0$ independent of R such that

$$T^* = \left(\frac{1}{C(R+1)} \right)^{\frac{1}{\theta}}. \tag{3.7}$$

Proof. This theorem is (essentially) proved in [15, Theorem 6.2], but the expression (3.7) is not made explicit there; we give a sketch. It is sufficient to prove that for T^* as in (3.7) the operator

$$\mathcal{M}_{T^*} v_t = S(t)x + \int_0^t S(t-s)F(v_s, \underline{Z}_s) ds$$

is a contraction on the set $\mathcal{B}_{T^*} := \{\sup_{0 \leq s \leq T^*} s^\gamma \|v_s\|_{C^\beta} \leq 1\}$, i.e. we need to show that \mathcal{M}_{T^*} maps \mathcal{B}_{T^*} into itself and that for $v, \tilde{v} \in \mathcal{B}_{T^*}$ we have $\sup_{0 \leq s \leq T^*} s^\gamma \|\mathcal{M}_{T^*} v_s - \mathcal{M}_{T^*} \tilde{v}_s\|_{C^\beta} \leq (1 - \lambda) \sup_{0 \leq s \leq T^*} s^\gamma \|v_s - \tilde{v}_s\|_{C^\beta}$ for some $\lambda > 0$. We only show the first property. First notice that using the explicit form of F (see (3.5))

$$\begin{aligned} \|\mathcal{M}_{T^*} v_t\|_{C^\beta} &\lesssim \|S(t)x\|_{C^\beta} + \int_0^t \|S(t-s)v_s^n\|_{C^\beta} ds + \sum_{j=0}^{n-1} \int_0^t \|S(t-s)v_s^j Z_s^{(n-j)}\|_{C^\beta} ds \\ &\lesssim t^{-\frac{\beta+\alpha_0}{2}} \|x\|_{C^{-\alpha_0}} + \int_0^t s^{-n\gamma} ds + \int_0^t (t-s)^{-\frac{\alpha+\beta}{2}} s^{-(n-j-1)\gamma} s^{-j\gamma} ds \\ &\lesssim t^{-\frac{\beta+\alpha_0}{2}} \|x\|_{C^{-\alpha_0}} + \int_0^t s^{-n\gamma} ds + \int_0^t (t-s)^{-\frac{\alpha+\beta}{2}} s^{-(n-1)\gamma} ds, \end{aligned}$$

where we use Proposition A.5 and we furthermore assume that $\alpha' < \gamma$. By (1.6) if we choose $\alpha > 0$ sufficiently small so that $\frac{\alpha+\beta}{2} + (n-1)\gamma < 1$ we get

$$\|\mathcal{M}_{T^*} v_t\|_{C^\beta} \lesssim t^{-\frac{\beta+\alpha_0}{2}} \|x\|_{C^{-\alpha_0}} + t^{1-n\gamma} + t^{1-\frac{\alpha+\beta}{2}-(n-1)\gamma}$$

and multiplying both sides by t^γ we obtain that

$$t^\gamma \|\mathcal{M}_{T^*} v_t\|_{C^\beta} \lesssim t^{\gamma-\frac{\beta+\alpha_0}{2}} R + t^{1-(n-1)\gamma} + t^{1-\frac{\alpha+\beta}{2}-(n-2)\gamma} \lesssim t^\theta (R+1).$$

Then, for $T^* \equiv T^*(R)$ as in (3.7) and every $t \leq T^*$ we get that

$$\sup_{0 \leq s \leq t} s^\gamma \|\mathcal{M}_{T^*} v_s\|_{C^\beta} \leq 1,$$

which implies that indeed \mathcal{M}_{T^*} maps \mathcal{B}_{T^*} into itself. □

The next proposition is a stability result which we use later on in Section 5. We first introduce some extra notation. Let $\{\underline{Z}^\varepsilon\}_{\varepsilon \in (0,1)}$ take values in $C^{n, -\alpha}(0; T)$ such that

$$\lim_{\varepsilon \rightarrow 0^+} \|\underline{Z}^\varepsilon - \underline{Z}\|_{\alpha; \alpha'; T} = 0.$$

Furthermore, let $F_\varepsilon = \hat{\Pi}_\varepsilon F$, where $\hat{\Pi}_\varepsilon$ is a linear smooth approximation such that the following properties hold for every $\lambda \in \mathbb{R}$:

- (i) $\|\hat{\Pi}_\varepsilon\|_{\mathcal{C}^\lambda \rightarrow \mathcal{C}^\lambda} \leq C$, for every $\varepsilon \in (0, 1)$.
(ii) For every $\delta > 0$ there exists $\theta = \theta(\lambda, \delta)$ such that

$$\|\hat{\Pi}_\varepsilon x - x\|_{\mathcal{C}^{\lambda-\delta}} \leq C\varepsilon^\theta \|x\|_{\mathcal{C}^\lambda}.$$

One can check that $\hat{\Pi}_\varepsilon = \sum_{-1 \leq \kappa < \log_2 \varepsilon^{-1}} \delta_\kappa$ is such a linear smooth approximation.

Denote by v^ε the corresponding mild solution of (3.3) with F replaced by F_ε , \underline{Z} by $\underline{Z}^\varepsilon$ and initial condition $x^\varepsilon = \hat{\Pi}_\varepsilon x$ (short time existence of v^ε is ensured by the same arguments as in the proof of [15, Theorem 6.1]). We then have the following proposition.

Proposition 3.4. *Let v be the unique solution to (3.3) on a closed interval $[0, T^*]$ (i.e. the solution does not explode at T^*). Then for every $\varepsilon \in (0, 1)$ there exists a unique solution v_ε to the approximate equation up to some (possibly infinite) explosion time T_ε^* . Furthermore, there exists $\varepsilon_0 > 0$ such that for every $\varepsilon < \varepsilon_0$, $T_\varepsilon^* \geq T^*$, and we have*

$$\lim_{\varepsilon \rightarrow 0^+} \sup_{0 \leq t \leq T_\varepsilon^* \wedge T^*} t^\gamma \|v_t - v_t^\varepsilon\|_{\mathcal{C}^\beta} = 0.$$

Proof. By (1.6) it is possible to find $\delta > 0$ such that

$$\frac{\delta}{2} + n\gamma < 1, \quad \frac{\alpha_0 + \delta + \beta}{2} + (n-1)\gamma < 1.$$

For $\varepsilon \in (0, 1)$ we notice that

$$v_t - v_t^\varepsilon = S(t)(x - x^\varepsilon) - \int_0^t S(t-s)(F(v_s, \underline{Z}_s) - F_\varepsilon(v_s^\varepsilon, \underline{Z}_s^\varepsilon)) ds$$

and using (A.7) and property (ii) of $\hat{\Pi}_\varepsilon$ we get

$$\begin{aligned} \|v_t - v_t^\varepsilon\|_{\mathcal{C}^\beta} &\lesssim t^{-\frac{\alpha_0 + \delta + \beta}{2}} \varepsilon^\theta \|x\|_{\mathcal{C}^{-\alpha_0}} + \int_0^t (t-s)^{\frac{\delta}{2}} \|v_s^n - \hat{\Pi}_\varepsilon(v_s^\varepsilon)^n\|_{\mathcal{C}^{\beta-\delta}} ds \\ &\quad + \int_0^t (t-s)^{-\frac{\alpha + \delta + \beta}{2}} \|R(v_s, \underline{Z}_s) - R_\varepsilon(v_s^\varepsilon, \underline{Z}_s^\varepsilon)\|_{\mathcal{C}^{-\alpha-\delta}} ds, \end{aligned}$$

where $R(v, \underline{Z}) = \sum_{j=0}^{n-1} v^j Z^{(n-j)}$ and $R_\varepsilon = \hat{\Pi}_\varepsilon R$. Using the triangle inequality as well as the properties (i) and (ii) of $\hat{\Pi}_\varepsilon$ we have that

$$\|v_s^n - \hat{\Pi}_\varepsilon(v_s^\varepsilon)^n\|_{\mathcal{C}^{\beta-\delta}} \lesssim \varepsilon^\theta \|(v_s^\varepsilon)^n\|_{\mathcal{C}^\beta} + \|v_s^n - (v_s^\varepsilon)^n\|_{\mathcal{C}^\beta}$$

and

$$\begin{aligned} \|R(v_s, \underline{Z}_s) - R_\varepsilon(v_s^\varepsilon, \underline{Z}_s^\varepsilon)\|_{\mathcal{C}^{-\alpha-\delta}} &\lesssim \varepsilon^\theta \|R(v_s, \underline{Z}_s)\|_{\mathcal{C}^{-\alpha}} + \|R(v_s, \underline{Z}_s) - R(v_s^\varepsilon, \underline{Z}_s^\varepsilon)\|_{\mathcal{C}^{-\alpha}} \\ &\quad + \|R(v_s^\varepsilon, \underline{Z}_s) - R(v_s^\varepsilon, \underline{Z}_s^\varepsilon)\|_{\mathcal{C}^{-\alpha}}. \end{aligned}$$

Let $M = \sup_{t \leq T^*} t^\gamma \|v_t\|_{\mathcal{C}^\beta}$, $N = \|\underline{Z}\|_{\alpha; \alpha'; T}$ and $\tau^\varepsilon = \inf\{t > 0, t \leq T_\varepsilon^* : t^\gamma \|v_t - v_t^\varepsilon\|_{\mathcal{C}^\beta} > 1\}$. Then, for every $t \leq \tau^\varepsilon \wedge T^*$, we have the bounds

$$\begin{aligned} \|(v_s^\varepsilon)^n\|_{\mathcal{C}^\beta} &\leq Cs^{-n\gamma}, \\ \|v_s^n - (v_s^\varepsilon)^n\|_{\mathcal{C}^\beta} &\leq Cs^{-n\gamma} \sup_{t \leq \tau^\varepsilon \wedge T^*} t^\gamma \|v_t - v_t^\varepsilon\|_{\mathcal{C}^\beta}, \end{aligned}$$

as well as

$$\begin{aligned} \|R(v_s, \underline{Z}_s)\|_{C^{-\alpha}} &\leq C s^{-(n-1)\gamma}, \\ \|R(v_s, \underline{Z}_s) - R(v_s^\varepsilon, \underline{Z}_s)\|_{C^{-\alpha}} &\leq C s^{-(n-1)\gamma} \sup_{t \leq \tau^\varepsilon \wedge T^*} t^\gamma \|v_t - v_t^\varepsilon\|_{C^\beta}, \\ \|R(v_s^\varepsilon, \underline{Z}_s) - R(v_s^\varepsilon, \underline{Z}_s^\varepsilon)\|_{C^{-\alpha}} &\leq C s^{-(n-1)\gamma} \|\underline{Z} - \underline{Z}^\varepsilon\|_{\alpha; \alpha'; T}, \end{aligned}$$

where the constant C depends on M and N . Thus

$$\begin{aligned} \|v_t - v_t^\varepsilon\|_{C^\beta} &\leq C \left(\varepsilon^\theta t^{-\frac{\alpha_0 + \delta + \beta}{2}} \|x\|_{C^{-\alpha_0}} + \varepsilon^\theta t^{1 - \frac{\delta}{2} - n\gamma} + \sup_{t \leq \tau^\varepsilon \wedge T^*} t^\gamma \|v_t - v_t^\varepsilon\|_{C^\beta} t^{1 - \frac{\delta}{2} - n\gamma} \right. \\ &\quad \left. + \varepsilon^\theta t^{1 - \frac{\alpha + \delta + \beta}{2} - (n-1)\gamma} + \sup_{t \leq \tau^\varepsilon \wedge T^*} t^\gamma \|v_t - v_t^\varepsilon\|_{C^\beta} t^{1 - \frac{\alpha + \delta + \beta}{2} - (n-1)\gamma} \right. \\ &\quad \left. + \|\underline{Z} - \underline{Z}^\varepsilon\|_{\alpha; \alpha'; T} t^{1 - \frac{\alpha + \delta + \beta}{2} - (n-1)\gamma} \right). \end{aligned}$$

Multiplying by t^γ and choosing $\tilde{T}^* \equiv \tilde{T}^*(M, N) > 0$ sufficiently small we can assure that

$$\sup_{t \leq \tilde{T}^*} t^\gamma \|v_t - v_t^\varepsilon\|_{C^\beta} \leq \varepsilon^\theta \|x\|_{C^{-\alpha_0}} + \|\underline{Z} - \underline{Z}^\varepsilon\|_{\alpha; \alpha'; T} + \varepsilon^\theta.$$

Iterating the procedure if necessary we find $N^* > 0$, independent of ε since $\tau^\varepsilon \wedge T^* \leq T^*$, and $C > 0$ such that

$$\sup_{t \leq \tau^\varepsilon \wedge T^*} t^\gamma \|v_t - v_t^\varepsilon\|_{C^\beta} \leq (N^* C + 1) (\varepsilon^\theta \|x\|_{C^{-\alpha_0}} + \|\underline{Z} - \underline{Z}^\varepsilon\|_{\alpha; \alpha'; T} + \varepsilon^\theta). \tag{3.8}$$

Let $\varepsilon_0 > 0$ such that for every $\varepsilon < \varepsilon_0$

$$\varepsilon^\theta \|x\|_{C^{-\alpha_0}} + \|\underline{Z} - \underline{Z}^\varepsilon\|_{\alpha; \alpha'; T} + \varepsilon^\theta < \frac{1}{(N^* C + 1)}.$$

Then for every $\varepsilon < \varepsilon_0$

$$\sup_{t \leq \tau^\varepsilon \wedge T^*} t^\gamma \|v_t - v_t^\varepsilon\|_{C^\beta} < 1$$

and the definition of τ^ε implies that $\tau^\varepsilon \wedge T^* = T^*$, which proves the first claim. For the second claim we just let $\varepsilon \rightarrow 0^+$ in (3.8). □

3.3. Testing the equation

Proposition 3.5 ([15, Proposition 6.8]). *Let $v \in C((0, T]; C^\beta)$ be a mild solution to (3.3). Then for all $s_0 > 0$ and all even integers $p \geq 2$*

$$\frac{1}{p} (\|v_t\|_{L^p}^p - \|v_{s_0}\|_{L^p}^p) = \int_{s_0}^t (-(p-1) \langle \nabla v_s, v_s^{p-2} \nabla v_s \rangle - \langle v_s, v_s^{p-1} \rangle - \langle F(v_s, \underline{Z}_s), v_s^{p-1} \rangle) ds, \tag{3.9}$$

for all $s_0 \leq t \leq T$. In particular, if we differentiate with respect to t ,

$$\frac{1}{p} \partial_t \|v_t\|_{L^p}^p = -(p-1) \langle \nabla v_t, v_t^{p-2} \nabla v_t \rangle - \langle v_t, v_t^{p-1} \rangle - \langle F(v_t, \underline{Z}_t), v_t^{p-1} \rangle, \tag{3.10}$$

for every $t \in (0, T)$.

Remark 3.6. This proposition involves spatial derivatives of v up to first order and the proof of (3.9) requires some time regularity on v . Our local existence theory implies that $v \in C((0, T]; \mathcal{C}^\beta)$ for some $\beta < 1$ (see (1.6)) due to the fact that we start (3.3) with initial condition in $\mathcal{C}^{-\alpha_0}$. This is the reason we state (3.9) for $s_0 > 0$. However one can prove that for fixed $t > 0$ v is almost a \mathcal{C}^2 function (see [15, Theorem 6.2]), as well as a Hölder continuous function from $(0, T]$ to $L^\infty(\mathbb{T}^2)$ (see [15, Proposition 6.5]) for some exponent strictly greater than $\frac{1}{2}$.

3.4. A priori estimates

Global existence of (3.3) for $x \in \mathcal{C}^\beta$ was already established in [15] based on a priori estimates of the L^p norm of v . Here we derive a stronger bound which does not depend on the initial condition x and we use later on to prove the main results of Sections 4 and 6.

Proposition 3.7. *Let $v \in C((0, T]; \mathcal{C}^\beta)$ be a solution of (3.3) with initial condition $x \in \mathcal{C}^{-\alpha_0}$ and $p \geq 2$ be an even integer. Then for every $0 < t \leq T$ and $\lambda = \frac{p+n-1}{p}$*

$$\|v_t\|_{L^p}^p \leq C \left[t^{-\frac{1}{\lambda-1}} \vee \left(\sum_{j,i} t^{-\alpha' p_i^j} \sup_{0 \leq r \leq t} (r^{\alpha' p_i^j} \|Z_r^{(n-j)}\|_{\mathcal{C}^{-\alpha}}^{p_i^j}) \right)^{\frac{1}{\lambda}} \right], \quad (3.11)$$

for some $p_i^j > 0$. In particular, the bound is independent from $\|x\|_{\mathcal{C}^{-\alpha_0}}$ and the randomness outside of the interval $[0, t]$.

Proof. Let

$$\alpha < \frac{1}{(p+n-1)(n-1)} \quad (3.12)$$

and recall that $F(v_s, \underline{Z}_s) = \sum_{j=0}^n v_s^j Z_s^{(n-j)}$. Thus

$$\langle F(v_s, \underline{Z}_s), v_s^{p-1} \rangle = \sum_{j=0}^n \langle v_s^{p+j-1}, Z_s^{(n-j)} \rangle = a_n \|v_s^{p+n-1}\|_{L^1} + \langle g_s, v_s^{p-1} \rangle,$$

where $g_s = \sum_{j=0}^{n-1} v_s^j Z_s^{(n-j)}$, and we rewrite (3.10) as

$$\frac{1}{p} \partial_s \|v_s\|_{L^p}^p = -((p-1) \|v_s^{p-2} |\nabla v_s|^2\|_{L^1} + a_n \|v_s^{p+n-1}\|_{L^1} + \|v_s^p\|_{L^1}) - \langle g_s, v_s^{p-1} \rangle, \quad (3.13)$$

for all $0 < s \leq t$, where we use that p is an even integer. Let

$$K_s := \|v_s^{p-2} |\nabla v_s|^2\|_{L^1}, \quad L_s := a_n \|v_s^{p+n-1}\|_{L^1}. \quad (3.14)$$

The idea is to control the terms of $\langle g_s, v_s^{p-1} \rangle$ by K_s and L_s .

We start with the leading term of $\langle g_s, v_s^{p-1} \rangle$, $\langle v_s^{p+n-2}, Z_s^{(1)} \rangle$. By Proposition A.8

$$\left| \langle v_s^{p+n-2}, Z_s^{(1)} \rangle \right| \lesssim \|v_s^{p+n-2}\|_{\mathcal{B}_{1,1}^\alpha} \|Z_s^{(1)}\|_{\mathcal{C}^{-\alpha}}. \quad (3.15)$$

Using (A.10)

$$\|v_s^{p+n-2}\|_{\mathcal{B}_{1,1}^\alpha} \lesssim \|v_s^{p+n-2}\|_{L^1}^{1-\alpha} \|v_s^{p+n-3} |\nabla v_s|\|_{L^1}^\alpha + \|v_s^{p+n-2}\|_{L^1}. \quad (3.16)$$

We handle each term of (3.16) separately. First we notice, using Jensen's inequality, that $\|v_s^{p+n-2}\|_{L^1} \lesssim L_s^{\frac{p+n-2}{p+n-1}}$. For the gradient term, using the Cauchy–Schwarz inequality we obtain

$$\|v_s^{p+n-3}|\nabla v_s|\|_{L^1} \lesssim \|v_s^{p+2(n-2)}\|_{L^1}^{\frac{1}{2}} K_s^{\frac{1}{2}}. \quad (3.17)$$

Recall the Sobolev inequality

$$\|f\|_{L^q} \lesssim (\|f\|_{L^2}^2 + \|\nabla f\|_{L^2}^2)^{\frac{1}{2}},$$

for every $q < \infty$ (see [16, Section 6], [8, Section 5.6] for Sobolev inequalities in the same spirit). In particular, for $q = \frac{2(p+2(n-2))}{p}$, we have that

$$\|v_s^{\frac{p}{2}}\|_{L^q}^{\frac{q}{2}} \lesssim \|v_s^{\frac{p}{2}}\|_{L^2}^{\frac{q}{2}} + \|\nabla(v_s)^{\frac{p}{2}}\|_{L^2}^{\frac{q}{2}},$$

which implies

$$\|v_s^{p+2(n-2)}\|_{L^1}^{\frac{1}{2}} \lesssim \|v_s^p\|_{L^1}^{\frac{1}{2} + \frac{n-2}{p}} + K_s^{\frac{1}{2} + \frac{n-2}{p}}, \quad (3.18)$$

where $\|v_s^p\|_{L^1}^{\frac{1}{2} + \frac{n-2}{p}} \lesssim L_s^{\frac{\frac{p}{2} + n - 2}{p+n-1}}$ by Jensen's inequality. Combining (3.16), (3.17) and (3.18)

$$\|v_s^{p+n-2}\|_{\mathcal{B}_{1,1}^\alpha} \lesssim K_s^{\frac{\alpha}{2}} L_s^{\frac{(p+n-2) - \frac{p}{2}\alpha}{p+n-1}} + K_s^{(1 + \frac{n-2}{p})\alpha} L_s^{\frac{(p+n-2)(1-\alpha)}{p+n-1}} + L_s^{\frac{p+n-2}{p+n-1}}. \quad (3.19)$$

By (3.12) we notice that

$$\frac{\alpha}{2} + \frac{(p+n-2) - \frac{p}{2}\alpha}{p+n-1} < 1$$

and

$$\left(1 + \frac{n-2}{p}\right)\alpha + \frac{(p+n-2)(1-\alpha)}{p+n-1} < 1,$$

thus we can find $\gamma_1, \gamma_2, \gamma_3, \gamma_4 < 1$ such that

$$\frac{\alpha}{2\gamma_1} + \frac{(p+n-2) - \frac{p}{2}\alpha}{(p+n-1)\gamma_2} = 1$$

and

$$\left(1 + \frac{n-2}{p}\right)\frac{\alpha}{\gamma_3} + \frac{(p+n-2)(1-\alpha)}{(p+n-1)\gamma_4} = 1.$$

In particular, we choose $\gamma_1 = \frac{(p+n-1)\alpha}{2}$, $\gamma_2 = \frac{(p+n-2) - \frac{p}{2}\alpha}{p+n-2}$, $\gamma_3 = \frac{(p+n-2)(p+n-1)\alpha}{p}$ and $\gamma_4 = (1-\alpha)$, apply the classical Young inequality to (3.19) and combine with (3.15) to obtain

$$|\langle v_s^{p+n-2}, Z_s^{(1)} \rangle| \lesssim (K_s^{\gamma_1} + L_s^{\gamma_2} + K_s^{\gamma_3} + L_s^{\gamma_4} + L_s^{\frac{p+n-2}{p+n-1}}) \|Z_s^{(1)}\|_{\mathcal{C}^{-\alpha}}.$$

Using Young's inequality once more, now in the form

$$a\zeta^\gamma \leq \gamma \frac{\zeta}{N^{\frac{1}{\gamma}}} + (1-\gamma)(Na)^{\frac{1}{1-\gamma}},$$

for $a = \|Z_s^{(1)}\|_{C^{-\alpha}}$, $\zeta \in \{K_s, L_s\}$, $N = (Cn)^\gamma$ and $\gamma \in \{\gamma_1, \dots, \gamma_5\}$, where $\gamma_5 = \frac{p+n-2}{p+n-1}$, we obtain the final bound

$$\langle v_s^{p+n-2}, Z_s^{(1)} \rangle \leq \frac{1}{n} \left(K_s + \frac{1}{2} L_s \right) + C \sum_{i=1}^5 \left(\|Z_s^{(1)}\|_{C^{-\alpha}}^{\frac{1}{1-\gamma_i}} \right), \quad (3.20)$$

where C is a constant depending only on γ_i , $i \in \{1, 2, \dots, 5\}$, and n .

For the remaining terms of $\langle g_s, v_s^{p-1} \rangle$ we need to estimate $\langle v_s^{p+j-1}, Z_s^{(n-j)} \rangle$, for all $0 \leq j \leq n-2$. Proceeding in the same spirit of calculations as above we first obtain that

$$\|v_s^{p+j-1}\|_{B_{1,1}^\alpha} \lesssim K_s^\alpha L_s^{\frac{(p+j-1)-\frac{p}{2}\alpha}{p+n-1}} + K_s^{(1+\frac{j-1}{p})\alpha} L_s^{\frac{(p+j-1)(1-\alpha)}{p+n-1}} + L_s^{\frac{p+j-1}{p+n-1}}.$$

We define the exponents $\gamma_1^j = \frac{(p+n-1)\alpha}{2}$, $\gamma_2^j = \frac{(p+j-1)-\frac{p}{2}\alpha}{p+n-2}$, $\gamma_3^j = \frac{(p+j-1)(p+j)\alpha}{p}$ and $\gamma_4^j = \frac{(p+j)(1-\alpha)}{p+n-1}$. Note that (3.12) implies that $\gamma_1^j, \gamma_2^j, \gamma_3^j, \gamma_4^j < 1$ and we also have that

$$\frac{\alpha}{2\gamma_1^j} + \frac{(p+j-1) - \frac{p}{2}\alpha}{(p+n-1)\gamma_2^j} = 1$$

and

$$\left(1 + \frac{j-1}{p}\right) \frac{\alpha}{\gamma_3^j} + \frac{(p+j-1)(1-\alpha)}{(p+n-1)\gamma_4^j} = 1.$$

Applying Young's inequality once more

$$\langle v_s^{p+j-1}, Z_s^{(n-j)} \rangle \lesssim (K_s^{\gamma_1^j} + L_s^{\gamma_2^j} + K_s^{\gamma_3^j} + L_s^{\gamma_4^j} + L_s^{\frac{p+j-1}{p+n-1}}) \|Z_s^{(n-j)}\|_{C^{-\alpha}}.$$

As before (see (3.20)), we obtain the bound

$$\langle v_s^{p+j-1}, Z_s^{(n-j)} \rangle \leq \frac{1}{n} \left(K_s + \frac{1}{2} L_s \right) + C \sum_{i=1}^5 \left(\|Z_s^{(n-j)}\|_{C^{-\alpha}}^{\frac{1}{1-\gamma_i^j}} \right), \quad (3.21)$$

for all $0 \leq j \leq n-2$, where $\gamma_5^j = \frac{p+j-1}{p+n-1}$. Thus, by (3.20) and (3.21),

$$\langle g_s, v_s^{p-1} \rangle \leq \left(K_s + \frac{1}{2} L_s \right) + C \sum_{j=0}^{n-1} \sum_{i=1}^5 \left(\|Z_s^{(n-j)}\|_{C^{-\alpha}}^{\frac{1}{1-\gamma_i^j}} \right), \quad (3.22)$$

where $\gamma_i^{n-1} = \gamma_i$, for all $i \in \{1, \dots, 5\}$.

Finally, for $p_i^j = \frac{1}{1-\gamma_i^j}$, combining (3.13) and (3.22) we obtain

$$\frac{1}{p} \partial_s \|v_s\|_{L^p}^p + \|v_s\|_{L^p}^p + (p-2)K_s + \frac{1}{2}L_s \leq C \sum_{j,i} \|Z_s^{(n-j)}\|_{C^{-\alpha}}^{p_i^j}.$$

Let $t > s$ and notice that by (3.4), for $r \in (s, t)$,

$$\sum_{j,i} \|Z_r^{(n-j)}\|_{C^{-\alpha}}^{p_i^j} \leq \sum_{j,i} r^{-\alpha' p_i^j} \sup_{s \leq r \leq t} (r^{\alpha' p_i^j} \|Z_r^{(n-j)}\|_{C^{-\alpha}}^{p_i^j})$$

for every $\alpha' > 0$. Thus for $r \in [s, t]$

$$\frac{1}{p} \partial_r \|v_r\|_{L^p}^p + \frac{1}{2} L_r \leq C \sum_{j,i} s^{-\alpha' p_i^j} \sup_{s \leq r \leq t} (r^{\alpha' p_i^j} \|Z_r^{(n-j)}\|_{C^{-\alpha}}^{p_i^j}).$$

By Jensen's inequality, for $\lambda = \frac{p+n-1}{p}$, we get that

$$\partial_r \|v_r\|_{L^p}^p + C_1 (\|v_r\|_{L^p}^p)^\lambda \leq C_2 \sum_{j,i} s^{-\alpha' p_i^j} \sup_{s \leq r \leq t} (r^{\alpha' p_i^j} \|Z_r^{(n-j)}\|_{C^{-\alpha}}^{p_i^j}),$$

and if we let $f(r) = \|v_r\|_{L^p}^p$, $r \geq s$, by Lemma 3.8

$$f(r) \leq \frac{f(s)}{(1 + (r - s)f(s)^{\lambda-1}(\lambda - 1)\tilde{C}_1)^{\frac{1}{\lambda-1}}} \vee \left(\frac{2C_2}{C_1} \sum_{j,i} s^{-\alpha' p_i^j} \sup_{s \leq r \leq t} (r^{\alpha' p_i^j} \|Z_r^{(n-j)}\|_{C^{-\alpha}}^{p_i^j}) \right)^{\frac{1}{\lambda}}, \tag{3.23}$$

where $\tilde{C}_1 = C_1/2$. In particular for $r = t$ and $s = t/2$ we have the bound

$$\|v_t\|_{L^p}^p \leq C \left[t^{-\frac{1}{\lambda-1}} \vee \left(\sum_{j,i} t^{-\alpha' p_i^j} \sup_{0 \leq r \leq t} (r^{\alpha' p_i^j} \|Z_r^{(n-j)}\|_{C^{-\alpha}}^{p_i^j}) \right)^{\frac{1}{\lambda}} \right],$$

which completes the proof. □

Lemma 3.8 (Comparison test). *Let $\lambda > 1$ and $f : [0, T] \rightarrow [0, \infty)$ differentiable such that*

$$f'(t) + c_1 f(t)^\lambda \leq c_2,$$

for every $t \in [0, T]$. Then for $t > 0$

$$f(t) \leq \frac{f(0)}{(1 + tf(0)^{\lambda-1}(\lambda - 1)\frac{c_1}{2})^{\frac{1}{\lambda-1}}} \vee \left(\frac{2c_2}{c_1} \right)^{\frac{1}{\lambda}} \leq t^{-\frac{1}{\lambda-1}} \left((\lambda - 1)\frac{c_1}{2} \right)^{-\frac{1}{\lambda-1}} \vee \left(\frac{2c_2}{c_1} \right)^{\frac{1}{\lambda}}.$$

Proof. Let $t > 0$. Then one of the following holds:

- (I) There exists $s_0 \leq t$ such that $f(s_0) \leq \left(\frac{2c_2}{c_1}\right)^{\frac{1}{\lambda}}$.
- (II) For every $s \leq t$, $f(s) > \left(\frac{2c_2}{c_1}\right)^{\frac{1}{\lambda}}$.

In the second case, using the assumption we have that for every $s \leq t$

$$f'(s) + \frac{c_1}{2} f(s)^\lambda \leq 0$$

and solving the above differential inequality on $[0, t]$ implies that

$$f(t) \leq \frac{f(0)}{(1 + tf(0)^{\lambda-1}(\lambda - 1)\frac{c_1}{2})^{\frac{1}{\lambda-1}}}.$$

In the first case, assume for contradiction that $f(t) > \left(\frac{2c_2}{c_1}\right)^{\frac{1}{\lambda}}$ and let

$$s^* = \sup \left\{ s < t : f(s) \leq \left(\frac{2c_2}{c_1}\right)^{\frac{1}{\lambda}} \right\}.$$

Then $f(s) > (\frac{2c_2}{c_1})^{\frac{1}{\lambda}}$, for every $s \in (s^*, t]$, while $f(s^*) = (\frac{2c_2}{c_1})^{\frac{1}{\lambda}}$ by continuity. However, the assumption implies

$$f'(s) + \frac{c_1}{2} f(s)^\lambda \leq 0$$

and in particular $f'(s) \leq 0$. But then

$$f(t) = f(s^*) + \int_{s^*}^t f'(s) ds \leq \left(\frac{2c_2}{c_1}\right)^{\frac{1}{\lambda}},$$

which is a contradiction. \square

The next theorem implies global existence of (3.3). Though it was already established in [15], we present it here for completeness.

Theorem 3.9. *For every initial condition $x \in \mathcal{C}^{-\alpha_0}$ and $\beta > 0$ as in (1.6) there exists a unique solution $v \in C((0, \infty); \mathcal{C}^\beta)$ of (3.3).*

Proof. Let $T > 0$. First fix any even integer $p \geq 2$ such that $L^p \hookrightarrow \mathcal{C}^{-\alpha_0}$ (for example $p \geq \frac{2}{\alpha_0}$ is enough; see also Proposition A.3 and (A.6)). Then the a priori estimate (3.11) (which depends only on $\|\mathbb{Z}\|_{\alpha; \alpha'; T}$) provides an a priori estimate on $\|v_t\|_{\mathcal{C}^{-\alpha_0}}$. Thus by Theorem 3.3 there exists $T^* \leq T$ bounded from below (by a constant depending only on the a priori estimate on $\|v_t\|_{\mathcal{C}^{-\alpha_0}}$) and a unique solution up to time T^* of (3.3). Using again Theorem 3.3 we construct a solution of (3.3) on $[T^*, 2T^* \wedge T]$ with initial condition v_{T^*} which satisfies the same a priori bounds depending on $\|\mathbb{Z}\|_{\alpha; \alpha'; T}$. We then proceed similarly until the whole interval $[0, T]$ is covered. To prove uniqueness we proceed as in the proof of Theorem [15, Theorem 6.2]. \square

Corollary 3.10. *For $x \in \mathcal{C}^{-\alpha_0}$ let $X(\cdot; x) = \mathbf{!}_{0, \cdot} + v$, where v is the solution to (3.2). Then for every $\alpha > 0$ and $p \geq 2$*

$$\sup_{x \in \mathcal{C}^{-\alpha_0}} \sup_{t \geq 0} (t^{\frac{p}{n-1}} \wedge 1) \mathbb{E} \|X(t; x)\|_{\mathcal{C}^{-\alpha}}^p < \infty. \quad (3.24)$$

Remark 3.11. Notice that the bound (3.24) does not follow immediately by taking the expectation of the a priori bound (3.11) on v_t . In fact the expectation of the supremum $\sup_{0 \leq r \leq t} (r^{\alpha'} p_i^j \|Z_s^{(n-j)}\|_{\mathcal{C}^{-\alpha}}^{p_j^i})$ on the right-hand side of this estimate is finite for every $t < \infty$ but it is not uniformly bounded in t . However, as (3.11) does not depend on the initial condition we can just restart (3.1) at time $t - 1$ for $t > 1$ and apply Proposition 3.7 for the restarted solution to obtain a bound which depends only on the randomness inside the interval $[t - 1, t]$. Given that the diagrams have the same law on intervals of the same size (see Proposition 2.3) we then obtain a bound which is independent of t .

Proof. Let $t > 1$ and notice that by Lemma 4.1 (see Section 4.2 for statement and proof) $X(t; x) = \mathbf{!}_{t-1, t} + \tilde{v}_{t-1, t}$ where $\tilde{v}_{t-1, r}$, $r \geq t - 1$, solves (3.2) with initial condition $X(t - 1; x)$ and

$$Z^{(n-j)} = \sum_{k=j}^n a_k \binom{k}{j} \mathbf{!}_{t-1, t-1+}^k,$$

for every $0 \leq j \leq n - 1$. Applying Proposition 3.7 on $\tilde{v}_{t-1, \cdot}$ we then have

$$\|\tilde{v}_{t-1, t}\|_{L^p}^p \lesssim 1 \vee \left(\sum_{j,i} \sup_{t-1 \leq r \leq t} ((r - (t - 1))^{\alpha'} p_i^j \|Z_r^{(n-j)}\|_{\mathcal{C}^{-\alpha}}^{p_j^i}) \right)^{\frac{1}{\lambda}}, \quad (3.25)$$

for every $p \geq 2$. To prove (3.24) we fix $\alpha > 0$ and using the embedding $L^p \hookrightarrow \mathcal{C}^{-\alpha}$ for $p \geq \frac{2}{\alpha}$ (see (A.6) and Proposition A.3) we first notice that for $t > 1$

$$\mathbb{E} \|X(t; x)\|_{\mathcal{C}^{-\alpha}}^p \lesssim \mathbb{E} \|\mathbf{!}_{t-1, t}\|_{\mathcal{C}^{-\alpha}}^p + \mathbb{E} \|\tilde{v}_{t-1, t}\|_{\mathcal{C}^{-\alpha}}^p \lesssim \mathbb{E} \|\mathbf{!}_{t-1, t}\|_{\mathcal{C}^{-\alpha}}^p + \mathbb{E} \|\tilde{v}_{t-1, t}\|_{L^p}^p.$$

Combining with (3.25) and given that for every $k \geq 1$ the law of $\mathbb{V}_{t-1,t+}$ does not depend on t we obtain that

$$\sup_{t \geq 1} \mathbb{E} \|X(t; x)\|_{\mathcal{C}^{-\alpha}}^p < \infty.$$

Finally, using (3.11) (and by possibly tuning down α' in the same equation) for $t \leq 1$ we get

$$\mathbb{E} \|X(t; x)\|_{\mathcal{C}^{-\alpha}}^p \lesssim \mathbb{E} \|\mathfrak{I}_{-\infty,t}\|_{\mathcal{C}^{-\alpha}}^p + \mathbb{E} \|v_t\|_{L^p}^p \lesssim 1 + t^{-\frac{p}{n-1}},$$

which completes the proof. □

4. Existence of invariant measures

4.1. Markov property

For $x \in \mathcal{C}^{-\alpha_0}$ we write $X(\cdot; x) = \mathfrak{I}_{0,\cdot} + v$ where v is the solution to (3.2) with initial condition x . We introduce a variant of the notation (3.5) and set

$$\tilde{F}(v, (\mathbb{V}_{0,\cdot}^k)_{k=1}^n) = \sum_{k=0}^n a_k \sum_{j=0}^k \binom{k}{j} v^j \mathbb{V}_{0,\cdot}^{k-j}. \tag{4.1}$$

We denote by $B_b(\mathcal{C}^{-\alpha_0})$ and $C_b(\mathcal{C}^{-\alpha_0})$ the spaces of bounded and continuous functions from $\mathcal{C}^{-\alpha_0}$ to \mathbb{R} , both endowed with the norm

$$\|\Phi\|_\infty := \sup_{x \in \mathcal{C}^{-\alpha_0}} |\Phi(x)|.$$

For every $\Phi \in B_b(\mathcal{C}^{-\alpha_0})$ and $t \in [0, \infty)$ we define the map $P_t : \Phi \mapsto P_t \Phi$ by

$$P_t \Phi(x) := \mathbb{E} \Phi(X(t; x)), \tag{4.2}$$

for every $x \in \mathcal{C}^{-\alpha_0}$.

In this section we prove that $\{X(t; \cdot) : t \geq 0\}$ is a Markov process with transition semigroup $\{P_t : t \geq 0\}$ with respect to the filtration $\{\mathcal{F}_t : t \geq 0\}$ defined in (2.1).

We first prove the following lemma.

Lemma 4.1. *Let $X(\cdot; x) = \mathfrak{I}_{0,\cdot} + v$. Then, for every $h > 0$,*

$$X(t+h; x) = \mathfrak{I}_{t,t+h} + \tilde{v}_{t,t+h},$$

where the remainder $\tilde{v}_{t,t+}$ solves (3.2) driven by the vector $(\mathbb{V}_{t,t+}^k)_{k=1}^n$ and initial condition $X(t; x)$, i.e.

$$\tilde{v}_{t,t+h} = S(h)X(t; x) - \int_0^h S(h-r) \tilde{F}(\tilde{v}_{t,t+r}, (\mathbb{V}_{t,t+r}^k)_{k=1}^n) dr.$$

Proof. Notice that for $h > 0$

$$X(t+h; x) = \mathfrak{I}_{0,t+h} + v_{t+h} = \mathfrak{I}_{t,t+h} + \tilde{v}_{t,t+h},$$

where

$$\tilde{v}_{t,t+h} = S(h)X(t; x) - \int_0^h S(h-r) \tilde{F}(v_{t+r}, (\mathbb{V}_{0,t+r}^k)_{k=1}^n) dr.$$

By (2.11) we have that

$$\begin{aligned}\tilde{F}(v_{t+r}, (\check{\nabla}_{0,t+r}^k)_{k=1}^n) &= \sum_{k=0}^n a_k \sum_{j=0}^k \binom{k}{j} v_{t+r}^j \check{\nabla}_{0,t+r}^{k-j}, \\ &= \sum_{k=0}^n a_k \sum_{i=0}^k \binom{k}{i} \tilde{v}_{t,t+r}^i \check{\nabla}_{t,t+r}^{k-i},\end{aligned}$$

where we use a binomial expansion of v_{t+r}^j and a change of summation. Hence

$$\tilde{F}(v_{t+r}, (\check{\nabla}_{0,t+r}^k)_{k=1}^n) = \tilde{F}(\tilde{v}_{t,t+r}, (\check{\nabla}_{t,t+r}^k)_{k=1}^n),$$

which completes the proof. \square

The fact that $\{X(t; \cdot) : t \geq 0\}$ is a Markov process is an immediate consequence of the following theorem.

Theorem 4.2. *Let $X(\cdot; x)$ be as in the lemma above with $x \in \mathcal{C}^{-\alpha_0}$. Then for every $\Phi \in B_b(\mathcal{C}^{-\alpha_0})$ and $t \geq 0$*

$$\mathbb{E}(\Phi(X(t+h; x)) | \mathcal{F}_t) = P_h \Phi(X(t; x)),$$

for all $h \geq 0$.

Proof. Let $h \geq 0$ and $\Phi \in B_b(\mathcal{C}^{-\alpha_0})$ and write

$$\mathcal{T}(X(t; x); h; (\check{\nabla}_{t,t+h}^k)_{k=1}^n)$$

to denote the solution of (3.2) at time h , driven by the vector $(\check{\nabla}_{t,t+h}^k)_{k=1}^n$ and initial condition $X(t; x)$. By Proposition 2.4 and [6, Proposition 1.12]

$$\mathbb{E}(\Phi(X(t+h; x)) | \mathcal{F}_t) = \bar{\Phi}(X(t; x)),$$

where for $w \in \mathcal{C}^{-\alpha_0}$

$$\bar{\Phi}(w) = \mathbb{E}\Phi(\mathcal{I}_{t,t+h} + \mathcal{T}(w; h; (\check{\nabla}_{t,t+h}^k)_{k=1}^n)).$$

Here we use the fact that $X(t; x)$ is \mathcal{F}_t -measurable and that the vector $(\check{\nabla}_{t,t+h}^k)_{k=1}^n$ is \mathcal{F}_t -independent (see Proposition 2.3). Given that $(\check{\nabla}_{t,t+h}^k)_{k=1}^n \stackrel{\text{law}}{=} (\check{\nabla}_{0,\cdot}^k)_{k=1}^n$ (see again Proposition 2.3) and the fact that (3.2) has a unique solution driven by any vector $\underline{Y} \in \mathcal{C}^{n, -\alpha}(0; T)$, for $T > 0$, and any initial condition $w \in \mathcal{C}^{-\alpha_0}$, we have that

$$\bar{\Phi}(w) = P_h \Phi(w),$$

which completes the proof if we set $w = X(t; x)$. \square

The theorem above implies that $\{P_t : t \geq 0\}$ is a semigroup. We finally prove that it is Feller.

Proposition 4.3. *Let $\Phi \in C_b(\mathcal{C}^{-\alpha_0})$. Then, for every $t \geq 0$, $P_t \Phi \in C_b(\mathcal{C}^{-\alpha_0})$.*

Proof. It suffices to prove that the solution to (3.2) is continuous with respect to its initial condition. Fix $T > 0$ and $x \in \mathcal{C}^{-\alpha_0}$. Let $y \in \mathcal{C}^{-\alpha_0}$ such that $\|x - y\|_{\mathcal{C}^{-\alpha_0}} \leq 1$ and

$$v_t = S(t)x - \int_0^t S(t-r) \tilde{F}(v_r, (\check{\nabla}_{0,r}^k)_{k=1}^n) dr,$$

$$u_t = S(t)y - \int_0^t S(t-r)\tilde{F}(u_r, (\mathcal{V}_{0,r}^k)_{k=1}^n) dr,$$

as well as $\tau = \inf\{t > 0 : t^\gamma \|v_t - u_t\|_{\mathcal{C}^\beta} > 1\}$ and

$$M = \sup_{t \leq T} t^\gamma \|v_t\|_{\mathcal{C}^\beta}, \quad N = \left\| (\mathcal{V}_{0,\cdot}^k)_{k=1}^n \right\|_{\alpha; \alpha'; T}.$$

Notice that

$$\tilde{F}(v_r, (\mathcal{V}_{0,r}^k)_{k=1}^n) - \tilde{F}(u_r, (\mathcal{V}_{0,r}^k)_{k=1}^n) = \sum_{k=0}^n a_k \sum_{j=0}^k \binom{k}{j} (u_r^k - v_r^k) \mathcal{V}_{0,r}^{k-j},$$

and by Propositions A.5 and A.7 we obtain that for all $T_* \leq T \wedge \tau$

$$\begin{aligned} \sup_{t \leq T_*} t^\gamma \|v_t - u_t\|_{\mathcal{C}^\beta} &\leq \sup_{t \leq T_*} t^\gamma \|v_t - u_t\|_{\mathcal{C}^\beta} \sum_{m=1}^n \lambda_m T_*^{\alpha_m} \\ &\quad + \|x - y\|_{\mathcal{C}^{-\alpha_0}} \sum_{m=n+1}^{2n} \lambda_m T_*^{\alpha_m}, \end{aligned}$$

where $\lambda_m \equiv \lambda_m(M, N, \|x\|_{\mathcal{C}^{-\alpha_0}})$ and $\alpha_m \in (0, 1]$. Choosing $T_* \equiv T_*(M, N, \|x\|_{\mathcal{C}^{-\alpha_0}}) \leq 1/2$ we obtain that

$$\sup_{t \leq T_*} t^\gamma \|v_t - u_t\|_{\mathcal{C}^\beta} \leq \|x - y\|_{\mathcal{C}^{-\alpha_0}}.$$

Iterating the procedure we find $N^* \in \mathbb{Z}_{\geq 0}$ and $C > 0$ such that

$$\sup_{t \leq T \wedge \tau} t^\gamma \|v_t - u_t\|_{\mathcal{C}^\beta} \leq (N^*C + 1) \|x - y\|_{\mathcal{C}^{-\alpha_0}},$$

for every $y \in \mathcal{C}^{-\alpha_0}$ such that $\|x - y\|_{\mathcal{C}^{-\alpha_0}} \leq 1$. At this point we should notice that for every $y \in \mathcal{C}^{-\alpha_0}$ such that $\|x - y\|_{\mathcal{C}^{-\alpha_0}} \leq 1/2(N^*C + 1)$ the above estimate implies that

$$\sup_{t \leq T \wedge \tau} t^\gamma \|v_t - u_t\|_{\mathcal{C}^\beta} \leq \frac{1}{2},$$

thus $T \wedge \tau = T$ because of the definition of τ . Hence, for all such $y \in \mathcal{C}^{-\alpha_0}$,

$$\sup_{t \leq T} t^\gamma \|v_t - u_t\|_{\mathcal{C}^\beta} \leq (N^*C + 1) \|x - y\|_{\mathcal{C}^{-\alpha_0}},$$

which implies convergence of u_t to v_t in \mathcal{C}^β for every $t \leq T$. Since T was arbitrary, the last implies continuity of the solution map of (3.2) with respect to its initial condition. The Feller property is then an immediate consequence of the above combined with the dominated convergence theorem. \square

4.2. Invariant measures

We denote by $\{P_t^* : t \geq 0\}$ the dual semigroup of $\{P_t : t \geq 0\}$ acting on the set of all probability Borel measures on $\mathcal{C}^{-\alpha_0}$ denoted by $\mathcal{M}_1(\mathcal{C}^{-\alpha_0})$. In the next proposition we prove existence of invariant measures of $\{P_t : t \geq 0\}$ as a semigroup acting on $C_b(\mathcal{C}^{-\alpha_0})$.

Proposition 4.4. *For every $x \in \mathcal{C}^{-\alpha_0}$ there exists a measure $\nu_x \in \mathcal{M}_1(\mathcal{C}^{-\alpha_0})$ and a sequence $t_k \nearrow \infty$ such that*

$$\frac{1}{t_k} \int_0^{t_k} P_s^* \delta_x ds \xrightarrow{w} \nu_x.$$

In particular the measure ν_x is invariant for the Markov semigroup $\{P_t : t \geq 0\}$ on $\mathcal{C}^{-\alpha_0}$.

Proof. For $t > 0$ and $\alpha > 0$ using Markov's and Jensen's inequality there exists a constant $C > 0$ such that

$$\mathbb{P}(\|X(t; x)\|_{\mathcal{C}^{-\alpha}} > K) \leq \frac{C}{K} (\mathbb{E}\|X(t; x)\|_{\mathcal{C}^{-\alpha}}^p)^{\frac{1}{p}},$$

for every $K > 0$ and $p \geq 2$. Thus

$$\begin{aligned} \int_0^t \mathbb{P}(\|X(s; x)\|_{\mathcal{C}^{-\alpha}} > K) \, ds &\leq \frac{C}{K} \int_0^t (\mathbb{E}\|X(s; x)\|_{\mathcal{C}^{-\alpha}}^p)^{\frac{1}{p}} \, ds \\ &\leq \frac{C}{K} \left[\int_0^1 s^{-\frac{1}{n-1}} \, ds + \int_1^t \, ds \right] \\ &= \frac{C}{K} t, \end{aligned}$$

where in the second inequality we use (3.24). If we let $R_t = \frac{1}{t} \int_0^t P_s^* \delta_x \, ds$, for $K_\varepsilon = \frac{C}{\varepsilon}$ we get

$$R_t(\{f \in \mathcal{C}^{-\alpha} : \|f\|_{\mathcal{C}^{-\alpha}} > K_\varepsilon\}) \leq \varepsilon.$$

Choosing $\alpha < \alpha_0$ we can ensure that $\{f \in \mathcal{C}^{-\alpha} : \|f\|_{\mathcal{C}^{-\alpha}} \leq K_\varepsilon\}$ is a compact subset of $\mathcal{C}^{-\alpha_0}$ since the embedding $\mathcal{C}^{-\alpha} \hookrightarrow \mathcal{C}^{-\alpha_0}$ is compact for every $\alpha < \alpha_0$ (see Proposition A.4 and (A.2)). This implies tightness of $\{R_t\}_{t \geq 0}$ in $\mathcal{C}^{-\alpha_0}$ and by the Krylov–Bogoliubov existence Theorem (see [7, Corollary 3.1.2]) there exist a sequence $t_k \nearrow \infty$ and a measure $\nu_x \in \mathcal{M}_1(\mathcal{C}^{-\alpha_0})$ such that $R_{t_k} \rightarrow \nu_x$ weakly in $\mathcal{C}^{-\alpha_0}$ and ν_x is invariant for the semigroup $\{P_t : t \geq 0\}$ in $\mathcal{C}^{-\alpha_0}$. \square

5. Strong Feller property

In this section we show that the Markov semigroup $\{P_t : t \geq 0\}$ satisfies the strong Feller property. The strong Feller property is to be expected when we deal with SPDEs where the noise forces every direction in Fourier space. However, the fact that the process X does not solve a self-contained equation forces us to translate everything onto the level of the remainder v . The most important step is to obtain a Bismut–Elworthy–Li formula (see Theorem 5.5) which captures enough information to provide a good control of the linearization of the remainder equation.

On the technical level, we work with a finite dimensional approximation X^ε for X . This choice and the fact that the equation is driven by white noise imply that the solution is Fréchet differentiable with respect to the (finite dimensional approximation of the) noise, so we can avoid working with Malliavin derivatives. This is expressed in Proposition 5.1 below, and in fact this proposition could even be established without splitting X^ε into v^ε and $\mathfrak{I}_{0,\cdot}^\varepsilon$. We make strong use of the splitting in Proposition 5.4 where the local solution theory is used to obtain deterministic bounds on v^ε and its linearization for small t provided that we control the diagrams $\mathfrak{V}_{0,\cdot}^{\varepsilon,n}$. This control is uniform in ε and enters crucially the proof of Proposition 5.8.

From now on we fix $0 < \alpha < \alpha_0$ sufficiently small. For $\varepsilon \in (0, 1)$ let $\Pi_\varepsilon[L^2(\mathbb{T}^2)]$ be the finite dimensional subspace of $L^2(\mathbb{T}^2)$ spanned by $\{e_m\}_{|m| < \frac{1}{\varepsilon}}$ (recall that we deal with real-valued functions and the symmetry condition (1.3) is always valid) and denote by Π_ε the corresponding orthogonal projection. We also let $\hat{\Pi}_\varepsilon$ be a linear smooth approximation taking values in $\Pi_\varepsilon[L^2(\mathbb{T}^2)]$ and having the properties (i) and (ii) introduced in the discussion before Proposition 3.4.

Let \mathfrak{R}^ε be the renormalization constant defined in (2.9) and consider a finite dimensional approximation of (3.1) given by

$$\begin{cases} dX^\varepsilon(t) = (\Delta X^\varepsilon(t) - X^\varepsilon(t) - \sum_{k=0}^n a_k \hat{\Pi}_\varepsilon \mathcal{H}_k(X^\varepsilon(t), \mathfrak{R}^\varepsilon)) dt + dW_\varepsilon(t, \cdot), \\ X^\varepsilon(0, \cdot) = \hat{\Pi}_\varepsilon x, \end{cases} \quad (5.1)$$

for some initial condition $x \in C^{-\alpha_0}$. Here $W_\varepsilon(t, z) = \sum_{|m| < \frac{1}{\varepsilon}} \hat{W}_m(t) e_m(z)$, where $(\hat{W}_m)_{m \in \mathbb{Z}^2}$ is a family of complex Brownian motions such that $\hat{W}_{-m} = \overline{\hat{W}_m}$ and independent otherwise. We furthermore assume that W_ε is defined on the same probability space Ω as ξ via the identity

$$\hat{W}_m(t) := \xi(\mathbf{1}_{[0,t]} \times e_m), \quad m \in \mathbb{Z}^2,$$

which also makes it adapted with respect to the filtration $(\mathcal{F}_t)_{t \geq 0}$. It is convenient to write $W_\varepsilon = G_\varepsilon(\hat{W}_m)_{m \in \mathbb{Z}^2 \cap [-d,d]^2}$ for $G_\varepsilon : C([0, \infty); \mathbb{R}^{(2d-1)^2}) \rightarrow C([0, \infty); \Pi_\varepsilon[L^2(\mathbb{T}^2)])$ such that

$$G_\varepsilon(\hat{W}_m)_{m \in \mathbb{Z}^2 \cap [-d,d]^2} = \sum_{|m| < \frac{1}{\varepsilon}} \hat{W}_m e_m$$

and $d = \lfloor \frac{1}{\varepsilon} \rfloor + \frac{1}{2}$. The Cameron–Martin space of W_ε is given by

$$\mathcal{CM} := W_0^{1,2}([0, \infty)) = \{w : \partial_t w \in L^2([0, \infty); \mathbb{R}^{(2d-1)^2}), w(0) = 0\}.$$

Last, we have the identity

$$\mathfrak{I}_{0,t}^\varepsilon = \sum_{|m| < \frac{1}{\varepsilon}} \int_0^t e^{-(1+4\pi^2|m|^2)(t-s)} d\hat{W}_m(s) e_m, \tag{5.2}$$

where $\mathfrak{I}_{0,\cdot}^\varepsilon$ is the finite dimensional approximation defined in Section 2.2.

For $v \in C^\beta$ and $\underline{Z} \in (C^{-\alpha})^n$, $\alpha < \beta$, we use the notation

$$\tilde{F}(v, \underline{Z}) = \sum_{k=0}^n a_k \sum_{j=0}^k \binom{k}{j} v^j Z^{(k-j)}$$

with the convention that $Z^{(0)} \equiv 1$. Recall that \tilde{F} (see (4.1)) is a variant of F in (3.5). Here and for the rest of this section \underline{Z} is a shortcut for $(\mathfrak{V}_{0,\cdot}^\varepsilon)_{k=1}^n$ (notice that this differs from the convention used in Section 3). We also let

$$\tilde{F}'(v, \underline{Z}) = \sum_{k=1}^n k a_k \sum_{j=0}^{k-1} \binom{k-1}{j} v^j Z^{(k-1-j)}.$$

Formally, \tilde{F}' stands for the derivative of $\sum_{k=0}^n a_k : X^k :$ with respect to X , with $: X^k :$ replaced by $\sum_{j=0}^k \binom{k}{j} v^j Z^{(k-j)}$.

From now on we also denote by \mathcal{D} the Fréchet derivative with respect to elements in $C([0, t]; \mathbb{R}^{(2d-1)^2})$ (i.e. with respect to the noise), for $t > 0$, and by D the Fréchet derivative with respect to elements in $C^{-\alpha_0}$ (i.e. with respect to the initial condition).

Existence and uniqueness of local in time solutions to (5.1) up to some random explosion time $T_\varepsilon^* > 0$ can be proven following the same method as in Section 3, i.e. using the ansatz $X^\varepsilon = \mathfrak{I}_{0,\cdot}^\varepsilon + v^\varepsilon$ and solving the PDE problem

$$\begin{cases} \partial_t v^\varepsilon = \Delta v^\varepsilon - v^\varepsilon - \hat{\Pi}_\varepsilon \tilde{F}(v^\varepsilon, \underline{Z}^\varepsilon), \\ v^\varepsilon(0, \cdot) = \hat{\Pi}_\varepsilon x, \end{cases} \tag{5.3}$$

where $\underline{Z}^\varepsilon = (\mathfrak{V}_{0,\cdot}^\varepsilon)_{k=1}^n$ (see Section 2.2 for definitions).

Notice that for fixed v , \tilde{F} is Fréchet differentiable with respect to any $\underline{Z} \in (C^{-\alpha})^n$ as a function taking values in $C^{-\alpha}$. Recall that $\mathfrak{V}_{0,\cdot}^\varepsilon = \mathcal{H}_k(\mathfrak{I}_{0,\cdot}^\varepsilon, \mathfrak{R}^\varepsilon)$, for every $1 \leq k \leq n$, so that the map

$$(v, Z^{(1)}) \mapsto S(t) \hat{\Pi}_\varepsilon x - \int_0^t S(t-s) \hat{\Pi}_\varepsilon \tilde{F}(v_s, (\mathcal{H}_k(Z^{(1)}, \mathfrak{R}^\varepsilon))_{k=1}^n) ds, \tag{5.4}$$

for $(v, Z^{(1)}) \in C([0, t]; \mathcal{C}^\beta) \times C([0, t]; \Pi_\varepsilon[L^2(\mathbb{T}^2)])$ and $t > 0$, is Fréchet differentiable as a composition of \tilde{F} with a linear operator shifted by a constant, since the mapping

$$C([0, t]; \Pi_\varepsilon[L^2(\mathbb{T}^2)]) \ni Z^{(1)} \mapsto (\mathcal{H}_k(Z^{(1)}, \mathfrak{N}^\varepsilon))_{k=1}^n \in C^{n, -\alpha}(0; t)$$

is Fréchet differentiable for any $\alpha > 0$, with respect to any $\|\cdot\|_{\alpha; \alpha'; t}$, for $\alpha' > 0$ fixed. Thus, for fixed $x \in \mathcal{C}^{-\alpha_0}$ and $\mathfrak{I}_{0,\cdot}^\varepsilon \in C([0, t]; \Pi_\varepsilon[L^2(\mathbb{T}^2)])$ the implicit function theorem for Banach spaces (see [23, Theorem 4E]) can be applied up to time $T_\varepsilon^* \equiv T_\varepsilon^*(x, \mathfrak{I}_{0,\cdot}^\varepsilon)$ where existence of v^ε is ensured. Hence, for $t \in (0, T_\varepsilon^*)$ there exists an open neighborhood $\mathcal{U}_{\mathfrak{I}_{0,\cdot}^\varepsilon} \subset C([0, t]; \Pi_\varepsilon[L^2(\mathbb{T}^2)])$ of $\mathfrak{I}_{0,\cdot}^\varepsilon$ such that the solution map $\mathcal{T}_t^{\varepsilon, x} : \mathcal{U}_{\mathfrak{I}_{0,\cdot}^\varepsilon} \rightarrow \mathcal{C}^\beta$ of (5.3) is Fréchet differentiable at $\mathfrak{I}_{0,\cdot}^\varepsilon$.

Using Itô's formula the stochastic integrals in (5.2) can be written as

$$\int_0^\cdot e^{-(1+4\pi^2|m|^2)(\cdot-s)} d\hat{W}_m(s) = \hat{W}_m(\cdot) - (1 + 4\pi^2|m|^2) \int_0^\cdot e^{-(1+4\pi^2|m|^2)(\cdot-s)} \hat{W}_m(s) ds. \quad (5.5)$$

Notice that the right-hand side in the above equation is well-defined if we replace $(\hat{W}_m)_{m \in \mathbb{Z}^2 \cap [-d, d]^2}$ by any $w \in C([0, t]; \mathbb{R}^{(2d-1)^2})$, therefore (5.5) is a continuous linear function on $C([0, t]; \mathbb{R}^{(2d-1)^2})$. Thus $\mathfrak{I}_{0,\cdot}^\varepsilon$ as a function from $C([0, t]; \mathbb{R}^{(2d-1)^2})$ to $C([0, t]; \Pi_\varepsilon[L^2(\mathbb{T}^2)])$ is Fréchet differentiable. Combining all the above we finally obtain Fréchet differentiability of v_t^ε from $C([0, t]; \mathbb{R}^{(2d-1)^2})$ to \mathcal{C}^β .

We let $\hat{W}_\varepsilon = (\hat{W}_m)_{|m| < \frac{1}{\varepsilon}}$ and for $w \in C([0, t]; \mathbb{R}^{(2d-1)^2})$ we write

$$\int_0^t S(t-s) G_\varepsilon dw(s) := \sum_{|m| < \frac{1}{\varepsilon}} \int_0^t e^{-(1+4\pi^2|m|^2)(t-s)} dw_m(s),$$

where the right-hand side is defined as in (5.5) with \hat{W}_m replaced by w_m .

In the next proposition we summarize the results of the previous discussion.

Proposition 5.1. Fix $x \in \mathcal{C}^{-\alpha_0}$, $\hat{W}_\varepsilon \in C([0, \infty); \mathbb{R}^{(2d-1)^2})$ and $\mathfrak{I}_{0,\cdot}^\varepsilon \equiv \mathfrak{I}_{0,\cdot}^\varepsilon(\hat{W}_\varepsilon) \in C([0, \infty); \Pi_\varepsilon[L^2(\mathbb{T}^2)])$ and let $T_\varepsilon^* \equiv T_\varepsilon^*(x, \mathfrak{I}_{0,\cdot}^\varepsilon) > 0$ be the explosion time of v^ε . Then for all $t < T_\varepsilon^*$ there exists an open neighborhood $\mathcal{O}_{\hat{W}_\varepsilon} \subset C([0, t]; \mathbb{R}^{(2d-1)^2})$ of \hat{W}_ε such that $X^\varepsilon(t; x) (= \mathfrak{I}_{0,t}^\varepsilon + v_t^\varepsilon)$ is Fréchet differentiable as a function from $\mathcal{O}_{\hat{W}_\varepsilon}$ to $\mathcal{C}^{-\alpha_0}$ and for any $w \in C([0, t]; \mathbb{R}^{(2d-1)^2})$ its directional derivative $\mathcal{D}X^\varepsilon(t; x)(w)$ is given in mild form as

$$\mathcal{D}X^\varepsilon(t; x)(w) = - \int_0^t S(t-s) \hat{\Pi}_\varepsilon[\tilde{F}'(v_s^\varepsilon, \underline{Z}_s^\varepsilon) \mathcal{D}X^\varepsilon(s; x)(w)] ds + \int_0^t S(t-s) G_\varepsilon dw(s). \quad (5.6)$$

Remark 5.2. We expect that $T_\varepsilon^* = +\infty$, which we already established in the limiting case $\varepsilon = 0$ in Section 3. However we only use the local solution theory to control the semigroup associated to $X^\varepsilon(t; x)$ (see Proposition 5.8), thus we do not insist on proving a global existence theorem. We then pass to the limit using the fact that $T_\varepsilon^* \rightarrow \infty$ (see the discussion above Remark 5.9).

Proof. The Fréchet differentiability of $X^\varepsilon(t; x)$ follows by the discussion above and (5.6) by differentiating (5.4). \square

For $h \in \mathcal{C}^{-\alpha_0}$, we let $h_\varepsilon = \hat{\Pi}_\varepsilon h$ and for $t \geq s$ we also consider the following linear equation,

$$\begin{cases} \partial_t J_{s,t}^\varepsilon h_\varepsilon = \Delta J_{s,t}^\varepsilon h_\varepsilon - J_{s,t}^\varepsilon h_\varepsilon - \hat{\Pi}_\varepsilon[\tilde{F}'(v_t^\varepsilon, \underline{Z}_t^\varepsilon) J_{s,t}^\varepsilon h_\varepsilon], \\ J_{s,s}^\varepsilon h_\varepsilon = h_\varepsilon. \end{cases} \quad (5.7)$$

Then $J_{0,t}^\varepsilon h_\varepsilon = \mathcal{D}X^\varepsilon(t; x)(h)$, i.e. it is the derivative of $X^\varepsilon(t; \cdot)$ in the direction h , and its existence for every $t \leq T_\varepsilon^*$ is ensured by a similar argument as the one discussed before Proposition 5.1.

At this point we should comment on the relation between (5.6) and (5.7). Given that (5.7) has a unique solution for every $h_\varepsilon \in \Pi_\varepsilon[L^2(\mathbb{T}^2)]$ up to time $t > 0$, then for $w \in \mathcal{CM}$, i.e. $w(0) = 0$ and $\partial_t w \in L^2([0, \infty); \mathbb{R}^{(2d-1)^2})$, by Duhamel’s principle

$$DX^\varepsilon(t; x)(w) = \int_0^t J_{s,t}^\varepsilon G_\varepsilon \partial_s w(s) ds, \tag{5.8}$$

where $J_{s,t}^\varepsilon : \mathcal{C}^{-\alpha_0} \rightarrow \mathcal{C}^\beta$ is the solution map of (5.7).

Remark 5.3. In the framework of Malliavin calculus $\mathcal{D}_s X^\varepsilon(t; x) = J_{s,t}^\varepsilon G_\varepsilon$ as an element of the dual of $L^2([0, \infty); \mathbb{R}^{(2d-1)^2})$ is the Malliavin derivative (see [18, Section 1.2]) in the sense that the latter coincides with the former when it acts on $X^\varepsilon(t; x)$. In our case, the presence of additive noise implies Fréchet differentiability with respect to the noise as an element in $C([0, t]; \mathbb{R}^{(2d-1)^2})$ (see Proposition 5.1), which is of course stronger than Malliavin differentiability with respect to the noise.

For $r \in [\frac{1}{4}, 1]$ (the precise value of r will be fixed below) and $0 < \alpha' < \alpha$ we consider the stopping times

$$\begin{aligned} \tau^{\varepsilon,r} &:= \inf\{t > 0 : \|\uparrow_{0,t}^\varepsilon\|_{\mathcal{C}^{-\alpha}} \vee t^{\alpha'} \|\mathbf{V}_{0,t}^\varepsilon\|_{\mathcal{C}^{-\alpha}} \vee \dots \vee t^{\alpha'(n-1)} \|\overset{\dots}{\mathbf{V}}_{0,t}^\varepsilon\|_{\mathcal{C}^{-\alpha}} > r\}, \\ \tau^r &:= \inf\{t > 0 : \|\uparrow_{0,t}\|_{\mathcal{C}^{-\alpha}} \vee t^{\alpha'} \|\mathbf{V}_{0,t}\|_{\mathcal{C}^{-\alpha}} \vee \dots \vee t^{(n-1)\alpha'} \|\overset{\dots}{\mathbf{V}}_{0,t}\|_{\mathcal{C}^{-\alpha}} > r\}. \end{aligned} \tag{5.9}$$

Let $\bar{B}_1(x)$ be the closed unit ball centered at x in $\mathcal{C}^{-\alpha_0}$. The next proposition provides local bounds on v^ε and $J_{0,\cdot}^\varepsilon$, given deterministic control on $\underline{Z}^\varepsilon$ (see also Theorem 3.3).

Proposition 5.4. *Let $x \in \mathcal{C}^{-\alpha_0}$ and let $R = 2\|x\|_{\mathcal{C}^{-\alpha_0}} + 1$. Then there exists a deterministic time $T^* \equiv T^*(R) > 0$, independent of ε , such that for all $t \leq T^* \wedge \tau^{\varepsilon,r}$ and initial conditions $y \in \bar{B}_1(x)$,*

$$\sup_{s \leq t} s^\gamma \|v_s^\varepsilon\|_{\mathcal{C}^\beta} \leq 1 \quad \text{and} \quad \sup_{s \leq t} s^\gamma \|J_{0,s}^\varepsilon h_\varepsilon\|_{\mathcal{C}^\beta} \leq 2\|h_\varepsilon\|_{\mathcal{C}^{-\alpha_0}},$$

for β, γ as in (1.6), uniformly in ε , for every $h_\varepsilon \in \hat{\Pi}_\varepsilon[L^2(\mathbb{T}^2)]$.

Proof. Let $t \leq \tau^{\varepsilon,r} \wedge T^*$ where $T^* \equiv T^*(R)$ is defined as in (3.7). We can also assume that $t \leq 1$. Then, from Theorem 3.3, we have that

$$\sup_{s \leq t} s^\gamma \|v_s^\varepsilon\|_{\mathcal{C}^\beta} \leq 1,$$

for every $y \in \bar{B}_1(x)$. Using Proposition A.5, (A.3) and (A.4) we get that

$$\|S(t-s)\hat{\Pi}_\varepsilon[\tilde{F}'(v_s^\varepsilon, \underline{Z}_s^\varepsilon)J_{0,s}^\varepsilon h_\varepsilon]\|_{\mathcal{C}^\beta} \lesssim (s^{-(n-1)\gamma} + (t-s)^{-\frac{\beta+\alpha}{2}} s^{-(n-2)\gamma}) \|J_{0,s}^\varepsilon h_\varepsilon\|_{\mathcal{C}^\beta}, \tag{5.10}$$

where we also use the fact that $\|\hat{\Pi}_\varepsilon f\|_{\mathcal{C}^{-\alpha}} \lesssim \|f\|_{\mathcal{C}^{-\alpha}}$, for every $f \in \mathcal{C}^{-\alpha}$. We are now ready to retrieve the appropriate bounds on the operator norm of $J_{0,\cdot}^\varepsilon$. For $h_\varepsilon \in \Pi_\varepsilon[L^2(\mathbb{T}^2)]$ we have in mild form,

$$J_{0,t}^\varepsilon h_\varepsilon = S(t)h_\varepsilon - \int_0^t S(t-s)\hat{\Pi}_\varepsilon[\tilde{F}'(v_s^\varepsilon, \underline{Z}_s^\varepsilon)J_{0,s}^\varepsilon h_\varepsilon] ds.$$

Thus for $s \leq t \leq \tau^{\varepsilon,r} \wedge T^*$ and $\alpha > 0$ sufficiently small (to ensure integrability of powers of s and $t-s$; see also (1.6)) by (5.10)

$$\|J_{0,s}^\varepsilon h_\varepsilon\|_{\mathcal{C}^\beta} \leq C s^{-\frac{\beta+\alpha_0}{2}} \|h_\varepsilon\|_{\mathcal{C}^{-\alpha_0}} + C (s^{1-n\gamma} + s^{1-\frac{\beta+\alpha}{2}-(n-1)\gamma}) \sup_{s \leq t} s^\gamma \|J_{0,s}^\varepsilon h_\varepsilon\|_{\mathcal{C}^\beta}.$$

Multiplying the above inequality by s^γ we get

$$\sup_{s \leq t} s^\gamma \|J_{0,s}^\varepsilon h_\varepsilon\|_{\mathcal{C}^\beta} \leq C t^{\gamma - \frac{\beta + \alpha_0}{2}} \|h_\varepsilon\|_{\mathcal{C}^{-\alpha_0}} + C t^\theta \sup_{s \leq t} s^\gamma \|J_{0,s}^\varepsilon h_\varepsilon\|_{\mathcal{C}^\beta},$$

for some $\theta \equiv \theta(\alpha, \beta, \gamma, n) > 0$. Using that $\gamma - \frac{\beta + \alpha_0}{2} > 0$ (see (1.6)) by possibly increasing the value of the constant C in (3.7) we finally obtain the bound

$$\sup_{s \leq t} s^\gamma \|J_{0,s}^\varepsilon h_\varepsilon\|_{\mathcal{C}^\beta} \leq 2 \|h_\varepsilon\|_{\mathcal{C}^{-\alpha_0}}, \quad (5.11)$$

which completes the proof. \square

We denote by $C_b^1(\mathcal{C}^{-\alpha_0})$ the set of continuously differentiable functions on $\mathcal{C}^{-\alpha_0}$. We furthermore let $\chi \in C^\infty(\mathbb{R})$ such that $\chi(\zeta) \in [0, 1]$, for every $\zeta \in \mathbb{R}$, and

$$\chi(\zeta) = \begin{cases} 1, & \text{if } |\zeta| \leq \frac{r}{2}, \\ 0, & \text{if } |\zeta| \geq r, \end{cases}$$

for r as in (5.9). For simplicity we also let $\|\cdot\|_t := \|\cdot\|_{\alpha, \alpha'; t}$, $t \geq 0$. Inspired by [17], we prove the following version of the Bismut–Elworthy–Li formula.

Theorem 5.5 (Bismut–Elworthy–Li Formula). *Let $x \in \mathcal{C}^{-\alpha_0}$, $\Phi \in C_b^1(\mathcal{C}^{-\alpha_0})$ and let $t > 0$. Let w be a process taking values in the Cameron–Martin space \mathcal{CM} with $\partial_s w$ adapted. Furthermore, assume that there exists a deterministic constant $C \equiv C(t) > 0$ such that $\|\partial_s w\|_{L^2([0,t]; \mathbb{R}^{(2d-1)^2})}^2 \leq C$ \mathbb{P} -almost surely. Then we have that*

$$\begin{aligned} \mathbb{E}(D\Phi(X^\varepsilon(t; x))(DX^\varepsilon(t; x)(w))\chi(\|\underline{Z}^\varepsilon\|_t)) &= \mathbb{E}\left(\Phi(X^\varepsilon(t; x)) \int_0^t \partial_s w(s) \cdot d\hat{W}_\varepsilon(s) \chi(\|\underline{Z}^\varepsilon\|_t)\right) \\ &\quad - \mathbb{E}(\Phi(X^\varepsilon(t; x))\partial_+ \chi(\|\underline{Z}^\varepsilon\|_t)(w)), \end{aligned} \quad (5.12)$$

where

$$\partial_+ \chi(\|\underline{Z}^\varepsilon\|_t)(w) = \partial_\zeta \chi(\|\underline{Z}^\varepsilon\|_t) \partial_+ \|\underline{Z}^\varepsilon\|_t(Q_w(\cdot), 2\uparrow_{0,\cdot}^\varepsilon Q_w(\cdot), \dots, n \overset{\varepsilon}{\curvearrowright}_{0,\cdot}^{n-1} Q_w(\cdot)), \quad (5.13)$$

$\partial_+ \|\cdot\|_t : C^{n, -\alpha}(0; t) \rightarrow C^{n, -\alpha}(0; t)^*$ is the one-sided derivative of $\|\cdot\|_t$ given by

$$\partial_+ \|\underline{Z}^\varepsilon\|_t(\underline{Y}) = \lim_{\delta \rightarrow 0^+} \frac{\|\underline{Z}^\varepsilon + \delta \underline{Y}\|_t - \|\underline{Z}^\varepsilon\|_t}{\delta},$$

for every direction $\underline{Y} \in C^{n, -\alpha}(0; t)$, and

$$Q_w(\cdot) := \int_0^\cdot S(\cdot - s) G_\varepsilon \partial_s w(s) ds.$$

Remark 5.6. It is worth mentioning that the usual Bismut–Elworthy–Li formula (see [17]) gives an explicit representation of derivatives with respect to the initial condition rather than the noise. Below we also prove such a representation (see (5.19)). However the core of our argument is (5.12) which is slightly more general than (5.19).

Remark 5.7. The presence of $\partial_+ \|\cdot\|_t$ in the theorem above is based on the fact that norms are not in general Fréchet differentiable functions. However, their one-sided derivatives always exist (see [6, Appendix D]) and they behave nicely in terms of the usual rules of differentiation.

Proof. Let $\delta > 0$ and $u = \partial_t w$, which is an $L^2([0, \infty); \mathbb{R}^{(2d-1)^2})$ function. For every $n \geq 1$, we define the shift $T_{\delta u}$ by

$$T_{\delta u} \overset{\varepsilon}{\mathcal{V}}_{0,t}^n = \sum_{k=0}^n \binom{n}{k} (\delta \mathcal{Q}_w(t))^k \overset{\varepsilon}{\mathcal{V}}_{0,t}^{n-k}$$

and we let $T_{\delta u} \underline{Z}^\varepsilon = (T_{\delta u} \overset{\varepsilon}{\mathcal{V}}_{0,\cdot}^k)_{k=1}^n$.

Let $X^{\varepsilon,\delta}(\cdot; x) = T_{\delta u} \overset{\varepsilon}{\mathcal{I}}_{0,\cdot} + v^{\varepsilon,\delta}$, where the remainder $v^{\varepsilon,\delta}$ solves the equation

$$\begin{cases} \partial_t v^{\varepsilon,\delta} = \Delta v^{\varepsilon,\delta} - v^{\varepsilon,\delta} - \hat{\Pi}_\varepsilon \tilde{F}(v^{\varepsilon,\delta}, T_{\delta u} \underline{Z}^\varepsilon), \\ v^{\varepsilon,\delta}(0, \cdot) = \hat{\Pi}_\varepsilon x. \end{cases}$$

As in [17], our aim is to construct a probability measure \mathbb{P}^δ such that the law of $T_{\delta u} \overset{\varepsilon}{\mathcal{I}}_{0,\cdot}$ under \mathbb{P}^δ is the same as the law of $\overset{\varepsilon}{\mathcal{I}}_{0,\cdot}$ under \mathbb{P} . That way we obtain the identity

$$\partial_{\delta+} \mathbb{E}_{\mathbb{P}^\delta} (\Phi(X^{\varepsilon,\delta}(t; x)) \chi(\|T_{\delta u} \underline{Z}^\varepsilon\|_t)) |_{\delta=0} = 0, \tag{5.14}$$

since $\overset{\varepsilon}{\mathcal{V}}_{0,\cdot}^k$ is a continuous function of $\overset{\varepsilon}{\mathcal{I}}_{0,\cdot}^\varepsilon$ for every $k \geq 2$, the solution map to (5.3) is a continuous function of the $\overset{\varepsilon}{\mathcal{V}}_{0,\cdot}^k$, and χ is a continuous function of $\overset{\varepsilon}{\mathcal{I}}_{0,\cdot}^\varepsilon$. Above $\partial_{\delta+}$ stands as a shortcut of the directional derivative of a function as $\delta \rightarrow 0^+$. We will then show below that the result follows by an expansion of the derivative in the above expression.

We start with the construction of \mathbb{P}^δ . Let $\mathbf{B}^\delta(r) := -\int_0^r \delta u(s) \cdot d\hat{W}_\varepsilon(s)$ where \cdot is the scalar product on $\mathbb{R}^{(2d-1)^2}$, and define the exponential process

$$\mathbf{A}^\delta(r) := \exp\left\{ \mathbf{B}^\delta(r) - \frac{1}{2} \int_0^r |\delta u(s)|^2 ds \right\}.$$

Notice that by the assumptions on w Novikov's condition is satisfied, i.e.

$$\mathbb{E} \exp\left\{ \frac{1}{2} \int_0^t |\delta u(s)|^2 ds \right\} < \infty,$$

thus by [20, Chapter 8, Proposition 1.15] \mathbf{A}^δ is a strictly positive martingale and we have that $\mathbb{E} \mathbf{A}^\delta(t) = 1$. We define \mathbb{P}^δ by its Radon–Nikodym derivative with respect to \mathbb{P}

$$\frac{d\mathbb{P}^\delta}{d\mathbb{P}} = \mathbf{A}^\delta(t).$$

By Girsanov's Theorem (see [20, Chapter 4, Theorem 1.4]) $\hat{W}_\varepsilon^\delta(r) := \hat{W}_\varepsilon(r) - [\hat{W}_\varepsilon(\cdot), \mathbf{B}^\delta(\cdot)]_r$, $r \leq t$, under \mathbb{P}^δ has the same law as \hat{W}_ε under \mathbb{P} , where $[\cdot, \cdot]_r$ stands for the quadratic variation at time r . We furthermore have that $[\hat{W}_\varepsilon(\cdot), \mathbf{B}^\delta(\cdot)]_r = -\int_0^r \delta u(s) ds$ as well as $\overset{\varepsilon}{\mathcal{I}}_{0,t}^\varepsilon = \int_0^t S(t-s) G_\varepsilon d\hat{W}_\varepsilon(s)$ and $T_{\delta u} \overset{\varepsilon}{\mathcal{I}}_{0,t}^\varepsilon = \int_0^t S(t-s) G_\varepsilon d\hat{W}_\varepsilon^\delta(s)$. Since the law of $\hat{W}_\varepsilon^\delta$ under \mathbb{P}^δ is the same as the law of \hat{W}_ε under \mathbb{P} , this is also the case for $T_{\delta u} \overset{\varepsilon}{\mathcal{I}}_{0,\cdot}^\varepsilon$ and $\overset{\varepsilon}{\mathcal{I}}_{0,\cdot}^\varepsilon$ (recall that $\overset{\varepsilon}{\mathcal{I}}_{0,\cdot}^\varepsilon$ is a continuous function of \hat{W}_ε , when the latter is seen as an element in $C([0, t]; \mathbb{R}^{(2d-1)^2})$ endowed with the supremum norm because of (5.5)). Thus \mathbb{P}^δ is the required measure and (5.14) in the form

$$\partial_{\delta+} \mathbb{E} (\Phi(X^{\varepsilon,\delta}(t; x)) \chi(\|T_{\delta u} \underline{Z}^\varepsilon\|_t) \mathbf{A}^\delta(t)) |_{\delta=0} = 0 \tag{5.15}$$

follows. Using the chain rule,

$$\partial_\delta \Phi(X^{\varepsilon,\delta}(x; t)) = D\Phi(X^{\varepsilon,\delta}(x; t)) (\partial_\delta X^{\varepsilon,\delta}(x; t))$$

and

$$\partial_\delta \mathbf{A}^\delta(t) = -\mathbf{A}^\delta(t) \left(\int_0^t u(s) \cdot d\hat{W}_\varepsilon(s) + \delta \int_0^t |u(s)|^2 ds \right).$$

For the directional derivative of $\chi(\|T_{\delta u}\underline{Z}^\varepsilon\|_t)$ at $\delta^+ = 0$ it suffices to check the existence of the limit

$$\lim_{\delta \rightarrow 0^+} \frac{\|T_{\delta u}\underline{Z}^\varepsilon\|_t - \|\underline{Z}^\varepsilon\|_t}{\delta}.$$

We claim that the above limit is the same as

$$\partial_+ \|\underline{Z}^\varepsilon\|_t(Y^\varepsilon) := \lim_{\delta \rightarrow 0^+} \frac{\|\underline{Z}^\varepsilon + \delta Y^\varepsilon\|_t - \|\underline{Z}^\varepsilon\|_t}{\delta},$$

where $Y^\varepsilon = (Q_w(\cdot), 2\overset{\varepsilon}{\uparrow}_0, Q_w(\cdot), \dots, n \overset{\varepsilon}{\triangleleft}_0, Q_w(\cdot))$. Using the fact that $\|\cdot\|_t$ is a norm, we have that

$$\frac{\|T_{\delta u}\underline{Z}^\varepsilon\|_t - \|\underline{Z}^\varepsilon\|_t}{\delta} = \frac{\|\underline{Z}^\varepsilon + \delta Y^\varepsilon\|_t - \|\underline{Z}^\varepsilon\|_t}{\delta} + \text{Error}_\delta,$$

where $\text{Error}_\delta \rightarrow 0$ as $\delta \rightarrow 0^+$. Subtracting $\partial_+ \|\underline{Z}^\varepsilon\|_t(Y^\varepsilon)$ from both sides of the above equation and letting $\delta \rightarrow 0^+$ we get

$$\limsup_{\delta \rightarrow 0^+} \left(\frac{\|T_{\delta u}\underline{Z}^\varepsilon\|_t - \|\underline{Z}^\varepsilon\|_t}{\delta} - \partial_+ \|\underline{Z}^\varepsilon\|_t(Y^\varepsilon) \right) \leq 0. \quad (5.16)$$

In a similar way we can prove that the reverse inequality of (5.16) is valid with the lim sup replaced by a lim inf, which makes $\partial_+ \|\underline{Z}^\varepsilon\|_t(Y^\varepsilon)$ the appropriate limit.

We now argue on how to pass the derivative inside the expectation in (5.15). The argument is similar to [17]. For any function $f = f(\delta)$ we introduce the difference operator $\Delta_\delta f(\cdot) = f(\delta) - f(0)$.

We first show that the family of random variables

$$\frac{\Delta_\delta(\Phi(X^{\varepsilon,\cdot}(t;x))\chi(\|T_u\underline{Z}^\varepsilon\|_t)\mathbf{A}^\cdot(t))}{\delta}, \quad \delta \in (0, 1], \quad (5.17)$$

are uniformly integrable. We first write

$$\begin{aligned} \Delta_\delta(\Phi(X^{\varepsilon,\cdot}(t;x))\chi(\|T_u\underline{Z}^\varepsilon\|_t)\mathbf{A}^\cdot(t)) &= \Delta_\delta\Phi(X^{\varepsilon,\cdot}(t;x))\chi(\|T_{\delta u}\underline{Z}^\varepsilon\|_t)\mathbf{A}^\delta(t) \\ &\quad + \Phi(X^{\varepsilon,\delta}(t;x))\Delta_\delta\chi(\|T_u\underline{Z}^\varepsilon\|_t)\mathbf{A}^\delta(t) \\ &\quad + \Phi(X^{\varepsilon,\delta}(t;x))\chi(\|T_{\delta u}\underline{Z}^\varepsilon\|_t)\Delta_\delta\mathbf{A}^\cdot(t), \end{aligned}$$

and then treat each term on the right-hand side separately. For the first term, we first use that $\Phi \in C_b^1(C^{-\alpha_0})$ which prompts us to bound $\|X^{\varepsilon,\delta}(t;x) - X^\varepsilon(t;x)\|_{C^{-\alpha_0}}$. By the mean value theorem we get

$$\|X^{\varepsilon,\delta}(t;x) - X^\varepsilon(t;x)\|_{C^{-\alpha_0}} \leq \int_0^\delta \|\mathcal{D}X^{\varepsilon,\lambda}(t;x)(w)\|_{C^{-\alpha_0}} d\lambda,$$

where $\mathcal{D}X^{\varepsilon,\lambda}(t;x)(w)$ solves (5.6) with $\underline{Z}^\varepsilon$ replaced by $T_{\lambda u}\underline{Z}^\varepsilon$. By (5.6) we get a bound on the quantity $\|\mathcal{D}X^{\varepsilon,\lambda}(t;x)(w)\|_{C^{-\alpha_0}}$ as soon as we have a bound on $\|T_{\lambda u}\underline{Z}^\varepsilon\|$. The presence of the smooth indicator function yields a bound on $\|T_{\delta u}\underline{Z}^\varepsilon\|$ which then by definition of the shift as well as the assumed boundedness of w yields a uniform bound on $\|T_{\lambda u}\underline{Z}^\varepsilon\|$ for all $0 \leq \lambda \leq 1$. Hence we obtain a bound of the form

$$|\Delta_\delta\Phi(X^{\varepsilon,\cdot}(t;x))\chi(\|T_{\delta u}\underline{Z}^\varepsilon\|_t)\mathbf{A}^\delta(t)| \leq C\delta\|D\Phi\|_\infty\mathbf{A}^\delta(t),$$

where the constant C depends on w , χ and t . Arguing in the same way we get for the second term

$$|\Phi(X^{\varepsilon,\delta}(t;x))\Delta_\delta\chi(\|T_u\underline{Z}^\varepsilon\|_t)\mathbf{A}^\delta(t)| \leq C\delta\|\Phi\|_\infty\mathbf{A}^\delta(t).$$

Finally, for the third term we get using the mean value theorem for $\Delta_\delta A^\cdot(t)$ that

$$|\Phi(X^{\varepsilon,\delta}(t; x))\chi(\|T_{\delta u}\underline{Z}^\varepsilon\|_t) \Delta_\delta A^\cdot(t)| \leq C \|\Phi\|_\infty \int_0^\delta |\partial_\lambda A^\lambda(t)| d\lambda.$$

All the above imply that for every $p \geq 1$

$$\mathbb{E} \left| \frac{\Delta_\delta(\Phi(X^{\varepsilon,\cdot}(t; x))\chi(\|T_{\cdot u}\underline{Z}^\varepsilon\|_t)A^\cdot(t))}{\delta} \right|^p \lesssim \sup_{\delta \in (0,1]} \mathbb{E} A^\delta(t)^p + \sup_{\delta \in (0,1]} \mathbb{E} |\partial_\delta A^\delta(t)|^p.$$

The key observation is now that $A^\delta(t)^p = A^{\delta p}(t) \exp\{\frac{p^2-p}{2} \int_0^t |\delta u(s)|^2 ds\}$, where $A^{\delta p}(t)$ is also an exponential martingale of expectation 1, while $\exp\{\frac{p^2-p}{2} \int_0^t |\delta u(s)|^2 ds\}$ is uniformly bounded in δ because of the almost sure bound on w . This implies that $\sup_{\delta \in (0,1]} \mathbb{E} A^\delta(t)^p$ is bounded for any $p \geq 1$. Recalling the identity

$$\partial_\delta A^\delta(t) = -A^\delta(t) \left(\int_0^t u(s) \cdot d\hat{W}_\varepsilon(s) + \delta \int_0^t |u(s)|^2 ds \right)$$

and using again the almost sure bound on w as well as the Cauchy–Schwarz inequality we have that

$$\mathbb{E} |\partial_\delta A^\delta(t)|^p \lesssim (\mathbb{E} A^\delta(t)^{2p})^{\frac{1}{2p}} \left(\left(\mathbb{E} \left| \int_0^t u(s) \cdot d\hat{W}_\varepsilon(s) \right|^{2p} \right)^{\frac{1}{2p}} + 1 \right).$$

The first term on the right-hand side of the above inequality is uniformly bounded in δ as we discussed earlier while the second term can be bounded uniformly in δ using the Burkholder–Davis–Gundy inequality (see [20, Chapter 4, Theorem 4.1]) and the almost sure bound on w . Hence

$$\mathbb{E} \left| \frac{\Delta_\delta(\Phi(X^{\varepsilon,\cdot}(t; x))\chi(\|T_{\cdot u}\underline{Z}^\varepsilon\|_t)A^\cdot(t))}{\delta} \right|^p < \infty,$$

uniformly in $\delta \in (0, 1]$, for every $p \geq 1$, which implies uniform integrability of (5.17).

Using Vitali’s convergence theorem (see [3, Theorem 4.5.4]), we can now pass the derivative inside the expectation and differentiate by parts to obtain the identity

$$\begin{aligned} &\mathbb{E}(D\Phi(X^{\varepsilon,\delta}(t; x))(\partial_\delta X^{\varepsilon,\delta}(t; x))\chi(\|T_{\delta u}\underline{Z}^\varepsilon\|_t)A^\delta(t))|_{\delta=0} \\ &= -\mathbb{E}(\Phi(X^{\varepsilon,\delta}(t; x))\chi(\|T_{\delta u}\underline{Z}^\varepsilon\|_t)\partial_\delta A^\delta(t))|_{\delta=0} \\ &\quad - \mathbb{E}(\Phi(X^{\varepsilon,\delta}(t; x))\partial_{\delta+}\chi(\|T_{\delta u}\underline{Z}^\varepsilon\|_t)(w)A^\delta(t))|_{\delta+=0}. \end{aligned}$$

The result follows since $\partial_\delta X^{\varepsilon,\delta}(x; t)|_{\delta=0} = \mathcal{D}X^\varepsilon(t; x)(w)$ and $\partial_\delta A^\delta(t)|_{\delta=0} = -\int_0^t u(s) \cdot d\hat{W}_\varepsilon(s)$. □

Let $\{P_t^\varepsilon : t \geq 0\}$ defined via the identity

$$P_t^\varepsilon \Phi(x) := \mathbb{E}\Phi(X^\varepsilon(t; x))\mathbf{1}_{\{t < T_\varepsilon^*(x)\}}$$

for every $\Phi \in C_b(C^{-\alpha_0})$, where we write $T_\varepsilon^*(x)$ for the explosion time of v^ε (see Proposition 5.1) dropping the dependence on $\mathfrak{I}_{0,\cdot}^\varepsilon$. We use (5.12) to prove the following proposition.

Proposition 5.8. *There exist a universal constant C and $\theta_1 > 0$ such that*

$$|P_t^\varepsilon \Phi(x) - P_t^\varepsilon \Phi(y)| \leq C \frac{1}{t^{\theta_1}} \|\Phi\|_\infty \|x - y\|_{C^{-\alpha}} + 2\|\Phi\|_\infty \mathbb{P}(t \geq \tau^{\varepsilon, \frac{t}{2}}) \tag{5.18}$$

for every $x \in C^{-\alpha_0}$, $y \in \bar{B}_1(x)$, $\Phi \in C_b^1(C^{-\alpha_0})$ and $t \leq T^* \equiv T^*(R)$ (defined in Proposition 5.4), where $R = 2\|x\|_{C^{-\alpha_0}} + 1$.

Proof. Let $\Phi \in C_b^1(C^{-\alpha})$ and $t \leq T^*$. Then

$$|P_t^\varepsilon \Phi(x) - P_t^\varepsilon \Phi(y)| = |\mathbb{E}[\Phi(X^\varepsilon(t; x))\mathbf{1}_{\{t < T_\varepsilon^*(x)\}} - \Phi(X^\varepsilon(t; y))\mathbf{1}_{\{t < T_\varepsilon^*(y)\}}]|$$

and the latter term is bounded by $I_1 + I_2$, where

$$\begin{aligned} I_1 &:= |\mathbb{E}[(\Phi(X^\varepsilon(t; x)) - \Phi(X^\varepsilon(t; y)))\chi(\|\underline{Z}^\varepsilon\|_t)]|, \\ I_2 &:= |\mathbb{E}[(\Phi(X^\varepsilon(t; x))\mathbf{1}_{\{t < T_\varepsilon^*(x)\}} - \Phi(X^\varepsilon(t; y))\mathbf{1}_{\{t < T_\varepsilon^*(y)\}})(1 - \chi(\|\underline{Z}^\varepsilon\|_t))]|. \end{aligned}$$

For the second term we have that $I_2 \leq 2\|\Phi\|_\infty \mathbb{P}(t \geq \tau^{\varepsilon, \frac{r}{2}})$ while by the mean value theorem we get that

$$\begin{aligned} I_1 &= \left| \mathbb{E} \left(\int_0^1 D\Phi(X^\varepsilon(t; x + \lambda(y-x)))(y-x) d\lambda \chi(\|\underline{Z}^\varepsilon\|_t) \right) \right| \\ &= \left| \int_0^1 \mathbb{E}(D\Phi(X^\varepsilon(t; x + \lambda(y-x)))(y-x)\chi(\|\underline{Z}^\varepsilon\|_t)) d\lambda \right|. \end{aligned}$$

For any $h_\varepsilon \in \Pi_\varepsilon[L^2(\mathbb{T}^2)]$ let w be such that $\partial_s w(s) = (\langle J_{0,s}^\varepsilon h_\varepsilon, e_m \rangle)_{|m| < \frac{1}{\varepsilon}}$ for $s \leq \tau^{\varepsilon, r}$ and 0 otherwise. Then $\partial_s w$ is an adapted process and by Proposition 5.4 there exists $C \equiv C(t) > 0$ such that $\|\partial_s w\|_{L^2([0,t]; \mathbb{R}^{(2d-1)^2})}^2 \leq C$, \mathbb{P} -almost surely, for every initial condition $z_\lambda = x + \lambda(y-x)$ (recall that $J_{0,\cdot}^\varepsilon$ depends on the initial condition and that $z_\lambda \in \bar{B}_1(x)$, for every $\lambda \in [0, 1]$, thus the estimates in Proposition 5.4 hold uniformly in λ). Furthermore, $\mathcal{D}X^\varepsilon(t; z_\lambda)(w) = tDX^\varepsilon(t; z_\lambda)(h_\varepsilon)$, for every $t \leq \tau^{\varepsilon, r}$, and as in [17] we can use (5.12) for this particular choice of w to obtain the following identity,

$$\begin{aligned} \mathbb{E}(D[\Phi(X^\varepsilon(t; z_\lambda))](h_\varepsilon)\chi(\|\underline{Z}^\varepsilon\|_t)) &= \frac{1}{t} \mathbb{E} \left(\Phi(X^\varepsilon(t; z_\lambda)) \int_0^t \langle J_{0,s}^\varepsilon h_\varepsilon, dW_\varepsilon(s) \rangle \chi(\|\underline{Z}^\varepsilon\|_t) \right) \\ &\quad - \frac{1}{t} \mathbb{E}(\Phi(X^\varepsilon(t; z_\lambda))\partial_+ \chi(\|\underline{Z}^\varepsilon\|_t)(w)), \end{aligned} \quad (5.19)$$

where we slightly abuse the notation since, as we already mentioned, the operator $J_{0,\cdot}^\varepsilon$ depends on the initial condition z_λ . In particular this is true for $h_\varepsilon = \hat{\Pi}_\varepsilon(y-x)$, hence

$$\begin{aligned} I_1 &\leq \frac{1}{t} \|\Phi\|_\infty \int_0^1 \mathbb{E} \left| \int_0^t \langle J_{0,s}^\varepsilon \hat{\Pi}_\varepsilon(y-x), dW_\varepsilon(s) \rangle \chi(\|\underline{Z}^\varepsilon\|_t) \right| d\lambda \\ &\quad + \frac{1}{t} \|\Phi\|_\infty \int_0^1 \mathbb{E} |\partial_+ \chi(\|\underline{Z}^\varepsilon\|_t)(w)| d\lambda. \end{aligned}$$

Estimating the first term above we get

$$\begin{aligned} \mathbb{E} \left| \int_0^t \langle J_{0,s}^\varepsilon \hat{\Pi}_\varepsilon(y-x), dW_\varepsilon(s) \rangle \chi(\|\underline{Z}^\varepsilon\|_t) \right| &\leq \mathbb{E} \left| \int_0^{t \wedge \tau^{\varepsilon, r}} \langle J_{0,s}^\varepsilon \hat{\Pi}_\varepsilon(y-x), dW_\varepsilon(s) \rangle \right| \\ &\leq \left(\mathbb{E} \int_0^{t \wedge \tau^{\varepsilon, r}} \|J_{0,s}^\varepsilon \hat{\Pi}_\varepsilon(y-x)\|_{L^2}^2 ds \right)^{\frac{1}{2}} \\ &\leq Ct^{\frac{1}{2}-\gamma} \|x-y\|_{C^{-\alpha_0}}, \end{aligned}$$

where we use a Cauchy–Schwarz inequality and Itô's isometry in the second step and Proposition 5.4 in the third step. Here we use crucially, that the deterministic bound on $J_{0,s}^\varepsilon$ provided in Proposition 5.4 holds uniformly in $\varepsilon > 0$ (and in λ). Using the explicit form (5.13) of $\partial_+ \chi(\|\underline{Z}^\varepsilon\|_t)$ we also have the uniform in λ bound

$$\mathbb{E} |\partial_+ \chi(\|\underline{Z}^\varepsilon\|_t)(w)| \leq Ct^{1-\gamma} \|x-y\|_{C^{-\alpha_0}},$$

since

$$\partial_+ \|\underline{Z}^\varepsilon\|_t(Q_w(\cdot), 2\uparrow_{0,\cdot}^\varepsilon Q_w(\cdot), \dots, n \overset{\varepsilon}{\downarrow}_{0,\cdot}^{n-1} Q_w(\cdot)) \leq C \|\underline{Z}^\varepsilon\|_t t^{1-\gamma} \|x - y\|_{C^{-\alpha_0}}$$

and the fact that $\|\underline{Z}^\varepsilon\|_t$ multiplied by $\partial_\zeta \chi(\|\underline{Z}^\varepsilon\|_t)$ is bounded by 1. Thus

$$I_1 \leq C \frac{1}{t^\gamma} \|\Phi\|_\infty \|x - y\|_{C^{-\alpha_0}}$$

and using both the bounds on I_1 and I_2 we get that for every $t \leq T^*$

$$|P_t^\varepsilon \Phi(x) - P_t^\varepsilon \Phi(y)| \leq C \frac{1}{t^\gamma} \|\Phi\|_\infty \|x - y\|_{C^{-\alpha_0}} + 2\|\Phi\|_\infty \mathbb{P}(t \geq \tau^{\varepsilon, \frac{r}{2}}),$$

which completes the proof. □

Given that the vector $(\overset{\varepsilon}{\downarrow}_{0,\cdot}^k)_{k=1}^n$ converges in law to $(\overset{\cdot}{\downarrow}_{0,\cdot}^k)_{k=1}^n$ on $C^{n,-\alpha}(0; T)$, for every $\alpha > 0$ and with respect to every norm $\|\cdot\|_{\alpha; \alpha'; T}$, for every $T > 0$, we have that $\tau^{\varepsilon, \frac{r}{2}}$ converges in law to $\tau^{\frac{r}{2}}$ when the mapping

$$\underline{Z} \mapsto \inf \left\{ t > 0 : \|\underline{Z}_t^{(1)}\|_{C^{-\alpha}} \vee t^{\alpha'} \|\underline{Z}_t^{(2)}\|_{C^{-\alpha}} \vee \dots \vee t^{(n-1)\alpha'} \|\underline{Z}_t^{(n)}\|_{C^{-\alpha}} > \frac{r}{2} \right\} \tag{5.20}$$

is \mathbb{P} -almost surely continuous on the path $(\overset{\cdot}{\downarrow}_{0,\cdot}^k)_{k=1}^n$. But if

$$L := \{r \in (0, 1] : \mathbb{P}((5.20) \text{ is discontinuous on } (\overset{\cdot}{\downarrow}_{0,\cdot}^k)_{k=1}^n) > 0\}$$

and $M : [0, \infty) \rightarrow [0, \infty)$ is the mapping

$$t \mapsto \|\uparrow_{0,t}\|_{C^{-\alpha}} \vee t^{\alpha'} \|\downarrow_{0,t}\|_{C^{-\alpha}} \vee \dots \vee t^{(n-1)\alpha'} \|\downarrow_{0,t}\|_{C^{-\alpha}},$$

then

$$L \subset \{r \in (0, 1] : \mathbb{P}(M \text{ has a local maximum at height } r) > 0\}$$

and the latter set is at most countable (see [14, proof of Theorem 6.1]), thus we can choose $r \in [\frac{1}{4}, 1]$ in (5.9) such that (5.20) is \mathbb{P} -almost surely continuous on $(\overset{\cdot}{\downarrow}_{0,\cdot}^k)_{k=1}^n$. This implies convergence in law of $\tau^{\varepsilon, \frac{r}{2}}$ to $\tau^{\frac{r}{2}}$, thus

$$\limsup_{\varepsilon \rightarrow 0^+} \mathbb{P}(t \geq \tau^{\varepsilon, \frac{r}{2}}) \leq \mathbb{P}(t \geq \tau^{\frac{r}{2}}).$$

Notice that global existence of v_t (see Theorem 3.9) implies global existence of $X(t; x)$ and in particular existence for every $t \leq T^*(R)$. Using Propositions 2.3 and 3.4, $\liminf_{\varepsilon \rightarrow 0^+} T_\varepsilon^* \geq T^*(R)$ and $\sup_{t \leq T_\varepsilon^* \wedge T^*(R)} \|X^\varepsilon(t; x^\varepsilon) - X(t; x)\|_{C^{-\alpha_0}} \rightarrow 0$ \mathbb{P} -almost surely, for every $x \in C^{-\alpha}$. By the dominated convergence theorem $P_t^\varepsilon \Phi(x)$ converges to $P_t \Phi(x)$, for every $\Phi \in C_b^1(C^{-\alpha_0})$, and we retrieve (5.18) for the limiting semigroup P_t , for every $t \leq T^*(R)$, in the form

$$|P_t \Phi(x) - P_t \Phi(y)| \leq C \frac{1}{t^{\theta_1}} \|\Phi\|_\infty \|x - y\|_{C^{-\alpha_0}} + 2\|\Phi\|_\infty \mathbb{P}(t \geq \tau^{\frac{r}{2}}). \tag{5.21}$$

Remark 5.9. The above argument can be modified to retrieve (5.21) without the knowledge of global existence for the limiting process. In this case, one can define the semigroup P_t by introducing a cemetery state for the process $X(t; x)$.

We finally prove the following theorem. Below we denote by $\|\mu_1 - \mu_2\|_{\text{TV}}$ the total variation distance of two probability measures $\mu_1, \mu_2 \in \mathcal{M}_1(C^{-\alpha_0})$ given by

$$\|\mu_1 - \mu_2\|_{\text{TV}} := \frac{1}{2} \sup_{\|\Phi\|_{\infty} \leq 1} |\mathbb{E}_{\mu_1} \Phi - \mathbb{E}_{\mu_2} \Phi|.$$

Theorem 5.10. *There exists $\theta \in (0, 1)$ and $\sigma > 0$ such that for every $x \in C^{-\alpha_0}$ and $y \in \bar{B}_1(x)$*

$$\|P_t^* \delta_x - P_t^* \delta_y\|_{\text{TV}} \leq C(1 + \|x\|_{C^{-\alpha_0}})^\sigma \|x - y\|_{C^{-\alpha_0}}^\theta,$$

for every $t \geq 1$. In particular, for every $t \geq 1$, P_t is locally uniformly θ -Hölder continuous with respect to the total variation norm in $C^{-\alpha_0}$.

Proof. Let $R = 2\|x\|_{C^{-\alpha_0}} + 1$. By a density argument (see [7, Lemma 7.1.5]) (5.21) is equivalent to

$$\|P_t^* \delta_x - P_t^* \delta_y\|_{\text{TV}} \leq \frac{C}{2} \frac{1}{t^{\theta_1}} \|x - y\|_{C^{-\alpha_0}} + \mathbb{P}(t \geq \tau^{\frac{r}{2}}),$$

for every $t \leq T^*$ and $y \in \bar{B}_1(x)$. Notice that

$$\mathbb{P}(t \geq \tau^{\frac{r}{2}}) \leq \mathbb{P}\left(\|\underline{Z}\|_{\alpha; \alpha'; t} > \frac{r}{2}\right)$$

and by Theorem 2.1

$$\mathbb{P}(\|\underline{Z}\|_{\alpha; \alpha'; t} > r) \leq C \frac{1}{r} t^{\theta_2},$$

for some $\theta_2 \in (0, 1)$. Since we can assume that $T^* \leq 1$, we have that

$$\|P_1^* \delta_x - P_1^* \delta_y\|_{\text{TV}} \leq \|P_{T^*}^* \delta_x - P_{T^*}^* \delta_y\|_{\text{TV}},$$

where

$$\|P_{T^*}^* \delta_x - P_{T^*}^* \delta_y\|_{\text{TV}} \leq \inf_{t \leq T^*} \left\{ C_1 \frac{1}{t^{\theta_1}} \|x - y\|_{C^{-\alpha_0}} + C_2 \frac{1}{r} t^{\theta_2} \right\}.$$

Let $f(t) := C_1 \frac{1}{t^{\theta_1}} \|x - y\|_{C^{-\alpha_0}} + C_2 \frac{1}{r} t^{\theta_2}$, $t > 0$, and notice that for

$$t_0 = \left(\frac{\theta_1 C_1 r \|x - y\|_{C^{-\alpha_0}}}{\theta_2 C_2} \right)^{\frac{1}{\theta_1 + \theta_2}},$$

$f(t_0) = \inf_{t > 0} f(t)$. If $t_0 \leq T^*$, then there exists $C \equiv C(\theta_1, \theta_2, r)$ such that

$$\|P_{T^*}^* \delta_x - P_{T^*}^* \delta_y\|_{\text{TV}} \leq f(t_0) = C \|x - y\|_{C^{-\alpha_0}}^{\frac{\theta_2}{\theta_1 + \theta_2}}.$$

Otherwise $t_0 \geq T^*$ and using

$$\begin{aligned} \|P_{T^*}^* \delta_x - P_{T^*}^* \delta_y\|_{\text{TV}} &\leq C_1 \frac{1}{(T^*)^{\theta_1}} \|x - y\|_{C^{-\alpha_0}} + C_2 \frac{1}{r} (T^*)^{\theta_2} \\ &\leq C_1 \frac{1}{(T^*)^{\theta_1}} \|x - y\|_{C^{-\alpha_0}} + C_2 \frac{1}{r} t_0^{\theta_2} \\ &= C_1 \frac{1}{(T^*)^{\theta_1}} \|x - y\|_{C^{-\alpha_0}} + \tilde{C}_2 \frac{1}{r} \|x - y\|_{C^{-\alpha_0}}^{\frac{\theta_2}{\theta_1 + \theta_2}} \end{aligned}$$

and the explicit estimate of T^* (see (3.7)) we get

$$\begin{aligned} \|P_{T^*}^* \delta_x - P_{T^*}^* \delta_y\|_{TV} &\leq \tilde{C}_1 (1 + R)^{3\frac{\theta_1}{\theta}} \|x - y\|_{C^{-\alpha_0}} + \tilde{C}_2 \frac{1}{r} \|x - y\|_{C^{-\alpha_0}}^{\frac{\theta_2}{\theta_1 + \theta_2}} \\ &\leq C (1 + R)^{3\frac{\theta_1}{\theta} + \frac{\theta_1}{\theta_1 + \theta_2}} \|x - y\|_{C^{-\alpha_0}}^{\frac{\theta_2}{\theta_1 + \theta_2}} \end{aligned}$$

for a constant $C \equiv C(\theta_1, \theta_2, r)$ and some $\theta > 0$ as in (3.7). Combining all the above we finally get

$$\|P_1^* \delta_x - P_1^* \delta_y\|_{TV} \leq C (1 + R)^{3\frac{\theta_1}{\theta} + \frac{\theta_1}{\theta_1 + \theta_2}} \|x - y\|_{C^{-\alpha_0}}^{\frac{\theta_2}{\theta_1 + \theta_2}},$$

which completes the proof. □

6. Exponential mixing of Φ_2^4

From now on we restrict ourselves in the case $n = 3$ (see Remark 6.2). In this section following [4] we first prove a support theorem for the solution to the Φ_2^4 equation. After that we combine this result with Corollary 3.10 and Theorem 5.10 and prove exponential convergence to a unique invariant measure with respect to the total variation norm.

6.1. A support theorem

We consider $\underline{Y} = (\bigtriangledown_{-\infty, \cdot})_{k=1}^3$ as an element of $C([0, T]; C^{-\alpha})^3$ endowed with the norm $\|\cdot\|_{\alpha; 0; T}$, for some $\alpha \in (0, 1)$, given by

$$\|\underline{Y}\|_{\alpha; 0; T} := \max_{k=1,2,3} \left\{ \sup_{t \leq T} \|\bigtriangledown_{0,t}\|_{C^{-\alpha}} \right\}.$$

Here we are allowed to use a non-weighted norm since there is no blow up of $\bigtriangledown_{-\infty, \cdot}$ at zero. We furthermore let

$$\mathcal{H}(T) := \left\{ h|_{[0,T]} : h(t) = \int_{-\infty}^t S(t-r) f(r) dr, t \geq 0, \text{ and } f \in L^2(\mathbb{R} \times \mathbb{T}^2) \right\}.$$

It is worth mentioning that $\mathcal{H}(T)$ consists of those L^2 -integrable space-time functions with zero initial datum and with one derivative in time and two derivatives in space in L^2 .

Lemma 6.1. *Let $\{C_m\}_{m \geq 1}$ be a sequence of positive numbers such that $C_m \leq C(m + 1)$. Then there exists a sequence of smooth functions $\{f_m\}_{m \geq 1}$ such that*

- (i) $f_m \in C^{-\alpha}$, for every $\alpha \in (0, 1)$.
- (ii) $\mathbb{E}|\langle f_m, e_l \rangle|^2 = C_m$ if $l = 2^m$ or $l = -2^m$ and 0 otherwise.
- (iii) For every $n = 1, 2, 3$, $H_n(f_m, C_m) \rightarrow 0$ in $C^{-\alpha}$, for every $\alpha \in (0, 1)$.

Proof. Let

$$f_m(z) := \frac{e^{2\pi i 2^m z_0 \cdot z} + e^{-2\pi i 2^m z_0 \cdot z}}{2^{1/2}} C_m^{1/2},$$

where $z_0 = (1, 1) \in \mathbb{Z}^2$, $z \in \mathbb{T}^2$. Then for $\kappa \geq -1$

$$\delta_\kappa f_m(z) = \frac{C_m^{1/2}}{2^{1/2}} \mathbf{1}_{\{m=\kappa\}} (e^{2\pi i 2^m z_0 \cdot z} + e^{-2\pi i 2^m z_0 \cdot z}),$$

$$\begin{aligned}\delta_\kappa f_m(z)^2 - C_m &= \frac{C_m}{2} \mathbf{1}_{\{m+1=\kappa\}} (e^{2\pi i 2^{m+1} z_0 \cdot z} + e^{-2\pi i 2^{m+1} z_0 \cdot z}), \\ \delta_\kappa f_m(z)^3 &= \frac{C_m^{3/2}}{2^{3/2}} [\chi_\kappa(2^m 3z_0) (e^{2\pi i 2^m 3z_0 \cdot z} + e^{-2\pi i 2^m 3z_0 \cdot z}) \\ &\quad + \mathbf{1}_{\{m=\kappa\}} 3 (e^{2\pi i 2^m z_0 \cdot z} + e^{-2\pi i 2^m z_0 \cdot z})].\end{aligned}$$

Notice here we have used the convenient fact that the particular choice of z_0 has the property that $\chi_\kappa(2^m z_0) = \mathbf{1}_{\{m=\kappa\}}$. Thus we have

$$\begin{aligned}\|f_m\|_{C^{-\alpha}} &\lesssim C_m^{1/2} 2^{-\alpha m}, \\ \|f_m^2 - C_m\|_{C^{-\alpha}} &\lesssim C_m 2^{-\alpha m}, \\ \|f_m^3 - 3C_m f_m\|_{C^{-\alpha}} &\lesssim C_m^{3/2} 2^{-\alpha m}.\end{aligned}$$

Given that $C_m \lesssim m + 1$ all the above quantities tend to 0 as $m \rightarrow \infty$, which completes the proof. \square

Remark 6.2. The sequence $\{f_m\}_{m \geq 1}$ introduced in the lemma above satisfies property (iii) for every odd n . For such n every term appearing in $H_n(f_m, C_m)$ is a multiple of $C_m^{k_1} e^{2^m k_2 z_0}$ for a $k_2 \neq 0$ and the fast (exponential) decay of $\|e^{2^m k_2 z_0}\|_{C^{-\alpha}}$ compensates the slow (polynomial) growth of $C_m^{k_1}$. However, for even n this property fails, because for such n the $H_n(f_m, C_m)$ contains a multiple of C_m^n which does not need to vanish. We suspect, that a first step in order to generalize Theorem 6.3 to the case of general n would be the construction of a sequence $\{f_m\}_{m \geq 1}$ with Fourier support on an annulus and such that

$$\int_{\mathbb{T}^2} f_m(z)^k dz = \mathcal{H}_k(0, C^m),$$

for every $k \geq 1$.

We now prove the following support theorem.

Theorem 6.3. Let $\mathbb{P}_{\underline{Y}}$ be the law of \underline{Y} in $C([0, T]; C^{-\alpha})^3$ endowed with the norm $\|\cdot\|_{\alpha; 0; T}$. Then

$$\text{supp } \mathbb{P}_{\underline{Y}} = \overline{\{(\mathcal{H}_k(h, \mathfrak{R}))_{k=1}^3 : h \in \mathcal{H}(T), \mathfrak{R} \geq 0\}}^{\|\cdot\|_{\alpha; 0; T}}.$$

Proof. For $h \in \mathcal{H}(T)$ and $\underline{Y} \in C^{3, -\alpha}(0; T)$ let T_h be the shift

$$T_h Y^{(k)} = \sum_{j=0}^k \binom{k}{j} h^j Y^{(k-j)}, \quad k = 1, 2, 3,$$

where we use again the convention that $Y^{(0)} \equiv 1$, and write $T_h \underline{Y} = (T_h Y^{(k)})_{k=1}^3$. Here we slightly abuse the notation since the action of T_h on $Y^{(k)}$ needs information on the lower order terms.

As in [4], it suffices to prove that $(0, -\mathfrak{R}, 0) \in \text{supp } \mathbb{P}_{\underline{Y}}$, for every $\mathfrak{R} \geq 0$. Then, given that shifts of the initial probability measure in the direction of the Cameron–Martin space generate equivalent probability measures, for every $h \in \mathcal{H}(T)$, $T_h(0, -\mathfrak{R}, 0) \in \text{supp } \mathbb{P}_{\underline{Y}}$, which completes the proof since by the definition of T_h the latter is equal to $(\mathcal{H}_k(h, \mathfrak{R}))_{k=1}^3$ (see also [4, Corollary 3.10]).

For $\lambda > 0$ and $\rho_{\lambda 2^m}(z) = \sum_{|\bar{m}| < \lambda 2^m} e_{\bar{m}}(z)$ we let

$$\mathfrak{I}_{-\infty, t}^m(z) := \mathfrak{I}_{-\infty, t}(\rho_{\lambda 2^m}(z - \cdot)), \quad \mathfrak{R}^m := \mathbb{E} \mathfrak{I}_{-\infty, t}^m(0)^2,$$

where $\mathfrak{I}_{-\infty, t}^m$ coincides with $\mathfrak{I}_{-\infty, t}^\varepsilon$ in Section 2.2 for $m = \frac{1}{\varepsilon}$. Notice that for $\mathfrak{R} \geq 0$ there exists $m_0 \equiv m_0(\mathfrak{R}) > 1$ such that $\mathfrak{R}^m - \mathfrak{R} > 0$, for every $m \geq m_0$ (recall that $\mathfrak{R}^m \sim \log m$). Thus if we set $C_m = 0$ for $m \leq m_0$ and $C_m = \mathfrak{R}^m - \mathfrak{R}$

otherwise, then $C_m \geq 0$ and $C_m \lesssim m + 1$. We consider f_m as in Lemma 6.1 for this particular choice of C_m and for $\lambda_m = 1 + 4\pi^2 2^{2m} |z_0|^2$ we let

$$h_m(t) := (1 - e^{-\lambda_m(t+1)}) f_m,$$

for $t \in [0, T]$. Then $h_m \in \mathcal{H}(T)$ since $h_m(t) = \frac{1}{\lambda_m} \int_{-1}^t S(t-r) f_m dr$ and we furthermore have the uniform in t estimates

$$\begin{aligned} \|h_m(t)\|_{C^{-\alpha}} &\leq \|f_m\|_{C^{-\alpha}}, \\ \|h_m(t)^2 - C_m\|_{C^{-\alpha}} &\leq \|f_m^2 - C_m\|_{C^{-\alpha}}^2 + 2e^{-\lambda_m} C_m, \\ \|h_m(t)^3\|_{C^{-\alpha}} &\leq \|f_m^3\|_{C^{-\alpha}}. \end{aligned}$$

Finally, we define

$$w_m := -\mathfrak{I}_{-\infty, \cdot}^m - h_m.$$

We prove that the following convergences hold in every stochastic L^p space of random variables taking values in $C([0, T]; C^{-\alpha})$,

$$T_{w_m} \mathfrak{I}_{-\infty, \cdot} \rightarrow 0, \quad T_{w_m} \mathfrak{V}_{-\infty, \cdot} \rightarrow -\mathfrak{R}, \quad T_{w_m} \mathfrak{V}_{-\infty, \cdot} \rightarrow 0.$$

By the same argument as in [4, Lemma 3.13] this implies the result. For the reader's convenience, we sketch the argument here. Since $w_m \in \mathcal{H}(T)$, by Lemma [4, Corollary 3.10] there exists a subset Ω' of Ω of probability one such that for every $\omega \in \Omega'$

$$(T_{w_m(\omega)} \mathfrak{I}_{-\infty, \cdot}(\omega), T_{w_m(\omega)} \mathfrak{V}_{-\infty, \cdot}(\omega), T_{w_m(\omega)} \mathfrak{V}_{-\infty, \cdot}(\omega)) \in \text{supp } \mathbb{P}_Y,$$

for every $m \geq 1$. Given that $\text{supp } \mathbb{P}_Y$ is closed under the norm $\|\cdot\|_{\alpha; 0; T}$, we can conclude that $(0, -\mathfrak{R}, 0) \in \text{supp } \mathbb{P}_Y$ as soon as the above convergence holds for a single element $\omega \in \Omega'$. The stochastic L^p convergence implies almost sure convergence along a subsequence which is sufficient.

The convergence of $T_{w_m} \mathfrak{I}_{-\infty, \cdot}$ to 0 is an immediate consequence of Proposition 2.3 and Lemma 6.1.

If we compute the corresponding shift for $\mathfrak{V}_{-\infty, t}$ we get

$$\begin{aligned} T_{w_m} \mathfrak{V}_{-\infty, t} &= \mathfrak{V}_{-\infty, t} + ((\mathfrak{I}_{-\infty, t}^m)^2 - \mathfrak{R}^m) - 2(\mathfrak{I}_{-\infty, t} \mathfrak{I}_{-\infty, t}^m - \mathfrak{R}^m) \\ &\quad + 2\mathfrak{I}_{-\infty, t}^m h_m(t) + \mathcal{H}_2(h_m(t), \mathfrak{R}^m), \end{aligned}$$

where we also add and subtract $2\mathfrak{R}^m$ where necessary. If we choose λ sufficiently small we can ensure that

$$\mathfrak{I}_{-\infty, t}^m \circ h_m(t) \equiv 0,$$

where $\mathfrak{I}_{-\infty, t}^m \circ h_m(t)$ is the resonant term define in (A.9). Using the Bony estimates (see Proposition A.6), Lemma 6.1 and the fact that $\mathfrak{I}_{-\infty, \cdot}^m$ is bounded in every stochastic L^p space taking values in $C([0, T]; C^{-\alpha})$ we get that $\mathfrak{I}_{-\infty, t}^m h_m(t) \rightarrow 0$. For the term

$$\mathfrak{V}_{-\infty, t} + ((\mathfrak{I}_{-\infty, t}^m)^2 - \mathfrak{R}^m) - 2(\mathfrak{I}_{-\infty, t} \mathfrak{I}_{-\infty, t}^m - \mathfrak{R}^m)$$

by Proposition 2.3 it suffices to compute the limit of $\mathfrak{I}_{-\infty, t} \mathfrak{I}_{-\infty, t}^m - \mathfrak{R}^m$. We only give a sketch of the proof since the idea is similar to the one in the proof of Proposition 2.3. Notice that for $m' > m$, $\mathbb{E} \mathfrak{I}_{-\infty, t}^{m'} \mathfrak{I}_{-\infty, t}^m = \mathfrak{R}^m$, thus using [18, Proposition 1.1.2] we have that

$$\mathfrak{I}_{-\infty, t}^{m'} \mathfrak{I}_{-\infty, t}^m - \mathfrak{R}^m = \mathfrak{I}_{-\infty, t}^{m'} \otimes \mathfrak{I}_{-\infty, t}^m,$$

where \otimes denotes the renormalized product given by

$$\begin{aligned} & \mathbb{V}_{-\infty,t}^{m'} \otimes \mathbb{V}_{-\infty,t}^m(z) \\ & := \int_{\{(-\infty,t] \times \mathbb{T}^2\}^{j+i}} \prod_{\substack{1 \leq j' \leq j \\ 1 \leq i' \leq i}} H_{m'}(t - r_{i'}, z - z_{i'}) H_m(t - r_{j'}, z - z_{j'}) \xi \left(\bigotimes_{k=1}^{i+j} dz_k, \bigotimes_{k=1}^{i+j} dr_k \right), \end{aligned}$$

for every $z \in \mathbb{T}^2$ and $i, j \geq 1$. In the same spirit as in the proof of Proposition 2.3 (see Appendix E) we can prove that

$$\lim_{m \rightarrow \infty} \lim_{m' \rightarrow \infty} \mathbb{E} \sup_{t \leq T} \|\mathfrak{I}_{-\infty,t}^{m'} \otimes \mathfrak{I}_{-\infty,t}^m - \mathfrak{V}_{-\infty,t}\|_{\mathcal{C}^{-\alpha}}^p = 0,$$

for every $p \geq 2$. Combining the above with the fact that $\sup_{t \leq T} \|h_m(t)^2 - (\mathfrak{R}^m - \mathfrak{R})\|_{\mathcal{C}^{-\alpha}}$ converges to 0, we obtain that $T_{w_m} \mathfrak{V}_{-\infty, \cdot} \rightarrow -\mathfrak{R}$.

For the term $T_{w_m} \mathfrak{V}_{-\infty,t}$, by adding and subtracting multiples of $\mathfrak{R}^m \mathfrak{I}_{-\infty,t}^m$ and \mathfrak{R}^m where necessary we have that

$$\begin{aligned} T_{w_m} \mathfrak{V}_{-\infty,t} &= \mathfrak{V}_{-\infty,t} - ((\mathfrak{I}_{-\infty,t}^m)^3 - 3\mathfrak{R}^m \mathfrak{I}_{-\infty,t}^m) - 3(\mathfrak{I}_{-\infty,t}^m \mathfrak{V}_{-\infty,t} - 2\mathfrak{R}^m \mathfrak{I}_{-\infty,t}^m) \\ &\quad + 3(\mathfrak{I}_{-\infty,t}^m (\mathfrak{I}_{-\infty,t}^m)^2 - 3\mathfrak{R}^m \mathfrak{I}_{-\infty,t}^m) \\ &\quad + 3h_m(t)(\mathfrak{V}_{-\infty,t} + ((\mathfrak{I}_{-\infty,t}^m)^2 - \mathfrak{R}^m) - 2(\mathfrak{I}_{-\infty,t}^m \mathfrak{I}_{-\infty,t}^m - \mathfrak{R}^m)) \\ &\quad + 3h_m(t)^2(\mathfrak{I}_{-\infty,t} - \mathfrak{I}_{-\infty,t}^m) + \mathcal{H}_3(h_m(t), \mathfrak{R}^m). \end{aligned}$$

For the terms $\mathfrak{I}_{-\infty,t}^m \mathfrak{V}_{-\infty,t} - 2\mathfrak{R}^m \mathfrak{I}_{-\infty,t}^m$, $\mathfrak{I}_{-\infty,t}^m (\mathfrak{I}_{-\infty,t}^m)^2 - 3\mathfrak{R}^m \mathfrak{I}_{-\infty,t}^m$ using again [18, Proposition 1.1.2] for $m' > m$ we have that

$$\begin{aligned} \mathfrak{I}_{-\infty,t}^m \mathfrak{V}_{-\infty,t} - 2\mathfrak{R}^m \mathfrak{I}_{-\infty,t}^m &= \mathfrak{I}_{-\infty,t}^m \otimes \mathfrak{V}_{-\infty,t}^{m'} + 2\mathfrak{R}^m (\mathfrak{I}_{-\infty,t}^{m'} - \mathfrak{I}_{-\infty,t}^m), \\ \mathfrak{I}_{-\infty,t}^m (\mathfrak{I}_{-\infty,t}^m)^2 - 3\mathfrak{R}^m \mathfrak{I}_{-\infty,t}^m &= \mathfrak{I}_{-\infty,t}^{m'} \otimes \mathfrak{V}_{-\infty,t}^m + \mathfrak{R}^m (\mathfrak{I}_{-\infty,t}^{m'} - \mathfrak{I}_{-\infty,t}^m). \end{aligned}$$

If we proceed again in the spirit of the proof of Proposition 2.3 (see Appendix E) we obtain that

$$\lim_{m \rightarrow \infty} \lim_{m' \rightarrow \infty} \mathbb{E} \sup_{t \leq T} \|\mathfrak{I}_{-\infty,t}^m \otimes \mathfrak{V}_{-\infty,t}^{m'} - \mathfrak{V}_{-\infty,t}\|_{\mathcal{C}^{-\alpha}}^p = 0,$$

$$\lim_{m \rightarrow \infty} \lim_{m' \rightarrow \infty} \mathbb{E} \sup_{t \leq T} \|\mathfrak{I}_{-\infty,t}^{m'} \otimes \mathfrak{V}_{-\infty,t}^m - \mathfrak{V}_{-\infty,t}\|_{\mathcal{C}^{-\alpha}}^p = 0,$$

$$\lim_{m \rightarrow \infty} \lim_{m' \rightarrow \infty} (\mathfrak{R}^m)^p \mathbb{E} \sup_{t \leq T} \|\mathfrak{I}_{-\infty,t}^{m'} - \mathfrak{I}_{-\infty,t}^m\|_{\mathcal{C}^{-\alpha}}^p = 0,$$

for every $p \geq 2$. It remains to handle the terms

$$h_m(t)(\mathfrak{V}_{-\infty,t} - (\mathfrak{I}_{-\infty,t}^m \mathfrak{I}_{-\infty,t}^m - \mathfrak{R}^m)), \quad (6.1)$$

$$h_m(t)((\mathfrak{I}_{-\infty,t}^m)^2 - \mathfrak{R}^m - (\mathfrak{I}_{-\infty,t}^m \mathfrak{I}_{-\infty,t}^m - \mathfrak{R}^m)) \quad (6.2)$$

and

$$h_m(t)^2(\mathfrak{I}_{-\infty,t} - \mathfrak{I}_{-\infty,t}^m). \quad (6.3)$$

We only show that (6.1) converges to 0 since (6.2) and (6.3) can be handled in a similar way. In particular due to Bony estimates (see Proposition A.6) it suffices to prove that the resonant term

$$h_m(t) \circ (\mathbf{V}_{-\infty,t} - (\mathfrak{I}_{-\infty,t} \mathfrak{I}_{-\infty,t}^m - \mathfrak{R}^m)) = \sum_{|\kappa_1 - \kappa_2| \leq 1} \delta_{\kappa_1} h_m(t) \delta_{\kappa_2} [\mathbf{V}_{-\infty,t} - (\mathfrak{I}_{-\infty,t} \mathfrak{I}_{-\infty,t}^m - \mathfrak{R}^m)]$$

converges to 0. Since the Fourier modes of h_m are localized at the points $2^m z_0$ and $-2^m z_0$ we have that

$$h_m(t) \circ (\mathbf{V}_{-\infty,t} - (\mathfrak{I}_{-\infty,t} \mathfrak{I}_{-\infty,t}^m - \mathfrak{R}^m)) = h_m(t) \sum_{i=-1,0,1} \delta_{m+i} [\mathbf{V}_{-\infty,t} - (\mathfrak{I}_{-\infty,t} \mathfrak{I}_{-\infty,t}^m - \mathfrak{R}^m)].$$

Let $\kappa \geq -1$ and $Y_m(t) = \mathbf{V}_{-\infty,t} - (\mathfrak{I}_{-\infty,t} \mathfrak{I}_{-\infty,t}^m - \mathfrak{R}^m)$. Then, for $i = -1, 0, 1$,

$$\begin{aligned} & \mathbb{E} \delta_\kappa [h_m(t_1) \delta_{m+i} Y_m(t_1)](z_1) \delta_\kappa [h_m(t_2) \delta_{m+i} Y_m(t_2)](\bar{z}_2) \\ &= \int_{\mathbb{T}^2 \times \mathbb{T}^2} C_{m,i}(t_1 - t_2, \bar{z}_1 - \bar{z}_2) \eta_\kappa(z_1 - \bar{z}_1) \eta_\kappa(z_2 - \bar{z}_2) h_m(t_1, \bar{z}_1) h_m(t_2, \bar{z}_2) d\bar{z}_1 d\bar{z}_2, \end{aligned}$$

where

$$C_{m,i}(t_1 - t_2, \bar{z}_1 - \bar{z}_2) = \mathbb{E} \delta_{m+i} [Y_m(t_1)](\bar{z}_1) \delta_{m+i} [Y_m(t_2)](\bar{z}_2).$$

For $m' > m$ using [18, Proposition 1.1.2] we have that $\mathfrak{I}_{-\infty,t}^{m'} \mathfrak{I}_{-\infty,t}^m - \mathfrak{R}^m = \mathfrak{I}_{-\infty,t}^{m'} \otimes \mathfrak{I}_{-\infty,t}^m$. Let $Y_{m,m'}(t) = \mathbf{V}_{-\infty,t} - \mathfrak{I}_{-\infty,t}^{m'} \otimes \mathfrak{I}_{-\infty,t}^m$ and notice that

$$\mathbb{E} \delta_{m+i} [Y_{m,m'}(t_1)](\bar{z}_1) \delta_{m+i} [Y_{m,m'}(t_2)](\bar{z}_2) = C \sum_{\substack{|l_1| > \lambda 2^{m'} \\ |l_2| > \lambda 2^m}} \prod_{j=1,2} \frac{1 - e^{-I_j |t_2 - t_1|}}{2 I_j} |\chi_{m+i}(l_1 + l_2)|^2 e_{l_1 + l_2}(\bar{z}_1 - \bar{z}_2),$$

for some constant C independent of m and m' and $I_j = 1 + 4\pi^2 |l_j|^2$. Then for every $\gamma \in (0, \frac{1}{2})$ by a change of variables

$$\begin{aligned} & \int_{\mathbb{T}^2 \times \mathbb{T}^2} C_{m,m',i}(t_1 - t_2, \bar{z}_1 - \bar{z}_2) \eta_\kappa(z_1 - \bar{z}_1) \eta_\kappa(z_2 - \bar{z}_2) h_m(t_1, \bar{z}_1) h_m(t_2, \bar{z}_2) d\bar{z}_1 d\bar{z}_2 \\ & \lesssim (m+1) |t_1 - t_2|^{2\gamma} \underbrace{\left(\sum_{\substack{l \in \mathcal{A}_{2^{m+i}} \\ l + 2^m z_0 \in \mathcal{A}_{2^\kappa}}} K^\gamma \star_{>\lambda 2^m}^2 K^\gamma(l) + \sum_{\substack{l \in \mathcal{A}_{2^{m+i}} \\ l - 2^m z_0 \in \mathcal{A}_{2^\kappa}}} K^\gamma \star_{>\lambda 2^m}^2 K^\gamma(l) \right)}_I, \end{aligned}$$

where $K^\gamma(l) = \frac{1}{(1+|l|^2)^{1-\gamma}}$ and $C_{m,m',i}$ is defined as $C_{m,i}$ with Y_m replaced by $Y_{m,m'}$. By Corollary C.3

$$I \lesssim \sum_{\substack{l \in \mathcal{A}_{2^{m+i}} \\ l + 2^m z_0 \in \mathcal{A}_{2^\kappa}}} \frac{1}{(1+|l|^2)^{1-2\gamma}} + \sum_{\substack{l \in \mathcal{A}_{2^{m+i}} \\ l - 2^m z_0 \in \mathcal{A}_{2^\kappa}}} \frac{1}{(1+|l|^2)^{1-2\gamma}},$$

thus for every $\varepsilon > 2\gamma$

$$I \lesssim 2^{2\varepsilon k} \sum_{l \in \mathbb{Z}^2} \frac{1}{(1+|l|^2)^{1-2\gamma}} \frac{1}{(1+|l+2^m z_0|^2)^\varepsilon}.$$

Using Corollary C.3 we obtain

$$\mathbb{E}\delta_\kappa[h_m(t_1)\delta_{m+i}Y_{m,m'}(t_1)](z_1)\delta_\kappa[h_m(t_2)\delta_{m+i}Y_{m,m'}(t_2)](z_2) \lesssim \frac{2^{2\varepsilon\kappa}(m+1)}{(1+|2^m z_0|^2)^{\varepsilon-2\gamma}}|t_1-t_2|^{2\gamma},$$

for every $\gamma \in (0, \frac{1}{2})$ and $\varepsilon > 2\gamma$. Using Nelson's estimate (B.3) for every $p \geq 2$, the usual Kolmogorov's criterion and the embedding $\mathcal{B}_{p,p}^{-\alpha+\frac{2}{p}} \hookrightarrow \mathcal{C}^{-\alpha}$ we finally obtain that

$$\lim_{m \rightarrow \infty} \lim_{m' \rightarrow \infty} \mathbb{E} \sup_{t \leq T} \|h_m(t) \circ (\mathbf{V}_{-\infty,t} - (\mathbf{I}_{-\infty,t} \mathbf{I}_{-\infty,t}^m - \mathfrak{R}^m))\|_{\mathcal{C}^{-\alpha}}^p = 0.$$

Convergence of $h_m(\mathbf{V}_{-\infty,\cdot} - (\mathbf{I}_{-\infty,\cdot} \mathbf{I}_{-\infty,\cdot}^m - \mathfrak{R}^m))$ to 0 then follows by Bony estimates (see Proposition A.6). \square

For $x \in \mathcal{C}^{-\alpha_0}$, $f \in L^2(\mathbb{R} \times \mathbb{T}^2)$ and $\mathfrak{R} \geq 0$, let $\mathcal{T}(x; f; \mathfrak{R})$ be the solution map of the equation

$$\begin{cases} \partial_t X = \Delta X - X - \sum_{k=0}^3 a_k \mathcal{H}_k(X, \mathfrak{R}) + f, \\ f(0, \cdot) = x. \end{cases} \quad (6.4)$$

The following corollary is an immediate consequence of Theorem 6.3.

Corollary 6.4. *Let $X(\cdot; x)$ be the solution to (3.2) for $n = 3$ and $x \in \mathcal{C}^{-\alpha_0}$ and denote by $\mathbb{P}_{X(\cdot; x)}$ its law in $C([0, T]; \mathcal{C}^{-\alpha_0})$. Then*

$$\text{supp } \mathbb{P}_{X(\cdot; x)} = \overline{\{\mathcal{T}(x; f; \mathfrak{R}) : f \in L^2(\mathbb{R} \times \mathbb{T}^2), \mathfrak{R} \geq 0\}}^{\mathcal{C}([0, T]; \mathcal{C}^{-\alpha_0})}.$$

Proof. See the proof of [4, Theorem 1.1]. \square

Using the above corollary we prove that for every $y \in \mathcal{C}^{-\alpha_0}$ and every $\varepsilon > 0$

$$\mathbb{P}(X(T; x) \in B_\varepsilon(y)) > 0. \quad (6.5)$$

To do so, it suffices to prove that for every $y \in C^\infty(\mathbb{T}^2)$ there exist $f \in L^2(\mathbb{T}^2)$ and $\mathfrak{R} \geq 0$ such that $\mathcal{T}(x; f; \mathfrak{R})(T) = y$. But if we set

$$X(t) = S(t)x + \frac{t}{T}(y - S(T)x),$$

for any choice of $\mathfrak{R} \geq 0$ and

$$f(t) = \sum_{k=0}^3 a_k \mathcal{H}_k(X(t), \mathfrak{R}) + \frac{1}{T}(y - S(T)x) - \frac{t}{T}(\Delta - I)(y - S(T)x)$$

we have that $X = \mathcal{T}(x; f; \mathfrak{R})$. Then the result follows by Corollary 6.4 and the fact that $C^\infty(\mathbb{T}^2)$ is dense in $\mathcal{C}^{-\alpha_0}$.

6.2. Convergence rate

We recall that for any coupling M of probability measures μ_1, μ_2 and F, G measurable functions with respect to the corresponding σ -algebras we have the identity

$$\int (F(x) - G(y))M(dx, dy) = \int \int (F(x) - G(y))\mu_1(dx)\mu_2(dy). \quad (6.6)$$

We finally combine the results of the previous sections to prove the following theorem.

Theorem 6.5. *Let $\{P_t : t \geq 0\}$ be the Markov semigroup (4.2) associated to the solution of (3.2) for $n = 3$. Then there exists $\lambda \in (0, 1)$ such that*

$$\|P_t^* \delta_x - P_t^* \delta_y\|_{\text{TV}} \leq 1 - \lambda, \quad (6.7)$$

for every $x, y \in \mathcal{C}^{-\alpha_0}$, $t \geq 3$.

Proof. Let $0 < \alpha < \alpha_0$ and for $R > 0$ consider the subset of $\mathcal{C}^{-\alpha_0}$

$$A_R := \{x \in \mathcal{C}^{-\alpha_0} : \|x\|_{\mathcal{C}^{-\alpha}} \leq R\}$$

which is compact since the embedding $\mathcal{C}^{-\alpha} \hookrightarrow \mathcal{C}^{-\alpha_0}$ is compact (see Proposition A.4). By Theorem 5.10 for every $a \in (0, 1)$ there exists $r \equiv r(a) > 0$ such that for every $x, y \in \bar{B}_r(0)$ and $t \geq 1$

$$\|P_t^* \delta_x - P_t^* \delta_y\|_{\text{TV}} \leq 1 - a.$$

By (6.5) for every $x \in A_R$

$$P_1(x; \bar{B}_r(0)) > 0,$$

which combined with the strong Feller property (which implies the continuity of $P_1(x; A)$ as a function of x for fixed measurable set A) and the fact that A_R is compact implies that there exists $b \equiv b(R) > 0$ such that

$$\inf_{x \in A_R} P_1(x; \bar{B}_r(0)) \geq b.$$

For $t \geq 0$ and $x, y \in A_R \setminus \bar{B}_r(0)$, let $\mathbb{P}_t^{x,y} \in \mathcal{M}_1(\mathcal{C}^{-\alpha_0} \times \mathcal{C}^{-\alpha_0})$ be the product coupling of $P_t(x)$ and $P_t(y)$ given by

$$\mathbb{P}_t^{x,y}(A \times B) = P_t(x; A)P_t(y; B),$$

for every measurable sets $A, B \subset \mathcal{C}^{-\alpha_0}$. Then, for $x, y \in A_R$, $t \geq 2$ and $\Phi \in C_b(\mathcal{C}^{-\alpha_0})$,

$$\begin{aligned} |P_t \Phi(x) - P_t \Phi(y)| &= |\mathbb{E}[P_{t-1} \Phi(X(1; x)) - P_{t-1} \Phi(X(1; y))]| \\ &= \left| \int [P_{t-1} \Phi(\tilde{x}) - P_{t-1} \Phi(\tilde{y})] \mathbb{P}_1^{x,y}(d\tilde{x}, d\tilde{y}) \right|, \end{aligned}$$

where in the first equality we use the Markov property and in the second (6.6). This implies that

$$\begin{aligned} \|P_t^* \delta_x - P_t^* \delta_y\|_{\text{TV}} &\leq \mathbb{P}_1^{x,y}((\bar{B}_r(0) \times \bar{B}_r(0))^c) + (1 - a) \mathbb{P}_1^{x,y}(\bar{B}_r(0) \times \bar{B}_r(0)) \\ &= 1 - a \mathbb{P}_1^{x,y}(\bar{B}_r(0) \times \bar{B}_r(0)) \\ &\leq 1 - ab^2. \end{aligned}$$

By (3.24) we can choose $R > 0$ sufficiently large such that

$$\inf_{x \in \mathcal{C}^{-\alpha_0}} \inf_{t \geq 1} \mathbb{P}(\|X(t; x)\|_{\mathcal{C}^{-\alpha}} \leq R) > \frac{1}{2}.$$

Then for any $x, y \in \mathcal{C}^{-\alpha_0}$ and $t \geq 3$, using the same coupling argument as above we get

$$\|P_t^* \delta_x - P_t^* \delta_y\|_{\text{TV}} \leq 1 - \frac{ab^2}{4},$$

which completes the proof if we set $\lambda = \frac{ab^2}{4}$. □

The following corollary contains our main result, the exponential convergence to a unique invariant measure.

Corollary 6.6. *There exists a unique invariant measure $\mu \in \mathcal{M}_1(\mathcal{C}^{-\alpha_0})$ for the semigroup $\{P_t : t \geq 0\}$ associated to the solution of (3.2) for $n = 3$ such that*

$$\|P_t^* \delta_x - \mu\|_{\text{TV}} \leq (1 - \lambda)^{\lfloor \frac{t}{3} \rfloor} \|\delta_x - \mu\|_{\text{TV}}, \quad (6.8)$$

for every $x \in \mathcal{C}^{-\alpha_0}$, $t \geq 3$.

Proof. We first notice that for $\mu_1, \mu_2 \in \mathcal{M}_1(\mathcal{C}^{-\alpha_0})$ and every $t \geq 0$ by (6.6) we have that

$$\|P_t^* \mu_1 - P_t^* \mu_2\|_{\text{TV}} \leq \frac{1}{2} \sup_{\|M\|_{\infty} \leq 1} \int \int |P_t \Phi(x) - P_t \Phi(y)| M(dx, dy),$$

for any coupling $M \in \mathcal{M}_1(\mathcal{C}^{-\alpha_0} \times \mathcal{C}^{-\alpha_0})$ of μ_1 and μ_2 . Thus by (6.7) for $t \geq 3$

$$\|P_t^* \mu_1 - P_t^* \mu_2\|_{\text{TV}} \leq (1 - \lambda) (1 - M(\{(x, x) : x \in \mathcal{C}^{-\alpha_0}\}))$$

and using the characterization of the total variation distance given by

$$\|\mu_1 - \mu_2\|_{\text{TV}} = \inf \{1 - M(\{(x, x) : x \in \mathcal{C}^{-\alpha_0}\}) : M \text{ coupling of } \mu_1 \text{ and } \mu_2\}$$

we get that

$$\|P_t^* \mu_1 - P_t^* \mu_2\|_{\text{TV}} \leq (1 - \lambda) \|\mu_1 - \mu_2\|_{\text{TV}}.$$

This implies that $\{P_t : t \geq 0\}$ has a unique invariant measure $\mu \in \mathcal{M}_1(\mathcal{C}^{-\alpha_0})$, since by Proposition [7, Proposition 3.2.5] any two distinct invariant measures are singular. Finally, for $x \in \mathcal{C}^{-\alpha_0}$ and $t \geq 3$

$$\|P_t^* \delta_x - \mu\|_{\text{TV}} \leq (1 - \lambda) \|P_{t-3}^* \delta_x - \mu\|_{\text{TV}},$$

which implies (6.8). □

Appendix A

The following three propositions can be found in [15, Section 3: pp. 11–12].

Proposition A.1. *Let $\alpha_1, \alpha_2 \in \mathbb{R}$, $p_1, p_2, q_1, q_2 \in [1, \infty]$. Then,*

$$\|f\|_{\mathcal{B}_{p_1, q_1}^{\alpha_1}} \leq C \|f\|_{\mathcal{B}_{p_1, q_1}^{\alpha_2}}, \quad \text{whenever } \alpha_1 \leq \alpha_2, \quad (\text{A.1})$$

$$\|f\|_{\mathcal{B}_{p_1, q_1}^{\alpha_1}} \leq \|f\|_{\mathcal{B}_{p_1, q_2}^{\alpha_1}}, \quad \text{whenever } q_1 \geq q_2, \quad (\text{A.2})$$

$$\|f\|_{\mathcal{B}_{p_1, q_1}^{\alpha_1}} \leq C \|f\|_{\mathcal{B}_{p_2, q_1}^{\alpha_1}}, \quad \text{whenever } p_1 \leq p_2, \quad (\text{A.3})$$

$$\|f\|_{\mathcal{B}_{p_1, q_1}^{\alpha_1}} \leq C \|f\|_{\mathcal{B}_{p_1, q_2}^{\alpha_2}}, \quad \text{whenever } \alpha_1 < \alpha_2. \quad (\text{A.4})$$

Proposition A.2. *Let $p \in [1, \infty]$. Then the space $\mathcal{B}_{p, 1}^0$ is continuously embedded in L^p and*

$$\|f\|_{L^p} \leq \|f\|_{\mathcal{B}_{p, 1}^0}. \quad (\text{A.5})$$

On the other hand, L^p is continuously embedded in $\mathcal{B}_{p, \infty}^0$ and

$$\|f\|_{\mathcal{B}_{p, \infty}^0} \leq C \|f\|_{L^p}. \quad (\text{A.6})$$

Proposition A.3. *Let $\alpha \leq \beta$ and $p, q \geq 1$ such that $p \geq q$ and $\beta = \alpha + d(\frac{1}{q} - \frac{1}{p})$. Then*

$$\|f\|_{\mathcal{B}_{p,\infty}^\alpha} \leq C \|f\|_{\mathcal{B}_{q,\infty}^\beta}.$$

The following proposition can be found in [2, Corollary 2.96] and it is generally true for Besov spaces over compact sets.

Proposition A.4. *Let $\alpha < \alpha'$. Then the embedding $\mathcal{B}_{\infty,\infty}^{\alpha'} \hookrightarrow \mathcal{B}_{\infty,1}^\alpha$ is compact.*

In the following proposition we describe the smoothing properties of the heat semigroup $(e^{t\Delta})_{t \geq 0}$ with generator Δ in space (see [15, Proposition 3.11]).

Proposition A.5. *Let $f \in \mathcal{B}_{p,q}^\alpha$. Then, for all $\beta \geq \alpha$,*

$$\|e^{t\Delta} f\|_{\mathcal{B}_{p,q}^\beta} \leq C t^{\frac{\alpha-\beta}{2}} \|f\|_{\mathcal{B}_{p,q}^\alpha}, \quad (\text{A.7})$$

for every $t \leq 1$.

For $f, g \in C^\infty(\mathbb{T}^d)$ we define the paraproduct $f \prec g$ and the resonant term $f \circ g$ by

$$f \prec g := \sum_{\iota < \kappa - 1} \delta_\iota f \delta_\kappa g, \quad (\text{A.8})$$

$$f \circ g := \sum_{|\iota - \kappa| \leq 1} \delta_\iota f \delta_\kappa g. \quad (\text{A.9})$$

We also let $f \succ g := g \prec f$. Notice that formally

$$fg = f \prec g + f \circ g + f \succ g.$$

We then have the following estimates due to Bony.

Proposition A.6. ([2, Theorems 2.82 and 2.85]) *Let $\alpha, \beta \in \mathbb{R}$ and $g \in \mathcal{C}^\beta$.*

- (i) *If $f \in L^\infty$, $\|f \prec g\|_{\mathcal{C}^\beta} \leq C \|f\|_{L^\infty} \|g\|_{\mathcal{C}^\beta}$.*
- (ii) *If $\alpha < 0$ and $f \in \mathcal{C}^\alpha$, $\|f \prec g\|_{\mathcal{C}^{\alpha+\beta}} \leq C \|f\|_{\mathcal{C}^\alpha} \|g\|_{\mathcal{C}^\beta}$.*
- (iii) *If $\alpha + \beta > 0$ and $f \in \mathcal{C}^\alpha$, $\|f \circ g\|_{\mathcal{C}^{\alpha+\beta}} \leq C \|f\|_{\mathcal{C}^\alpha} \|g\|_{\mathcal{C}^\beta}$.*

The above proposition allows us to define the product of a distribution and a function in a canonical way under certain regularity assumptions (see [15, Corollary 3.21]).

Proposition A.7. *Let $f \in \mathcal{C}^\alpha$ and $g \in \mathcal{C}^\beta$, where $\alpha < 0 < \beta$, $\alpha + \beta > 0$. Then fg can be uniquely defined as an element in \mathcal{C}^α such that*

$$\|fg\|_{\mathcal{C}^\alpha} \leq C \|f\|_{\mathcal{C}^\alpha} \|g\|_{\mathcal{C}^\beta}.$$

Regarding the inner product on $L^2(\mathbb{T}^d)$ we have the following extension result (see [15, Proposition 3.23]).

Proposition A.8. *Let $p, q \geq 1$ and p', q' their conjugate exponents. Then, for every $0 \leq \alpha < 1$, the $L^2(\mathbb{T}^d)$ inner product can be uniquely extended to a continuous bilinear form on $\mathcal{B}_{p,q}^\alpha \times \mathcal{B}_{p',q'}^{-\alpha}$ such that*

$$|\langle f, g \rangle| \leq C \|f\|_{\mathcal{B}_{p,q}^\alpha} \|g\|_{\mathcal{B}_{p',q'}^{-\alpha}},$$

for all $(f, g) \in \mathcal{B}_{p,q}^\alpha \times \mathcal{B}_{p',q'}^{-\alpha}$.

Finally we have the following gradient estimate for functions of positive regularity (see [15, Proposition 3.25]).

Proposition A.9. *Let $f \in \mathcal{B}_{1,1}^\alpha$, $\alpha \in (0, 1)$. Then*

$$\|f\|_{\mathcal{B}_{1,1}^\alpha} \leq C(\|f\|_{L^1}^{1-\alpha} \|\nabla f\|_{L^1}^\alpha + \|f\|_{L^1}). \quad (\text{A.10})$$

Appendix B

Definition B.1. Let $\{\xi(\phi)\}_{\phi \in L^2(\mathbb{R} \times \mathbb{T}^d)}$ be a family of centered Gaussian random variables on a probability space $(\Omega, \mathcal{F}, \mathbb{P})$ such that

$$\mathbb{E}(\xi(\phi)\xi(\psi)) = \langle \phi, \psi \rangle_{L^2(\mathbb{R} \times \mathbb{T}^d)},$$

for all $\phi, \psi \in L^2(\mathbb{R} \times \mathbb{T}^d)$. Then ξ is called a space-time white noise on $\mathbb{R} \times \mathbb{T}^d$.

The existence of such a family of random variables on some probability space $(\Omega, \mathcal{F}, \mathbb{P})$ is assured by Kolmogorov's extension theorem and by definition we can check that it is linear, i.e. for all $\lambda, \nu \in \mathbb{R}$, $\phi, \psi \in L^2(\mathbb{R} \times \mathbb{T}^d)$ we have that $\xi(\lambda\phi + \nu\psi) = \lambda\xi(\phi) + \nu\xi(\psi)$ \mathbb{P} -almost surely (see [18, Chapter 1]). We interpret $\xi(\phi)$ as a stochastic integral and write

$$\int_{\mathbb{R} \times \mathbb{T}^d} \phi(t, x) \xi(dt, dx) := \xi(\phi),$$

for all $\phi \in L^2(\mathbb{R} \times \mathbb{T}^d)$. We use this notation, but stress that ξ is almost surely not a measure and that the stochastic integral is only defined on a set of measure one which may depend on the specific choice of ϕ .

We also define multiple stochastic integrals (see [18, Chapter 1]) on $\mathbb{R} \times \mathbb{T}^d$ for all symmetric functions f in $L^2((\mathbb{R} \times \mathbb{T}^d)^n)$, for some $n \in \mathbb{N}$, i.e. functions such that $f(z_1, z_2, \dots, z_n) = f(z_{i_1}, z_{i_2}, \dots, z_{i_n})$ for any permutation (i_1, i_2, \dots, i_n) of $(1, 2, \dots, n)$. Here z_j is an element of $\mathbb{R} \times \mathbb{T}^d$, for all $j \in \{1, 2, \dots, n\}$. For such a symmetric function f we denote its n th iterated stochastic integral by

$$I_n(f) := \int_{(\mathbb{R} \times \mathbb{T}^d)^n} f(z_1, z_2, \dots, z_n) \xi(dz_1 \otimes dz_2 \otimes \dots \otimes dz_n).$$

The following theorem can be found in [18, Theorem 1.1.2].

Theorem B.2. *Let \mathcal{F}_ξ be the σ -algebra generated by the family of random variables $\{\xi(\phi)\}_{\phi \in L^2(\mathbb{R} \times \mathbb{T}^d)}$. Then every element $X \in L^2(\Omega, \mathcal{F}_\xi, \mathbb{P})$ can be written in the following form*

$$X = \mathbb{E}(X) + \sum_{n=1}^{\infty} I_n(f_n),$$

where $f_n \in L^2((\mathbb{R} \times \mathbb{T}^d)^n)$ are symmetric functions, uniquely determined by X .

The above theorem implies that $L^2(\Omega, \mathcal{F}_\xi, \mathbb{P})$ can be decomposed into a direct sum of the form $\bigoplus_{n \geq 0} S_n$, where $S_0 := \mathbb{R}$ and

$$S_n := \{I_n(f) : f \in L^2((\mathbb{R} \times \mathbb{T}^d)^n) \text{ symmetric}\}, \quad (\text{B.1})$$

for all $n \geq 1$. The space S_n is called the n th homogeneous Wiener chaos and the element $I_n(f_n)$ the projection of X onto S_n .

Given a symmetric function $f \in L^2((\mathbb{R} \times \mathbb{T}^d)^n)$, we have the isometry

$$\mathbb{E}(I_n)^2 = n! \|f\|_{L^2((\mathbb{R} \times \mathbb{T}^d)^n)}^2. \tag{B.2}$$

Furthermore, by Nelson’s estimate (see [18, Section 1.4]) for every $n \geq 1$ and $Y \in S_n$,

$$\mathbb{E}|Y|^p \leq (p - 1)^{\frac{n}{2}} (\mathbb{E}|Y|^2)^{\frac{p}{2}}, \tag{B.3}$$

for every $p \geq 2$.

Appendix C

Definition C.1. For symmetric kernels $K_1, K_2 : \mathbb{Z}^2 \rightarrow (0, \infty)$ we denote by $K_1 \star K_2$ the convolution given by

$$K_1 \star K_2(m) := \sum_{l \in \mathbb{Z}^2} K_1(m - l) K_2(l)$$

and for $N \in \mathbb{N}$ we let

$$K_1 \star_{\leq N} K_2(m) := \sum_{|l| \leq N} K_1(m - l) K_2(l),$$

as well as

$$K_1 \star_{> N} K_2 := (K_1 \star K_2) - (K_1 \star_{\leq N} K_2).$$

We are interested in symmetric kernels K for which there exists $\alpha \in (0, 1]$ such that

$$K(m) \leq C \frac{1}{(1 + |m|^2)^\alpha}.$$

In the spirit of [9, Lemma 10.14] we have the following lemma.

Lemma C.2. Let $\alpha, \beta \in (0, 1]$ such that $\alpha + \beta - 1 > 0$ and let $K_1, K_2 : \mathbb{Z}^2 \rightarrow (0, \infty)$ be symmetric kernels such that

$$K_1(m) \leq C \frac{1}{(1 + |m|^2)^\alpha}, \quad K_2(m) \leq C \frac{1}{(1 + |m|^2)^\beta}.$$

If $\alpha < 1$ or $\beta < 1$ then

$$K_1 \star K_2(m) \leq C \frac{1}{(1 + |m|^2)^{\alpha + \beta - 1}},$$

$$K_1 \star_{> N} K_2(m) \leq C \begin{cases} \frac{1}{(1 + |m|^2)^{\alpha + \beta - 1}}, & \text{if } |m| \geq N, \\ \frac{1}{(1 + |N|^2)^{\alpha + \beta - 1}}, & \text{if } |m| < N \end{cases}$$

and if $\alpha = \beta = 1$

$$K_1 \star K_2(m) \leq C \frac{\log |m| \vee 1}{1 + |m|^2},$$

$$K_1 \star_{> N} K_2(m) \leq C \begin{cases} \frac{\log |m| \vee 1}{1 + |m|^2}, & \text{if } |m| \geq N, \\ \frac{\log |N| \vee 1}{1 + |N|^2}, & \text{if } |m| < N. \end{cases}$$

Proof. We only prove the estimates for $K_1 \star K_2$. The corresponding estimates for $K_1 \star_{>N} K_2$ can be proven in a similar way. We consider the following regions of \mathbb{Z}^2 ,

$$A_1 = \left\{ l : |l| \leq \frac{|m|}{2} \right\},$$

$$A_2 = \left\{ l : |l - m| \leq \frac{|m|}{2} \right\},$$

$$A_3 = \left\{ l : \frac{|m|}{2} \leq |l| \leq 2|m|, |l - m| \geq \frac{|m|}{2} \right\},$$

$$A_4 = \{ l : |l| > 2|m| \}.$$

For every $l \in A_1$ we notice that $|m - l| \geq \frac{3|m|}{4}$, which implies that

$$\begin{aligned} \sum_{l \in A_1} K_1(m-l)K_2(l) &\lesssim \frac{1}{(1+|m|^2)^\alpha} \sum_{l \in A_1} K_2(l) \\ &\lesssim \begin{cases} \frac{(1+|m|^2)^{\beta-1}}{(1+|m|^2)^\alpha}, & \text{if } \beta < 1, \\ \frac{\log |m| \vee 1}{(1+|m|^2)^\alpha}, & \text{if } \beta = 1. \end{cases} \end{aligned}$$

By symmetry we get that

$$\sum_{l \in A_2} K_1(m-l)K_2(l) \lesssim \begin{cases} \frac{(1+|m|^2)^{\alpha-1}}{(1+|m|^2)^\beta}, & \text{if } \alpha < 1, \\ \frac{\log |m| \vee 1}{(1+|m|^2)^\beta}, & \text{if } \alpha = 1. \end{cases}$$

For the summation over A_3 we notice that

$$\sum_{l \in A_3} K_1(m-l)K_2(l) \lesssim \frac{1+|m|^2}{(1+|m|^2)^{\alpha+\beta}}.$$

Finally, for $l \in A_4$ we have that $|m - l| \geq \frac{|l|}{2}$, which implies that

$$\sum_{l \in A_4} K_1(m-l)K_2(l) \lesssim \sum_{|l| > 2|m|} \frac{1}{(1+|l|^2)^{\alpha+\beta}} \lesssim \frac{1}{(1+|m|^2)^{\alpha+\beta}}.$$

Combining all the above we thus obtain the appropriate estimate on $K_1 \star K_2(m)$. \square

Because we are interested in nested convolutions of the same kernel we introduce the following recursive notation

$$K \star^1 K = K, \quad K \star^n K = K \star (K \star^{n-1} K),$$

for every $n \geq 2$, with the obvious interpretation for $K \star_{\leq N}^n K$ and $K \star_{> N}^n K$. We then have the following corollary, the proof of which is omitted since it is an immediate consequence of Lemma C.2.

Corollary C.3. *Let K be a symmetric kernel as above for some $\alpha \in (\frac{n-1}{n}, 1]$. If $\alpha < 1$ then*

$$\begin{aligned} K \star^n K(m) &\leq C \frac{1}{(1+|m|^2)^{n\alpha-(n-1)}}, \\ K \star_{>N}^n K(m) &\leq C \begin{cases} \frac{1}{(1+|m|^2)^{n\alpha-(n-1)}}, & \text{if } |m| \geq N, \\ \frac{1}{(1+|N|^2)^{n\alpha-(n-1)}}, & \text{if } |m| < N \end{cases} \end{aligned}$$

and if $\alpha = 1$

$$K \star^n K(m) \leq C \frac{1}{(1 + |m|^2)^{1-\varepsilon}},$$

$$K \star_{>N} K(m) \leq C \begin{cases} \frac{1}{(1+|m|^2)^{1-\varepsilon}}, & \text{if } |m| \geq N, \\ \frac{1}{(1+N^2)^{1-\varepsilon}}, & \text{if } |m| < N \end{cases}$$

for every $\varepsilon \in (0, 1)$.

Appendix D

Proof of Theorem 2.1. Let $\phi_1, \phi_2 \in L^2(\mathbb{T}^2)$ and notice that for $t_1, t_2 > -\infty$ by (B.2)

$$\mathbb{E} \overset{\circ}{\nabla}_{-\infty, t_1}^n(\phi_1) \overset{\circ}{\nabla}_{-\infty, t_2}^n(\phi_2) = n! \int_{\mathbb{T}^2} \int_{\mathbb{T}^2} \phi_1(z_1) \phi_2(z_2) \left(\int_{-\infty}^{t_1 \wedge t_2} H(t_1 + t_2 - 2r, z_1 - z_2) dr \right)^n dz_1 dz_2, \tag{D.1}$$

where we also use the semigroup property

$$\int_{\mathbb{T}^2} H(t_1 - r, z_1 - z) H(t_2 - r, z_2 - z) dz = H(t_1 + t_2 - 2r, z_1 - z_2).$$

For $I_m = 1 + 4\pi^2|m|^2, m \in \mathbb{Z}^2$, we rewrite (D.1) as

$$\mathbb{E} \overset{\circ}{\nabla}_{-\infty, t_1}^n(\phi_1) \overset{\circ}{\nabla}_{-\infty, t_2}^n(\phi_2) = n! \sum_{\substack{m_i \in \mathbb{Z}^2 \\ i=1,2,\dots,n \\ m=m_1+\dots+m_n}} \prod_{i=1}^n \frac{e^{-I_{m_i}|t_1-t_2|}}{2I_{m_i}} \hat{\phi}_1(m) \overline{\hat{\phi}_2(m)},$$

and if we replace ϕ_1, ϕ_2 by $\eta_\kappa(z_1 \cdot \cdot), \eta_\kappa(z_2 \cdot \cdot)$ respectively, for $\kappa \geq -1, z_1, z_2 \in \mathbb{T}^2$, we have that

$$\mathbb{E} \delta_\kappa \overset{\circ}{\nabla}_{-\infty, t_1}^n(z_1) \delta_\kappa \overset{\circ}{\nabla}_{-\infty, t_2}^n(z_2) = n! \sum_{\substack{m_i \in \mathbb{Z}^2 \\ i=1,2,\dots,n \\ m=m_1+\dots+m_n}} \prod_{i=1}^n \frac{e^{-I_{m_i}|t_1-t_2|}}{2I_{m_i}} |\chi_\kappa(m)|^2 e_m(z_1 - z_2).$$

By a change of variables we finally obtain

$$\mathbb{E} \delta_\kappa \overset{\circ}{\nabla}_{-\infty, t_1}^n(z_1) \delta_\kappa \overset{\circ}{\nabla}_{-\infty, t_2}^n(z_2) \approx n! \sum_{m_1 \in \mathcal{A}_{2^\kappa}} \sum_{\substack{m_i \in \mathbb{Z}^2 \\ i=2,\dots,n}} \prod_{i=1}^n \frac{e^{-I_{m_i-m_{i-1}}|t_1-t_2|}}{2I_{m_i-m_{i-1}}} e_{m_1}(z_1 - z_2),$$

with the convention that $m_0 = 0$. Let $K^\gamma(m) = \frac{1}{(1+|m|^2)^{1-\gamma}}$, for $\gamma \in [0, 1)$, and write $K^\gamma \star^n K^\gamma$ to denote the n th iterated convolution of K^γ with itself (see Definition C.1). If we let $z_1 = z_2 = z$, for $t_1 = t_2 = t$ we get an estimate of the form

$$\mathbb{E} \delta_\kappa \overset{\circ}{\nabla}_{-\infty, t}^n(z)^2 \lesssim \sum_{m \in \mathcal{A}_{2^\kappa}} K^0 \star^n K^0(m)$$

while for $t_1 \neq t_2$ and every $\gamma \in (0, 1)$

$$\mathbb{E} (\delta_\kappa \overset{\circ}{\nabla}_{-\infty, t_1}^n(z) - \delta_\kappa \overset{\circ}{\nabla}_{-\infty, t_2}^n(z))^2 \lesssim |t_1 - t_2|^{n\gamma} \sum_{m \in \mathcal{A}_{2^\kappa}} K^\gamma \star^n K^\gamma(m).$$

By Corollary C.3

$$\mathbb{E} \delta_\kappa \overset{\circ}{\nabla}_{-\infty, t}^n(z)^2 \lesssim \sum_{m \in \mathcal{A}_{2^\kappa}} \frac{1}{(1 + |m|^2)^{1-\varepsilon}},$$

for every $\varepsilon \in (0, 1)$, and

$$\mathbb{E} (\delta_\kappa \overset{\circ}{\nabla}_{-\infty, t_1}^n(z) - \delta_\kappa \overset{\circ}{\nabla}_{-\infty, t_2}^n(z))^2 \lesssim |t_1 - t_2|^{n\gamma} \sum_{m \in \mathcal{A}_{2^\kappa}} \frac{1}{(1 + |m|^2)^{1-n\gamma}}.$$

Using the fact that $m \in \mathcal{A}_{2^\kappa}$ we have that for every $\kappa \geq -1$

$$\begin{aligned} \mathbb{E} \delta_\kappa \overset{\circ}{\nabla}_{-\infty, t}^n(z)^2 &\lesssim 2^{2\lambda_1 \kappa}, \\ \mathbb{E} (\delta_\kappa \overset{\circ}{\nabla}_{-\infty, t_1}^n(z) - \delta_\kappa \overset{\circ}{\nabla}_{-\infty, t_2}^n(z))^2 &\lesssim |t_1 - t_2|^{n\gamma} 2^{2\lambda_2 \kappa} \end{aligned}$$

for every $\lambda_1 > 0$ and every $\gamma \in (0, \frac{1}{n})$, $\lambda_2 > n\gamma$, while for every $p \geq 2$ by Nelson's estimate (B.3) we finally get

$$\begin{aligned} \mathbb{E} \delta_\kappa \overset{\circ}{\nabla}_{-\infty, t}^n(z)^p &\lesssim 2^{p\lambda_1 \kappa}, \\ \mathbb{E} (\delta_\kappa \overset{\circ}{\nabla}_{-\infty, t_1}^n(z) - \delta_\kappa \overset{\circ}{\nabla}_{-\infty, t_2}^n(z))^p &\lesssim |t_1 - t_2|^{n\frac{p}{2}\gamma} 2^{p\lambda_2 \kappa}. \end{aligned}$$

The result then follows from [15, Lemma 5.2, Lemma 5.3], the usual Kolmogorov's criterion and the embedding $\mathcal{B}_{p,p}^{-\alpha + \frac{2}{p}} \hookrightarrow \mathcal{C}^{-\alpha}$, for $\alpha > \frac{2}{p}$. \square

Appendix E

Proof of Proposition 2.3. For all $n \geq 1$, using the formula

$$\mathcal{H}_n(X + Y, C) = \sum_{k=0}^n \binom{n}{k} X^k \mathcal{H}_{n-k}(Y, C)$$

we have

$$\overset{\circ}{\nabla}_{s,t}^n{}^\varepsilon = \sum_{k=0}^n \binom{n}{k} (-1)^k (S(t-s))_{-\infty, s}^{\varepsilon}{}^k \mathcal{H}_{n-k}(\overset{\circ}{\nabla}_{-\infty, t}^\varepsilon, \mathfrak{R}^\varepsilon).$$

Thus it suffices to prove convergence only for $\overset{\circ}{\nabla}_{-\infty, t}^n{}^\varepsilon$, $n \geq 1$.

By [18, Proposition 1.1.4] for $t_1, t_2 > -\infty$ and $z_1, z_2 \in \mathbb{T}^2$

$$\mathbb{E} \overset{\circ}{\nabla}_{-\infty, t_1}^n{}^\varepsilon(z_1) \overset{\circ}{\nabla}_{-\infty, t_2}^n{}^\varepsilon(z_2) = n! (\mathbb{E} \overset{\circ}{\nabla}_{-\infty, t_1}^\varepsilon(z_1) \overset{\circ}{\nabla}_{-\infty, t_2}^\varepsilon(z_2))^n.$$

Using (D.1) we get

$$\mathbb{E} \overset{\circ}{\nabla}_{-\infty, t_1}^n{}^\varepsilon(z_1) \overset{\circ}{\nabla}_{-\infty, t_2}^n{}^\varepsilon(z_2) = n! \sum_{\substack{|m_i| \leq \frac{1}{\varepsilon} \\ i=1, 2, \dots, n \\ m=m_1+\dots+m_n}} \prod_{i=1}^n \frac{e^{-I_{m_i}|t_1-t_2|}}{2I_{m_i}} e_m(z_1 - z_2),$$

and by a change of variables the above implies that for $\kappa \geq -1$

$$\mathbb{E} \delta_\kappa \nabla_{-\infty, t_1}^\varepsilon(z_1) \delta_\kappa \nabla_{-\infty, t_2}^\varepsilon(z_2) \approx n! \sum_{m_1 \in \mathcal{A}_{2\kappa}} \sum_{\substack{|m_i| \leq \frac{1}{\varepsilon} \\ i=2, \dots, n}} \prod_{i=1}^n \frac{e^{-I_{m_i - m_{i-1}} |t_1 - t_2|}}{2I_{m_i - m_{i-1}}} e_{m_1}(z_1 - z_2). \tag{E.1}$$

In a similar way

$$\mathbb{E} \delta_\kappa \nabla_{-\infty, t_1}^\varepsilon(z_1) \delta_\kappa \nabla_{-\infty, t_2}^\varepsilon(z_2) \approx n! \sum_{m_1 \in \mathcal{A}_{2\kappa}} \sum_{\substack{|m_i| \leq \frac{1}{\varepsilon} \\ i=2, \dots, n}} \prod_{i=1}^n \frac{e^{-I_{m_i - m_{i-1}} |t_1 - t_2|}}{2I_{m_i - m_{i-1}}} e_{m_1}(z_1 - z_2) \tag{E.2}$$

and for $K^\gamma(m) = \frac{1}{(1+|m|^2)^{1-\gamma}}$ combining (E.1) and (E.2) for $z_1 = z_2 = z$ and $t_1 = t_2 = t$ we have that

$$\mathbb{E} (\delta_\kappa \nabla_{-\infty, t}^\varepsilon(z) - \delta_\kappa \nabla_{-\infty, t}^\varepsilon(z))^2 \lesssim \sum_{m \in \mathcal{A}_{2\kappa}} K^0 \star_{> \frac{1}{\varepsilon}}^n K^0(m),$$

while for $t_1 \neq t_2$ and every $\gamma \in (0, 1)$

$$\begin{aligned} & \mathbb{E} [(\delta_\kappa \nabla_{-\infty, t_1}^\varepsilon(z) - \delta_\kappa \nabla_{-\infty, t_1}^\varepsilon(z)) (\delta_\kappa \nabla_{-\infty, t_2}^\varepsilon(z) - \delta_\kappa \nabla_{-\infty, t_2}^\varepsilon(z))] \\ & \lesssim |t_1 - t_2|^{n\gamma} \sum_{m \in \mathcal{A}_{2\kappa}} K^\gamma \star_{> \frac{1}{\varepsilon}}^n K^\gamma(m). \end{aligned}$$

Proceeding as in the proof of Theorem 2.1 (see Appendix D) and using Corollary C.3 we obtain that

$$\mathbb{E} (\delta_\kappa \nabla_{-\infty, t}^\varepsilon(z) - \delta_\kappa \nabla_{-\infty, t}^\varepsilon(z))^2 \lesssim 2^{2\lambda_1 \kappa} \frac{1}{(1 + \frac{1}{\varepsilon^2})^{\lambda_1/2}}$$

for every $\lambda_1 \in (0, 1)$, and

$$\begin{aligned} & \mathbb{E} [(\delta_\kappa \nabla_{-\infty, t_1}^\varepsilon(z) - \delta_\kappa \nabla_{-\infty, t_1}^\varepsilon(z)) (\delta_\kappa \nabla_{-\infty, t_2}^\varepsilon(z) - \delta_\kappa \nabla_{-\infty, t_2}^\varepsilon(z))] \\ & \lesssim |t_1 - t_2|^{n\gamma} 2^{2\lambda_2 \kappa} \frac{1}{(1 + \frac{1}{\varepsilon^2})^{\lambda_2 - n\gamma}}, \end{aligned}$$

for every $\gamma \in (0, \frac{1}{n})$, $\lambda_2 > n\gamma$. The result then follows by Nelson’s estimate (B.3) combined with the usual Kolmogorov’s criterion, as well as the embedding $\mathcal{B}_{p,p}^{-\alpha + \frac{2}{p}} \hookrightarrow \mathcal{C}^{-\alpha}$, for $\alpha > \frac{2}{p}$. □

Acknowledgements

The authors would like to thank Martin Hairer, Jonathan Mattingly and Philipp Schoenbauer for helpful discussions. PT is supported by ESPRC as part of the MASDOC DTC at the University of Warwick, Grant No. EP/HO23364/1. HW is supported by the Royal Society through the University Research Fellowship UF140187.

References

[1] S. Albeverio and M. Röckner. Stochastic differential equations in infinite dimensions: Solutions via Dirichlet forms. *Probab. Theory Related Fields* **89** (3) (1991) 347–386. [MR1113223](#)
 [2] H. Bahouri, J. Y. Chemin and R. Danchin. *Fourier Analysis and Nonlinear Partial Differential Equations*. Springer, Heidelberg, 2011. [MR2768550](#)
 [3] V. I. Bogachev. *Measure Theory*. Springer-Verlag, Berlin, Heidelberg, 2007. [MR2267655](#)

- [4] K. Chouk and P. K. Friz. Support theorem for a singular SPDE: the case of gPAM. Preprint, 2016. Available at [arXiv:1409.4250v3](#).
- [5] G. Da Prato and A. Debussche. Strong solutions to the stochastic quantization equations. *Ann. Probab.* **31** (4) (2003) 1900–1916. [MR2016604](#)
- [6] G. Da Prato and J. Zabczyk. *Stochastic Equations in Infinite Dimensions*. Cambridge University Press, Cambridge, 1992. [MR1207136](#)
- [7] G. Da Prato and J. Zabczyk. *Ergodicity for Infinite Dimensional Systems*. Cambridge University Press, Cambridge, 1996. [MR1417491](#)
- [8] L. C. Evans. *Partial Differential Equations*. AMS, Providence, 2010.
- [9] M. Hairer. A theory of regularity structures. *Invent. Math.* **198** (2) (2014) 269–504. [MR3274562](#)
- [10] M. Hairer and J. Mattingly. The strong Feller property for singular stochastic PDEs. Preprint, 2016. Available at [arXiv:1610.03415v2](#).
- [11] M. Hairer, J. Mattingly and M. Scheutzow. Asymptotic coupling and a weak form of Harris’ theorem with applications to stochastic delay equations. *Probab. Theory Related Fields* **149** (1) (2011) 223–259. [MR2773030](#)
- [12] M. Hairer, A. M. Stuart and J. Voss. Analysis of SPDEs arising in path sampling, Part II: The nonlinear case. *Ann. Appl. Probab.* **17** (5/6) (2007) 1657–1706. [MR2358638](#)
- [13] G. Jona-Lassinio and P. K. Mitter. On the stochastic quantization of field theory. *Comm. Math. Phys.* **101** (3) (1985) 409–436. [MR0815192](#)
- [14] J. C. Mourrat and H. Weber. Convergence of the two-dimensional dynamic Ising–Kac model to ϕ_2^4 . Preprint, 2014. Available at [arXiv:1410.1179](#).
- [15] J. C. Mourrat and H. Weber. Global well-posedness of the dynamic ϕ^4 in the plane. Preprint, 2015. Available at [arXiv:1501.06191](#).
- [16] E. D. Nezza, G. Palatucci and E. Valdinoci. Hitchhiker’s guide to the fractional Sobolev spaces. *Bull. Sci. Math.* **136** (2012) 521–573. [MR2944369](#)
- [17] J. R. Norris. Simplified Malliavin calculus. *Sémin. Probab. (Strasbourg)* **20** (1986) 101–130. [MR0942019](#)
- [18] D. Nualart. *The Malliavin Calculus and Related Topics*. Springer-Verlag, Berlin, 2006. [MR2200233](#)
- [19] G. Parisi and S. C. Wu. Perturbation theory without gauge fixing. *Sci. Sin.* **24** (4) (1981) 483–496. [MR0626795](#)
- [20] D. Revuz and M. Yor. *Continuous Martingales and Brownian Motion*. Springer-Verlag, Berlin, 1999. [MR1725357](#)
- [21] M. Röckner, R. Zhu and X. Zhu. Restricted markov uniqueness for the stochastic quantization of $\mathcal{P}(\phi)$ and its applications. *J. Funct. Anal.* **272** (2017) 4263–4303. [MR3626040](#)
- [22] M. Röckner, R. Zhu and X. Zhu. Ergodicity for the stochastic quantization problems on the 2D-torus. *Comm. Math. Phys.* **352** (2017) 1061–1090. [MR3631399](#)
- [23] E. Zeidler. *Applied Functional Analysis: Main Principles and Their Applications*. Springer-Verlag, New York, 1995. [MR1347692](#)