



UNIVERSITY OF LEEDS

This is a repository copy of *Detection of a novel gammaherpesvirus (genus Rhadinovirus) in wild muntjac deer in Northern Ireland*.

White Rose Research Online URL for this paper:  
<http://eprints.whiterose.ac.uk/113711/>

Version: Accepted Version

---

**Article:**

McKillen, J, Hogg, K, Lagan, P et al. (5 more authors) (2017) Detection of a novel gammaherpesvirus (genus Rhadinovirus) in wild muntjac deer in Northern Ireland. *Archives of Virology*, 162 (6). pp. 1737-1740. ISSN 0304-8608

<https://doi.org/10.1007/s00705-017-3254-z>

---

© 2017, Springer-Verlag Wien. This is an author produced version of a paper published in *Archives of Virology*. Uploaded in accordance with the publisher's self-archiving policy. The final publication is available at Springer via <https://doi.org/10.1007/s00705-017-3254-z>.

**Reuse**

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.



[eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk)  
<https://eprints.whiterose.ac.uk/>

## Detection of a novel gammaherpesvirus (genus *Rhadinovirus*) in wild muntjac deer in Northern Ireland

John McKillen<sup>1\*</sup>, Kayleigh Hogg<sup>2</sup>, Paula Lagan<sup>1</sup>, Cheryl Ball<sup>1</sup>, Simon Doherty<sup>1</sup>, Neil Reid<sup>2,3</sup>, Lisa Collins<sup>4</sup>  
Jaimie Dick<sup>2,3</sup>

<sup>1</sup>Veterinary Sciences Division, Agri-Food and Biosciences Institute, Belfast, BT4 3SD

<sup>2</sup>School of Biological Sciences, Queen's University Belfast, MBC, 97 Lisburn Road, Belfast, BT9 7BL

<sup>3</sup>Institute for Global Food Security (IGFS), Queen's University Belfast, 18-30 Malone Road, Belfast, BT9 5BN

<sup>4</sup>University of Lincoln, Brayford Pool, Lincoln, Lincolnshire, LN6 7TS

\*Corresponding author. Tel.: +44 (0)2890 525752; fax. +44 (0)2890 525823

E-mail address: [john.mckillen@afbini.gov.uk](mailto:john.mckillen@afbini.gov.uk)

### Abstract

This study represents the initial part of an investigation into the potential for non-native, wild, free-living muntjac deer (*Muntiacus reevesi*) to carry viruses that could be a threat to livestock. A degenerate PCR assay was used to screen a range of tissues from muntjac deer culled in Northern Ireland for the presence of herpesviral nucleic acids. This was followed by sequencing of PCR amplicons and phylogenetic analysis. We report the detection of a novel gammaherpesvirus most closely related to a type 2 ruminant rhadinovirus from mule deer. It remains to be determined if this new virus is pathogenic to deer or presents a risk to food security through the susceptibility of domestic livestock.

### Key words

Muntjac deer, *Muntiacus reevesi*, Rhadinovirus, gammaherpesvirus, degenerate PCR

Deer, especially introduced species, are of major concern with regards to impacts on biodiversity, forestry and agriculture [1]. Deer also affect humans directly, causing fatalities and injury through deer-vehicle collisions [2]. Consequently, expansion of non-native, introduced and native deer populations will present considerable management challenges in the future [3].

The rapidity of the spread of invasive Reeves' muntjac (*Muntiacus reevesi*) Ogilby, 1839 is currently of great concern due to deliberate translocations and natural dispersal from their initial inoculation site(s) in the

south-east of Great Britain [3]. The first verified record of muntjac deer in the wild in Northern Ireland was confirmed during June 2009 as a result of a road traffic accident near Newtownards, Co. Down [4]. This is the first known introduction of a new large mammal species in Northern Ireland since the introduction of sika deer (*Cervus nippon* Temminck, 1838) in 1870. The second record was a culled animal (referred to hereafter as ‘Animal 1’) shot in the grounds of Mount Stewart, Co. Down, in June 2011. In February 2013, another animal was shot in the same locality (referred to hereafter as ‘Animal 2’). Furthermore, in the spring of 2013 and 2014 a camera trapping survey confirmed the presence of a small muntjac population within this area [5]. In April 2013, eleven positive detections of muntjac were made at seven separate locations in this area using camera traps.

Animal 1 was a healthy male aged 55-57 weeks of age and Animal 2 was a healthy female aged approximately 15 weeks of age. Carcasses were submitted to Veterinary Sciences Division, Agri-Food and Biosciences Institute for post mortem examination. Tissues were collected from both animals and processed to produce a 10% homogenate. Tissues taken included the kidney, heart, spleen, lung, bronchial lymph node, thyroid, caecum, faeces, oesophagus, colon, small intestine, pancreas, liver, abomasum, mediastinal lymph node, mesenteric lymph node, trachea, rumen contents and blood clot. Total nucleic acids were isolated from the homogenates using the MagNA Pure LC liquid handling system and the MagNA Pure Total Nucleic Acid Isolation Kit (Roche Diagnostics Ltd., U.K.) according to the manufacturer's instructions. These were subject to SYBR Green I real-time PCR using Quantitect SYBR Green PCR master mix (Qiagen, Crawley, U.K.). PCR was carried out with nested degenerate primers designed to detect herpesviral DNA at the family level [6].

These degenerate primers were used for two purposes; 1) Diagnostic PCR and 2) PCR for phylogenetic analysis. For diagnostic PCR the first round was carried out using forward primers DFA and ILK with reverse primer KG1. The second round was carried out with forward primer ILK and reverse primer IYG; expected amplicon size was 215-315 base pairs. For the PCR for phylogenetic analysis the first round was carried out using forward primer DFA with reverse primer KG1. The second round was carried out with forward primer DFA and reverse primer IYG; the expected amplicon size was ~500 base pairs. A selection of samples found positive by the diagnostic PCR were subject to PCR for phylogenetic analysis. PCR reactions were subject to agarose gel electrophoresis and amplicons were excised and purified. Suspected herpesvirus positive samples were sent for commercial sequencing. Resultant sequence data were first identified by BlastN analysis [7]. Sequences were trimmed and contigs generated using Geneious [8] software before alignment and generation of phylogenetic

trees using MEGA6 [9]. Geneious was also used to compare percentage nucleotide identities between the muntjac viral sequences and known herpesviral sequences from GenBank.

Post-mortem examination showed no significant gross pathological lesions for either animal. Laboratory examination of Animal 1 did not show the presence of any significant bacterial or parasitic infections including bovine tuberculosis (data not shown). Amplicons of the expected size generated by the diagnostic PCR were detected in the colon, mediastinal lymph node, lung and blood clot for Animal 1 and in the bronchial lymph node in Animal 2. Sequencing and subsequent BlastN analysis of short 175 bp sequences determined that they were herpesviral sequences most similar to those found in a number of deer species. The sequences were identical in both animals. Amplicons of the expected size generated by the PCR for phylogenetic analysis were detected in the mediastinal lymph node of animal 1 and in the bronchial lymph node in Animal 2. Sequencing and subsequent BlastN analysis of a 500 bp sequence from the bronchial lymph node of Animal 2 determined that these were herpesviral sequences most closely related to the DNA dependant DNA polymerase gene of type 2 ruminant rhadinovirus of mule deer Family *Gammaherpesvirinae*, Genus *Rhadinovirus*. Phylogenetic analysis (Figure 1) of this sequence compared to homologous herpesviral sequences from GenBank confirmed this relationship with type 2 rhadinovirus of mule deer. Analysis of nucleotide similarity showed 82.2% similarity between Animal 2 and the type 2 rhadinovirus of mule deer (accession no. HM014314). Other members of the Family *Gammaherpesvirinae*, genus *Macavirus* showed nucleotide similarity of approximately 65% with Animal 2 whereas human herpesvirus and bovine herpesvirus 1 (Family *Alphaherpesvirinae*) showed lower similarity (Table 1).

The introduction of invasive species poses a great threat to conservation in Northern Ireland. There is a significant risk of them introducing novel pathogens to native wildlife that are immunologically naive. In addition invasive species pose a similar threat to domesticated food animals and a zoonotic threat to human populations. Deer are known to be susceptible to bovine, caprine and ovine viruses and *vice versa*. A recent study from Japan showed native wild and American farmed sika deer to have antibodies against bovine parainfluenza virus type 3, bovine respiratory syncytial virus, bovine adenovirus type 7, bovine coronavirus, bovine viral diarrhoea virus (BVDV) and bovine herpesvirus type 1 [10]. Antibodies to the recently emerged Schmallenberg virus which causes foetal abnormalities, have been reported in red deer (*Cervus elaphus*), roe deer (*Capreolus capreolus*) and muntjac deer in Austria, Belgium and England [11]. BVDV has been detected in a wide range of wild and captive deer in Europe and North America including roe deer, red deer, fallow deer (*Dama dama*), elk (*Cervus elaphus nelsoni*), white-tailed deer (*Odocoileus virginianus*), and mule deer

(*Odocoileus hemionus*) [12]. In the Republic of Ireland, an industry-led compulsory national BVDV eradication programme commenced in 2013 ([www.bvdfree.ie](http://www.bvdfree.ie)), while the compulsory phase of the Northern Ireland eradication programme began in 2016 ([www.animalhealthni.com/BVD.aspx](http://www.animalhealthni.com/BVD.aspx)). In order for muntjac deer to be considered as a reservoir for re-introduction of BVDV to cattle after eradication efforts, the animals must be susceptible to infection and able to shed the virus, the virus must be capable of being maintained in the population and there must be sufficient contact with domestic livestock [13]. It is not known if muntjac deer fulfil these criteria for BVDV or other pathogens, but an expanding muntjac population should be monitored closely to assess the level of risk posed.

It remains to be determined if the novel rhadinovirus detected in this study is pathogenic to deer or is a potential threat to food security if livestock are susceptible. In the animals examined the virus was present at very low levels and was only detected after two rounds of PCR in healthy animals. A member of the genus *Rhadinovirus* known as bovine herpesvirus-4 is implicated in post-partum metritis [14] in cattle and a number of primate viruses are associated with cancer in immunosuppressed individuals such as [15] Kaposi sarcoma herpesvirus (human herpesvirus 8) [16]. The genus *Macavirus*, another genus in the *Gammaherpesvirinae* subfamily [17] includes alcephaline herpes viruses 1 and ovine herpes virus 2, which cause malignant catarrhal fever.

So while the risks associated with viruses in invasive muntjac deer in Ireland are not fully understood, if the muntjac population becomes established it could potentially interact with livestock in terms of the evolution, spread and maintenance of viral infections. Camera trap and anecdotal evidence strongly suggest that muntjac are becoming established in Ireland and it is known that muntjac can form successful populations from a limited number of founding females [18]. As such there is significant value in monitoring the virological status of these invasive animals. As our phylogenetic tree is based only on a small fragment of a single gene further sequencing would be necessary to provide a definitive classification of this gammaherpesvirus and metagenomic analysis of additional samples using next generation sequencing would be a powerful method of further virus discovery.

### **Acknowledgements**

This study was commissioned and funded by the Natural Heritage Research Partnership (NHRP) between the Northern Ireland Environment Agency (NIEA) and *Quercus*, Queen's University Belfast (QUB). The joint NIEA Client Officers were Drs Michael Meharg and Declan Looney.

## Compliance with ethical standards

Funding: This study funded by the Natural Heritage Research Partnership (NHRP).

Conflict of Interest: The authors declare that they have no conflict of interest.


Ethical approval: This article does not contain any studies with human participants or animals performed by any of the authors.

## References

1. Putman RJ, Moore NP (1998) Impact of deer in lowland Britain on agriculture, forestry and conservation habitats. *Mammal Review* 28:141-164
2. Putman RJ (1997) Deer and Road Traffic Accidents: Options for Management. *Journal of environmental management* 51:43-57
3. Ward AI (2005) Expanding ranges of wild and feral deer in Great Britain. *Mammal Review* 35:165-173
4. Dick JTA, Freeman M, Provan J, Reid N (2010) First record of free-living Reeves' muntjac deer (*Muntiacus reevesi* Ogilby 1839) in Northern Ireland. *Irish Naturalists' Journal* 31:151
5. Hogg K, Freeman, M., Burdekin, O., Lennon, J., Reid, N. & Dick, J.T.A. (2014) Muntjac deer (*Muntiacus reevesi*) survey 2013-14 of the Ards Peninsula, Northern Ireland.
6. VanDevanter DR, Warrener P, Bennett L, Schultz ER, Coulter S, Garber RL, Rose TM (1996) Detection and analysis of diverse herpesviral species by consensus primer PCR. *Journal of clinical microbiology* 34:1666-1671
7. Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ (1990) Basic local alignment search tool. *Journal of molecular biology* 215:403-410
8. Kearse M, Moir R, Wilson A, Stones-Havas S, Cheung M, Sturrock S, Buxton S, Cooper A, Markowitz S, Duran C, Thierer T, Ashton B, Meintjes P, Drummond A (2012) Geneious Basic: an integrated and extendable desktop software platform for the organization and analysis of sequence data. *Bioinformatics* 28:1647-1649
9. Tamura K, Stecher G, Peterson D, Filipski A, Kumar S (2013) MEGA6: Molecular Evolutionary Genetics Analysis version 6.0. *Mol Biol Evol* 30:2725-2729
10. Yokoi K, Okazaki H, Inahara K, Hatama S (2009) Prevalence of eight bovine viruses in sika deer (*Cervus nippon yezoensis*) in Japan. *The Veterinary record* 165:754-755
11. Barlow A, Green P, Banham T, Healy N (2013) Serological confirmation of SBV infection in wild British deer. *The Veterinary record* 172:429
12. Glawischnig W, Schoepf K, Matt M (2010) Monitoring for bovine viral diarrhoea virus in Austrian red deer (*Cervus elaphus elaphus*) by using ear-notch samples. *Journal of wildlife diseases* 46:1269-1273
13. Passler T, Ditchkoff SS, Walz PH (2016) Bovine Viral Diarrhoea Virus (BVDV) in White-Tailed Deer (*Odocoileus virginianus*). *Frontiers in microbiology* 7:945
14. Donofrio G, Franceschi V, Capocéfalo A, Cavarani S, Sheldon IM (2009) Isolation and characterization of bovine herpesvirus 4 (BoHV-4) from a cow affected by post partum metritis and cloning of the genome as a bacterial artificial chromosome. *Reproductive biology and endocrinology : RB&E* 7:83
15. Russell GC, Stewart JP, Haig DM (2009) Malignant catarrhal fever: a review. *Veterinary journal* 179:324-335

16. Cesarman E (2011) Gammaherpesvirus and lymphoproliferative disorders in immunocompromised patients. *Cancer letters* 305:163-174
17. Davison AJ, Eberle R, Ehlers B, Hayward GS, McGeoch DJ, Minson AC, Pellett PE, Roizman B, Studdert MJ, Thiry E (2009) The order Herpesvirales. *Archives of virology* 154:171-177
18. Freeman MS, Beatty GE, Dick JTA, Reid N, Provan J (2016) The paradox of invasion: Reeves' muntjac deer invade the British Isles from a limited number of founding females. *Journal of Zoology* 298 54–63

**Figure 1.** Phylogenetic tree showing the relationship between NI muntjac deer herpesvirus sequence (500 base pairs of the DNA dependant DNA polymerase gene) and homologous sequences from GenBank. The evolutionary history was inferred using the UPGMA method. The tree is drawn to scale, with branch lengths in the same units as those of the evolutionary distances used to infer the phylogenetic tree. The evolutionary distances were computed using the Maximum Composite Likelihood method and are in the units of the number of base substitutions per site. All positions containing gaps and missing data were eliminated. Test of phylogeny was by the bootstrap method with 500 replications. Analyses were conducted using MEGA6.





	NI Muntjac 2	Type 2 ruminant rhadinovirus of mule deer	Ovine herpesvirus 2	Alcelaphine herpesvirus 1	Bovine herpesvirus 6	Alcelaphine herpesvirus 2	Human herpesvirus 1	Bovine herpesvirus type 1.2
NI Muntjac 2	-	82.2	63.1	65.4	65.7	64.4	45.9	29.6
Type 2 ruminant rhadinovirus of mule deer	82.2	-	63.1	64.2	64.1	64.2	44.5	30.9
Ovine herpesvirus 2	63.1	63.1	-	75.5	67.5	75.9	50.8	33.6
Alcelaphine herpesvirus 1	65.4	64.2	75.5	-	67.6	84.8	48	31.1
Bovine herpesvirus 6	65.7	64.1	67.5	67.6	-	68.8	48	30.9
Alcelaphine herpesvirus 2	64.4	64.2	75.9	84.8	68.8	-	46.9	29.8
Human herpesvirus 1	45.9	44.5	50.8	48	48	46.9	-	37.8
Bovine herpesvirus type 1.2	29.6	30.9	33.6	31.1	30.9	29.8	37.8	-

**Table 1.** Percentage sequence identity between 500 base pairs of DNA dependant DNA polymerase gene sequence from Northern Ireland muntjac rhadinovirus and other homologous sequences from GenBank. Type 2 ruminant rhadinovirus of mule deer, accession no. HM014314; ovine herpesvirus 2, accession no. DQ198083, a causative agent of sheep-associated malignant catarrhal fever; alcelaphine herpesvirus 1, accession no. NC002531, a causative agent of wildebeest-associated malignant catarrhal fever; bovine herpesvirus 6, accession no. NC024303; alcelaphine herpesvirus 2, accession no. NC024382; human herpesvirus 1, accession no. JQ780693; bovine herpesvirus 1.2, accession no. KM258881.