

Complete Genome Sequence of *Ehrlichia mineirensis*, a Novel Organism Closely Related to *Ehrlichia canis* with a New Host Association

Alejandro Cabezas-Cruz,^a Erich Zwegarth,^{b,c} Marzena Broniszweska,^b Lygia M. F. Passos,^d Múcio Flávio Barbosa Ribeiro,^e Marina Manrique,^f Raquel Tobes,^f José de la Fuente^{g,h}

Center for Infection and Immunity of Lille (CII), INSERM, CNRS UMR, Université de Lille U1019 8204, Institut Pasteur de Lille, Lille, France^a; Comparative Tropical Medicine and Parasitology, Ludwig-Maximilians-Universität München, Munich, Germany^b; Department of Veterinary Tropical Diseases, Faculty of Veterinary Science, University of Pretoria, Onderstepoort, South Africa^c; Departamento de Medicina Veterinária Preventiva, Escola de Veterinária-UFMG, Belo Horizonte, Minas Gerais, Brazil^d; Departamento de Parasitología, ICB-UFMG, Belo Horizonte, Brazil^e; Oh No Sequences! Research Group, Era7 Bioinformatics, Granada, Spain^f; SaBio, Instituto de Investigación en Recursos Cinegéticos IREC, CSIC-UCLM-JCCM, Ronda de Toledo, Spain^g; Department of Veterinary Pathobiology, Center for Veterinary Health Sciences, Oklahoma State University, Stillwater, Oklahoma, USA^h

A.C.-C. and E.Z. contributed equally to this work.

We report here the complete genome sequencing of *Ehrlichia mineirensis*, an *Ehrlichia* organism that was isolated from the hemolymph of *Rhipicephalus microplus*–engorged females. *E. mineirensis* is the best characterized *Ehrlichia* isolate from a novel cattle-related clade closely related to the monocytotropic pathogen *E. canis*.

Received 4 December 2014 Accepted 9 December 2014 Published 22 January 2015

Citation Cabezas-Cruz A, Zwegarth E, Broniszweska M, Passos LMF, Ribeiro MFB, Manrique M, Tobes R, de la Fuente J. 2015. Complete genome sequence of *Ehrlichia mineirensis*, a novel organism closely related to *Ehrlichia canis* with a new host association. *Genome Announc* 3(1):e01450-14. doi:10.1128/genomeA.01450-14.

Copyright © 2015 Cabezas-Cruz et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 3.0 Unported license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to José de la Fuente, jose_delafuente@yahoo.com.

Ehrlichia species are the etiological agents of emerging tick-borne human zoonoses that inflict serious and fatal infections in companion animals and livestock. *Ehrlichia* species are tick-borne Gram-negative alphaproteobacteria that belong to the family Anaplasmataceae. Five *Ehrlichia* species are recognized, three of which can cause human ehrlichiosis (*E. canis*, *E. chaffeensis*, and *E. ewingii*) (1, 2). The complete genome of *E. chaffeensis*, *E. ruminantium*, *E. canis*, and *E. muris* were previously reported (3–6).

E. mineirensis was isolated from the hemolymph of *Rhipicephalus microplus*–engorged females and was characterized as a new species within the *Ehrlichia* genus (7). The organism has been maintained in the laboratory by continuous passage in the IDE8 tick cell line, where the ultrastructure was characterized (8, 9). Recently, we reported evidence that *E. mineirensis* evolved from a highly variable clade of *E. canis* under adaptive diversifying selection (10). For genome sequencing, the bacteria were grown in IDE8 cells, purified by Percoll density-gradient centrifugation (11), and the total DNA was extracted using TRI Reagent (Sigma, St. Louis, MO, USA). A next-generation sequencing (NGS) library was made starting from 800 ng of DNA using the NEB Next kit (New England Biolabs, Ipswich, MA, USA). The final library had a mean insert size of 516 bp. The library was then titrated by qPCR, denatured, equilibrated, and diluted for sequencing in MiSeq (Illumina). A total of 7.5 million pass-filter quality reads, 2 × 150-bp in length, were generated, which showed an average quality score above Q30 in more than 95% of the bases. The *de novo* assembly was performed with all the reads using SPADES (12). Prior to the assembly the overlapping reads were merged with FLASH (13). The contigs were aligned with Mauve (14) to the *E. canis* Jake genome (GenBank accession no. CP000107.1), detecting the pres-

ence of some contigs from *Ixodes* ticks. After massive BLAST comparison, only those contigs with similarity to Anaplasmataceae sequences were selected for annotation resulting in 182 contigs and 1,414,066 bases with an N_{50} of 32,220.

The genome of *E. mineirensis* consists of 1,414,066 bp, with 30.36% G+C content. The origin of replication (*oriC*), predicted by similarity to the *E. canis* Jake *oriC* region defined in the DoriC database (<http://tubic.tju.edu.cn/doric/info1.php?ac=ORI10030069>), seems to be placed in contig ehr000001 in the intergenic space upstream from the divergent genes encoding for uroporphyrinogen decarboxylase (EC 4.1.1.37) and cytochrome *c* oxidase, subunit III (EC 1.9.3.1), and 3 genes downstream of the *DnaJ* gene.

The genome of *E. mineirensis* was annotated using the BG7 system (15, 16) and resulted in 944 genes, including protein-coding sequences (CDSs), RNA genes, and pseudogenes. Of them, 322 genes encode proteins with enzymatic activity (with defined EC code), 51 encode for membrane proteins, 55 are involved in DNA repair, and 144 are related with oxidoreduction processes.

The availability of the *E. mineirensis* genome will allow comparative analysis to *E. canis* and *E. ruminantium* in studying the evolution of host specificity of *Ehrlichia* spp.

Nucleotide sequence accession numbers. The *E. mineirensis* genome sequence has been deposited in GenBank under the accession numbers CDGH01000001 through CDGH01000187.

ACKNOWLEDGMENTS

This research was supported by Spanish National Research Program grant BFU2011-23896 and by EU FP7 ANTIGONE project number 278976. R.T. and M.M. were partially funded by Spanish Ministerio de Ciencia e Innovación grant IPT-2011-0735-010000. A.C.C. was supported by a

grant from the Ministère de l'Éducation Supérieure et de la Recherche of France.

REFERENCES

- Perez M, Bodor M, Zhang C, Xiong Q, Rikihisa Y. 2006. Human infection with *Ehrlichia canis* accompanied by clinical signs in Venezuela. *Ann N Y Acad Sci* 1078:110–117. <http://dx.doi.org/10.1196/annals.1374.016>.
- Thomas RJ, Dumler JS, Carlyon JA. 2009. Current management of human granulocytic anaplasmosis, human monocytic ehrlichiosis and *Ehrlichia ewingii* ehrlichiosis. *Expert Rev Anti. Infect Ther* 7:709–722. <http://dx.doi.org/10.1586/eri.09.44>.
- Dunning Hotopp JC, Lin M, Madupu R, Crabtree J, Angiuoli SV, Eisen JA, Eisen J, Seshadri R, Ren Q Wu M, Utterback TR, Smith S, Lewis M, Khouri H, Zhang C, Niu H, Lin Q Ohashi N, Zhi N, Nelson W, Brinkac LM, Dodson RJ, Rosovitz MJ, Sundaram J, Daugherty SC, Davidsen T, Durkin AS, Gwinn M, Haft DH, Selengut JD, Sullivan SA, Zafar N, Zhou L, Benahmed F, Forberger H, Halpin R, Mulligan S, Robinson J, White O, Rikihisa Y. 2006. Comparative genomics of emerging human ehrlichiosis agents. *PLoS Genet* 2:e21. <http://dx.doi.org/10.1371/journal.pgen.0020021>.
- Collins NE, Liebenberg J, de Villiers EP, Brayton KA, Louw E, Pretorius A, Faber FE, van Heerden H, Josemans A, van Kleef M, Steyn HC, van Strijp MF, Zwegarth E, Jongejan F, Maillard JC, Berthier D, Botha M, Joubert F, Corton CH, Thomson NR, Allsopp MT, Allsopp BA. 2005. The genome of the heartwater agent *Ehrlichia ruminantium* contains multiple tandem repeats of actively variable copy number. *Proc Natl Acad Sci U S A* 102:838–843. <http://dx.doi.org/10.1073/pnas.0406633102>.
- Mavromatis K, Doyle CK, Lykidis A, Ivanova N, Francino MP, Chain P, Shin M, Malfatti S, Larimer F, Copeland A, Detter JC, Land M, Richardson PM, Yu XJ, Walker DH, McBride JW, Kyrpides NC. 2006. The genome of the obligately intracellular bacterium *Ehrlichia canis* reveals themes of complex membrane structure and immune evasion strategies. *J Bacteriol* 188:4015–4023. <http://dx.doi.org/10.1128/JB.01837-05>.
- Thirumalapura NR, Qin X, Kuriakose JA, Walker DH. 2014. Complete genome sequence of *Ehrlichia muris* strain as145t, a model monocyto-tropic *Ehrlichia* strain. *Genome Announc* 2(1):e01234-13. <http://dx.doi.org/10.1128/genomeA.01234-13>.
- Cruz AC, Zwegarth E, Ribeiro MF, da Silveira JA, de la Fuente J, Grubhoffer L, Valdés JJ, Passos LM. 2012. New species of *Ehrlichia* isolated from *Rhipicephalus (Boophilus) microplus* shows an ortholog of the *E. canis* major immunogenic glycoprotein gp36 with a new sequence of tandem repeats. *Parasit Vectors* 5:291. <http://dx.doi.org/10.1186/1756-3305-5-291>.
- Zwegarth E, Schöl H, Lis K, Cabezas Cruz A, Thiel C, Silaghi C, Ribeiro MFB, Passos LMF. 2013. *In vitro* culture of a novel genotype of *Ehrlichia* sp. from Brazil. *Transbound Emerg Dis* 60:86–92. <http://dx.doi.org/10.1111/tbed.12122>.
- Cabezas-Cruz A, Vancová M, Zwegarth E, Ribeiro MF, Grubhoffer L, Passos LM. 2013. Ultrastructure of *Ehrlichia mineirensis*, a new member of the *Ehrlichia* genus. *Vet Microbiol* 167:455–458. <http://dx.doi.org/10.1016/j.vetmic.2013.08.001>.
- Cabezas-Cruz A, Valdés JJ, de la Fuente J. 2014. The glycoprotein TRP36 of *Ehrlichia* sp. UFMG-EV and related cattle pathogen *Ehrlichia* sp. UFMT-BV evolved from a highly variable clade of *E. canis* under adaptive diversifying selection. *Parasit Vectors* 7:584. <http://dx.doi.org/10.1186/s13071-014-0584-5>.
- Lis K, Najm N, de la Fuente J, Fernández de Mera I, Zwegarth E, Pfister K, Passos LM. 2014. Use of Percoll gradients to purify *Anaplasma marginale* (Rickettsiales: Anaplasmataceae) from tick cell cultures. *Ticks Tick Borne Dis* 5:511–515. <http://dx.doi.org/10.1016/j.ttbdis.2014.03.006>.
- Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Pribelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol* 19:455–477. <http://dx.doi.org/10.1089/cmb.2012.0021>.
- Magoč T, Salzberg SL. 2011. FLASH: fast length adjustment of short reads to improve genome assemblies. *Bioinformatics* 27:2957–2963. <http://dx.doi.org/10.1093/bioinformatics/btr507>.
- Rissman AI, Mau B, Biehl BS, Darling AE, Glasner JD, Perna NT. 2009. Reordering contigs of draft genomes using the Mauve Aligner. *Bioinformatics* 25:2071–2013. <http://dx.doi.org/10.1093/bioinformatics/btp356>.
- Tobes R, Pareja-Tobes P, Manrique M, Pareja-Tobes E, Kovach E, Alekhin A, Pareja E. 2015. Gene calling and bacterial genome annotation with BG7. *Methods Mol Biol* 1231:177–189. http://dx.doi.org/10.1007/978-1-4939-1720-4_12.
- Pareja-Tobes P, Manrique M, Pareja-Tobes E, Pareja E, Tobes R. 2012. BG7: a new approach for bacterial genome annotation designed for next generation sequencing data. *PLoS One* 7:e49239. <http://dx.doi.org/10.1371/journal.pone.0049239>.