



$\gamma\delta$ T lymphocytes in the diagnosis of human T cell receptor immunodeficiencies

Beatriz Garcillán^{1†}, Ana V. M. Marin^{1†}, Anaïs Jiménez-Reinoso¹, Alejandro C. Briones¹, Miguel Muñoz-Ruiz¹, María J. García-León², Juana Gil³, Luis M. Allende⁴, Eduardo Martínez-Naves¹, María L. Toribio² and José R. Regueiro^{1*}

¹ Department of Immunology, Complutense University School of Medicine and Hospital 12 de Octubre Health Research Institute, Madrid, Spain

² Centro de Biología Molecular Severo Ochoa, Consejo Superior de Investigaciones Científicas and Universidad Autónoma, Madrid, Spain

³ Division of Immunology, Hospital General Universitario, Gregorio Marañón and Health Research Institute, Madrid, Spain

⁴ Division of Immunology, Hospital Universitario 12 de Octubre and Health Research Institute, Madrid, Spain

*Correspondence: regueiro@med.ucm.es

[†]Joint first authors.

Edited by:

Julie Dechanet-Merville, Centre National de la Recherche Scientifique, France

Reviewed by:

Sylvain Latour, Centre National de la Recherche Scientifique, France

Paul Fisch, University Freiburg Medical Center, Germany

Keywords: TCR, immunodeficiency, diagnosis, $\gamma\delta$, T lymphocyte

INTRODUCTION

Human T cell receptor (TCR) immunodeficiencies (TCRID) are rare autosomal recessive disorders caused by mutations affecting TCR, CD3, or CD247 chains, which share developmental, functional, and TCR expression defects (1). Their rapid diagnosis is fundamental for patient survival and early hematopoietic stem cell transplantation. Here, we propose that studying $\gamma\delta$ T cells, which are often neglected, can be helpful for a timely diagnosis. We thus offer a diagnostic flowchart and some lab tricks based on published cases.

$\gamma\delta$ T CELL AND TCR PHYSIOPATHOLOGY

$\gamma\delta$ T lymphocytes are a minor subset (1–10%) of human peripheral blood T cells. Most (>70%) are CD4⁻CD8⁻ [double negative (DN)], some (30%) are CD8⁺CD4⁻ and very few (<1%) are CD4⁺CD8⁻ [CD8⁺ or CD4⁺ single positive (SP), respectively]. Most $\gamma\delta$ T cells in adults express V δ 2/V γ 9 TCR variable regions (65–90%), the rest being mostly V δ 1⁺, some V δ 3⁺ or V δ 5⁺, all with different V γ chains (2). As peripheral blood $\gamma\delta$ T cells are scarce, their over-representation is more conspicuous than their under-representation, which is very rarely reported and normally associated to a single subset, such as V δ 2⁺ in

granulomatosis (3) or aging (4). Indeed, no selective $\gamma\delta$ T cell immunodeficiency (ID) has been reported to date, although absence of $\gamma\delta$ T cells has been described together with other lymphocyte derangements in rare primary ID (5). The clinical significance of increased $\gamma\delta$ T cells, defined as >10% of peripheral blood T lymphocytes (6), requires clarification in several diseases including infection, autoimmunity, cancer, and primary ID.

The human $\gamma\delta$ TCR (Figure 1A inset) is an octameric protein complex composed of three heterodimers (TCR γ /TCR δ , CD3 γ /CD3 ϵ , and CD3 δ /CD3 ϵ) and a single CD247 homodimer (also termed ζ/ζ). The complex can be abbreviated as $\gamma\delta$ TCR/ $\gamma\epsilon\delta\epsilon\zeta\zeta$. The TCR γ /TCR δ heterodimer contains variable regions, which allow for antigen recognition, while the other three dimers are invariant and are required for surface TCR expression and for intracellular propagation of the recognition signal (7). Therefore, defects in any chain would expectedly impact $\gamma\delta$ TCR expression and $\gamma\delta$ T cell selection and function.


$\gamma\delta$ T CELLS IN TCRID

$\alpha\beta$ T cells have been extensively studied in TCRID. In contrast, $\gamma\delta$ T cells have been frequently ignored, in part due to their scarcity but also to the lack of markers

other than the TCR to identify them when TCR expression is reduced, as is the case in TCRID. Although their functions are still debated, we believe that their accurate study (relative numbers, Figure 1A, TCR expression, Figure 1B, and main subsets) can help to diagnose TCRID, as reviewed below and summarized in a practical diagnostic flowchart in Figure 1C.

TCR α deficient patients showed combined ID and autoimmune features due to a selective block in $\alpha\beta$ T cell development, as TCR α is part of the TCR α /TCR β ($\alpha\beta$ TCR, Figure 1A inset) antigen-binding heterodimer (9). In contrast, the $\gamma\delta$ TCR was unaffected, as demonstrated by normal surface expression (Figure 1B), which allowed for normal absolute but increased relative numbers of $\gamma\delta$ T cells (Figure 1A). This is unique among TCRID and thus a useful feature in the diagnostic flowchart (Figure 1C). Such $\gamma\delta$ T cells were proposed to be in part protective from infections in the two reported patients. Indeed, $\gamma\delta$ T cells are involved in immune responses against a variety of pathogens including virus, bacteria, and parasites, whereas still other act as antigen-presenting cells (10) or B cell helper cells (11). Their beneficial effects *in vivo* have found recent unexpected recognition in haploidentical allogeneic hematopoietic cell transplantation after depletion of $\alpha\beta$ T and B cells (12),

Abbreviations: DN, double negative; DP, double positive; ID, immunodeficiency; mAb, monoclonal antibody; SCID, severe combined immunodeficiency; SP, single positive; TCR, T cell receptor; TCRID, T cell receptor immunodeficiencies.



which showed that $\gamma\delta$ T cells did not cause graft vs. host disease and may have helped with host immune maintenance and recovery. The fact is that, compared to other complete TCRID, symptoms in both TCR α deficient patients appeared rather late (6 and 15 months of age) and transplantation took place very late (6–7 years of age).

Similar to TCR α deficient patients, patients with partial CD3 δ deficiency (CD3 δ * in **Figure 1**) due to a leaky splicing mutation showed strongly reduced $\alpha\beta$ T cell numbers and normal absolute but high relative numbers of $\gamma\delta$ T cells (**Figure 1A**), although with low surface TCR expression [(13) and **Figure 1B**]. In contrast to TCR α deficiency, partial CD3 δ deficiency showed early severe combined ID (SCID) features and required very early transplantation (before 2 years of age), thus their $\gamma\delta$ T cells were not protective, perhaps as a consequence of their impaired TCR expression and function (13). Unexpectedly, partial CD3 δ deficiency caused a stronger impact in $\gamma\delta$ (**Figure 1B**) than in $\alpha\beta$ TCR surface expression (25 vs. 55% relative to controls (13)). A detailed study of their $\gamma\delta$ T cells showed an enrichment in a subset of otherwise rare CD4⁺ $\gamma\delta$ T cells, which exhibited an activated phenotype and were refractory to further TCR stimulation (14). This CD4-expressing $\gamma\delta$ T cell subset seems to be pathognomonic for partial CD3 δ deficiency, since: (i) it has been ascertained in three of three tested patients with this condition and (ii) it was not found in other TCRID (14, 15). Its developmental origin deserves further comment. $\alpha\beta$ and $\gamma\delta$ T cells differentiate within the thymus from a late DN common progenitor (16). In humans, development of most $\gamma\delta$ T cells seems to mimic that of $\alpha\beta$ T cells: from DN progenitors through a CD4⁺CD8⁺ double positive (DP) pathway (17, 18), to DN and either CD4⁺ or CD8⁺ SP populations. DN and CD8⁺ SP $\gamma\delta$ T cells are minor intrathymic subsets but become the major $\gamma\delta$ T cell subsets in the periphery, while CD4⁺ SP are the main intrathymic subset, followed by DP (19). Notably, the last two subsets can be found in peripheral blood in pathological conditions, and most bone marrow and peripheral blood $\gamma\delta$ T cells from patients with $\gamma\delta$ T cell acute lymphoblastic leukemia are either CD4⁺ SP or DP (20). Thus, we believe that the 10-fold enrichment of CD4⁺ SP $\gamma\delta$ T cells observed

in patients with partial CD3 δ deficiency is due to low TCR-dependence for positive selection of CD4⁻ $\gamma\delta$ T cells and disrupted negative selection of CD4⁺ $\gamma\delta$ T cells (14).

CD3 γ deficient patients, most of which showed mild ID (21), had normal numbers of polyclonal peripheral blood $\gamma\delta$ T cells [absolute and relative, (22) and **Figure 1A**] with low surface TCR [around 30% of control levels (22) and **Figure 1B**], similarly to their $\alpha\beta$ T cell counterparts (23), likely with an abnormal $\gamma\delta$ TCR/ $\delta\epsilon\delta\epsilon\zeta\zeta$ stoichiometry. Despite their high homology, the invariant CD3 γ and δ chains show different roles in human vs. mouse $\gamma\delta$ T cell development. Indeed, CD3 γ -deficient mice exhibited a severe $\gamma\delta$ T cell developmental block (24).

$\gamma\delta$ T cells were studied in only two of three reported CD247 deficient patients (21). The patients showed SCID features and reduced absolute and relative $\gamma\delta$ T cells numbers (**Figure 1A**). Surface $\gamma\delta$ TCR expression was also reduced (4% vs. healthy controls, **Figure 1B**). The number of $\alpha\beta$ T cells was only slightly reduced despite their similarly reduced surface TCR, with all reported cases showing reduced numbers of CD4⁺ T cells but normal or high numbers of CD8⁺ T cells (21, 25–27).

A single patient with partial CD3 ϵ deficiency (28–30) showed very low surface $\alpha\beta$ TCR expression (10% of normal levels, **Figure 1B**, CD3 ϵ^*), mild ID, normal CD8⁺, and reduced CD4⁺ ($\alpha\beta$) T cells, but no $\gamma\delta$ T cells as determined with the anti-TCR δ -1 monoclonal antibody (mAb) (**Figure 1A**). We have however considered for **Figure 1C** that surface $\gamma\delta$ TCR expression might have been similar to $\alpha\beta$ TCR expression.

Three studied CD3 δ deficient patients [out of 16 reported, all with severe T cell lymphopenia and SCID (31, 32)], showed a few circulating CD3⁺ T cells, which were DN but $\gamma\delta$ TCR⁻ (33). $\gamma\delta$ TCR⁺ cells were indeed undetectable by flow cytometry in peripheral blood or by immunohistochemistry in the thymus, lymph nodes, spleen, or gut. However, gene microarray analysis and protein expression of patient thymocytes showed increased levels of TCR γ and TCR δ transcripts and proteins (33), which could be interpreted as presence and thus significant selection of $\gamma\delta$ T cells unable however to leave the thymus, perhaps due to insufficient surface TCR compared to partial CD3 δ deficiency.

Finally, $\gamma\delta$ T cells have not been studied in SCID patients with complete CD3 ϵ deficiency (31). Nevertheless, given their severe T cell lymphopenia, we can safely presume for **Figure 1C** that they were absent.

In summary, the proportion of $\gamma\delta$ T cells within total T lymphocytes (**Figure 1A**) and the level of surface $\gamma\delta$ vs. $\alpha\beta$ TCR expression (**Figure 1B**), as well as the severity of lymphopenia (T⁻B⁺NK⁺ or T[±]B⁺NK⁺ phenotype), can be used to generate a practical TCRID diagnostic flowchart (**Figure 1C**). For instance, if an infant has SCID and no T cells but normal B and NK cell numbers (T⁻B⁺NK⁺ phenotype) and other causes have been ruled out, CD3 δ or CD3 ϵ deficiency should be considered (**Figure 1C**). In contrast, if some T cells are present (T[±]B⁺NK⁺ phenotype) and $\gamma\delta$ TCR expression is low, TCR α deficiency can be ruled out. If CD16 expression by NK cells is normal, CD247 deficiency can be excluded, and the presence or absence of high absolute numbers of CD4⁺ $\gamma\delta$ T cells will rule out CD3 γ or partial CD3 δ deficiency, respectively.

LAB TRICKS TO IDENTIFY $\alpha\beta$ AND $\gamma\delta$ T CELLS IN TCRID

When surface TCR expression is low, $\alpha\beta$ T cells can be identified by the expression of CD4 or CD8 $\alpha\beta$ (i.e., CD8^{bright}) within the lymphoid subset (23), whereas $\gamma\delta$ T cells are identified only by expression of the $\gamma\delta$ TCR. We have reported that most CD3⁺ cells within normal DN lymphocytes are $\gamma\delta$ T cells (34), and this may also help in certain TCRID.

Despite their reduced numbers and surface TCR expression, an appropriate multicolor flow cytometry approach can help to identify $\gamma\delta$ T cells in TCRID. To avoid underestimation due to low TCR surface expression, we recommend: (i) the use of bright fluorochromes such as PE, PE-Cy5.5, PE-Cy7, or APC, rather than FITC, (ii) an appropriate choice of CD3 mAb such as UCHT-1, F101.01, or S4.1 due to their high signal-to-noise ratio in TCRID, (iii) two-color stainings with CD3 and $\gamma\delta$ TCR mAb, which can also help to single out $\gamma\delta$ T cells as a DP subset, and (iv) to avoid mixing $\alpha\beta$ TCR and CD3 mAb, as they sometimes compete (UCHT-1, for instance).

CD4 and CD8 expression by $\gamma\delta$ T cells should also be tested to rule out partial CD3 δ deficiency (see above). CD4, CD8,

$\gamma\delta$ TCR (IMMU510 or 11F2), and CD3 (UCHT-1 or S4.1) is a useful combination, to this end. Lastly, intracellular stainings for invariant TCR chains has been shown to be useful to identify T cells expressing very low surface TCR, such as those with CD247 (21) or partial CD3 δ deficiency (14).

CONCLUSION AND PERSPECTIVES

Human $\gamma\delta$ T lymphocytes are still puzzling in terms of development, function, and TCR stoichiometry in ways that mouse models do not wholly recapitulate. Human TCRID share defects in T cell development and function and in TCR expression. While their $\alpha\beta$ T cells have been studied in detail, $\gamma\delta$ T cells have been frequently ignored, in part due to their scarcity and to the lack of appropriate markers to identify them when TCR expression is reduced. Here, pooling published studies, we proposed some technical tricks to identify $\gamma\delta$ T cells in TCRID patients and made the point that their careful analysis can help to inform a rapid differential diagnosis using a flowchart, with clinical benefit.

ACKNOWLEDGMENTS

Supported by grants from MINECO (SAF 2011-24235, BES-2012-055054, SAF2014-54708-R, and SAF2014-53563-REDT), CAM (S2010/BMD-2316), ISCIII (RD08-0075-0002 and PI12/02761), and Lair (2012/0070). We thank M. J. Recio and E. Fernández-Malavé (Universidad Complutense) for helpful discussions, M. Jurado (Community Transfusion Centre of Madrid) and Craig W. Reynolds (National Cancer Institute, National Institutes of Health, Frederick, USA) for critical reagents and all patients, families, and clinicians for their cooperation.

REFERENCES

1. Regueiro JR, Recio MJ. T-cell-receptor complex deficiency. 3 ed. In: Ochs HD, Smith CIE, Puck JM, editors. *Primary Immunodeficiency Diseases, a Molecular and Genetic Approach*. New York: Oxford University Press (2013). p. 156–62.
2. Kalyan S, Kabelitz D. Defining the nature of human gammadelta T cells: a biographical sketch of the highly empathetic. *Cell Mol Immunol* (2013) **10**(1):21–9. doi:10.1038/cmi.2012.44
3. Fazio J, Quabius ES, Muller A, Adam-Klages S, Wesch D, Sebens S, et al. Vdelta2 T cell deficiency in granulomatosis with polyangiitis (Wegener's granulomatosis). *Clin Immunol* (2013) **149**(1):65–72. doi:10.1016/j.clim.2013.06.003
4. Michishita Y, Hirokawa M, Guo YM, Abe Y, Liu J, Ubukawa K, et al. Age-associated alteration of

- gammadelta T-cell repertoire and different profiles of activation-induced death of Vdelta1 and Vdelta2 T cells. *Int J Hematol* (2011) **94**(3):230–40. doi:10.1007/s12185-011-0907-7
5. Pannicke U, Baumann B, Fuchs S, Henneke P, Rensing-Ehl A, Rizzi M, et al. Deficiency of innate and acquired immunity caused by an IKBKB mutation. *N Engl J Med* (2013) **369**(26):2504–14. doi:10.1056/NEJMoa1309199
 6. Granel B, Camoin L, Serratrice J, de Roux-Serratrice C, Brunet C, Pache X, et al. Retrospective study of 55 patients with circulating blood T gamma/delta lymphocytosis. *Rev Med Interne* (2002) **23**(2):137–43. doi:10.1016/S0248-8663(01)00529-X
 7. Call ME, Pyrdol J, Wucherpfennig KW. Stoichiometry of the T-cell receptor-CD3 complex and key intermediates assembled in the endoplasmic reticulum. *EMBO J* (2004) **23**(12):2348–57. doi:10.1038/sj.emboj.7600245
 8. Wen L, Peakman M, Mieli-Vergani G, Vergani D. Elevation of activated gamma delta T cell receptor bearing T lymphocytes in patients with autoimmune chronic liver disease. *Clin Exp Immunol* (1992) **89**(1):78–82. doi:10.1111/j.1365-2249.1992.tb06881.x
 9. Morgan NV, Goddard S, Cardno TS, McDonald D, Rahman F, Barge D, et al. Mutation in the TCRalpha subunit constant gene (TRAC) leads to a human immunodeficiency disorder characterized by a lack of TCRalphabeta+ T cells. *J Clin Invest* (2011) **121**(2):695–702. doi:10.1172/JCI41931
 10. Casetti R, Martino A. The plasticity of gamma delta T cells: innate immunity, antigen presentation and new immunotherapy. *Cell Mol Immunol* (2008) **5**(3):161–70. doi:10.1038/cmi.2008.20
 11. Brandes M, Willmann K, Lang AB, Nam KH, Jin C, Brenner MB, et al. Flexible migration program regulates gamma delta T-cell involvement in humoral immunity. *Blood* (2003) **102**(10):3693–701. doi:10.1182/blood-2003-04-1016
 12. Pulsipher MA. Haplo is the new black. *Blood* (2014) **124**(5):675–6. doi:10.1182/blood-2014-06-579714
 13. Gil J, Busto EM, Garcillán B, Chean C, Garcia-Rodriguez MC, Diaz-Alderete A, et al. A leaky mutation in CD3D differentially affects alpha-beta and gammadelta T cells and leads to a Tab-Tgd+B+NK+ human SCID. *J Clin Invest* (2011) **121**(10):3872–6. doi:10.1172/JCI44254
 14. Garcillán B, Mazariegos MS, Fisch P, Res PC, Munoz-Ruiz M, Gil J, et al. Enrichment of the rare CD4(+) gammadelta T-cell subset in patients with atypical CD3delta deficiency. *J Allergy Clin Immunol* (2014) **133**(4):1205–8. doi:10.1016/j.jaci.2013.10.002
 15. Martínez-Martínez L, Cedeño R, Marin AVM, Rubiales MV, Jiménez-Reinoso A, Recio MJ, et al. A founder splicing mutation in CD3D in a Tab-Tgd+B+NK+SCID pedigree of Ecuadorian descent. *J Clin Immunol* (2014) **34**(Suppl 2):S440. doi:10.1007/s10875-014-0101-9
 16. García-Peydro M, de Yébenes VG, Toribio ML. Sustained Notch1 signaling instructs the earliest human intrathymic precursors to adopt a gammadelta T-cell fate in fetal thymus organ culture. *Blood* (2003) **102**(7):2444–51. doi:10.1182/blood-2002-10-3261
 17. Offner F, Van Beneden K, Debacker V, Vanhecke D, Vandekerckhove B, Plum J, et al. Phenotypic and functional maturation of TCR gammadelta cells in the human thymus. *J Immunol* (1997) **158**(10):4634–41.
 18. Van Coppennolle S, Vanhee S, Verstichel G, Snauwaert S, van der Spek A, Velghe J, et al. Notch induces human T-cell receptor gammadelta+ thymocytes to differentiate along a parallel, highly proliferative and bipotent CD4 CD8 double-positive pathway. *Leukemia* (2012) **26**(1):127–38. doi:10.1038/leu.2011.324
 19. Groh V, Porcelli S, Fabbi M, Lanier LL, Picker LJ, Anderson T, et al. Human lymphocytes bearing T cell receptor gamma/delta are phenotypically diverse and evenly distributed throughout the lymphoid system. *J Exp Med* (1989) **169**(4):1277–94. doi:10.1084/jem.169.4.1277
 20. Langerak AW, Wolvers-Tettero IL, van den Beemd MW, van Wering ER, Ludwig WD, Hahlen K, et al. Immunophenotypic and immunogenotypic characteristics of TCRgd+ T cell acute lymphoblastic leukemia. *Leukemia* (1999) **13**(2):206–14. doi:10.1038/sj.leu.2401276
 21. Marin AVM, Garcillán B, Jiménez-Reinoso A, Muñoz-Ruiz M, Briones AC, Fernández-Malave E, et al. Human congenital T cell receptor disorders. *LymphoSign J* (2014) **2**(1). doi:10.14785/lpsn-2014-0012
 22. Siegers GM, Swamy M, Fernandez-Malave E, Minguet S, Rathmann S, Guardo AC, et al. Different composition of the human and the mouse gammadelta T cell receptor explains different phenotypes of CD3gamma and CD3delta immunodeficiencies. *J Exp Med* (2007) **204**(11):2537–44. doi:10.1084/jem.20070782102207c
 23. Recio MJ, Moreno-Pelayo MA, Kilic SS, Guardo AC, Sanal O, Allende LM, et al. Differential biological role of CD3 chains revealed by human immunodeficiencies. *J Immunol* (2007) **178**(4):2556–64. doi:10.4049/jimmunol.178.4.2556
 24. Haks MC, Krimpenfort P, Borst J, Kruisbeek AM. The CD3gamma chain is essential for development of both the TCRalphabeta and TCRgammadelta lineages. *EMBO J* (1998) **17**(7):1871–82. doi:10.1093/emboj/17.7.1871
 25. Rieux-Laucat F, Hivroz C, Lim A, Mateo V, Pellier I, Selz F, et al. Inherited and somatic CD3zeta mutations in a patient with T-cell deficiency. *N Engl J Med* (2006) **354**(18):1913–21. doi:10.1056/NEJMoa053750
 26. Roberts JL, Lauritsen JP, Cooney M, Parrott RE, Sajaroff EO, Win CM, et al. T-B+NK+ severe combined immunodeficiency caused by complete deficiency of the CD3zeta subunit of the T-cell antigen receptor complex. *Blood* (2007) **109**(8):3198–206. doi:10.1182/blood-2006-08-043166
 27. Recio MJ, Aydogmus C, Marin AVM, Haskologlu S, Jiménez-Reinoso A, Cipe F, et al. Complete deficiency of CD247, the zeta chain of the T-cell antigen receptor and the NK Fc gamma receptor III. *J Clin Immunol* (2014) **34**(Suppl 2):S463. doi:10.1007/s10875-014-0101-9
 28. Thoenes G, Soudais C, le Deist F, Griscelli C, Fischer A, Lisowska-Groszpiere B. Structural analysis of low TCR-CD3 complex expression in T cells of an immunodeficient patient. *J Biol Chem* (1992) **267**(1):487–93.
 29. Soudais C, de Villartay JP, Le Deist F, Fischer A, Lisowska-Groszpiere B. Independent mutations of the human CD3-epsilon gene resulting in a T cell receptor/CD3 complex immunodeficiency. *Nat Genet* (1993) **3**(1):77–81. doi:10.1038/ng0193-77
 30. Le Deist F, Thoenes G, Corado J, Lisowska-Groszpiere B, Fischer A. Immunodeficiency with low expression of the T cell receptor/CD3 complex. Effect on T lymphocyte activation. *Eur J Immunol* (1991) **21**(7):1641–7. doi:10.1002/eji.1830210709
 31. de Saint Basile G, Geissmann F, Flori E, Uring-Lambert B, Soudais C, Cavazzana-Calvo M, et al. Severe combined immunodeficiency caused by deficiency in either the delta or the epsilon subunit of CD3. *J Clin Invest* (2004) **114**(10):1512–7. doi:10.1172/JCI200422588
 32. Marcus N, Takada H, Law J, Cowan MJ, Gil J, Regueiro JR, et al. Hematopoietic stem cell transplantation for CD3delta deficiency. *J Allergy Clin Immunol* (2011) **128**(5):1050–7. doi:10.1016/j.jaci.2011.05.031
 33. Dadi HK, Simon AJ, Roifman CM. Effect of CD3delta deficiency on maturation of alpha/beta and gamma/delta T-cell lineages in severe combined immunodeficiency. *N Engl J Med* (2003) **349**(19):1821–8. doi:10.1056/NEJMoa031178
 34. Muñoz-Ruiz M, Perez-Flores V, Garcillán B, Guardo AC, Mazariegos MS, Takada H, et al. Human CD3gamma, but not CD3delta, haploinsufficiency differentially impairs gammadelta versus alphabeta surface TCR expression. *BMC Immunol* (2013) **14**:3. doi:10.1186/1471-2172-14-3

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Received: 03 November 2014; accepted: 09 January 2015; published online: 29 January 2015.

Citation: Garcillán B, Marin AVM, Jiménez-Reinoso A, Briones AC, Muñoz-Ruiz M, García-León MJ, Gil J, Allende LM, Martínez-Naves E, Toribio ML and Regueiro JR (2015) γδ T lymphocytes in the diagnosis of human T cell receptor immunodeficiencies. *Front. Immunol.* **6**:20. doi: 10.3389/fimmu.2015.00020

This article was submitted to *T Cell Biology*, a section of the journal *Frontiers in Immunology*.

Copyright © 2015 Garcillán, Marin, Jiménez-Reinoso, Briones, Muñoz-Ruiz, García-León, Gil, Allende, Martínez-Naves, Toribio and Regueiro. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.